Chapter Twelve ◆ TERRESTRIAL AND FRESHWATER ECOLOGY AND BIODIVERSITY

INTRODUCTION

- 12.1 This chapter sets out a detailed assessment of the likely significant effects of the Proposed Development (as described in ES Chapter 3: *Project Description* (Document Reference 6.1.3)) on terrestrial and freshwater features of nature conservation value. In particular, it considers the likely effects of the Proposed Development on the Important Ecological Features (IEFs) identified within the Project Site or its potential zone of influence¹.
- 12.2 The Project Site comprises two parts including the 'Kent Project Site', which includes land on the Swanscombe Peninsula and the Ebbsfleet Valley on the south side of the River Thames centred approximately at Ordnance Survey Grid Reference (OSGR) TQ 606 758, and the 'Essex Project Site', which includes land to the east of the A1089 Ferry Road and the Tilbury Ferry Terminal centred approximately at OSGR TQ 643 752. The Project Site lies partly within three local planning authority areas: Dartford Borough and Gravesham Borough for the Kent Project Site; and Thurrock Council for the Essex Project Site. Collectively these two parts of the entire Development Consent Order DCO boundary are referred to as 'the Project Site'.
- 12.3 The IEFs included within the assessment are described in detail in Appendix 12.1 *Ecology Baseline Report* (Document Reference 6.2.12.1) and summarised within the 'Baseline Conditions' section of this Chapter. Aquatic features in the Thames Estuary intertidal and subtidal environment are dealt with separately in ES Chapter 13: *Marine Ecology* (Document Reference 6.1.13).
- 12.4 The assessment has been prepared with reference to the Chartered Institute of Ecology and Environmental Management's (CIEEM) 'Guidelines for Ecological Impact Assessment in the UK and Ireland' (CIEEM, 2018)², hereafter referred to as 'the EcIA Guidelines'. This chapter has been prepared and reviewed by experienced senior EDP Ecologists and full members of CIEEM.
- 12.5 This chapter describes:
 - the methods used for the assessment;

² CIEEM (2018) Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, Freshwater, Coastal and Marine version 1.1. Chartered Institute of Ecology and Environmental Management, Winchester.



12 - 1

¹ Zone of Influence - the area over which ecological features may be affected by biophysical changes as a result of the proposed development

- a summary of the baseline conditions currently existing at the Project Site and in its surroundings;
- the likely direct and indirect effects arising from the Proposed Development during construction and operation;
- the <u>mitigation</u> measures <u>required proposed</u> to avoid, <u>or mitigate or compensate</u> likely significant negative effects;
- the potential opportunities to provide enhancements over the existing situation;
- the assessment of residual effects taking into account the additional mitigation measures proposed;
- <u>the measures proposed to compensate for those residual effects, which cannot be</u> avoided or mitigated; and
- the assessment of potential cumulative and in-combination effects on ecological receptors.
- 12.6 This document supersedes the original ES Chapter, which was submitted with the DCO Application in late 2020. The key updates to the original ES Chapter and the supporting documentation are as follows:
 - Updates to the baseline conditions and evaluation of important ecological features to reflect the notification of Swanscombe Peninsula SSSI (notified on 11 March 2021), which covers a large proportion of the Kent Project Site;
 - Additional baseline information collected through site surveys since the original submission;
 - Further clarification in respect of predicted direct and indirect effects upon important ecological features;
 - Amendments to the proposals for on-site mitigation of impacts upon non-SSSI notified features, where these have potential to result in negative effects on SSSI notified features;
 - Greater distinction made between avoidance and mitigation, and compensation;
 - More detail in respect of proposed off-site habitat creation measures to mitigate (in the case of potential effects on off-site designated sites) or compensate (in the case of effects on on-site features) for negative effects on-site.



This Chapter should be read in conjunction with the following ES Appendices and Figures.

Appendices

- Appendix 12.1: *Ecology Baseline Report* (Document Reference 6.2.12.1), which includes full details of survey methods, results and a full set of associated drawings;
- Appendix 12.2: Biodiversity Net Gain Impact Assessment (Document Reference 6.2.12.2), which presents the results of a formal calculation of the net change in biodiversity value resulting from the Proposed Development using a standardised Metric;
- Appendix 12.3: Ecological Mitigation and Management Framework (Document Reference 6.2.12.3), which describes in further detail the <u>on-site</u> ecological avoidance , <u>mitigation</u>, <u>compensation</u> and <u>enhancementand mitigation</u> —measures proposed during the construction and operational stages of the Proposed Development;
- Appendix 12.4: Shadow Habitats Regulations Assessment (Document Reference 6.2.12.4), which sets out a detailed assessment of likely significant effects on European Sites resulting from the Proposed Development both in the absence of, and then with the inclusion of, mitigation, and, as a last resort, compensation measures;
- Appendix 12.5: Consultation responses to the 2020 EIA Scoping request (Relevant to Terrestrial and Freshwater Ecology; Document Reference 6.2.12.5), includes copies of consultation responses and how the points raised have been addressed in the ES;
- Appendix 12.6: Statutory consultee responses to the Preliminary Environmental Information Report (Relevant to Terrestrial and Freshwater Ecology; Document Reference 6.2.12.6), includes copies of consultation responses and how the points raised have been addressed in the ES;
- Appendix 12.7: Non-statutory consultee responses to the Preliminary Environmental Information Report (Relevant to Terrestrial and Freshwater Ecology; Document Reference 6.2.12.7), includes copies of consultation responses and how the points raised have been addressed in the ES;
- Appendix 12.8: Water Framework Directive (Screening) Assessment: River Ebbsfleet (Document Reference 6.2.12.8), provides information in respect of an assessment of potential effects arising upon the River Ebbsfleet and associated aquatic communities with due regard to objectives of the Water Framework Directive;
- Appendix 12.9: Arboricultural Impact Assessment (Document Reference 6.2.12.9), evaluates the existing tree stock and identifies trees suitable for retention, and assesses the impacts on the existing stock from the Proposed Development;



- Appendix 12.10: General Principles for Offsite Ecological Mitigation Ecological
 <u>Compensation Framework</u> (Document Reference 6.2.12.10), describes the general
 principles to be adhered to in the delivery of off site ecological mitigation;
- Appendix 12.11: Artificial Lighting Environmental Impact Assessment (Document Reference 6.2.12.11), provides details on the artificial lighting strategy for the Proposed Development together with the design criteria, recommendations and mitigation measures to avoid or mitigate undue light pollution or adverse impact upon the natural environment;
- Landscape Strategy (Document Reference 6.2.11.7), provides the overall landscape and green infrastructure design for the Proposed Development covering a number of elements such as accessibility, habitat creation and ecology, hydrology and public facilities, and provides the basis for fully a detailed Soft Landscaping Scheme (SLS) to be prepared post-consent;
- Landscape Management Plan (Document Reference 6.2.11.8), provides the strategy for delivering landscape management, maintenance and monitoring within the site wide landscape, to ensure that the ecological habitats retained, created or enhanced provide long term benefits to wildlife throughout the operational period of the Proposed Development; and
- Public Rights of Way Assessment (Document Reference 6.2.11.9), considers the impacts resulting from the Proposed Development on existing Public Rights of Way (PRoW) and provides mitigation and/or enhancement measures to the PRoW network.

Figures

- Figure 12.1: Project Site Areas (Document Reference 6.3.12.1);
- Figure 12.2: Statutory Designations Plan (Document Reference 6.3.12.2);
- Figure 12.3: Non-Statutory Designations Plan (Document Reference 6.3.12.3);
- Figure 12.4: Extended Phase 1 Habitat Survey (Document Reference 6.3.12.4);
- Figure 12.5: Rare Plants Species Population Plan (Document Reference 6.3.12.5);
- Figure 12.6: River Ebbsfleet River Corridor Survey Plan (Document Reference 6.3.12.6);
- Figure 12.7: River Ebbsfleet River Habitat Survey Sections and Results (Document Reference 6.3.12.7);



- Figure 12.8: Breeding Bird Survey Plan (April 2020; Document Reference 6.3.12.8);
- Figure 12.9: Breeding Bird Survey Plan (May 2020; Document Reference 6.3.12.9);
- Figure 12.10: Breeding Bird Survey Plan (June 2020; Document Reference 6.3.12.10);
- Figure 12.11: Breeding Bird Survey Plan (July 2020; Document Reference 6.3.12.11);
- Figure 12.12: Preliminary Roost Assessment of Trees (Document Reference 6.3.12.12);
- Figure 12.13: Preliminary Roost Assessment of Buildings (Document Reference 6.3.12.13);
- Figure 12.14: Static Detector Locations and Bat Transect Routes (Document Reference 6.3.12.14);
- Figure 12.15: Bat Transect Survey Results (Document Reference 6.3.12.15);
- Figure 12.16: Bat Activity Results per Species (Document Reference 6.3.12.16);
- Figure 12.17: Bat activity heat maps (Document Reference 6.3.12.17);
- Figure 12.18: Dormouse Survey Nest Tube Locations and Previous Records (Document Reference 6.3.12.18);
- Figure 12.19: Dormouse Survey Results 2020 (Document Reference 6.3.12.19);
- Figure 12.20: Summary of Dormouse Survey Results 2020 (Document Reference 6.3.12.20);
- Figure 12.21: Water Vole Survey Results (Document Reference 6.3.12.21);
- Figure 12.22: Otter Survey Results (Document Reference 6.3.12.22);
- Figure 12.23: Harvest Mouse Survey Plan (Document Reference 6.3.12.23);
- Figure 12.24: GCN Survey Area (Document Reference 6.3.12.24);
- Figure 12.25: Reptile Survey Plan and Previous Records (Document Reference 6.3.12.25);
- Figure 12.26: Reptile Survey Results (Document Reference 6.3.12.26);



- Figure 12.27: Aquatic Macroinvertebrate Sample Locations (standing waterbodies; Document Reference 6.3.12.27);
- Figure 12.28: Aquatic Macroinvertebrate Survey Results (River Ebbsfleet) Spring 2020 (Document Reference 6.3.12.28);
- Figure 12.29: Aquatic Macroinvertebrate Survey Results (River Ebbsfleet) Autumn 2020 (Document Reference 6.3.12.29);
- Figure 12.30: Site Location Plan and Core Count Zones for Low and High Tide Surveys (Document Reference 6.3.12.30);
- Figure 12.31: Statutory Designations Recognised for Their Importance to Overwintering Birds (Document Reference 6.3.12.31);
- Figure 12.32: Winter Bird Survey Results November 2019 (Document Reference 6.3.12.32);
- Figure 12.33: Winter Bird Survey Results December 2019 (Document Reference 6.3.12.33);
- Figure 12.34: Winter Bird Survey Results January 2020 (Document Reference 6.3.12.34)
- Figure 12.35: Winter Bird Survey Results February 2020 (Document Reference 6.3.12.35);
- Figure 12.36: Winter Bird Survey Results March 2020 (Document Reference 6.3.12.36);
- Figure 12.37: Vantage Point Survey Results November 2019 (Document Reference 6.3.12.37);
- Figure 12.38: Vantage Point Survey Results December 2019 (Document Reference 6.3.12.38);
- Figure 12.39: Vantage Point Survey Results January 2020 (Document Reference 6.3.12.39);
- Figure 12.40: Vantage Point Survey Results February 2020 (Document Reference 6.3.12.40);



- Figure 12.41: Vantage Point Survey Results March 2020 (Document Reference 6.3.12.41);
- Figure 12.42: Biodiversity Net Gain Impact Assessment Existing Habitats (Document Reference 6.3.12.42);
- Figure 12.43: Biodiversity Net Gain Impact Assessment Proposed Habitats (Document Reference 6.3.12.43);
- Figure 12.44: Ecology Mitigation Strategy: Species Measures (Document Reference 6.3.12.44);
- Figure 12.45: Light Mitigation Strategy for Biodiversity (Document Reference 6.3.12.45);
- Figure 12.46: Locations of Other Nearby Dormouse Licenced Sites (Document Reference 6.3.12.46);
- Figure 12.47: Dormouse Habitat Suitability Assessment (Document Reference 6.3.12.47);
- Figure 12.48: Dormouse Habitat Impacts Plan (Document Reference 6.3.12.48);
- Figure 12.49: Desk Study Results Water Vole (Document Reference 6.3.12.49);
- Figure 12.50: Existing Hydrology (Document Reference 6.3.12.50);
- Figure 12.51: Predicted Impacts Hydrology (Document Reference 6.3.12.51);
- Figure 12.52: Mitigation and Enhancements Hydrology (Document Reference 6.3.12.52);
- Figure 12.53: Water Vole Receptor Location (Document Reference 6.3.12.53);
- Figure 12.54: Water Vole Receptor Design (Document Reference 6.3.12.54);
- Figure 12.55: Artificial Otter Holt Location and Design (Document Reference 6.3.12.55);
- Figure 12.56: Tree Constraints Plan (Document Reference 6.3.12.56); and
- Figure 12.57: Tree Retention and Removal Plan (Document Reference 6.3.12.57).

METHODOLOGY AND DATA SOURCES



The assessment will follow the methodology provided in the EcIA Guidelines. These Guidelines confirm that the 'EcIA is a process of identifying, quantifying and evaluating the potential effects of development-related or other proposed actions on habitats, species and ecosystems'. The assessment methodology has been agreed through consultation on EIA Scoping and the Preliminary Environmental Information Report.

Consultation

EIA Scoping

A Scoping Opinion was subsequently received from the SoS in July 2020, and the comments received have been used to inform the assessment. **Table 12.1** includes the Planning Inspectorate's comments from the 2020 EIA Scoping Opinion in relation to ecology, and the actions taken.

12.912.10 The comments received from statutory consultation bodies formally consulted by the Inspectorate in preparing the 2020 Scoping Opinion, and how the comments have been taken into consideration, are provided in Appendix 12.5 (Document Reference 6.2.12.5).

Table 12.1: Planning Inspectorate's comments from the 2020 EIA Scoping Opinion in relation to Ecology (July 2020).

PINS ID	Applicant's proposed matters to scope out	Inspectorate's comments	Action taken
4.5.1	Effects on fish populations within the Thames Estuary, Swanscombe Marshes and the Ebbsfleet Stream	The Scoping Report seeks to scope out these effects on the grounds that survey information shows very few species within these areas. This appears to contradict the statement in paragraph 12.73 of the Scoping Report which states that surveys are proposed. Accordingly, the Inspectorate does not agree to scope this matter out. The ES should assess impacts to fish populations in the Thames Estuary, Swanscombe Marshes and the Ebbsfleet Stream where significant	An Ecology Briefing Note 'Further Information in Respect of an EIA Scoping Opinion' (Report ref: edp5988_r019) was submitted to EA on 13 August 2020 to provide further information on potential effects on a fish population within the river Thames, Swanscombe Marshes and River Ebbsfleet, and thus assess the requirement for further survey effort of each waterbody with respect to fish. The EA 'Response to LR Ecology Briefing Note' (Ref: KT/2020/127432/01-L01) received 15 September 2020 (see Appendix 12.5; Document Reference 6.2.12.5) confirmed additional fish surveys of



PINS	Applicant's	Inspectorate's comments	Action taken
ID	proposed		
	matters to		
	scope out	effects are likely to occur. The	the River Ebbsfleet were not
		Applicant should make effort to agree the approach to the assessment with relevant consultation bodies including the Environment Agency.	required, with the existing baseline considered adequate to inform potential effects. Further surveys of the Swanscombe Marsh waterbodies were, however, progressed during September 2020 to confirm presence/absence of a notable fish population, particularly given the
			potential impacts in this area. The surveys confirmed the absence of a notable fish population.
			It has been agreed with the MMO that subtidal fish surveys will not be required in the Thames Estuary. Inter-tidal fish surveys of the River Thames have been progressed, the results of which have informed Chapter 13: Marine ecology and biodiversity (Document Reference: 6.1.13) and the Water Framework Directive (WFD) assessment (Appendix 13.7; Document Reference 6.2.13.7).
			Although assessments of the River Ebbsfleet and Swanscombe Marshes have not identified any significant fish assemblages such that potential significant effects are unlikely, correspondence with the EA have confirmed a requirement to assess the potential for development to result in deterioration of each waterbody, and therefore each waterbody's potential to support a fish population in future. This
			assessment has now been completed (see Appendix 12.7; Document
4.5.2	Selection of	The response from relevant	Reference 6.2.12.7) The assessment has been updated to
4.3.2	Jeiection of	The response montrelevant	The assessment has been upuated to



PINS	Applicant's	Inspectorate's comments	Action taken
ID	proposed		
	matters to		
	Important Ecological Features	consultation bodies identifies a number of receptors, including additional European sites and Sites of Special Scientific Interest, nonstatutory wildlife sites and ancient woodland which are not included in the assessment. The assessment in the ES should be comprehensive with regards to all relevant receptors likely to experience significant effects. The Applicant should make effort to agree the relevant receptors for the	include additional receptors with potential to experience significant effects, as identified in the relevant consultation responses. This includes the inclusion of the Mucking Flats and Marshes SSSI, which is the underpinning SSSI for the Thames Estuary and Marshes SPA and Ramsar Site in Essex, as requested by Natural England. Furthermore, North Downs Woodland SAC, and the underpinning Wouldham to Detling Escarpment SSSI, are now included within the assessment of potential air quality effects, as advised by Natural England. Open Mosaic Habitat on
		assessment with consultation	Previously Developed Land (OMHPDL)
		bodies.	has been included as an IEF, as referred to by Thurrock Council.
4.5.3	Baseline conditions	Relevant consultation bodies have raised concerns regarding the proposed baseline data for the assessment in the Scoping Report. In particular concerns are raised in relation to the winter bird surveys required for the assessment of effects on designated sites. The Applicant should make effort to agree the scope and nature of baseline data required to support the assessment with relevant consultation bodies.	The ES and shadow HRA (Appendix 12.4; Document Reference 6.2.12.4) submitted along with the DCO application is informed by a comprehensive suite of wintering, passage and breeding bird survey data collated during winter 2019/2020 through to spring 2020. In addition, this recent data is supplemented by a suite of 'baseline' surveys completed during 2012/2013 on behalf of London Resort Company Holdings (LRCH) Ltd. Two seasons of bird survey data is submitted along with the DCO application, albeit not in consecutive years.
			The wetland habitats within the Kent Project Site have not changed significantly in the intervening years between the 2012/2013 baseline surveys and the recent 2019/2020 surveys, and the data collected is



PINS ID	Applicant's proposed matters to scope out	Inspectorate's comments	Action taken
			more or less consistent across that time period. As set out in the ES the wintering wader/wildfowl assemblage using the Kent Project Site has been valued at the International level owing to its association with the nearby Ramsar and SPAs. The assemblage, which is treated as an Important Ecological Feature (IEF) in the Ecological Impact Assessment, has therefore being valued at the highest level, and data from additional surveys will not affect this valuation or significantly alter the impact assessment and mitigation measures being developed.
			To further assess the ornithological interest at the Project Site a comprehensive desk study, based on Wetland Bird Survey (WeBS) data and local record centre data, has been updated in 2020.
			A meeting was held to discuss these matters with Natural England on 10 August 2020, and further justification on the scope of surveys was provided to Natural England within an Ecology Briefing Note (edp5988_r022) issued 21 August 2020 (see Appendix 12.5; Document Reference 6.2.12.5). At the time of writing, no further comment has been made by Natural England in response to the Ecology Briefing Note.
4.5.4	Temporal scope	The Scoping Report states that the assessment of ecological effects will be undertaken in the context of how the predicted baseline conditions might change between the	The anticipated future baseline conditions in the absence of development are set out in this ES Chapter. Changes in baseline conditions between the surveys and start of construction activities are



PINS ID	Applicant's proposed matters to scope out	Inspectorate's comments	Action taken
		surveys and the start of construction activities. The Scoping Report does not explain how this would be done, how it relates to the future baseline conditions referred to in paragraph 11.68 or if any additional survey work is proposed after consent prior to construction beginning. The ES should explain and justify any use of a future baseline.	likely to be very minor over that relatively short time period, however such changes would be identified through updated surveys proposed as part of the pre-construction mitigation measures, and the avoidance/mitigation strategy adjusted if necessary.

Preliminary Environmental Information Report (PEIR)

- <u>12.1012.11</u> As part of its pre-application duties, the Applicant commenced a statutory public consultation in July 2020. A Preliminary Environmental Information Report (PEIR) was prepared in support of this process to assist consultees in understanding the potential environmental effects of the Proposed Development and to enable consultees to develop an informed view of the project, ahead of a DCO application being made.
- 12.11 12.12 Relevant comments received on the PEIR from statutory and non-statutory consultees, and how these have been addressed, are provided in Appendix 12.6 and Appendix 12.7 (Document References 6.2.12.6 and 6.2.12.7) respectively.

Consultation Meetings/Correspondence

12.12

<u>12.13</u> In addition to the above, consultation has been held with parties with an interest or remit in biodiversity. **Table 12.2** provides a summary of the meetings and/or email/telephone correspondence with relevant consultees.

Table 12.2: Summary of consultation meetings and correspondence

Consultee	Details
Natural England, Environment	Environmental Liaison Group meeting on 18 May
Agency, Kent County Council,	2020. Introductory meeting to re-introduce consultees
Dartford Borough Council,	to the proposals and the scope of survey and
Gravesham Borough Council,	assessment to be undertaken to inform the DCO
Marine Management Organisation,	application.
Ebbsfleet Development	Green Infrastructure and Biodiversity Workshop, to
Corporation, Gravesham Borough	provide consultees an opportunity to ask the project's



Consultee	Details
Council, Kent County Council,	ecology and landscape consultants questions about
Natural England, Environment	the approach to survey, assessment and design.
Agency,	
Natural England (NE)	Prior to DCO Application Submission Email
	correspondence regarding terrestrial invertebrate
	survey methodology in late May and June 2020;
	 Meeting on 10 August 2020 to discuss NE
	response to EIA Scoping request;
	Ecology Briefing Note 'Natural England
	Consultation' (Report ref: edp5988_r022) and
	'Consultation Issues Tracker' submitted to NE on 21
	August 2020 in response to matters discussed on 10
	August 2020 and in response to NE's EIA Scoping
	comments. Briefing note provided further information
	on how the mitigation hierarchy has been applied and
	further justification on the robustness of the wintering
	bird survey data;
	• 'Marsh management' meeting on 24 August
	2020 along with the Environment Agency to discuss
	surface water drainage proposals, and proposed
	wetland creation, enhancement and management;
	 Second meeting on 24 August 2020 to discuss
	response to EIA Scoping request;
	 Meeting on 01 October 2020 to discuss NE
	response to PEIR;
	 Meeting on 16 October 2020 to discuss
	protected species licensing (water vole, otter, bat); and
	 Meeting on 20 October 2020 to discuss
	protected species licensing (dormouse).
	• Email correspondence regarding terrestrial
	invertebrate survey methodology in late May and June 2020;
	 Meeting on 10 August 2020 to discuss NE response to EIA Scoping request;



Consultee	Details
	 Ecology Briefing Note 'Natural England Consultation'
	(Report ref: edp5988 r022) and 'Consultation Issues
	Tracker' submitted to NE on 21 August 2020 in
	response to matters discussed on 10 August 2020 and
	in response to NE's EIA Scoping comments Briefing
	note provided further information on how the
	mitigation hierarchy has been applied and further
	justification on the robustness of the wintering bird
	survey data;
	• 'Marsh management' meeting on 24 August 2020
	along with the Environment Agency to discuss surface
	water drainage proposals, and proposed wetland
	creation, enhancement and management;
	 Second meeting on 24 August 2020 to discuss
	response to EIA Scoping request;
	 Meeting on 01 October 2020 to discuss NE response
	to PEIR;
	Marking and Contains 2020 to discuss and advantage
	• Meeting on 16 October 2020 to discuss protected
	species licensing (water vole, otter, bat); and
	 Meeting on 20 October 2020 to discuss protected
	species licensing (dormouse).
	Post DCO Application Submission
	Meeting on 24 February 2021 to acknowledge
	acceptance of the DCO Application by PINS and
	request feedback on mitigation strategies shared with
	NE prior to submission, in particular for invertebrates;
	 Meeting on 24 March 2021 to discuss recent
	notification of Swanscombe Peninsula SSSI and
	implications for on-site mitigation and off-site
	compensation, and to agree approach to further
	consultation prior to examination;
	a Corios of topic based resetings informed by NE/
	Series of topic-based meetings, informed by NE's Relevant Representations on the DCO Application
	Relevant Representations on the DCO Application (dated 31 March 2021), during April and May 2021 as
	(dated 31 March 2021), during April and May 2021 as follows:
	IOHOWS.
	 Breeding and wintering birds (22 April)
<u> </u>	



Consultee	Details
	 Vascular plants (22 April)
	O Invertebrates (28 April)
	 Protected species licensing (29 April)
	O HRA and EIA (06 May)
	 Meeting on 07 March 2022 to discuss scope of
	updated documents
Environment Agency (EA)	Prior to DCO Application Submission Email
0 , , ,	correspondence regarding aquatic invertebrate survey
	methodology in late May 2020, and meeting 09 July
	2020 to discuss requirements for Water Framework
	Directive assessment and Ecological Impact
	Assessment;
	, 18666
	• 'Marsh management' meeting on 24 August
	2020 along with Natural England to discuss surface
	water drainage proposals, and proposed wetland
	creation, enhancement and management;
	ereation, emancement and management,
	Ecology Briefing Note 'Further Information in
	Respect of an EIA Scoping Opinion' (Report ref:
	edp5988 r019) submitted to EA on 13 August 2020 to
	provide further information on potential effects on a
	fish population within the river Thames, Swanscombe
	Marshes and River Ebbsfleet, and requirements for
	water quality and Water Framework Directive
	assessment; and
	assessment, and
	EA 'Response to LR Ecology Briefing Note' (Ref:
	KT/2020/127432/01 L01) received 15 September 2020
	confirming additional fish survey on River Ebbsfleet
	not required.
	Email correspondence regarding aquatic invertebrate
	survey methodology in late May 2020, and meeting
	09 -July 2020 to discuss requirements for Water
	Framework Directive assessment and Ecological
	Impact Assessment;
	a (March management) mosting at 24 August 2020
	• 'Marsh management' meeting on 24 August 2020
	along with Natural England E to discuss surface water
	drainage proposals, and proposed wetland creation,
	enhancement and management;



Consultee	Details
	 Ecology Briefing Note 'Further Information in Respect of an EIA Scoping Opinion' (Report ref: edp5988 r019) submitted to EA on 13 August 2020 to provide further information on potential effects on a fish population within the river Thames, Swanscombe Marshes and River Ebbsfleet, and requirements for water quality and Water Framework Directive assessment; and EA 'Response to LR Ecology Briefing Note' (Ref: KT/2020/127432/01-L01) received 15 September 2020 confirming additional fish survey on River Ebbsfleet not required.
Kent County Council	 Meeting on 13 August 2020 to discuss 'biodiversity elements' of the Proposed Development with Helen Forster (Senior Biodiversity Officer); Email to Helen Forster on 14 September, 16_September and 01 October 2020 requesting information on dormouse mitigation strategies for approved developments in local area, along with information on existing dormouse monitoring/recording schemes.
Kent Wildlife Trust	 Meeting to discuss ecological mitigation proposals on 24 June 2020 and 23 July 2020 with Richard Bloor and Nicky Britton-Williams (Wilder Towns Officer).
Buglife	Introductory meeting on 22 July 2020 with Jamie Robbins (Project Manager) to discuss scope of invertebrate surveys, approach to evaluation, and mitigation/compensation measures.

Identification of the Resource/Baseline Conditions

- 12.13 12.14 This assessment is based upon a review of the data obtained from previous ecology surveys, desk-based reviews and update ecology surveys commencing in winter 2019/2020 and continuing throughout 2020 undertaken between late 2019 and mid-2021, as discussed further below.
- <u>12.1412.15</u> Collectively, findings from the above have been used to identify the important ecological features/receptors (IEFs) within the Project Site and its potential zone of influence, which has in turn been used to inform the masterplanning and mitigation strategy and form the basis for the assessment of potential effects.

Previous Ecology Assessment



- <u>12.15</u>12.16 The Kent Project Site has previously been subject to a suite of ecological baseline surveys as set out in various reports prepared in 2012 which were summarised within the 2014 EIA Scoping Report received by the Secretary of State in November 2014.
- 12.16 12.17 The 2012 reports are appended to EDP's Ecology Baseline Report (Appendix 12.1; Document Reference 6.2.12.1), and include:
 - 2012 Desk Study and Phase I Habitat Survey Report (CBA, 2012);
 - 2012 Botanical Survey Report (CBA, 2012);
 - 2012 Amphibian Survey Report (CBA, 2012);
 - 2012 Breeding Birds Survey Report (CBA, 2012);
 - 2012 Terrestrial Invertebrate Survey Report (CBA, 2012);
 - 2012 Terrestrial Invertebrate Survey Supplementary Report (Spiders [Araneae] and related groups) (CBA, 2012); and
 - 2012/13 Wintering Birds Survey Report (CBA, 2013).
- 12.17 12.18 In addition, further surveys were undertaken on the Kent Project Site throughout 2015 and 2016, the findings of which are provided in full in the following reports:
 - Phase 1 and Botanical Survey Report (CBA February 2016);
 - Wintering Bird Survey Report (Corylus Ecology April 2016);
 - Common Bird Survey Report (Corylus Ecology April 2016);
 - Bat Activity Report 2015 (Corylus Ecology June 2016);
 - Dormouse Report (Corylus Ecology February 2016);
 - 2015 Harvest Mouse Survey Report (CBA February 2016);
 - 2015 Water Vole Survey Report (CBA February 2016);
 - 2015 Badger Survey Report (CBA February 2016);
 - 2015 Amphibian Survey Report (CBA February 2016);
 - 2015 & 2016 Reptile Survey Report (CBA August 2016);



- Invertebrate Survey and Assessment of the London Paramount Entertainment Resort 2015 (Edwards Ecological Services, 2015);
- An ecological survey of the waterbodies and wetlands on and around the Swanscombe Peninsula, Kent (Aseda, 2016); and
- A targeted ecological survey of selected waterbodies and wetlands on the Swanscombe peninsula, Kent (Aseda, 2016);
- Fish survey of Swanscombe Marshes (Colclough and Coates, 2015); and
- Fish survey of the Ebbsfleet Stream (Colclough and Coates, 2015).
- 12.1812.19 These reports are appended in full, and the key findings summarised, within Appendix 12.1 (Document Reference 6.2.12.1).

2019/-2020-2021 Update Ecology Surveys

- 12.1912.20 The update ecological information obtained for the Project Site by EDP is set out in detail within Appendix 12.1 (Document Reference 6.2.12.1). The appendix details the full survey methodologies that have been employed, any findings from surveys completed and analysed and any limitations. It seeks to identify the IEFs within the Project's ecological zone of influence and defines the zone of influence for different receptors.
- 12.2012.21 A summary of the update ecology surveys undertaken across the Project Site in between 2019 and 2020-2021 is provided below:
 - Desk study (April 2020);
 - Extended Phase 1 Habitat Survey (May 2020);
 - Botanical survey (June 2020);
 - Wintering bird surveys (winter 2019/2020 and winter 2020/2021);
 - Breeding bird surveys (April to July 2020) and passage bird surveys (April, August and September 2020) and nocturnal long-eared owl survey July 2021);
 - Bat surveys, including bat roost surveys and bat activity surveys (May to September October -2020) and winter hibernation surveys December 2020 to February 2021);
 - Dormouse surveys (April to October 2020);
 - Badger survey (May 2020 and October 2020);



- Otter surveys (June and August 2020 across the Kent Project Site, with Botany Marsh West surveyed later in July and September 2020, and updated between March and July 2021);
- Water vole surveys (June, August and September 2020 across the Swanscombe Peninsula, with Botany Marsh West surveyed later in September 2020, and updated between March and July 2021);
- Harvest mouse surveys (October 2020);
- Great crested newt survey (April 2020);
- Reptile surveys (April to September 2020);
- Terrestrial invertebrate surveys, including an invertebrate scoping study (April 2020) and detailed invertebrate sampling (May to August 2020);
- Aquatic invertebrate surveys, including an invertebrate scoping study (April 2020) and detailed invertebrate sampling (May, July and August 2020);
- River corridor survey and river habitat survey (May 2020); and
- Freshwater fish surveys of suitable waterbodies across Swanscombe Marshes (September 2020).
- 42.2112.22 All surveys have been undertaken with reference to best practice guidance where available. Any limitations in the survey work are detailed in Appendix 12.1 (Document Reference 6.2.12.1) and summarised below under 'Assumptions and Limitations'.

Assessment Methodology

12.2212.23 The Assessment Methodology adopted by this chapter follows that set out in the EcIA Guidelines, as explained in the following sections.

Ecological Zone of Influence or Geographical Scope (Study Area)

The Ecological Zone of Influence (EZoI) is an area defined by the assessment in which there may be receptors subject to effects as part of the Proposed Development, both those which may occur as a result of land-take and habitat loss and those which may occur through disturbance such as noise. Such receptors are likely to include designated sites, notable habitats and protected species, and these could be affected directly, e.g. construction works affecting a receptor within the Project Site such as removal of a tree occupied by roosting bats, or indirectly, e.g. a designated site located down river of a development being affected by increased storm run-off, etc.



12.2412.25 The EZoI was determined through:

- A review of existing baseline conditions in comparison with that proposed by the Proposed Development;
- Consideration of the proposed activities during construction and operation;
- Desk study information including an examination of mapping data;
- Consultation responses; and
- Findings of the survey work.
- <u>12.25</u>12.26 The EZoI is defined as the areas/resources that may be affected by the biophysical changes caused by activities associated with the Proposed Development. The definition of the EZoI used in this assessment is taken from the EcIA Guidelines (CIEEM 2018) described above.
- <u>12.26</u>12.27 Due to the scale and nature of the proposals, the EZoI includes all land within the DCO Order Limits. In addition, the study area for the desk study extends beyond the DCO Order Limits in order to take account of potential for effects to occur within a wider geographical area.
- <u>12.27</u> The following desk study search radii, taken from the boundary of the DCO Order Limits, -were employed and are considered to be sufficient to cover the ecological zone of influence of the Proposed Development:
 - International statutory designations (15km);
 - National statutory designations (5km);
 - Non-statutory locally designated sites (2km);
 - Annex II bat species³ records (6km); and
 - All other protected/notable species records (2km).
- The field surveys undertaken to inform the assessment cover the Project Site, but are tailored to the individual habitats present on either the Kent Project Site or the Essex Project Site and their suitability to support protected species orf species of conservation concern. Further details are provided in Appendix 12.1 (Document Reference 6.2.12.1).

³ Bat species listed in Annex II of the *EC Habitats Directive*, namely Greater horseshoe, Lesser horseshoe, Barbastelle and Bechstein's bats



Ecology and Biodiversity Evaluation

- 42.2912.30 An evaluation of IEFs has been made with reference to the EcIA Guidelines. A summary of the evaluation approach is provided below.
- <u>12.3012.31</u> The guidelines advocate an approach to valuing features that involves professional judgement based on available guidance and information, together with advice from experts, who know the locality of the project and/or the distribution and status of the species or features that are being considered.
- 12.31 12.32 The EcIA Guidelines recommend that the value or potential value of an ecological resource or feature should be determined within a defined geographical context, and the EcIA Guidelines provide a geographical range ('frame of reference') that can be adapted. The geographical frame of reference used in this assessment is as follows:
 - International value (SACs, SPAs, Ramsar sites);
 - National value (within England), e.g. Sites of Special Scientific Interest (SSSIs) and National Nature Reserves (NNRs);
 - Regional value (within south-east England);
 - County value (within Kent or Essex), e.g. Local Nature Reserves, Local Wildlife Sites, atypical and diverse species assemblages with good population sizes;
 - District value (within the Borough of Dartford, Borough of Gravesend or Thurrock Council), e.g. where species rich/extensive/atypical examples are present – moderate population sizes or species assemblages with moderate to high diversity;
 - Local value (within Swanscombe and Greenhithe Civil Parish, Bean Civil Parish, or Southfleet Civil Parrish, or the towns of Northfleet, Gravesend or Tilbury), e.g. common and widespread species with relatively moderate populations and relatively limited diversity;
 - Site value (the Project Site and immediate environs): small areas of common habitats such as species-poor grassland and scrub – common and widespread species with small populations and limited diversity; and
 - Negligible value (typically applied to areas of bare open ground/built development/areas of hardstanding).



Designated Sites

- 12.32 Some sites have already been assigned a level of nature conservation value through designation, and the EcIA Guidelines recommend that the reasons for this designation need to be taken into account in the assessment. Such designations include:
 - Internationally important sites such as SACs, SPAs and Ramsar sites;
 - Nationally important sites such as SSSIs and NNRs;
 - Regional/County important sites such as LNRs; and
 - County/District important sites such as LWSs.
- <u>12.33</u>12.34 Where a feature has value at more than one designation level, its overriding value is that of the highest level.

Habitats

The EcIA Guidelines recommend that the value of areas of habitat and plant communities should be measured against published selection criteria where available. Examples include criteria for Local Wildlife Site selection in Kent⁴ and criteria for Priority Habitats/Habitats of Principal Importance in England, although the value of specific Priority Habitats must then be judged on a case by case basis. Where areas of a habitat or plant community do not meet the necessary criteria for designation at a specific level, the EcIA Guidelines recommend that the ecologist may consider the local context if appropriate. It also emphasises the importance of not under-estimating habitats in suboptimal condition where there is potential for restoration.

Species

- 12.35 12.36 The guidance deals with species that need to be assessed because they are of biodiversity value, rather than because they are legally protected (although some species may be legally protected as well as being of biodiversity value).
- <u>12.36</u>12.37 In assigning value to a species, it is necessary to consider its distribution and status, including a consideration of trends based on available historical records. The valuation of populations should make use of any relevant published evaluation criteria.

Assessment of Potential Impacts

<u>12.37</u>12.38 The EcIA Guidelines advocate an approach to assessing likely impacts that involves professional judgement based on available guidance and information.

⁴ Kent Wildlife Trust (2015) "Local Wildlife Sites in Kent. Criteria for Selection and Delineation". Adopted by the Kent Nature Partnership October 2015, Version 1.5.



- 12.38 12.39 The assessment of the potential impacts of the Proposed Development takes into account both on-site impacts and those that may occur to adjacent and more distant ecological features. Impacts can be permanent/temporary, direct or indirect, positive or negative and can include:
 - Direct loss of wildlife habitats;
 - Fragmentation and isolation of habitats;
 - Disturbance to species from noise, light or other visual stimuli;
 - Changes to key habitat features; and/or
 - Changes to the local hydrology, water quality and/or air quality.
- 12.3912.40 The significance of a negative effect (or a positive effect) is the product of the magnitude of the impact and the value or sensitivity of the nature conservation features affected. In order to characterise the impacts on each feature, the following parameters are taken into account:
 - The magnitude of the impact;
 - The spatial extent over which the impact would occur;
 - The temporal duration of the impact;
 - Whether the impact is reversible and over what timeframe; and
 - The timing and frequency of the impact.

Significance of Effects

- 12.4012.41 There is no agreed absolute method for assessing the significance of negative or positive impacts on nature conservation features within the EcIA Guidelines. In addition, since the purpose of an EIA is to focus on likely significant effects, it is not reasonable to expect the assessment to include every ecological feature that may be affected, since effects are unlikely to be significant where features of low (Site level or below) value or sensitivity are, for example, subject to low or short-term impacts. On this basis, the assessment therefore focuses on ecological features that are considered, based on professional judgement, experience and contextual information, to be protected and/or of Local nature conservation value or above.
- 12.4112.42 However, this does not mean that effects upon features of less than Local level nature conservation value have been discounted. Certain species and habitats that may not constitute IEFs based upon their nature conservation value, may still warrant



consideration during the design of the development (and any mitigation identified) on the basis of their legal protection, their implications for policies and plans, or other issues, such as animal welfare.

- 12.42 12.43 The EcIA Guidelines also recommend that where ecosystem service provision (benefits that people derive from the natural environment) might be affected as a result of a project's ecological effects, this should be recognised, and the relevant data collected during the EcIA to inform separate specialist assessments of social and economic value. This can enable the social and economic implications of ecological changes to be taken into account.
- 12.43 12.44 The assessment identifies those positive and negative impacts which would be 'significant', based on effects that either support or undermine the conservation objectives of the ecological feature or biodiversity in general. Significant effects encompass impacts on structure and function of defined sites, habitats or ecosystems and the conservation status of habitats and species (including extent abundance and distribution)'. Such significant effects are qualified with reference to an appropriate geographic scale and based on the best available scientific evidence. Where it is not possible to robustly justify that no significant effect will occur, a significant effect is assumed.
- 12.4412.45 The integrity of 'designated' sites is described as follows and is taken from the EcIA Guidelines. It has been used in this assessment to determine whether the impacts of the Proposed Development on a designated site are likely to be significant:

'Significant effects encompass impacts on structure and function of defined sites and ecosystems. The following need to be determined:

- for designated sites is the project and associated activities likely to undermine the site's conservation objectives of the site, or positively or negatively affect the conservation status of species or habitats for which the site is designated, or may it have positive or negative effects on the condition of the site or its interest/qualifying features?'
- 12.45 The conservation status of habitats and species within a defined geographical area is described in the EcIA Guidelines as follows and has been used in this chapter to determine whether the impacts of the Proposed Development on non-designated habitats and species are likely to be significant:

'Consideration of conservation status is important for evaluating the effects of impacts on individual habitats and species and assessing their significance:

• habitats - conservation status is determined by the sum of the influences acting on the habitat that may affect its extent, structure and functions as well as its distribution and its typical species within a given geographical area;



- species conservation status is determined by the sum of influences acting on the species concerned that may affect its abundance and distribution within a given geographical area.'
- <u>12.46</u>12.47 On the basis of the above, and within this assessment, ecological effects are described as either:
 - Significant or not significant;
 - Direct and/or indirect;
 - Permanent or temporary; and
 - Negative or positive.
- <u>12.47</u>12.48 Mitigation measures have been taken into account during the assessment of effects, such that the residual impact assessment reflects the completed scheme. These measures include those required to achieve the minimum standard of established practice plus additional measures to further reduce the effects of the scheme. The assessment takes into account the likely success of the mitigation.
- <u>12.48</u> The significance of the likely impacts upon IEFs has been assessed both before and after consideration of additional mitigation measures. The latter represents the assessment of the residual impacts of the project.
- <u>12.49</u>12.50 In relation to EIA and assessment of significant effects, the EcIA Guidelines highlight that:
 - 'A significant effect does not necessarily equate to an effect so severe that consent for the project should be refused planning permission. For example, many projects with significant negative ecological effects can be lawfully permitted following EIA procedures.'
- <u>12.50</u>12.51 In addition to determining the significance of an impact on any ecological features, this chapter also identifies any legal requirements in relation to wildlife.

Temporal Scope

12.5112.52 The assessment of potential ecological effects resulting from the Proposed Development will be undertaken in the context of how the predicted baseline conditions within the EZoI might change between the surveys and the start of construction activities. It is anticipated a phased work programme for the Proposed Development will commence in 2022, with Gate One opening in 2024, Gate Two in 2029, with full maturity being achieved by 2038.

Cumulative Effects



- 12.52 12.53 Cumulative effects generally occur where there may be simultaneous or prolonged similar effects on the same habitats or species populations as a result of two or more developments of the same type and scale, or where the consideration of other schemes would increase an effect identified. Where other schemes are relevant, these are considered in conjunction with the Proposed Development.
- 12.5312.54 Those cumulative impacts that can be foreseen as a result of the Proposed Development in conjunction with the identified schemes, and any possible mitigation measures required, are assessed in 'Cumulative and in-combination assessment' section of this chapter.

Assumptions and Limitations

12.5412.55 The vast majority of surveys have been undertaken in suitable weather conditions at optimum times of year with reference to best practice guidance. Limitations specific to each of the field survey methodologies are detailed within the appropriate sections of Appendix 12.1 (Document Reference 6.2.12.1) and are summarised below.

Access to Botany Marsh West

12.5512.56 Whilst landowner permission was granted in spring 2020 to undertake environmental DNA (eDNA) testing for the presence of great crested newts (Triturus cristatus), permission to access 'Botany Marsh West' for other surveys was not granted until mid-July 2020. Therefore, no additional survey work was undertaken within the interior of the fields here prior to this, including habitat surveys or wintering bird surveys, breeding bird surveys, and bat surveys. There is some possibility therefore that some species occupying the interior of the fields, such as snipe or skylark, may have gone unrecorded but these species were recorded elsewhere on the Kent Project Site by EDP in 2020 and therefore assumed potentially present in Botany Marsh West. Furthermore, prior to July 2020 surveys were conducted on similar habitat close to, or adjacent to, the marsh and so it is reasonably assumed that any species confirmed present in suitable, connected habitat near to/adjacent to the marsh are also present in these areas. Following receipt of permission for ecology surveys, the following surveys were completed: Extended Phase 1 Habitat Survey and botanical survey (July 2020), terrestrial and aquatic invertebrate sampling (July and August 2020), water vole surveys (1 visit July 2020, and 1 visit September 2020), harvest mouse survey (October 2020), and fish survey of the ditch network (end-September 2020).

Access to Waterbodies

12.5612.57 During the course of the eDNA surveys to confirm the presence/absence of great crested newts, access was not possible to every single one of the ponds and ditches on or adjacent to the Project Site. However, a large number of waterbodies within the Kent Project Site were surveyed and all eDNA analysis returned a negative result (i.e. no evidence of the presence of great crested newt DNA), and there were no recent records returned during the desk study. On this basis, it is considered a fair assumption that great



crested newts are unlikely to be present on the Kent Project Site. Furthermore, due to the lack of suitable habitat, great crested newts are considered highly unlikely to be present on the Essex Project Site.

- 12.5712.58 The lack of access to some water bodies also posed a limitation to the otter and water vole survey. However, presence of water vole within the Kent Project Site was confirmed during the first survey in June 2020, and it is assumed that water vole could utilise other parts of the ditch network due to the presence of suitable habitat. On a precautionary basis therefore, the assessment assumes that water vole could be present in the other suitable water bodies on the Kent Project Site. Similarly, with regard to the presence of otter, an otter was sighted within Black Duck Marsh in March 2020, and it is assumed that otters may be present in other parts of the Kent Project Site such as along the River Ebbsfleet and the ditch network due to the presence of suitable habitat.
- The lack of access to some water bodies posed a limitation to freshwater fish surveys undertaken across Swanscombe marshes, whilst high alkalinity levels in some waterbodies reduced the effectiveness of electrofishing operation. However, it is considered that the results of surveys undertaken can be applied to those waterbodies to which access was not possible, given the relative uniformity of habitat conditions, topography and flow. The findings of the survey are, furthermore, consistent with previous survey effort undertaken during 2015 and a habitat suitability assessment, in respect of fish.

Access to Buildings/Structures

- Limits, of which 117 were assessed as having negligible potential to support roosting bats, with 23 buildings found to have some potential, and 26 buildings (16% of the total) which could not be adequately assessed due to access restrictions including permission not been granted by the landowner or buildings being inaccessible due to dense scrub. In light of the 'lockdown' and social distancing measures imposed as a result of the COVID-19 virus global pandemic, internal access to buildings has not been possible for all buildings. For these 26 buildings where this is a lack of information, a precautionary approach to the assessment of potential effects has been followed in this chapter, with precautionary mitigation measures detailed within the 'Bat Mitigation Strategy' enclosed within the EMMF (Document reference: 6.2.12.3), as advised by Natural England in their Discretionary Advice Service letter of 9th October 2020 (copy of which is enclosed as Annex EDP 13 to the EMMF (Document reference: 6.2.12.3)).
- 12.6012.61 Tunnels TU/014A and TU/015 within Craylands Pit (see Appendix 12.1; Document Reference 6.2.12.1) were not surveyed beyond their initial inspection due to health and safety constraints relating to access and night-time surveys. As such no summer roosting or autumn swarming bat surveys have been undertaken on these tunnels. In addition, there was no access for an autumn swarming survey on tunnel TU/012 in August due to positioning of the boarding across the entrance, this was later moved to allow access for the microphone. Given access constraints to the tunnels it was not always possible to



position static detectors during the autumn swarming <u>and winter hibernation</u> surveys to ensure only bats from within the tunnel are recorded. As such a number of recordings may be made from foraging bats outside the tunnel entrances. This has been taken into account during the interpretation of survey findings.

Access to Bamber Pit

- <u>12.61</u>12.62 Due to access and security issues, bat transects and static detector deployment could not take place in Bamber Pit in July and August 2020. However, the remainder of the Kent Project Site was surveyed, including areas close to Bamber Pit, and Bamber Pit was surveyed in May, June and September 2020. It is considered unlikely that the overall value of the bat assemblage has been underestimated as a result of missing the July and August 2020 survey.
- Due to access and security issues, the reptile refugia within Bamber Pit were not checked after the second survey visit in May 2020, until these issues were resolved in September 2020, with the remaining visits then completed in September 2020. A total of seven survey visits were able to be conducted. Reptile surveys were therefore completed within the recognised survey period of April to September, and reptiles were found during the visits, thus confirming presence within these areas of the Kent Project Site. The reptile refugia within and around Botany Marsh were all checked in September 2020, which is considered an optimal month for recording reptiles.

Damage to Reptile Refugia

12.63 12.64 Following the first reptile survey in May 2020, some of the artificial refugia in the north of the Kent Project Site were moved and damaged due to mowing activity. However, reptiles were still recorded in this area, and replacement refugia were deployed in June 2020 and remained in place for the following checks, allowing for an accurate presence/absence survey and population estimate.

General Limitations

- 12.6412.65 Owing to the seasonality of some species, as well as the ability for some species to quickly colonise sites (such as badger), the absence of evidence of any particular species from within the Project Site should not be taken as conclusive evidence that the species is not present or that it will not be present in the future.
- 12.6512.66 Overall, it is considered that the results of the Phase 1 survey and additional Phase 2 surveys undertaken in 2019 and 2020 are robust and reliable for the identification of the habitats and the presence or absence of legally protected/notable species within the Project Site, and sufficient to identify the IEFs requiring further consideration in the EcIA.
- 12.6612.67 The scope of surveys undertaken at the Essex Project Site is considered a robust level of survey effort based on the nature of the habitats present, being largely developed



land of negligible intrinsic ecological value. The surveys undertaken at the Essex Project Site are not considered to pose a limitation to the robustness of the EcIA.

RELEVANT LAW, POLICY AND GUIDANCE

Legislative and Policy Context

<u>12.68</u> There are a number of different legislative instruments that are relevant to the assessment of potential effects of the construction and operation of the London Resort. The key international legislation that has been considered is set out within **Table 12.3** <u>Table 12.3: A Summary of Legislation and Conventions of Relevance to the Proposed Development.</u>

12.6712.69 -

Table 12.3: A Summary of Legislation and Conventions of Relevance to the Proposed Development.

Title	Summary and Relevance	
International		
Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC) (the 'Habitats Directive')	Provides a framework for the conservation and management of natural habitats, wild fauna (except birds) and flora in Europe. Its aim is to maintain or restore natural habitats and wild species at a favourable conservation status. The relevant provisions of the Directive are the identification and classification of Special Areas of Conservation (SACs) (Article 4) and procedures for the protection of SACs (Article 6). SACs are identified based on the presence of natural habitat types listed in Annex I and populations of the species listed in Annex II. The Directive requires national Governments to establish SACs, and to have in place mechanisms to protect and manage them. SACs are also termed Natura 2000 sites, and those that are covered by tidal water (continuously or intermittently) are also termed 'European Marine Site' (EMS) - although this is not a statutory site designation.	
Birds Directive - Council Directive 2009/147/EC on the Conservation of Wild Birds	This Directive provides a framework for the conservation and management of wild birds in Europe. The most relevant provisions of the Directive are the identification and classification of Special Protection Areas (SPAs) for rare or vulnerable species listed in Annex I of the Directive and for all regularly occurring migratory species (required by Article 4). It also establishes a general scheme of protection for all wild birds (required by Article 5). The Directive requires national Governments to establish SPAs and to have in place mechanisms to protect and manage them. The SPA protection procedures originally set out in Article 4 of the Birds Directive have been replaced by the Article 6 provisions of the Habitats Directive.	



Title	Summary and Relevance
Ramsar Convention on Wetlands of International Importance (1972)	In accordance with Government advice in both England and Wales, Ramsar sites (internationally important wetlands) must be given the same consideration as European sites, so they are afforded the same protection as those under the Habitats Directive - Council Directive 92/43/EEC on the Conservation of Natural Habitats and of Wild Fauna and Flora.
Water Framework Directive (WFD) (2000/60/EC)	The WFD establishes a framework for the management and protection of Europe's water resources. It is implemented in England and Wales through the Water Environment (Water Framework Directive) (England and Wales) Regulations 2017 (the Water Framework Regulations). Central to the WFD is the philosophy to make water bodies better through sustainable development for the joint benefits of aquatic habitats and the human environment.
	Ecological status is an expression of the quality of the structure and functioning of surface water ecosystems as indicated by the condition of a number of 'quality elements'. These include biological, hydromorphological and chemical indicators. The development and implementation of strategic long-term River Basin Management Plans (RBMPs) is a key requirement of the WFD. They include a programme of measures outlining the on-going monitoring and management actions required for water bodies to achieve future objectives.
	Proposed developments or activities that have the potential to affect the water environment require a WFD Assessment. In this context, compliance with the WFD means prevention of deterioration (of ecological status, chemical status and supporting element status) and prevention of ability to achieve future targets. However, WFD Article 4.7 provides legislation for exemption conditions that could allow implementation of schemes that cause deterioration in ecological status, for example for reasons of overriding public interest.
EU Alien Invasive Species Regulation (Regulation No 1143/2014)	Regulation (EU) 1143/2014 on invasive alien species (the IAS Regulation) entered into force on 1 January 2015, fulfilling Action 16 of Target 5 of the EU 2020 Biodiversity Strategy. The regulation is designed to establish a framework for action to prevent, minimise and mitigate the adverse impacts of invasive non-native species on biodiversity and ecosystem services, and focuses on a list of invasive alien species of EU concern, which has been drawn up with Member States using risk assessments and scientific evidence. Selected species are banned from the EU, meaning it will not be possible to import, buy, use, release or sell them. The proposal is for three types of intervention: prevention; early warning and rapid response; and management.



Title	Summary and Relevance
Convention on the	This Convention was adopted in Bern, Switzerland in 1979, and came
Conservation of	into force in 1982. The principal aims of the Convention are to ensure
European Wildlife	conservation and protection of all wild plant and animal species and
and Natural Habitats	their natural habitats (listed in Appendices I and II of the Convention),
(the 'Berne	to increase co-operation between contracting parties, and to afford
Convention')	special protection to the most vulnerable or threatened species
	(including migratory species).
Convention on the	This Convention was adopted in Bonn, Germany in 1979 and came into
Conservation of	force in 1985. Contracting Parties work together to conserve migratory
Migratory Species of	species and their habitats by providing strict protection for endangered
Wild Animals (the	migratory species (listed in Appendix 1 of the Convention), concluding
'Bonn Convention')	multilateral agreements for the conservation and management of
	migratory species which require or would benefit from international co-
	operation (listed in Appendix 2 of the Convention), and by undertaking
	co-operative research activities.
Convention on	The Convention focuses on the conservation of all species and
Biological Diversity	ecosystems and, therefore, provides protection to all biodiversity. The
1992	Convention requires the development of national strategies, plans or
	programmes for the conservation and sustainable use of biodiversity, its
	sustainable use, and equitable sharing of benefits arising from the
	utilisation of natural resources (i.e. the Ecosystem Approach). In
	accordance with this, the UK developed Biodiversity Action Plans
	(BAPs).
National	
The Conservation of	The Conservation of Habitats and Species Regulations 2017 (the
Habitats and Species	'Habitats Regulations') consolidate the Conservation of Habitats and
Regulations (2017)	Species Regulations 2010 with subsequent amendments. The
	Regulations transpose Council Directive 92/43/EEC, on the conservation
	of natural habitats and of wild fauna and flora (EC Habitats Directive),
	into national law. They also transpose elements of the EU Wild Birds
	Directive in England and Wales, and make provision for the protection
	and management of sites, including the control of potentially damaging
	operations that may affect designated sites. The Regulations came into
	force on 30 November 2017.



Title	Summary and Relevance
Wildlife and	This Act is the principal mechanism for the legislative protection of
Countryside Act	wildlife in Britain. It consolidates and amends existing national
1981	legislation to implement the Convention on the Conservation of
	European Wildlife and Natural Habitats (Bern Convention) and Council
	Directive 79/409/EEC on the conservation of wild birds (Birds Directive)
	in Great Britain. The Act provides for the designation of Sites of Special
	Scientific Interest (SSSIs), which are selected as the best national
	examples of habitat types, sites with notable species and sites of
	geological importance. Various species of marine animals are also
	protected from being killed, injured or disturbed under provisions in
	Schedule 5 of the WCA 1981. It is also the principal UK legislation
	dealing with non-native species.
National	The NERC Act 2006 makes provision for bodies concerned with the
Environment and	natural environment and rural communities, amends protection for
Rural Communities	some designated wildlife areas, and amends the law relating to rights of
(NERC) Act 2006	way. Section 41 of the Act required the Secretary of State to publish a
	list of habitats and species of principal importance for the conservation
	of biodiversity in England. This list (the S41 list) includes habitats and
	species which have been identified as requiring action in the UK
	Biodiversity Action Plan (UK BAP), in line with the 1992 Convention on
	Biological Diversity.
The Protection of	Legislation primarily aimed at preventing cruel treatment and
Badgers Act 1992	persecution of badgers. Makes it an offence to kill, injure or take a
	badger and intentionally or recklessly damage or destroy a badger sett,
	or obstruct access to it.
The Water	The WRA regulates water resources, water quality, pollution and flood
Resources Act 1991	defence. The policing of this act is the responsibility of the Environment
(WRA)	Agency; under the act is an offence to cause or knowingly permit any
	poisonous, noxious or polluting material, or any solid waste to enter
	any controlled water.
Salmon and	Makes it an offence to discharge effluents which may damage fish, their
Freshwater Fisheries	food or their spawning grounds.
Act 1975 (as	
amended)	
Water Act 2014	Part 3 of the Water Act 2014 focusses on the Environmental Permitting
	regime in relation to water abstraction and pollution prevention and
Th 14/-1 -	control (enabling single rather than multiple permit applications).
The Water	Key transposing Directions that set out the environmental standards to
Framework	be used for the second cycle of river basin plans.
Directive (Standards	
and Classification)	
Directions (England	
and Wales) 2015	



Policy

<u>12.68</u>12.70 There are a number of national and local policies that are of relevance to the London Resort ecological assessment process (summarised in **Table 12.4**).

Table Error! No text of specified style in document.12. Error! No text of specified style in document.4: Summary of national and local policies of relevance to the London Resort.

Title	Summary and Relevance
National	
National Policy Statement (NSP) for National Networks (2014)	Whilst there is no NPS for business and commercial NSIP project, the extent that the Proposed Development includes transport and highways infrastructure means that regard will be had to the NPS on National Networks, including:
	 Environmental and Social impacts (NPS paragraphs 3.2 to 3.5);
	 Climate Change Adaptation (NPS paragraphs 4.36 to 4.47);
	 Pollution Control and Other Environmental Protection Regimes (NPS paragraphs 4.48 to 4.56);
	 Biodiversity and ecological conservation (NPS paragraphs 5.20 to 5.38);
	Flood Risk and Mitigation (NPS paragraphs 5.90 to 5.115); and
	Water Quality and Resources (NPS paragraphs 5.219 to 5.231).
National Planning Policy Framework, 2019 2021	The revised latest National Planning Policy Framework (NPPF) was published in 2019 July 2021. The NPPF is a material consideration that must be taken into account in the determination of planning applications. The NPPF requires that an overall approach is taken to sustainable development, incorporating social, economic and environmental dimensions which should not be considered in isolation. A section of the NPPF document addresses 'Conserving and enhancing the natural environment'. Amongst other objectives this section indicates when determining planning applications, that local planning authorities should aim to contribute to and enhance the natural and local environment by applying a number of outlined principles. Planning Practice Guidance on the natural environment supports the NPPF by explaining the government's planning policies for England, in regard to (amongst others) the protection and enhancement of biodiversity, ecosystem and green infrastructure, and how these are expected to be applied.



Title	Summary and Relevance
National	
UK Post-2010 Biodiversity Framework	The UK Post-2010 Biodiversity Framework has succeeded the UK Biodiversity Action Plan (UK BAP). The Framework demonstrates how the work of the four countries and the UK contributes to achieving the Aichi Biodiversity Targets, and identifies the activities required to complement the country biodiversity strategies in achieving the targets. Although the UK BAP has been superseded, the list of UK BAP priority habitats and species remains a useful reference for local authority decision-makers and forms the basis of the Section 41 list of the NERC Act.
Local	
Thurrock Council Local Plan	The Thurrock Borough Local Plan 1997 sets out the Council's policies and proposals for the way in which land, buildings and infrastructure should be developed. By law, although the end date of the Borough Local Plan has passed, its policies are a material consideration when deciding planning applications. The Council are in the process of developing a new Local Plan for Thurrock, however, this is in the early stages of development and is not anticipated to be adopted until 2021. The development plan for Thurrock comprises the following documents: Thurrock Core Strategy and Policies for Management of Development and Policies Map, as amended, adopted January 2015 and Thurrock Borough Local Plan Saved Policies, Site Allocations and Annexes, February 2012. Policies relevant to this chapter include: Policy CSTP18: Green Infrastructure; and
	Policy CSTP19: Biodiversity.
Dartford Borough Local Plan	The Dartford Borough Local Plan 2011 sets out the Council's policies and proposals for the way in which land, buildings and infrastructure should be developed. The development plan for Dartford comprises the following documents: Dartford Core Strategy, adopted September 2011 and Dartford Development Policies Plan, adopted July 2017. Policies relevant to this chapter include: Policy CS14: Green Space.
Gravesham Borough Local Plan	The Gravesham Borough Local plan 2014 Plan sets out the Council's policies and proposals for the way in which land, buildings and
	infrastructure should be developed.



Title	Summary and Relevance
National	
	The development plan for Gravesham comprises the following documents: Gravesham Local Plan Core Strategy and Local Plan Policies Map, adopted September 2014 and Local Plan Review, undertaken in September 2019.
	Policies relevant to this chapter include: • Policy CS12: Green Infrastructure.
	Following the Regulation 18 (Stage 1) Consultation of the Gravesham Borough Council emerging Local Plan, the Council are undertaking a Regulation 18 (Stage 2) consultation to seek views on the updated policies in the Development Management Policies Document. Proposed 'draft' policies relevant to this chapter include:
	Policy GI 6: Biodiversity.

Guidance

12.6912.71 The key guidance document used to inform this assessment is the EclA Guidelines (CIEEM 2018) described above. Other relevant guidance can be found in Table 12.Table 12.5 below.

Table 12.5: A Summary of Guidance followed for the Proposed Development.

Title	Summary and Relevance	
National		
Advice Note 7:	This Advice Note explains the Environmental Impact	
Environmental Impact	Assessment (EIA) process set out in the Infrastructure Planning	
Assessment: Preliminary	(Environmental Impact Assessment) Regulations 2017 (the EIA	
Environmental	Regulations). In particular the note addresses the procedures	
Information, Screening and	for EIA screening and scoping; notification and consultation;	
Scoping (PINS, 2020)	matters relating to the production of Preliminary	
	Environmental Information (PEI) and the preparation of	
	Environmental Statements (ES).	
Advice Note 9: Rochdale	This Advice Note addresses the use of the 'Rochdale Envelope'	
Envelope (PINS, 2018)	approach under the Planning Act 2008, as amended (PA2008).	



Title	Summary and Relevance
Advice Note 10: Habitats Regulations Assessment relevant to nationally significant infrastructure projects (PINS, 2017)	When preparing applications for NSIPs under the PA2008, Applicants should consider the potential effects of the application on protected habitats. If an NSIP, when taken alone or with existing and known future projects, is likely to affect a European site and/or a European Marine site, the Applicant must provide a report with the application showing the site(s) that may be affected together with sufficient information to enable the competent authority to make an Appropriate Assessment, if required. This Advice Note provides advice for Applicants in relation to the preparation of that report, and the PA2008 processes relating to HRA.
Advice Note 12: Transboundary Impacts and Process (PINS, 2018)	This Advice Note explains the legal context for transboundary impacts, and the NSIP transboundary process (including special arrangements) that will be followed by the Inspectorate on behalf of the Secretary of State during the pre-application, examination and recommendations stages of a DCO application. The Advice Note also explains how Applicants, States and the public can participate more generally in the DCO process where transboundary impacts occur.
Advice Note 17: Cumulative effects assessment relevant to nationally significant infrastructure projects (PINS, 2019)	A range of public sector and industry-led guidance is available on Cumulative Effects Assessment (CEA) but at present there is no single, agreed industry standard method. Consequently, the approach to CEA varies between applications. This Advice Note sets out a staged process that Applicants for the London Resort DCO are adopting in CEA for NSIPs.
Advice Note 18: The Water Framework Directive (PINS, 2017)	The purpose of this Advice Note is to alert Applicants to the requirements of the WFD and 2017 Regulations, as applicable to NSIPs under the PA2008. This Advice Note explains the information that the Inspectorate considers an Applicant must provide with their NSIP application in order to clearly demonstrate that the WFD and the 2017 Regulations have been appropriately considered.
The Design Manual for Roads and Bridges (DMRB) Volume 11: Environmental Assessment (and updates) (Highways Agency <i>et al</i> . 2008)	This Advice Note provides guidance for determining the significance of environmental effects, including for cumulative effects, and for the management of those effects.
British Standards Institute (2013) BS 42020 – Biodiversity – Code of Practice for Planning and Development	Provides recommendations and guidance to ecological consultants, planning applicants and planning authority decision makers, to ensure that ecological considerations are given the appropriate weight at each stage of the planning process, and are sufficiently informed by scientific assessment.



Title	Summary and Relevance	
The IUCN Red list	The International Union for Conservation of Nature (IUCN) has	
	compiled a Red list of threatened species that are facing a high	
	risk of global extinction.	
Guidance for survey	A range of guidance documentation was referred to for the	
methodologies	design and completion of project-specific surveys. Relevant	
	guidance is referred to in the survey reports.	

BASELINE CONDITIONS

- 12.7012.72 This section summarises the baseline ecological conditions determined through the course of desk-based and field-based investigations described above. In particular, this section identifies and evaluates those ecological features/receptors that lie within the Project Site's potential zone of influence and are pertinent in the context of the Proposed Development.
- 12.7112.73 Full results of the desk-based searches and field surveys undertaken, along with their findings, are provided within Appendix 12.1 (Document Reference 6.2.12.1). The findings are summarised below along with a final list of 'Important Ecological Features' to be considered within the EcIA. References to parts of the Project Site use commonly used names as shown on Figure 12.1 (Document Reference 6.2.12.1).

Statutory Designations

- <u>12.72</u> Statutory designations represent the most significant ecological receptors, being of recognised importance at an international and/or <u>national levelNational level</u>. International designations include Special Protection Areas (SPAs), Special Areas of Conservation (SACs) and Ramsar Sites. National designations include Sites of Special Scientific Interest (SSSIs) and National Nature Reserves (NNRs).
- <u>12.73</u> <u>12.75</u> Statutory designations occurring within the potential zone of influence of the Project Site are detailed below and illustrated in Figure 12.2 (Document Reference 6.2.12.2).

International Designations (SPAs/SACs/Ramsar)

12.7412.76 No part of the Project Site is covered by any international statutory designations. However, there are four statutory designations of international importance within 15km of the Project Site as described below. Reasons for their designation are provided in full within Appendix 12.1 (Document Reference 6.2.12.1).

Thames Estuary and Marshes SPA/Ramsar



- 12.75 12.77 The Thames Estuary and Marshes SPA is located c.3.3km east of the Essex Project Site, and c.6.0km east of the Kent Project Site. The corresponding Thames Estuary and Marshes Ramsar site is located 2.8km east of the Essex Project Site and 4.8km from the Kent Project Site. It comprises extensive intertidal mudflats that are visible at low tide. Additionally, there is saltmarsh and complex channel systems.
- Directive (Council Directive 79/409/EEC) as it supports populations of European importance over winter of the regularly occurring Annex 1 species avocet (*Recurvirostra avosetta*) and hen harrier (*Circus cyaneus*). This Site also qualifies as an SPA under Article 4.2 of the EU Birds Directive as it supports populations of European importance of regularly occurring migratory species including ringed plover (*Charadrius hiaticula*), grey plover (*Pluvialis squatarola*), dunlin (*Calidris alpina alpine*), knot (*Calidris canutus*), blacktailed godwit (*Limosa limosa islandica*) and common redshank (*Tringa totanus*). This SPA site also supports an internationally important assemblage of waterfowl as stated in Article 4.2 of the Directive, which includes gadwall (*Anas strepera*), shoveler (*Anas clypeata*), tufted duck (*Aythya fuligula*), and pochard (*Aythya ferina*).
- as it supports more than 20 British Red Data Book invertebrates and populations of the GB Red Book endangered least lettuce (Lactuca saligna), as well as the vulnerable slender hare's-ear (Bupleurum tenuissimum), divided sedge (Carex divisa), sea barley (Hordeum marinum), Borrer's saltmarsh-grass (Puccinellia fasciculata), and dwarf eelgrass (Zostera noltei). It also qualifies under Criterion 5 for its internationally important assemblage of waterfowl, and Criterion 6 for its internationally important numbers of over-wintering waterfowl.

Medway Estuary and Marshes SPA/Ramsar

- <u>12.78</u>12.80 The Medway Estuary and Marshes SPA/Ramsar is located c.13.1km south-east of the Essex Project Site, and c.16.4km east of the Kent Project Site.
- Directive (79/409/EEC) by supporting populations of European importance over winter of avocet and bewick's swan (*Cygnus columbianus bewickii*). It also qualifies under Article 4.2 of the Directive (79/409/EEC) by supporting populations of European importance of the following species over winter: black-tailed godwit, redshank, curlew (*Numenius arquata*), dark-bellied brent goose (*Branta bernicla bernicla*), dunlin, greenshank (*Tringa nebularia*), grey plover, knot, oystercatcher (*Haematopus ostralegus*), pintail (*Anas acuta*), ringed plover, shelduck (*Tadorna tadorna*), shoveler, teal (*Anas crecca*), turnstone (*Arenaria interpres*), and wigeon (*Anas penelope*). This SPA site also supports an internationally important assemblage of waterfowl as stated in Article 4.2 of the Directive, which includes red-throated diver (*Gavia stellata*), great crested grebe (*Podiceps cristatus*), cormorant (*Phalacrocorax carbo*), Bewick's swan, dark-bellied brent goose, shelduck, wigeon, teal, mallard (*Anas platyrhynchos*), pintail, shoveler, pochard, oystercatcher, avocet, ringed



plover, grey plover, lapwing (*Vanellus vanellus*), knot, dunlin, black-tailed godwit, curlew, redshank, greenshank and turnstone.

as it supports a number of species of rare plants and animals including several nationally scarce plants, and a total of at least twelve British Red Data Book species of wetland invertebrates. It also qualifies under Criterion 5 (formerly 3a) as it supports an international important waterfowl assemblage (greater than 20,000 birds), and it qualifies under Criterion 6 (formerly 3c), over winter, the site regularly supports internationally important populations of dark-bellied brent goose, dunlin, grey plover, knot, pintail, redshank, ringed plover and shelduck.

North Downs Woodland SAC

- 12.81 The North Downs Woodland SAC is located c.9.7km south-east of the Essex Project Site, and 8km south-east of the Kent Project Site. It is designated based on the following qualifying features:
 - Annex I habitats that are a primary reason for selection:
 - 9130 Asperulo-Fagetum beech forests; and
 - o 91J0 Taxus baccata woods of the British Isles * Priority feature.

Peters Pit SAC

- <u>12.82</u>12.84 Peters Pit SAC is located 13.8km south-east of the Essex Project Site, and 12.8km south-east of the Kent Project Site. It is designated for the following qualifying features:
 - Annex II species that are a primary reason for selection of this site:
 - Great crested newt for which this is considered to be one of the best areas in the United Kingdom.
- 42.8312.85 Both Thames Estuary and Marshes SPA/Ramsar and Medway Estuary and Marshes SPA/Ramsar support migratory birds and bird populations that have potential to use areas of 'functionally linked' land⁵ near the designations. Two distinct areas within the Kent Project Site that appear to be functionally linked directly to the estuary assemblage, and therefore potentially to nearby Ramsar/SPA/SSSI designations, are Botany Marsh West and Black Duck Marsh, which are locally important areas at dawn (rest)/high tide (refuge) for small numbers of several target species.

⁵ 'Functionally linked land' refers to land (or sea) beyond the boundary of a European site which might provide a (potentially important) role in maintaining or restoring a protected population at favourable conservation status through the role or function the land (or sea) provides, e.g. providing habitat for refuge, rest or foraging



- 12.8412.86 Therefore, both sites are at potential risk of adverse effects as a result of loss/damage/disturbance of functionally linked land within Botany Marsh and Black Duck Marsh and the Proposed Development. Both designations will therefore be considered Important Ecological Features scoped into the EcIA and are further considered within the Shadow Habitat Regulations Assessment (HRA) submitted along with the DCO application (see Appendix 12.4; Document Reference 6.2.12.4).
- 12.85
 12.87 In relation to the North Downs Woodland SAC, there is potential for traffic generated air quality impacts as a result of the Proposed Development both alone and incombination with other developments, as identified in Natural England's consultation response to the EIA Scoping (see Appendix 12.5; Document Reference 6.2.12.5) and PEIR (see Appendix 12.6; Document Reference 6.2.12.6). The SAC is therefore considered further as an IEF within the EcIA, and potential effects are 'screened' within the HRA (Document Reference 6.2.12.4).
- <u>12.86</u>12.88 Peters Pit SAC is located 13.8km away from the Proposed Development and it is considered highly unlikely that the Proposed Development will have any adverse effects on this designation, which is subsequently scoped out of the EcIA and not considered further. Within their consultation response to the EIA Scoping and PEIR, Natural England have not raised any objection to the scoping out of this designation.
- <u>12.87</u> Furthermore, it should be acknowledged that the scope of the HRA for the Tilbury2 development was limited to the Thames Estuary and Marshes SPA/Ramsar stating that:

'All other European nature conservation designations are located at least 9km from the Site, at which distance they are considered to be beyond the range of potential significant effects. No other designations have emerged through the scoping process, and therefore no other European nature conservation designations are given any further consideration...'

National Designations (SSSIs and NNRs)

12.90 In total, ten-11eleven SSSIs have been identified as being IEFs at risk of adverse effects from the Proposed Development. This includes the Swanscombe Peninsula SSSI, which occupies the majority of the Kent Project Site, and includes the former Bakers Hole SSSI, and a number of SSSIs within the EZoI of the Proposed Development as described further below.

Swanscombe Peninsula SSSI These include a number of SSSIs within the theoretical 5km EZoI, however some of the SSSIs within this zone have been excluded from the EcIA based on their reasons for designation and/or lack of effect-receptor pathways. Several SSSIs located beyond the 5km EZoI have been included in the EcIA owing to their location within one of the higher order IEFs identified above (i.e. SAC, SPA, Ramsar Site) and/or due to effect-receptor pathways having been identified over greater distances. This is summarised in Table 12.6 below.



- 12.91 Swanscombe Peninsula SSSI was notified by Natural England on 11 March 2021.
- 12.92 The Swanscombe Peninsula SSSI notification document⁶ describes the site as "a complex of open mosaic habitats on previously developed land and traditional estuarine habitats located near and within the River Thames, Kent. Habitats include chalk pits, free-draining grassland, scrub, wetlands, grazing marsh, mudflats and saltmarsh."
- 12.93 The SSSI boundary includes within it an existing SSSI (Bakers Hole, notified in 1989), and an enlarged SSSI area to take account of the biological interest on the peninsula. The total SSSI boundary equates to an area of 264.10ha.
- 12.94 Bakers Hole SSSI was originally designated for its geological/palaeontological interest and no additional reasons have been put forward for its designation. Bakers Hole SSSI now forms part of the Swanscombe Peninsula SSSI, however, impacts upon the SSSI from a geological perspective are addressed within Chapter 14: Cultural Heritage and Archaeology (Document reference: 6.1.14).
- 12.95 The enlarged SSSI area, to be named Swanscombe Peninsula SSSI, covers the majority of the Kent Project Site and has been notified for the following biological interest:
 - Vascular plants populations of the following plants: divided sedge (*Carex divisa*), yellow vetchling (*Lathyrus aphaca*), slender hare's-ear (*Bupleurum tenuissimum*), Bithynian vetch (*Vicia bithynica*) and round-leaved wintergreen (*Pyrola rotundifolia subsp. maritima*);
 - Invertebrates assemblages of invertebrates associated with bare sand and chalk, open short swards, open water on disturbed mineral sediments and saltmarsh and transitional brackish marsh; and
 - Breeding birds two diverse assemblages of breeding birds; one associated with lowland open waters and their margins, lowland fen and lowland damp grassland; and the other with lowland scrub.

LRCH Ltd have received correspondence from Natural England informing them of an area of land within the DCO boundary that is being considered for notification as a SSSI. Given the early stage of this process, this pre-notification is not considered any further as part of this assessment. The Applicant is working closely with Natural England as further details of the SSSI notification process emerge.

Other SSSIs within theoretical EZOI

12.96 In addition to the above, there are also a number of other SSSIs within the theoretical 5km EZoI, however, some of the SSSIs within this zone have been excluded from the EcIA based

^{6 &#}x27; Swanscombe Peninsula SSSI Kent – Notification under Section 28C of the Wildlife and Countryside Act 1981.' Natural England, published 11 March 2021.



on their reasons for designation and/or lack of effect-receptor pathways. Several SSSIs located beyond the 5km EZoI have been included in the EcIA owing to their location within one of the higher order IEFs identified above (i.e. SAC, SPA, Ramsar Site) and/or due to effect-receptor pathways having been identified over greater distances. This is summarised in **Table 12.6** below.

Table 12.6: Consideration of SSSIs for inclusion in the EcIA.

SSSI	Approximate Distance and Direction from the Project Sites*	Include in or exclude from EcIA?	Reasoning
Swanscombe Peninsula SSSI	Covers the majority of the KPS, and 3km W of ExPS.	<u>Include</u>	Located within KPS with direct impacts upon biological features identified.
			Impacts upon the SSSI from a geological perspective are addressed within Chapter 14: Cultural Heritage and Archaeology (Document reference: 6.1.14).
Bakers Hole SSSI	Within KPS, and 3km S of ExPS.	Exclude	SSSI designated for its geological rather than ecological importance and will be addressed within Chapter 14: Cultural Heritage and Archaeology (Document reference: 6.1.14).
Darenth Woods SSSI	Adjacent to the south-west corner of the KPS and 6.2km south-west of the ExPS.	Include	Within theoretical EZoI and effect-receptor pathways identified.
Swanscombe Skull Site SSSI and NNR	0.5km south of KPS and 4.2km south-west of ExPS.	Exclude	SSSI designated for its geological rather than ecological importance and will be addressed if appropriate within Chapter 14: Cultural Heritage and Archaeology (Document reference: 6.1.14).



SSSI	Approximate Distance and Direction from the Project Sites*	Include in or exclude from EcIA?	Reasoning
West Thurrock Lagoon and Marshes SSSI	450m west of KPS and 4.1km west of the ExPS.	Include	Within theoretical EZoI and effect-receptor pathways identified.
Lion Pit SSSI	1.4km north of KPS and 3.4km west of the ExPS.	Exclude	SSSI designated for its geological rather than ecological importance and will be addressed if appropriate within Chapter 14: Cultural Heritage and Archaeology (Document reference: 6.1.14).
Grays Thurrock Chalk Pit SSSI	1.9km north of KPS and 2.5km north of the ExPS.	Exclude	Within theoretical EZoI but no effect-receptor pathways identified.
Hangmans Wood and Denholes SSSI	1.9km north of ExPS and 3.3km north-east of the KPS.	Exclude	Within theoretical EZoI but no effect-receptor pathways identified.
Globe Pit SSSI	2.2km north-east of KPS and 1.1km north of ExPS.	Exclude	SSSI designated for its geological rather than ecological importance and will be addressed if appropriate within Chapter 14: Cultural Heritage and Archaeology (Document reference: 6.1.14).
South Thames Estuary and Marshes SSSI	2.5km east of ExPS and 4.7km east of the KPS.	Include	Within theoretical EZoI, effect-receptor pathways identified and included by virtue of its location within the higher order IEF (Thames Estuary and Marshes SPA).
Purfleet Chalk Pits SSSI	3.6km north-west of KPS and 6.1km west of the ExPS.	Exclude	sssi designated for its geological rather than ecological importance and will be addressed if appropriate within Chapter 14: Cultural Heritage and Archaeology (Document



Mucking Flats and Marshes SSSI	Approximate Distance and Direction from the Project Sites* 4.1km east of ExPS and 7.5km east of the KPS.	Include in or exclude from EcIA?	reference: 6.1.14). Within theoretical EZoI, effect-receptor pathways identified and included by virtue of its location within the higher order IEF
Shorne and Ashenbank Woods SSSI	4.8km south-east of both Project	Include	(Thames Estuary and Marshes SPA). On outer limit of theoretical EZOI but offect recentor pathways.
Inner Thames Marshes	Sites. 5.5km north-west	Include	but effect-receptor pathways identified. Outside of theoretical EZoI but
SSSI	of KPS and 8.3km north-west of the ExPS.	include	effect-receptor pathways identified.
Cobham Woods SSSI	6.7km south-east of the KPS and 7.5km south-east of the ExPS.	<u>Include</u>	Outside of theoretical EZoI but effect-receptor pathways identified.
Great Crabbles Wood SSSI	7.5km south-east of the KPS and 6.9km south-east of the ExPS.	<u>Include</u>	Outside of theoretical EZoI but effect-receptor pathways identified.
Wouldham to Detling Escarpment SSSI	13.0km south-east of the ExPS, and 12.4km SE of KPS.	Include	Outside of theoretical EZoI but effect-receptor pathways identified by virtue of its location within the higher order IEF (North Downs Woodland SAC).
Medway Estuary and Marshes SSSI	13.0km south-east of the ExPS and 14.7km east of the KPS.	Include	Outside of theoretical EZoI but effect-receptor pathways identified by virtue of its location within the higher order IEF (Medway Estuary and Marshes SPA).
Cobhan Woods SSSI	6.7km south east of the KPS and 7.5km south east of the ExPS.	Include	Outside of theoretical EZOI but effect receptor pathways identified.



SSSI	Approximate	Include in	Reasoning
	Distance and	or exclude	
	Direction from the	from EcIA?	
	Project Sites*		
Great Cabbles Wood	7.5km south-east	Include	Outside of theoretical EZoI but
SSSI	of the KPS and		effect-receptor pathways
	6.9km south-east		identified.
	of the ExPS.		Tachtinea.

^{*}KPS = Kent Project Site; ExPS = Essex Project Site

Non-statutory Designations

12.8812.97 Non-statutory designations are also commonly referred to in planning policies as 'local sites', although in fact these designations are typically considered to be important at a county level. In Kent and Essex, such designations are referred to as Local Wildlife Sites (LWSs). Additional non-statutory designations which should be considered in the assessment include Local Nature Reserves (LNRs) and Ancient Semi-natural Woodland (ASNW) where these are not covered by other designations.

12.8912.98 In total, five LWSs have been identified as being IEFs at risk of adverse effects from the Proposed Development. These include two LWSs within the Kent Project Site and a further three LWSs within the theoretical 2km EZol, however a large proportion of the LWSs within this zone (as shown on Figure 12.3; Document Reference 6.3.12.3) have been excluded from the EcIA based on their reasons for designation, spatial separation and/or lack of effect-receptor pathways. This is summarised in **Table 12.7** below.

Table 12.7: Consideration of LWSs for inclusion in the EcIA.

LWS	Approximate Distance and Direction from the Project Sites*	Include in or exclude from EcIA?	Reasoning
GR19.	Within the KPS and	Include	Within the Project Site with
Botany Marshes	2.2km west of the		potential for direct effects.
	ExPS.		
GR05.	Partly within the KPS	Include	Within theoretical EZoI with
Ebbsfleet	and 2.2km south-east		potential for direct effects.
Marshes,	of the ExPS.		
Northfleet			
DA13.	28m south of the KPS	Include	Within theoretical EZoI and
Alkerden Lane Pit	and 3.7km west of the		effect-receptor pathways
	ExPS.		identified.



LWS	Approximate	Include in or	Reasoning
	Distance and	exclude from	incusorining
	Direction from the	EcIA?	
	Project Sites*	ECIA:	
DA1.	Adjacent to the	Exclude	LWS designated for its
Bluewater Quarry	western edge of KPS		geological rather than
	and 6.1km west of the		ecological importance and will
	ExPS.		be addressed if appropriate
			within Chapter 14: Cultural
			Heritage and Archaeology
			(Document reference: 6.1.14).
DA12.	1.3km south-west of	Include	Within theoretical EZoI with
Disused Hospital	KPS and 7.7km south-		potential for indirect effects
Grounds,	west of ExPS.		due to changes in air quality.
Mabledon			
DA01.	1.6km south-west of	Exclude	Within theoretical EZoI but no
Green Street	KPS and 6.9km south-		effect-receptor pathways
Common	west of the ExPS.		identified
Th37.	25m to the east of the	Include	Within theoretical EZoI and
Tilbury Marshes	ExPS and 3.3km east		effect-receptor pathways
	of the KPS.		identified.
Th39.	732m north-east of	Exclude	Within theoretical EZoI but no
Lytag Brownfield	ExPS and 4.2km east		effect-receptor pathways
	of the KPS.		identified.
Th40.	900m east of ExPS	Exclude	Within theoretical EZoI but no
Tilbury Centre	and 4.3km east of the		effect-receptor pathways
	KPS.		identified.
Th15.	1.8km north-west of	Exclude	Within theoretical EZoI but no
West Thurrock	KPS and 4.8km west		effect-receptor pathways
Reedbed	of the ExPS.		identified.
Th18.	1.4km north-west of	Exclude	Within theoretical EZoI but no
West Thurrock	KPS and 4.2km west		effect-receptor pathways
Lagoon	of the ExPS.	Finalizata	identified.
Th22.	2.0km north-west of KPS and 4.1km north-	Exclude	Within theoretical EZoI but no
Grenville Road			effect-receptor pathways
Grasslands Th23.	west of ExPS. 1.7km north-west of	Exclude	identified. Within theoretical EZoI but no
Anchor Field	KPS and 4.0km west	Exclude	effect-receptor pathways
Andrior rielu	of the ExPS.		identified.
Th24.	2km north of KPS and	Exclude	Within theoretical EZoI but no
Mill Wood and	3.8km north-west of	LACIOUC	effect-receptor pathways
Cliff	ExPS.		identified.
Th28.	1.7km north-west of	Exclude	Within theoretical EZoI but no
Lion Gorge	KPS and 3.4km north-		effect-receptor pathways
	west of ExPS.		identified.
	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1		



LWS	Approximate Distance and Direction from the Project Sites*	Include in or exclude from EcIA?	Reasoning
Th31.	1.8km north of KPS	Exclude	Within theoretical EZoI but no
Grays Pit	and 2.3km north-west		effect-receptor pathways
Extensions	of ExPS.		identified.
Th38.	1.8km north of ExPS	Exclude	Within theoretical EZoI but no
Broom Hill	and 4.0km east of		effect-receptor pathways
	KPS.		identified.
Th49.	1.8km east of ExPS	Exclude	Within theoretical EZoI but no
Goshems Farm	and 5.1km east of		effect-receptor pathways
	KPS.		identified.

^{*}KPS = Kent Project Site; ExPS = Essex Project Site

Habitats/Flora

<u>12.9012.99</u> A full description of the habitats present within the Project Site is set out within Appendix 12.1: *Ecology Baseline Report* (Document Reference 6.2.12.1) and illustrated on the 'Extended Phase 1 Habitat Survey' plan (Figure 12.4; Document Reference 6.3.12.4). A summary of the habitats found and described onsite is provided in **Table 12.8**.

Table 12.8: Summary of the Distribution and Value of Habitats within the Project Site.

Habitat/Flora	Distribution within the	Intrinsic Ecological Importance
	Project Site	
Rare plants	Populations of 13 nationally	National, nationally scarce.
	scarce species were found	Populations of the plants divided
	on Swanscombe Peninsula in	sedge (Carex divisa), yellow vetchling
	the Kent Project Site 2016.	(Lathyrus aphaca), slender hare's-ear
	Eight were recorded in 2020.	(Bupleurum tenuissimum), Bithynian
		vetch (Vicia bithynica) and round-
		leaved wintergreen (Pyrola
		rotundifolia subsp. maritima) are
		listed as a reason for notification of
		the Swanscombe Peninsula SSSI.
Broad-leaved semi-	Mainly in the South along	Local, owing to connectivity, good
natural woodland	the A2(T) corridor, some	canopy species diversity, ground flora
	along the eastern boundary.	diversity and status as a Priority
		Habitat ⁷ .

Priority habitats and species include those habitats and species which are 'of principal importance for the purpose of conserving biodiversity' under Section 41 of the Natural Environment and Rural Communities Act, 2006



12 - 47

Habitat/Flora	Distribution within the	Intrinsic Ecological Importance
	Project Site	
Broad-leaved	Many areas throughout the	Site, as small in extent and immature
plantation woodland	Project Site.	planting.
Scrub	Many areas throughout the	Local , due to large extent on the Kent
	Project Site.	Project Site and structure, despite
		being a common habitat.
Tall ruderal	Small extent in station	Site, owing to small extent and low
	quarter south.	species diversity.
Improved grassland	Small area in A2(T) corridor	Site , owing to low species diversity.
Poor semi-improved	Botany marsh east,	Local , species diversity is low across
grassland	Broadness grasslands and	most of extent, but forms large area of
	former landfill (west of	continuous habitat.
	Ebbsfleet International	
	Station).	
	Includes areas within the	
	overall OMHPDL.	
Semi-improved	Neutral grassland in	Local to District , some areas of lower
neutral and	Broadness grassland area	botanical diversity but supports
calcareous grassland	and to the west of Black	nationally scarce plant species. One
	Duck Marsh, calcareous	small area of MG1d grassland in
	grassland in Craylands Pit,	Broadness Grassland with increased
	Black Duck Marsh and north	value.
	of Tiltman Avenue.	
	Includes areas within the	
Coastal/Floodalaia	overall OMHPDL.	District a large area of Driegity
Coastal/Floodplain	Botany Marsh West and part	District, a large area of Priority
Grazing Marsh	in Botany Marsh East.	Habitat ⁸ coastal/floodplain grazing
Open mosaic on	Large areas across the Kent	marsh. District, a large area of Priority
previously developed	Project Site where there is a	Habitat ⁹ within the Kent Project Site.
land	verifiable history of industrial	The areas are not floristically diverse
ianu	disturbance. Comprises an	but have the potential to support
	'umbrella' habitat including a	important invertebrate assemblages.
	number of several individual	po. tane invertebrate assemblages.
	habitat components.	

⁹ UK Biodiversity Action Plan Priority Habitat Descriptions Open Mosaic Habitats on Previously Developed Land (Updated July 2010) From: UK Biodiversity Action Plan; Priority Habitat Descriptions. BRIG (ed. Ant Maddock) 2008.



⁸ UK Biodiversity Action Plan Priority Habitat Descriptions Coastal and Floodplain Grazing Marsh From: UK Biodiversity Action Plan; Priority Habitat Descriptions. BRIG (ed. Ant Maddock) 2008

Habitat/Flora	Distribution within the Project Site	Intrinsic Ecological Importance
Amenity grassland and shrub	Small area concentrated around Northfleet and roadand trackside verges.	Negligible , owing to low species diversity, intensive management.
Marshy grassland	Small area in station quarter south.	Site owing to small extent and limited species diversity.
Waterbodies (ponds, temporary pools, standing water and ditches)	Scattered throughout the Project Site, ditches forming two main networks at Black Duck and Botany Marshes.	District - many ponds are leachate treatment lagoons or contaminated by leachate. The remaining ponds and ditches support little vegetation other than reeds. Ditch network forms part of a large marsh area including Botany Marshes LWS and adjacent grazing marsh and is considered of district level District level.
Swamp (reedbed)	Black duck Marsh and Channel Tunnel Rail Link (CTRL) Wetland form extensive reedbed, smaller areas exist within Botany Marsh East and along ditch network.	County - Priority Habitat forming a large area of the Project Site in close proximity to the River Thames. Priority Habitat.
Bare ground	Mostly around Manor Way industrial Estate.	Negligible, no species diversity and surrounded by buildings.
Hardstanding and buildings	Mostly around Manor Way industrial Estate, Station Quarter North and the Essex Project Site.	Negligible, man-made structures.
River Ebbsfleet	Along the eastern boundary of the Kent Project Site.	Local , as this river corridor and its associated riparian habitat provide a green corridor.

12.8, the Essex Project Site comprises predominantly hardstanding, industrial land, whilst the Kent Project Site supports a range of habitats including, intertidal sediment, saltmarsh, wetlands, including running water (the River Ebbsfleet), open water (ponds), reedbed/swamp and ditch networks, a range of grasslands and open mosaic habitats, arable, scrub, woodland, chalk cliffs/exposures, buildings and bare ground. The saltmarsh and intertidal habitat are described and evaluated in ES Chapter 13: Marine Ecology (Document Reference 6.1.13). Those habitats within the Project Site considered to be of local, or greater, value are taken forward as IEFs in the EcIA.

<u>12.92</u><u>12.101</u> In addition to the identification of specific important habitats within the EcIA the contribution made by all existing habitats to the overall biodiversity value of the Project



Site is fully accounted for within Appendix 12.2: *Biodiversity Net Gain Assessment* (Document Reference 6.2.12.2).

Swanscombe Peninsula SSSI

- 12.102 A number of the individual habitats on the Kent Project Site, as described in Table 12.8, form part of the Swanscombe Peninsula SSSI (considered an IEF of national importance).

 None of the individual habitats within the SSSI are identified as reasons for the notification, and therefore impacts upon these habitats themselves is not strictly covered through the assessment of effects upon the higher level IEF of the SSSI. Nevertheless, the mosaic of habitats present is clearly important to sustaining the assemblages of breeding birds and invertebrates that are reasons for notification, and therefore for the purpose of this EcIA individual habitats are described, mapped and assessed independently (and as part of an open mosaic of habitats), irrespective of their inclusion within land notified as a SSSI.
- 12.103 Furthermore, there are instances where habitats which form part of the SSSI also occur on areas of land outside of the SSSI boundary, and therefore these habitats are also treated as individual habitat IEFs, such that impacts on the 'non-SSSI' areas can also be adequately assessed.
- 12.104 For the assessment of potential effects on habitats described later in this report, the quantities of habitat loss/damaged is provided for the total area of the individual habitat (e.g. the loss of scrub habitat includes the total area of scrub habitat, regardless of its position in/out of the SSSI boundary).
- 12.105 The assessment of potential effects upon the rare plant populations recorded on the Kent Project Site is undertaken as part of the assessment of effects upon Swanscombe Peninsula SSSI in regard to those plant populations, which are reason for notification of the SSSI (namely populations of the plants divided sedge (*Carex divisa*), yellow vetchling (*Lathyrus aphaca*), slender hare's-ear (*Bupleurum tenuissimum*), Bithynian vetch (*Vicia bithynica*) and round-leaved wintergreen (*Pyrola rotundifolia subsp. maritima*)). In addition, not all of the rare or scarce plants recorded on the Kent Project Site are listed as reasons for notification of the Swanscombe Peninsula SSSI, and therefore a separate assessment of these 'non-SSSI' plant populations is undertaken.

Open Mosaic Habitat on Previously Developed Land

12.93 12.106 A number of the individual habitats on the Kent Project Site, as described in **Table**12.8, collectively form part of a much wider 'Open Mosaic Habitat on Previously Developed Land' (OMHPDL) (considered a Priority habitat). This includes areas of bare ground, ephemeral vegetation, open grassland (poor semi-improved grassland and semi-improved neutral and calcareous grassland) and patches of scrub (less than 2500m²)¹⁰,

Large blocks of dense scrub (larger than 2500m2 or smaller, adjacent blocks making up a similar area) were excluded



where there is verifiable history of industrial disturbance, and where the habitat characteristics as described within the Priority Habitat description¹¹ could be identified using professional judgement and knowledge of the Project Site. The extent and boundary of the OMHPDL as defined by the 2020 survey by EDP is shown on Figure 12.4 (Document Reference 6.3.12.4).

12.9412.107 For the purpose of the habitat assessment within the EcIA, the individual habitat components which form part of the open mosaic habitat are described and mapped individually in order to illustrate the extent and spatial distribution of the individual habitats which occur across the Kent Project Site, and to enable a robust assessment of individual habitats, including where this is required for the assessment of effects on species (e.g. assessing the impact of scrub loss on dormice). However, in recognition of their presence within the 'open mosaic habitat on previously developed land', these individual habitats are collectively grouped into an 'umbrella' IEF which is 'open mosaic habitat on previously developed land'.

12.9512.108 In instances where habitats which form part of the OMHPDL also occur outside of this area (e.g. areas of open grassland that occur in areas considered to be OMHPDL and areas not considered to be), these habitats are also treated as individual habitat IEFs, such that impacts on the 'non-OMHPDL' areas can also be adequately assessed. For the assessment of potential effects on species/species assemblages described later in this report, the quantities of habitat loss/damaged is provided for the total area of the individual habitat (e.g. the loss of suitable scrub habitat for dormice includes the total area of suitable scrub habitat, regardless of its position in/out of the OMHPDL).

Ancient Woodland

12.9612.109 There are a number of areas of ancient woodland present along the A2 corridor, to the south of the Kent Project Site. Their location is shown on Figure 12.56: Tree Constraints Plan (Document reference: 6.3.12.56). As described within Appendix 12.9: Arboricultural Impact Assessment (Document reference: 6.2.12.9), ancient woodland is defined as an area which has been continuously wooded since at least 1600 AD¹² and includes ancient semi- natural woodland and plantation on ancient woodland. 'Wooded continuously' does not mean there has been continuous tree cover across the whole area. Not all trees in the woodland must be old. Open space, both temporary and permanent, is also an important component of ancient woodland.

<u>12.97</u>12.110 As described in paragraph 175c of the NPPF, ancient woodland is recognised in planning terms as an irreplaceable habitat:

¹² https://www.gov.uk/guidance/ancient-woodland-and-veteran-trees-protection-surveys-licences#history



12 - 51

¹¹ UK Biodiversity Action Plan Priority Habitat Descriptions Open Mosaic Habitats on Previously Developed Land (Updated July 2010) From: UK Biodiversity Action Plan; Priority Habitat Descriptions. BRIG (ed. Ant Maddock) 2008

"Development resulting in the loss or deterioration of irreplaceable habitats (such as ancient woodland and ancient or veteran trees) should be refused, unless there are wholly exceptional reasons and a suitable compensation strategy exists".

- 12.9812.111 Of those areas of ancient woodland shown on Figure 12.56 (Document reference 6.3.12.56), Darenth Woods is designated as a SSSI (as described in Table 12.6 above), and therefore considered of national importance commensurate with its statutory designation.
- 12.9912.112 The remaining areas of ancient woodland within the EZoI of the Proposed Development are not recognised as either SSSI or LWS. Within Kent, ancient woodlands are relatively well represented, and not all examples of ancient woodland would be considered of county/ national value.
- <u>12.100</u>12.113 For the purpose of the EcIA, ancient woodland is considered an IEF of at least county value.

Species

12.10112.114 As set out previously, information on protected and/or notable species within or near to the Project Site has been collected through a desk study and a range of field surveys. The findings of these investigations to date are set out in full within Appendix 12.1: *Ecology Baseline Report* (Document Reference 6.2.12.1) and summarised below.

Birds

Wintering Birds

- 12.10212.115 A combined total of up to 44 species were recorded during 2012/13, 2014/15 and 2019/20 intertidal and high tide surveys. Of the 30 Ramsar/SPA/SSSI qualifying species mentioned in the Thames Estuary and Marshes and Medway Estuary and Marshes designation citations, a total of 22 have been recorded at either low or high tide during the winter 2019/2020. Of the 22 Ramsar/SPA qualifying species which have stated peak population counts, EDP recorded an overall total of 12 over the course of the winter 2019/20 high and low tide surveys, with the numbers recorded during surveys at either low or high tide between 0.07% and 8.66% of the peak population counts stated in the citations.
- 12.116 Over winter 2020/2021, intertidal surveys recorded no further species over those recorded previously within the Kent Project Site. Activity levels were broadly consistent with those recorded in previous years.
- 12.117 Intertidal surveys undertaken at the Essex Project Site during 2020/2021 recorded smaller numbers of species, possibly due to the limited intertidal zone and high levels of existing disturbance from the Port of Tilbury. A total of eleven11 species were recorded in the intertidal zone, with a further seven7 species either flying over or passing along the



<u>channel of the river. Peak counts were also lower, with only 3 species surpassing 1% of the WeBS total for the Thames estuary (teal, redshank and little grebe).</u>

- 12.10312.118 As described in detail in Appendix 12.1 (Document Reference 6.2.12.1), in 2019/2020 it was concluded that although the Project Site itself is not regarded to have value at the International level, the wintering wader/wildfowl assemblage present within inland areas of the Kent Project Site itself must be valued at the International level given their status as functionally linked to the estuary assemblage.
- 12.10412.119 Additionally, 28 other terrestrial species (non-wader, non-wildfowl species) of conservation concern were also recorded in generally low to moderate numbers, typically relating to individuals or small flocks of each species recorded on one or two survey visits, but also including a high diversity and moderate abundance of several species listed on Schedule 1 of the Wildlife & Countryside Act (WCA) and species of Conservation Concern. Therefore, the terrestrial wintering bird assemblage present within the Kent Project Site should be considered separately and is of County Importance.
- 12.10512.120 The assessment of the wintering bird assemblage has been based on the completion of a comprehensive suite of wintering bird surveys collated during winter 2019/2020. In addition, this recent data is supplemented by a suite of 'baseline' surveys completed by Chris Blandford Associates (CBA) during 2012/2013, provided in full in Appendix 12.1 (Document Reference 6.2.12.1). The DCO application is therefore informed by two seasons of bird survey data, albeit not in consecutive years.
- 12.10612.121 It is considered that the survey data gathered is sufficient and robust to inform the DCO application. The wetland habitats within the Kent Project Site have not changed significantly in the intervening years between the 2012/2013 baseline surveys and the recent 2019/2020 surveys, and the data collected is more or less consistent across that time period. As set out above the wintering wader/wildfowl assemblage using the Kent Project Site has been valued at the International level owing to its association with the nearby Ramsar and SPAs. The assemblage, which is treated as an IEF, has therefore been valued at the highest level, and data from additional surveys will not affect this valuation or significantly alter the impact assessment and mitigation measures being developed.
- 12.10712.122 Justification for the scope of surveys was provided to Natural England within an Ecology Briefing Note (edp5988_r022) issued 21 August 2020 (see Appendix 12.5; Document Reference 6.2.12.5). At the time of writing, no further comment has been made by Natural England in response to the Ecology Briefing Note.

Breeding Birds

<u>12.10812.123</u> As described in detail in Appendix 12.1 (Document Reference 6.2.12.1), the overall bird abundance was very high across the Project Site. A total of 99 species were recorded during the breeding bird surveys. Of those species recorded during the survey, 33 were confirmed as breeding within the survey area, 26 are considered to have probably bred,



18 to have possibly bred and the remaining 22 species are considered to be non-breeding species.

Schedule 1 Species

- 12.10912.124 Fifteen species that are included on Schedule 1 of the Wildlife and Countryside Act 1981 (as amended) were recorded during the 2020 breeding bird surveys. Of these, greylag goose (Anser anser) (2-16 pairs), bearded tit (Panurus biarmicus; 3-5 pairs) and Cetti's warbler (Cettia cetti) (51-87 pairs) are confirmed breeding species; marsh harrier (Circus aeruginosus) (single pair) probably bred within the Kent Project Site; and spotted crake (Porzana porzana), little ringed plover (Charadrius dubius), barn owl (Tyto alba; one pair), kingfisher (Alcedo atthis) and peregrine falcon (Falco peregrinus) possibly bred. The remaining six species; whimbrel (Numenius phaeopus), greenshank (Tringa nebularia), mediterranean gull (Larus melanocephalus), little tern (Sternula albifrons), redwing (Turdus iliacus) and black redstart (Phoenicurus ochruros) are considered to be non-breeding species.
- A single spotted crake was recorded on the track by Pond P8 on the Kent Project Site in June by an ecologist who was completing invertebrate surveys. This species was not recorded during the breeding bird survey or during the nocturnal spotted crake survey, which was completed in late June after the incidental sighting. The Kent Project Site contains habitat that is suitable breeding habitat for this species so, based on the European Bird Census Council (EBBC) criteria, it is considered to be a possible breeding species, although based on the results of the survey effort it is more than likely that the incidental record relates to a migrant bird.

Green List Species

42.11112.126 Green listed species were present in large numbers across all areas of the Kent Project Site surveyed. The most commonly recorded green listed species were wren (*Troglodytes troglodytes*), with a peak count of 87-155 pairs, whitethroat (*Sylvia communis*) with 85-130 pairs individuals, reed warbler (*Acrocephalus scirpaceus*) with 70-133 pairs, robin (*Erithacus rubecula*) with 47-103 pairs, blackcap (*Sylvia atricapilla*) with 57-133 pairs, blackbird (*Turdus merula*) with 61-116 pairs, chiffchaff (*Phylloscopus collybita*) with 43-73 pairs, and most notably Cetti's warbler, a WCA Schedule 1 species, with 51-87 pairs. It is considered very likely that these species are breeding within the Kent Project Site. Peregrine falcon (*Falco peregrinus*), another Schedule 1 species, was also recorded hunting within the Kent Project Site.

<u>Amber List Species</u>

12.11212.127 Twenty-nine Amber List Bird Species of Conservation Concern were recorded on or over the Kent Project Site during the 2020 breeding bird surveys including greylag goose, mute swan, shelduck, shoveler, gadwall (*Mareca Strepera*), mallard, teal (*Anas crecca*), swift (*Apus apus*), stock dove (*Columba oenas*), spotted crake, oystercatcher, redshank, greenshank, black-headed gull (*Chroicocephalus ridibundus*),



Mediterranean gull, common gull (*Larus canus*), great black-backed gull (*Larus marinus*), yellow-legged gull (*Larus michahellis*), lesser black-backed gull (*Larus fuscus*), little tern, common tern (*Sterna hirundo*), marsh harrier, kingfisher, kestrel (*Falco tinnunculus*), house martin (*Delichon urbicum*), willow warbler (*Phylloscopus trochilus*), dunnock (*Prunella modularis*), bullfinch (*Pyrrhula pyrrhula*) and reed bunting (*Emberiza schoeniclus*).

- 12.11312.128 Of the above species greylag goose, spotted crake, greenshank, little tern, marsh harrier and kingfisher are included on Schedule 1 and have been discussed above.
- 12.11412.129 In terms of wetland birds, it is considered that the Kent Project Site supports a probable breeding pair of mute swan, up to eleven possible breeding pairs of shelduck, up to four pairs of possibly breeding shoveler, up to six probable breeding pairs of gadwall, between fourteen and seventeen probable breeding pairs of mallard, and one or two breeding pairs of oyster catcher. A single teal was recorded in April in Botany Marsh West; this species is considered to have possibly bred.
- <u>12.11512.130</u> Redshank, black-headed gull, common gull, great black-backed gull, yellow-legged gull, lesser black-backed gull and common tern are all considered to be non-breeding birds and registrations of these species are limited to individual or low numbers of birds or of birds flying over the Kent Project Site.
- 12.11612.131 In terms of non-wetland birds, swift and house martin (*Delichon urbica*) were recorded foraging but are not considered to have bred on the Kent Project Site, although there is suitable nesting habitat within the industrial area on the Swanscombe Peninsula on the Kent Project Site.
- <u>12.11712.132</u> The Kent Project Site is considered to support up to 10 breeding pairs of stock dove, one to three pairs of probably breeding kestrel, a single breeding pair of willow warbler, 45 to 84 breeding pairs of dunnock, 3 to 5 breeding pairs of bullfinch and seven to 14 breeding pairs of reed bunting.

Red List Species

12.11812.133 Seventeen Red List Birds of Conservation Concern were recorded on the Kent Project Site during 2020 including pochard, cuckoo (*Cuculus canorus*), lapwing (possibly bred, 1 pair), whimbrel, herring gull (*Larus argentatus*; non breeding), skylark (9-13 possibly breeding pairs), grasshopper warbler (*Locustella naevia*; 12-15 probably breeding pairs), starling (*Sturnus vulgaris*) (likely just foraging), redwing, song thrush (*Turdus philomelos*) (up to 67 breeding pairs), mistle thrush (*Turdus viscivorus*; three breeding pairs), nightingale (*Luscinia megarhynchos*; 3 to 4 probably breeding pairs), black redstart, house sparrow (*Passer domesticus*; up to 8 probably breeding pairs), yellow wagtail (*Motacilla flava*; not breeding), grey wagtail (*Motacilla cinereal*; 1 breeding pair) and linnet (*Carduelis cannabina*; up to 39 breeding pairs).



12.11912.134 Cuckoo are considered to have probably breed on the Kent Project site as adults were recorded between April and June 2020. The Kent Project Site supports suitable host species including dunnock and reed warbler.

Evaluation

12.135 The breeding bird assemblage found within the Kent Project Site is listed as a reason for notification of the Swanscombe Peninsula SSSI. The supporting information to the SSSI notification published by Natural England on 11 March 2021¹³ summarises the breeding bird assemblage as follows:

<u>"two diverse assemblages of breeding birds: one associated with lowland open waters and their margins, lowland fen and lowland damp grassland; the other with lowland scrub."</u>

12.136 The Swanscombe Peninsula SSSI notification document¹⁴ provides a fuller description of the breeding bird assemblage, which is a reason for notification, as follows:

"The wetland areas of the site support an assemblage of breeding birds of lowland damp grassland, lowland open waters and their margins and lowland fen. Species associated with the fen and swamp habitats of Black Duck Marsh and the CTRL wetlands include bearded tit Panurus biarmicus, marsh harrier Circus aeruginosus and the elusive water rail Rallus aquaticus. Wetland habitats across the site support reed bunting Emberiza schoeniclus, sedge warbler Acrocephalus schoenobaenus and reed warbler. The wetland mosaic with scrub supports Cetti's warblers Cettia cetti.

Botany Marsh West is a surviving fragment of grazing marsh, providing damp grassland habitat for lapwing Vanellus vanellus. Little egret Egretta garzetta and grey heron Ardea cinerea utilise a number of wetland habitats and forage within the intertidal habitats of the adjacent River Thames. The water bodies support breeding waterfowl including pochard Aythya ferina, mute swan Cygnus olor and little grebe Tachybaptus ruficollis.

The chalk pits and areas of scrub support an assemblage of breeding birds of lowland scrub. Grasshopper warbler Locustella naevia can be found in open habitats on Broadness, while nightingale Luscinia megarhynchos favours the denser scrub areas of Botany Marsh East and the chalk pits. Linnets Linaria cannabina and lesser whitethroats Curruca curruca can be found in the former landfill tips and areas north of the CTRL wetlands."

<u>12.137 Overall The two</u>, the breeding bird assemblages found within the Kent Project Site, as described above, are therefore considered in 2020 is considered to be of Regional National value, whilst. As the two breeding bird assemblages identified on the Kent Project Site are reasons for notification of the SSSI, the assessment of potentially

¹⁴ 'Swanscombe Peninsula SSSI Kent – Notification under Section 28C of the Wildlife and Countryside Act 1981.'
Natural England, published 11 March 2021.



¹³ 'Swanscombe Peninsula SSSI Kent – Supporting Information'. Natural England, published 11 March 2021.

significant effects upon the assemblages is included within the assessment of effects upon the higher tier IEF of the Swanscombe Peninsula SSSI.

12.120 The the Kent Project Site is also considered to be of National Importance for breeding pochard.

Passage Birds

- 12.12112.138 Full results of the passage bird surveys are discussed in Appendix 12.1 (Document Reference 6.2.12.1). In summary, the surveys recorded a total of thirty-seven species, including the following species listed as qualifying features of the Thames Estuary and Marshes SPA: ringed plover (*Charadrius hiaticula*), dunlin (*Calidris alpina*) and redshank. As described in full in Appendix 12.1 (Document Reference 6.2.12.1) the numbers of individual birds from each of these species recorded during the surveys constitute less than 1% of the SPA population and are not considered significant.
- 12.12212.139 In terms of abundance and diversity of birds recorded, this was significantly reduced from that found along the estuary front throughout winter, with the most abundant birds being black-headed gulls and mallard. Three Peregrines were recorded flying over on 15 April 2020.

Bats

Bat Roosting – Trees

- <u>12.12312.140</u> During the visual assessment for roosting bats on 04 June 2020, no bats or evidence of bats was found but a total of 19 trees within the Kent Project Site were identified as offering potential to support roosting bats. Following the aerial tree climbing surveys, no evidence of roosting bats was found in any of the trees and a further tree with moderate potential was identified and surveyed.
- <u>12.12412.141</u> Therefore, in total nine trees have been identified on the Kent Project Site as having high potential, five as moderate potential and six as having low potential. No evidence of bat roosting has been recorded in any of the trees.
- 12.12512.142 There are no trees with bat roosting potential within the Essex Project Site.

Bat Roosting – Buildings

12.12612.143 A total of 150 buildings were identified from maps/aerial photographs within the Project Site. A preliminary 'rapid assessment' of these buildings from public footpaths and roads, was undertaken on 01 May 2020. The purpose of this survey was to determine whether any buildings could be scoped out as having negligible potential to support roosting bats, in order to focus the efforts of further external and internal inspections and presence/absence surveys as required. The assessment was undertaken by two Natural



- England licensed bat workers. Further visits to the Project Site were made on 07 July 2020, 15 July 2020 and 17 July 2020 to complete external building assessment.
- <u>12.127</u><u>12.144</u> Following these assessments 101 buildings were assessed as having negligible potential to support roosting bats, due to their construction, or were no longer present. These buildings were therefore not subject to any further level of survey.
- <u>12.128</u>12.145 A total of 23 buildings were found to have some level of potential to support roosting bats during the rapid assessment, with 10 assessed as having *Low potential*, 10 assessed as having *Moderate potential* and three assessed as having *High potential*.
- <u>12.12912.146</u> An individual soprano pipistrelle (Pipistrellus pygmaeus) was recorded entering B67 during the survey on 27 August 2020. An individual common pipistrelle (Pipistrellus pipistrellus) was recorded entering B32 on 17 September 2020. It is considered B67 and B32 each support a summer day roost for individual bat(s) and it is likely the buildings are only occasionally used as other surveys on the buildings recorded no bats emerging.
- 12.13012.147 No emergences or re-entries have been detected from any other buildings surveyed.
- 12.13112.148 There remain a further 26 buildings that could not be accessed during 2020. Precautionary mitigation measures, including the need for pre-commencement surveys, are provided within the 'Bat Mitigation Strategy' enclosed within the EMMF (Document reference: 6.2.12.3).

Bat Roosting – Tunnels (Summer)

- 12.13212.149 The assessment of roosting potential undertaken in August 2020 noted 10 tunnels with suitability for roosting, swarming and hibernating bats. For summer roosting; two tunnels had moderate potential and two had low. For autumn swarming; five had moderate potential, four low and one negligible. For winter hibernation; nine had low potential and one had negligible potential. Full details are included in Appendix 12.1 (Document Reference 6.2.12.1).
- <u>12.133</u>12.150 No emergences or re-entries have been detected from the tunnels indicating no tunnels are used as summer roosts

Bat Roosting – Tunnels (Autumn Swarming)

12.13412.151 Static detectors deployed at the entrance of the tunnels in August and September recorded low levels of bat activity. Due to access constraints for health and safety reasons it was not always possible to position statics so that recordings were from solely within the tunnels themselves. As such it is difficult to determine absolutely whether behaviour can be attributed to autumn swarming or general foraging.



- 12.13512.152 A number of the tunnels returned no records of bats or low numbers of recordings of an assemblage typical of the area including common pipistrelle, soprano pipistrelle, noctule (*Nytalus noctula*), long-eared bat (Plecotus auratus) and Myotis bats. Of the tunnels surveyed, tunnel T7 and tunnel T16 recorded larger than average numbers of Myotis sp. calls. There were 14 Myotis recordings made between midnight and 1am on 25 September at tunnel T7 but no bats were recorded at tunnel T7 during the August or October deployments.
- 12.13612.153 There were 42 Myotis recordings made between 10.30pm and midnight on 01 September 2020 were made at the south end of T16. Conversely, there were no Myotis calls recorded at the northern end of T16 during this time and, nor was there a distinct repeated peak of activity within the target period in August or October.
- <u>12.137</u>12.154 The results do not indicate autumn swarming behaviour by any species at the tunnels surveyed.

Bat Roosting – Tunnels (Winter)

- 12.13812.155 The preliminary roost assessment of tunnels has identified that 1 of the tunnels has moderate potential to support hibernating bats, 7 tunnels have low potential and 1 has negligible potential. A further tunnel, TU/016, has negligible potential with the exception of a single large crack at the tunnel entrance which was considered to offer low potential.
- 12.13912.156 The Proposed Development will not result in any direct impacts upon any of the tunnels with the exception of tunnels TU/016 (negligible/low potential) and TU/018 (low potential) which will be used for access between the transport interchange and staff accommodation.
- 12.157 Static detectors were deployed inside four4 tunnels and at the entrance of four4 tunnels in December 2020, January 2021 and February 2021. Due to access constraints for health and safety reasons it was not always possible to position static detectors so that recordings were from solely within the tunnels themselves.
- 12.158 Tunnel TU/07 recorded 38 Myotis sp. passes between December 2020 and February 2021,

 1 common pipistrelle pass (19/01/21), and 1 soprano pipistrelle pass (13/02/21). This
 tunnel is considered to be a confirmed hibernation roost. Four tunnels (TU/011, TU/013A,
 TU/014A and TU/016) returned low numbers of recordings of common pipistrelle and
 soprano pipistrelle bats. It is unlikely that pipistrelle bat hibernation roosts were present
 in three3 of these tunnels but possible that common pipistrelle and soprano pipistrelle
 hibernation roosts were present in Tunnel TU/016 due to the time the bat passes were
 recorded before sunrise and the bat detector was deployed inside the tunnel, eliminating
 the possibility of the detector recording bats passing by outside of the tunnel.
- 12.159 Based on the results of the surveys and assessment of the conservation status of the bat species present, the roosting bat assemblage is considered to be of District level



- importance, with the exception of the roosting bat assemblage present in Tunnels TU/07 and TU/016, which are considered to be of County level importance for a small number of hibernating bats (common and rarer *Myotis* species).
- 12.140 Winter hibernation surveys of these two tunnels are being undertaken between December 2020 to February 2021, the results will be submitted as an addendum during the Examination of the DCO application once the surveys are complete.

Bat Foraging/Commuting Activity

- 12.14112.160 A bat assemblage of at least eight species has been recorded within the Kent Project Site. This assemblage includes one nationally rare bat; Nathusius' pipistrelle (Pipistrellus nathusii) and three Kent Red Data Book species; noctule, Leisler's bat (Nyctalus leisleri) and serotine (*Eptesicus serotinus*) (the latter also being a Kent BAP species), however, recordings from these species are few with 82.6-89.3% of activity resulting from common pipistrelle. Further details can be found in Appendix 12.1.
- 12.14212.161 The total amount of activity was lowest along the A2(T) corridor and highest in Botany Marsh, Black Duck Marsh, the Swanscombe Peninsula and the NE tip. This is not surprising as the grassland and wetland habitat on these high activity areas provides good foraging habitat for bats. Land north of Tiltman Avenue recorded the highest average number of species.
- 12.14312.162 The abundance and diversity of bat species recorded at the Kent Project Site is considered to be fairly typical of an urban edge site in south-east England with common and widespread generalist species such as common pipistrelle bats accounting for the vast majority of foraging and commuting activity. However, a number of rarer bat species were recorded using the Kent Project Site in low numbers including Nathusius' pipistrelle. Serotine bats are also recorded, which are considered rare in Kent.
- 12.14412.163 The overall bat assemblage using the Kent Project Site, taking into consideration the presence of rare and uncommon species (albeit only present in low numbers), is considered to be of District level value.
- <u>12.145</u>12.164 Owing to the nature of the habitats present, being predominantly developed land with little natural vegetation present, the Essex Project Site is considered of negligible value to foraging/commuting bats.
- 12.14612.165 In their Discretionary Advice Service letter of 9th October 2020 (copy of which is enclosed as Annex EDP 13 to the EMMF (Document reference: 6.2.12.3)), Natural England have identified the potential for the Kent Project Site to be used by winter foraging bats, due to the nature of the habitats present and the proximity to the River Thames. Whilst no dedicated winter survey work was considered necessary to inform the EcIA, nor requested by consultees during the EIA Scoping Opinion received in July 2020 or through the PEIR consultation in July 2020, on a precautionary basis, the potential effects of the Proposed Development on potential winter foraging habitats is included.



Dormouse

Essex Project Site

- <u>12.147</u>12.166 Essex Field Club (EFC) returned one dormouse record from Tilbury Marshes, dated 2009, located c. 1.5km east of the Essex Project Site.
- 12.14812.167 The Essex Project Site comprises predominantly hardstanding, being occupied by a large area used for vehicle storage, and buildings associated with Tilbury Ferry Terminal. It supports a small linear strip of scrub adjacent to seasonally wet ditches, which is isolated, and not considered suitable to support dormouse (*Muscardinus avellanarius*). Based on the lack of suitable dormouse habitat, it is considered highly unlikely that dormice are present within the Essex Project Site and on this basis, surveys for dormice were not undertaken. The level of survey effort at the Essex Project Site was not raised by consultees during consultation on the EIA Scoping (see Appendix 12.5; Document Reference 6.2.12.5) or PEIR (see Appendix 12.6; Document Reference 6.2.12.6).

Kent Project Site

- 12.14912.168 During the 2020 update desk study Kent and Medway Biological Records Centre (KMBRC) returned 12 records of dormouse, including three records dating from 2017 near the Bluewater Shopping Centre. The closest of these was 250m west of the Kent Project Site. Another record from 2011 originated from a similar area between the Bluewater Shopping Centre and the A296. The other records were all over ten years old, none of which originated from within the Project Site.
- 12.15012.169 In addition, the 2016 Dormouse report by Corylus Ecology showed dormouse records from the woodland running east-west just to the north of the A2(T) and within Darenth Woods SSSI (an ancient semi-natural woodland).
- 12.15112.170 Nest tube surveys to confirm the presence/likely absence of dormouse have been completed within suitable habitat throughout the Kent Project Site between April and November 2020. The results are provided in full detail in Appendix 12.1 (Document Reference 6.2.12.1), along with a habitat suitability assessment in the Dormouse Mitigation Strategy enclosed within Appendix 12.3 (Document Reference 6.2.12.3), are summarised below. The following should be read in conjunction with Figure 12.19 (Document Reference 6.3.12.19) which shows the distribution of dormouse activity across each month surveyed, and Figure 12.20 (Document Reference 6.3.12.20) which summarises the survey findings and shows the locations of any evidence of likely breeding.
- 12.15212.171 During the deployment of the tubes in April 2020, 3 individual dormice were found in old tubes that had remained on the Kent Project Site from previous surveys. Following this, during the May survey visit a total of 8 dormice and 4 nests found. During the August visit, a total of 6 adult and 13 juvenile dormice were found along with 23 nests. In September across all tubes, a total of 26 adult and 15 juvenile dormice were found across the Kent Project Site along with 39 nests. In October, across all tubes, a total of 14 adult



and 7 juvenile dormice were recorded, along with 51 nests. In November, 5 adult and 2 juvenile dormice were found, along with 67 nests.

- 12.15312.172 In summary, surveys have confirmed that dormice are breeding/very likely breeding within the southern half of the Kent Project Site, including within the Sportsground, Bamber Pit, in the Former Landfill and in Station Quarter South. The habitats supporting breeding/likely breeding within these areas comprise unmanaged, dense, continuous scrub of moderate quality, which offers a suitable foraging resource throughout the active season. No evidence of breeding has been recorded on the Swanscombe Peninsula itself (taken to be the land north of Tiltman Avenue, London Road and Galley Hill Road).
- 12.15412.173 The Swanscombe Peninsula supports large areas of moderate quality dense/continuous scrub offering year-round foraging due to the variety of woody species present, in addition to areas of low quality, species-poor scrub scattered over areas of rough, tussocky grassland. Dormouse surveys have confirmed the presence of individual (non-breeding) dormice on the Swanscombe Peninsula (north of Tiltman Avenue, London Road and Galley Hill Road), which is being used for foraging purposes during the summer months. Within parts of the Kent Project Site on the Swanscombe Peninsula, including Botany Marsh, Black Duck Marsh, NE tip and broadness grassland, only adults and nests have been found, with no evidence of breeding.
- 12.15512.174 As such, it is considered that the Kent Project Site supports suitable foraging habitat for dormouse across the Swanscombe Peninsula, alongside some, albeit suboptimal breeding/hibernation habitat within Station Quarter South, the former landfill, Bamber Pit and the Sports ground. The dormouse population within the Kent Project Site is considered of importance at the District level and is taken forward as an IEF.

Harvest Mouse

- 12.15612.175 In 2020, one harvest mouse (Micromys minutus) record was returned by KMBRC dating from 1963 located within the peninsula on the Kent Project Site. Three records were returned from EFC records with only one from the last 10 years and none from the Project Site.
- 12.15712.176 The 2020 surveys confirmed the continued presence of harvest mice on the Kent Project Site, with 11 full harvest mouse nests and a further three partial nests found as shown on Figure 12.23. The majority of the nests were found on Broadness Grasslands with a nest on the NE Tip and another nest in Botany Marsh.
- 12.15812.177 A presence/absence surveys based on nest searches are unable to estimate population size, however the population is considered of at least local level value.



Badger

<u>12.15912.178</u> In 2020, KMBRC and EFC returned many records of badger. None of the records originate from within the Project Site. No evidence of badger (*Meles meles*) has been found during the Extended Phase 1 Survey or any other survey visits so far in 2020. It is not considered likely that badgers are using the Project Site and they will not be taken forward as an IEF.

Otter

- 12.16012.179 The 2020 desk study returned no records of otter (Lutra lutra) from KMBRC or EFC.
- 12.16112.180 A habitat suitability assessment completed across the Project Site (as presented in full in the Otter Mitigation Strategy within Appendix 12.3 (Document Reference 6.2.12.3)) confirmed that none of the habitats within the Essex Project Site are suitable for otter, due their being developed and heavily industrialised with the banks of the River Thames comprising either steep wharfs or concrete banks, and no inland water courses.
- 12.16212.181 By comparison, the Swanscombe Peninsula on the Kent Project Site provides suitable otter foraging habitat with a mosaic of interconnected drainage ditches, reedbed and swamp, fringed by coastal saltmarsh and tidal creeks connected to the River Thames. Further assessment of the suitability of the individual parts of the Swanscombe Peninsula, including Black Duck Marsh, the CTRL wetland, Botany Marsh West, Botany Marsh East and the River Ebbsfleet, is provided in the Otter Mitigation Strategy within Appendix 12.3 (Document Reference 6.2.12.3).
- <u>12.163</u>12.182 An otter was sighted within Black Duck Marsh during the March 2020 wintering bird survey, and the habitat within the reedbeds, marshes and River Ebbsfleet is suitable. The population is considered likely to be of at least Local value.

Water Vole

- 12.16412.183 In 2020, KMBRC returned 25 records for water vole (*Arvicola amphibus*), the most recent record originates from 2016 for a location 5.0km east of the Kent Project Site. The other records all predate 2005 with the majority originating from within the marshes in the north of the Kent Project Site. EFC returned six records of water vole with the most recent from 2009 from Thurrock Park.
- 12.16512.184 As described above in relation to otter, the habitats on the Essex Project Site are not considered suitable for water vole. Large areas of the Swanscombe Peninsula provide suitable water vole foraging habitat due to the mosaic of wetland habitats present, including a connected network of ditches. Further information on the suitability of habitats on the Kent Project Site for water voles is provided in the Water Vole Mitigation Strategy enclosed within Appendix 12.3 (Document Reference 6.2.12.3).



- <u>12.166</u>12.185 The first water vole survey in June 2020 identified water vole signs in Botany Marsh East including extensive signs of feeding and many latrines. In addition, pockets of water vole signs were recorded along the western boundary of Botany Marsh West.
- 12.16712.186 No further water vole signs were recorded in August 2020 however a single water vole latrine was identified on one of the rafts in Black Duck Marsh, and a single latrine was found within the ditch running along the edge of the northeast tip. No signs of water vole were found during either of the hand-search surveys on Botany Marsh West. Furthermore, no signs of water vole were recorded along the River Ebbsfleet.
- 12.187 Surveys of the Site in April 2021 found fresh evidence of water vole in Botany Marsh East only. No signs of water vole were recorded during a visual search completed during deployment of the water vole rafts by boat in June 2021, however, two droppings were recorded on a raft in the lagoon immediately north of the HS1 portal, within the CTRL wetland. No droppings or other evidence of water vole presence was recorded anywhere else within CTRL wetland or Black Duck Marsh during the July 2021 surveys.
- 12.188 The survey results would imply that there are populations present in three areas of the Kent Project Site; Botany Marsh (East and West), CTRL Wetland and Black Duck Marsh. A breeding population of water vole would meet the criteria for designation as an LWS in Kent.—
- 12.189 The population on Black Duck Marsh is isolated. There are no water courses linking this marsh with other areas of the Kent Project Site. The water voles on Botany Marsh east and west and those on the NE Tip are linked. The population as a whole, occupying the Swanscombe Peninsula, is considered to be of District level value. The locations of all water vole signs are shown on Figure 12.21 (Document Reference (6.3.12.21).
- 12.168 The population on Black Duck Marsh is isolated. There are no water courses linking this marsh with other areas of the Kent Project Site. The water voles on Botany Marsh east and west and those on the NE Tip are linked. The population as a whole, occupying the Swanscombe Peninsula, is considered to be of District level value. The locations of all water vole signs are shown on Figure 12.21 (Document Reference (6.3.12.21). A breeding population of water vole would meet the criteria for designation as an LWS in Kent. However, the small amount of evidence gathered suggests that the population is of no greater value than at the local to district level. It is considered likely that the water vole population on the Peninsula is isolated and self supporting with little movement of water vole into or out of the area.

Other Mammals

12.16912.190 KMBRC returned 62 records for hedgehog (*Erinaceus europaeus*) of which 10ten records dated from within the last decade and EFC returned seven records with only three from the last decade. None originated from within the Project Site.



- <u>12.17012.191</u> Ten records for common shrew (*Sorex araneus*) were returned by KMBRC with the majority originating from within the Project Site. Three old records were returned from EFC not from within the Project Site. Six records for pygmy shrew (*Sorex minutus*) were returned by KMBRC of these one originated from within the Project Site dating from 1975.
- <u>12.17112.192</u> Four records of stoat (*Mustela erminea*) and five records of weasel (*Mustela nivalis*) were returned by KMBRC. Three of the weasel records originate from within the Project Site on the Swanscombe Peninsula. EFC retuned a single weasel record from Tilbury.
- 12.17212.193 Two records of brown hare (*Lepus europaeus*) were returned by KMBRC, and one from EFC. All originated from locations outside of the Project Site and the records were old records. Whilst all of the above species could potentially be supported by the Kent Project Site, populations are considered likely to be of less than local importance. None of these species are likely to be supported by the Essex Project Site, based on the lack of suitable habitat present. Other mammals will not be taken forward as an IEF.

Great Crested Newt

- <u>12.173</u>12.194 KMBRC returned 29 records of great crested newts of which only one was from within the last decade. EFC returned seven records with only two from the last decade. That record dated from 2012 and was from 800m south of the Kent Project Site, to the south of Bean. No records were returned from within the Project Site.
- <u>12.17412.195</u> All eDNA results were returned negative. Therefore, great crested newts are considered unlikely to be present and breeding within the Project Site and will not be taken forward as an IEF.

Other Amphibian Assemblage

- 12.17512.196 In 2020, KMBRC and EFC returned many records for palmate newts (*Lissotriton helveticus*), smooth newt (*Lissotriton vulgaris*) common toad (*Bufo bufo*), marsh frog (*Pelophylax ridibundus*) and common frog (*Rana temporaria*). Only one smooth newt record was from the Project Site; however, there were incidental records of smooth newt and marsh frog from the Swanscombe Peninsula, smooth newt from Botany Marsh East and smooth newt and common toad from Bamber Pit during the 2015 surveys.
- <u>12.176</u>12.197 The extent of the wetland habitat and waterbodies within the Kent Project Site has the potential to support large numbers of amphibians and records suggest that smooth and palmate newt, common toad and common frog are present. On a precautionary basis, it is therefore considered that the Kent Project Site supports an amphibian assemblage, valued at the local to <u>district levelDistrict level</u>.

Reptiles



- 12.17712.198 In 2020, KMBRC returned many reptile records. The majority of records were from the Old Malbeon Hospital which is located 1.5km west of the Kent Project Site. There were 189 records for common lizard (*Zootoca vivipara*) (nine of which were from within the Kent Project Site), 104 records of slow worm (*Anguis fragilis*) (two from within the Kent Project Site) and 22 records of grass snake (*Natrix helvetica helvetica*) (three from within the Kent Project Site) returned.
- <u>12.178</u>12.199 KMBRC returned 53 records of adder (*Vipera berus*) but none of these were from within the Project Site. EFC returned less records of all four species, none of which were from within the Project Site.
- 12.17912.200 As described in full detail in Appendix 12.1 (Document Reference 6.2.12.1), surveys have confirmed the presence of an *exceptional* grass snake population (peak survey count = 11 individuals), and *exceptional* common lizard population (peak survey count = 21 individuals) on Swanscombe Peninsula, including land within Black Duck Marsh, Botany Marsh, Broadness Grassland, CTRL wetland, NE tip, and SW tip.
- 12.18012.201 Due to the presence of barriers to dispersal such as existing roads, built development or topographical barriers afforded by the chalk spines and steep pits, the reptile population recorded within some parts of the Kent Project Site are considered isolated. Within these isolated areas, surveys have confirmed the following populations present:
 - Craylands Pit *exceptional* slow-worm population (peak survey count = 39 individuals), and *good* common lizard population (peak survey count = 5 individuals);
 - Bamber Pit good slow-worm population (peak survey count = 14 individuals), low common lizard population (peak survey count = 3 individuals), and low grass snake population (peak survey count = 1 individuals;
 - Sports Ground *low* common lizard population (peak survey count = 2 individuals);
 - Former landfill *low* slow-worm population (peak survey count = 2 individuals), *good* common lizard population (peak survey count = 9 individuals), and *low* grass snake population (peak survey count = 1 individual);
 - Station Quarter North low common lizard population (peak survey count = 1 individual); and
 - Station Quarter South *low* slow-worm population (peak survey count = 3 individuals), *exceptional* common lizard population (peak survey count = 23 individuals), and *low* grass snake population (peak survey count = 2 individuals).
- <u>12.181</u>12.202 For the purpose of the EcIA the reptile populations recorded are considered collectively as an IEF of district value.



<u>12.182</u>12.203 Reptiles are not considered to be present on the Essex Project Site due to the lack of suitable habitat.

Invertebrates

- <u>12.18312.204</u> The Essex Project Site does not support any habitat with potential to support a significant assemblage of terrestrial and aquatic invertebrates.
- <u>12.18412.205</u> The Kent Project Site contains a large complex of habitats offering very diverse array of different micro-habitats and, accordingly, it supports a diverse range of terrestrial and aquatic invertebrate species.
- 12.18512.206 Rather than one particular habitat being of key importance, the value of the Kent Project Site to invertebrates lies in its complex mosaic of habitats in which a range of different successional stages and are represented and in which other environmental conditions such as water/moisture levels and salinity vary significantly. These conditions are in large part the result of a long history of modification and disturbance by industrial activity which continues on the site to the present day. The mosaic of habitats formed upon previously disturbed or made ground, which cover large portions of the Swanscombe Peninsula and the disused chalk pits in the Ebbsfleet Valley, meet the definition of Open Mosaic Habitats on Previously Developed Land (OMH) which are known to support particularly diverse invertebrate populations.

<u>12.186</u>12.207 The habitat mosaics of particular importance to invertebrates are as follows:

- Dry habitats on made ground and/or hardstanding with well drained generally nutrient poor thin soils and supporting a mosaic of bare ground, early colonising/ephemeral vegetation, grassland and scrub; and
- Fresh water and brackish wetland habitats predominantly comprising saltmarsh reedbed, and marshy grassland but including open water (ponds, ditches and streams).
- 12.18712.208 The previous surveys and assessment of the invertebrate population at the Project Site in 2012- and 2015 considered terrestrial and aquatic invertebrate populations as somewhat separate entities and the previous were are divided on these lines. However, in reality there is continuum between wet and dry conditions, and many species of conservation importance rely on habitats at the transition between the two. The 2020 invertebrate survey and assessment recognises transitional as well as purely wet or dry biotopes and habitat types. Therefore, whilst the aquatic invertebrate survey findings will play a specific and separate role in assessing the water quality within the Project Site's waterbodies, the evaluation of the invertebrate population, the assessment of impacts and the strategy for avoidance and mitigation of impacts will consider the invertebrate assemblage as a whole.



- <u>12.188</u>12.209 From the combined 2020 survey area a total of 1,446 invertebrate species were recorded, comprising 1,304 derived from terrestrial sampling methods and 142 from aquatic sampling.
- 12.18912.210 In total, 204 species of recognised conservation status in the UK were recorded from the 2020 survey. These included 10 species listed as 'Species of Principal importance' and two 'research only' species under Section 41 of the NERC Act (2006); as well as 33 species listed in one of the pre-1994 or post-2001 IUCN red data book categories as being RDB3 or 'Near Threatened', or rarer and 159 species currently classed as Nationally Scarce in the UK.
- 12.19012.211 Evaluation of the 2020 survey results indicates that the whole Kent Project Site supports an invertebrate population of National Importance; whilst the majority of sites within the survey area have been found to support representative invertebrate assemblages of National Importance, usually the aquatic elements of the freshwater habitats and the more wooded areas were of somewhat lower conservation value. However, the interdependence of species requiring a combination of one or more habitat means that the value of wooded and wetland elements in relation to open ground habitats should not be disregarded.
- 12.19112.212 From evaluation of the 2020 survey results on a sub-site level; ten of the 17 sample areas were found to support invertebrate assemblages of National Importance; five sample areas were considered to support assemblages of Regional importance; one sample area was assessed as supporting an assemblage of County Importance; and one sample area (Tilbury Docks, verges) was considered to support an assemblage of District importance at most.

Evaluation

12.213 The invertebrate assemblages found within the Kent Project Site is listed as a reason for notification of the Swanscombe Peninsula SSSI. The supporting information to the SSSI notification published by Natural England on 11 March 202115 summarises the invertebrate interest as follows:

"assemblages of invertebrates chiefly associated with bare sand and chalk; open short swards; open water on disturbed mineral sediments; and saltmarsh and transitional brackish marsh".

12.214 The Swanscombe Peninsula SSSI notification document¹⁶ provides a more detailed description of the four invertebrate assemblages present, which is a reason for notification, as follows:

¹⁶ 'Swanscombe Peninsula SSSI Kent – Notification under Section 28C of the Wildlife and Countryside Act 1981.'
Natural England, published 11 March 2021.



¹⁵ 'Swanscombe Peninsula SSSI Kent – Supporting Information'. Natural England, published 11 March 2021.

"Extensive areas of open mosaic habitat on previously developed land and semi-natural habitat including chalk pits, free-draining grassland, scrub, wetlands, grazing marsh and saltmarsh support four diverse assemblages of invertebrates.

Brownfield areas within the SSSI support assemblages of species associated with bare sand and chalk, and open short swards. These assemblages are rich in bee and wasp species which use the open substrates for nesting, prey collection and basking, and the rich wild flower resource for nectar and pollen. Priority Species (Section 41) within the assemblage include the Critically Endangered distinguished jumping spider Sitticus distinguendus, rare five-banded weevil-wasp Cerceris quinquefasciata and Nationally Scarce brown-banded carder bee Bombus humilis.

Wetland areas, primarily Black Duck Marsh and ponds resulting from construction of the Channel Tunnel Rail Link (CTRL), support assemblages of species associated with open water on disturbed mineral sediments, and saltmarsh and transitional brackish marsh. These wetlands support 84 species of water beetle. This represents over a quarter of the UK water beetle fauna. Notable species include the nationally scarce Enochrus halophilus, and the nationally rare great silver water Beetle Hydrophilus piceus."

- 12.215 The invertebrate assemblages present are recognised as being of national importance, and importance and treated as an IEF of national value within this EcIA. For the purpose of this assessment, the invertebrate assemblages are grouped collectively within as an IEF, but impacts and mitigation are considered on each assemblage separately where relevant.
- 12.216 As the four invertebrate assemblages identified on the Kent Project Site are reasons for notification of the Swanscombe Peninsula SSSI, the assessment of potentially significant effects upon the assemblages is included within the assessment of effects upon the higher tier IEF of the Swanscombe Peninsula SSSI.

Freshwater Fish

Previous Surveys

- 12.19212.217 A fish survey of the River Ebbsfleet initially undertaken by Coclough and Coates Aquatic Consultants in 2015 (see Appendix 12.1; Document Reference 6.2.12.1) captured modest populations of mature roach and perch. The fish surveys identified no evidence of active recruitment to the fishery, such that a population present within the River Ebbsfleet is not self-sustaining in the long-term. It was considered that fish populations within the River Ebbsfleet could either be wash outs from local fisheries or may have been introduced by anglers.
- <u>12.193</u>12.218 A fish survey of waterbodies across Swanscombe Marshes in August 2015 recorded three-spined stickleback in small numbers, in isolated locations in the eastern complex of Swanscombe Marshes and on the bottom edge of Botany Marshes. No fish were captured anywhere in the western complex of Swanscombe Marshes despite suitable conditions.



Overall, survey effort identified a poor head of fish within waterbodies comprising Swanscombe marshes, likely attributed to poor water quality and anaerobic bed conditions in addition to saline intrusion and unstable water levels observed during 2015.

12.19412.219 A desk study request to the Environment Agency confirmed there are no Environment Agency monitoring stations along the River Ebbsfleet and no historical data with respect to an assessment of fish communities. This is with the exception of a single fish survey undertaken by the Environment Agency during 2007 (adjacent to Ebbsfleet International Station) during which no fish were captured.

2020 Surveys

- 12.19512.220 Further fish surveys of waterbodies comprising Swanscombe Marshes (where access was possible) confirmed the continued absence of a significant fish assemblage, likely attributed to poor water quality, anaerobic bed conditions in addition to saline intrusion. No fish were captured at any of the survey sites or observed during survey effort whilst the ditch network across Botany Marsh was predominantly dry.
- <u>12.196</u>12.221 Particularly given limitations to survey effort, it is assumed that several of the wet drains and ponds have potential to support some fish species, such as three spined stickleback as previously recorded during 2015.
- 12.19712.222 With respect to the River Ebbsfleet, fish populations are considered to be constrained by the availability of suitable habitat whilst the presence of significant culverts at both the upstream and downstream extent of the River Ebbsfleet are considered a significant barrier to the fish movement and migration. As such, a freshwater fish community within waterbodies across the Project Site are considered to be of no more than Site importance.
- 12.19812.223 The freshwater fish assemblage is not an IEF as the value is considered likely to be at a Site level only however, due to consultation responses from the Environment Agency (see Appendix 12.5; Document Reference 6.2.12.5), the freshwater fish assemblage will be taken forward and considered within the EcIA to inform a 'no deterioration assessment' of on-site waterbodies.

Summary of Important Ecological Features

<u>12.199</u>12.224 Based on the baseline ecological information described above (and presented in full in Appendix 12.1; Document Reference 6.2.12.1), a number of IEFs have been identified and are summarised in **Table 12.9**.



Table 12.9: Important Ecological Features to be Considered Within the EcIA.

Important Ecological Feature	Key Attributes	Nature Conservation Value
Designations		
Thames Estuary and Marshes SPA/Ramsar	Extensive intertidal mudflats with saltmarsh and channel systems. Internationally importance assemblage of birds and wintering populations of many wader species.	International
Medway Estuary and Marshes SPA/Ramsar/SSSI	Single tidal system with the Swale and joins the southern part of the Thames Estuary between the Isle of Grain and Sheerness. Internationally importance of assemblage of birds and wintering populations of many wader species. The Medway Estuary and Marshes form the largest area of intertidal habitats which have been identified as of value for nature conservation in Kent and are representative of the estuarine habitats found on the North Kent coast. A complex of mudflats and saltmarsh is present with in places grazing marsh behind the sea walls which is intersected by dykes and fleets. The area holds internationally important populations of wintering and passage birds and is also of importance for its breeding birds. An outstanding assemblage of plant species also occurs on the site.	International
North Downs Woodland SAC	Semi-natural dry grasslands and scrubland facies: on calcareous substrates (Festuco-Brometalia) for which the area is considered to support a significant presence. Asperulo-Fagetum beech forests for which this is considered to be one of the best areas in the United Kingdom. Taxus baccata woods of the British Isles for which this is considered to be one of the best areas in the	International



Important	Key Attributes	Nature
Ecological Feature		Conservation
		Value
<u>Swanscombe</u>	Swanscombe Peninsula SSSI is a corridor of	<u>National</u>
Peninsula SSSI	habitats connecting Ebbsfleet Valley with the	
	southern shore of the River Thames between	
	<u>Dartford and Gravesend. Industrial processes such</u>	
	as engineering, power generation, landfill and	
	dredging have left a legacy of low nutrient and	
	often toxic substrates, which have developed into	
	bare open ground habitats with low scrub cover.	
	The peninsula peninsula also supports wetland,	
	grazing marsh and saltmarsh habitats. These	
	habitats, coupled with a mild climate, provide ideal	
	conditions for certain species and assemblages of	
	plants, invertebrates and breeding birds. The SSSI	
	is of interest for the following biological interest	
	features:	
	Vascular plants - populations of the plants	
	divided sedge (<i>Carex divisa</i>), yellow vetchling	
	(Lathyrus aphaca), slender hare's-ear	
	(Bupleurum tenuissimum), Bithynian vetch	
	(Vicia bithynica) and round-leaved wintergreen	
	(Pyrola rotundifolia subsp. maritima);	
	 Invertebrates - assemblages of invertebrates 	
	associated with bare sand and chalk, open short	
	swards, open water on disturbed mineral	
	sediments and saltmarsh and transitional	
	brackish marsh; and	
	Breeding birds - two diverse assemblages of	
	breeding birds, one associated with lowland	
	open waters and their margins, lowland fen and	
	lowland damp grassland, the other with lowland	
	scrub.	
Darenth Woods	Some of the most valuable areas of ancient semi-	National
SSSI	natural woodland in north-west Kent with rare	
	woodland types.	



Important Ecological Feature	Key Attributes	Nature Conservation
		Value
Medway Estuary	Component SSSI to the Medway Estuary and	National
and Marshes SSSI	Marshes SPA/Ramsar. Forms the largest area of	
	intertidal habitats that have been identified as of	
	value for nature conservation in Kent, and are	
	representative of the estuarine habitats found on	
	the North Kent coast. A complex of mudflats and	
	saltmarsh is present with in places grazing marsh	
	behind the sea walls, which is intersected by dykes	
	and fleets.	
Shorne and	A complex of ancient and plantation woodland,	<u>National</u>
Ashenbank Woods	and includes a variety of stand-types associated	
<u>SSSI</u>	with Tertiary gravels, clays and sands. The site	
	supports an important and diverse invertebrate	
	fauna.	
Inner Thames	Largest remaining expanse of wetland bordering	National
Marshes SSSI	the upper reaches of the Thames Estuary.	
	Diverse bird interest especially the variety of	
	breeding birds and the numbers of wintering	
	wildfowl, waders, finches and birds of prey.	
South Thames	Component SSSI to the Thames Estuary and	National
Estuary and	Marshes SPA/Ramsar.	
Marshes SSSI		
	Extensive mosaic of grazing marsh, saltmarsh,	
	mudflats and shingle characteristic of the estuarine	
	habitats of the north Kent marshes. Freshwater	
	pools and some areas of woodland provide	
	additional variety and complement the estuarine	
	habitats. Supports outstanding numbers of	
	waterfowl, total counts regularly over 20,000.	
West Thurrock	One of the most important sites for wintering	National
Lagoon and	waders and wildfowl on the Inner Thames Estuary.	
Marshes SSSI	Extensive intertidal mudflats together with a large	
	and secure high tide roost, attracts waders in	
	nationally important numbers, with significant	
	populations of other bird species. The adjacent	
	Stone Ness saltmarsh is noted for the size and	
	character of its high marsh plant community.	



Important Ecological Feature	Key Attributes	Nature Conservation Value
Mucking Flats and Marshes SSSI	Mucking Flats and Marshes is the underpinning SSSI for the Thames Estuary and Marshes SPA. It comprises an extensive stretch of Thames mudflats and saltmarsh, together with sea wall grassland. Wintering wildfowl and waders reach both nationally and internationally important numbers on the mudflats, roosting and feeding on adjacent saltmarsh and disused silt lagoons. The mudflats form the largest intertidal feeding	National
	area for wintering wildfowl and waders west of Canvey Island on the north bank of the Thames	
Cobham Woods SSSI	Woodland and old parkland representative of acidic Thanet Sands and chalk soils. An outstanding assemblage of plants is present at this site, which is also of importance for its breeding birds. Includes an area of arable land, which is notable for the number of uncommon annual plant species.	National
Great Crabbles	Representative of woods on North West Kent	National
Wood SSSI	Tertiary sediments. Mostly mixed coppice under oak standards, with sweet chestnut as the dominant species. A number of scarce plants occur, including lady orchid and man orchid.	
Wouldham to Detling Escarpment SSSI	Component SSSI of the North Downs Woodland SAC. Designated for chalk grassland, woodland and invertebrate interest.	National
Botany Marshes LWS	Reedbed and potential for ditch & grazing marsh restoration. Reedbed and grazing marsh are of principal importance in England. Also supports three species of reptile, water vole, otter and is of value to birds.	County
Ebbsfleet Marshes, Northfleet LWS	Range of habitats including reedbed, calcareous stream, lake, scrub, woodland, calcareous and neutral grassland. Protected species have been recorded including reptiles and great crested newts.	County
Alkerden Lane Pit LWS	Contains nationally scarce plants and Kent's largest population of green-flowered helleborine (<i>Epipactis phyllanthes</i>). Also contains round leaved wintergreen (<i>Pyrola rotundifolia</i>) and several species of nationally rare and scarce invertebrates.	County



Important Ecological Feature	Key Attributes	Nature Conservation Value
Disused Hospital Grounds, Mabledon LWS	Designated for its chalk habitats including chalk grassland. The designation contains mixed scrub. The habitats within the designation support	County
Tilbury Marshes LWS	reptiles and lepidoptera. Diverse saltmarsh flora. Good grazing-marsh flora. An important invertebrate habitat destroyed by development, but some of the key species may survive on these remaining fragments.	County
Habitats/Flora		
Rare plants Broad leaved Semi Natural Woodland	Populations of 13 nationally scarce species were found in 2016. Eight species were recorded in 2020, along with an additional two2 species not recorded previously. Populations of the plants divided sedge (Carex divisa), yellow vetchling (Lathyrus aphaca), slender hare's-ear (Bupleurum tenuissimum), Bithynian vetch (Vicia bithynica) and round-leaved wintergreen (Pyrola rotundifolia subsp. maritima) are listed as a reason for notification of the Swanscombe Peninsula SSSI. The assessment of effects upon these notified features of the Swanscombe Peninsula SSSI is undertaken as part of the assessment of effects upon the SSSI. Woodland with good canopy species and ground flora species diversity. Connects to other woodlands in wider area. Meets criteria for Priority	National
Ancient woodland	habitat. Considered an irreplaceable habitat in planning terms, with Darenth Woods also designated as a SSSI. Remaining areas of ancient woodland not designated as SSSI or LWS. Collectively, considered of county value.	County
Scrub	Extensive mature and colonising scrub forming a corridor of woody habitats between the A2(T) and the River Thames.	Local
Semi-improved grassland	Including areas of species-poor semi-improved grassland and areas of semi-improved neutral, and calcareous grassland (with relict areas of more species-rich grassland of NVC MG1d and CG2 but not extensive or fine examples).	Local to District



Important Ecological Feature	Key Attributes	Nature Conservation Value
Coastal/Floodplain Grazing Marsh	Botany Marsh West - Priority Habitat ¹⁷ coastal/floodplain grazing marsh. Would qualify as a LWS.	District
Open mosaic on previously developed land	Large areas across the Kent Project Site where there is a verifiable history of industrial disturbance, and which fulfil the Priority Habitat description.	District
	Comprises an 'umbrella' IEF which includes a range of individual habitat types, including areas bare ground, ephemeral vegetation, open grassland (poor semi-improved grassland and semi-improved neutral and calcareous grassland) and patches of scrub (less than 2500m²).	
Waterbodies (ponds, temporary pools, standing water and ditches)	Extensive ditch network around the peninsula with associated ponds. Ditch network forms part of a large marsh area including Botany Marshes LWS and adjacent grazing marsh and is considered of district level District level. Some ponds, within Broadness grassland particularly, are contaminated by leachate from the nearby cement production facility and are of negligible ecological value.	District
Swamp (reedbed)	Three main areas in Black Duck Marsh, CTRL wetland and Botany Marsh, the latter of which is partially designated as a LWS. The other areas could qualify as LWSs and all qualify as Priority Habitat.	County
River Ebbsfleet	Acts as a wildlife corridor and is linked to reed bed and woodland habitats. Moderate water quality.	Local
Species	·	
Wintering waterfowl and wading bird assemblage	Supports many of the species associated with the nearby SPA/Ramsar <u>Sites</u> (albeit only a small proportion of the total populations).	International
Wintering terrestrial bird assemblage	28 species of conservation concern recorded in low to moderate numbers.	County

UK Biodiversity Action Plan Priority Habitat Descriptions Coastal and Floodplain Grazing Marsh From: UK Biodiversity Action Plan; Priority Habitat Descriptions. BRIG (ed. Ant Maddock) 2008



Important	Key Attributes	Nature
Ecological Feature		Conservation
D 11 D1 1		Value
Breeding Bird	99 species recorded of which 29 were listed on the	Regional National
Assemblage	Amber list of Birds of Conservation Concern, and	
	17 on the Red list. Of the 99 species, 77 have been	
	confirmed breeding or possibly breeding, and the	
	other 22 species confirmed as non-breeding.	
	Two distinct breeding bird assemblages are present	
	on the Kent Project Site, including one associated	
	with lowland open waters and their margins,	
	lowland fen and lowland damp grassland; and one	
	associated with lowland scrub. The wetland	
	breeding bird assemblage includes a nationally	
	significant population of pochard (7-10 pairs	
	present, which equates to between 0.99% and	
	1.4% of the national breeding population).	
	The two breeding bird assemblages found on the	
	Kent Project Site are listed as a reason for	
	notification of the Swanscombe Peninsula SSSI.	
	The assessment of potentially significant effects	
	upon the assemblages is included within the	
	assessment of effects upon the higher tier IEF of the	
	Swanscombe Peninsula SSSI.	
Pochard breeding	Possibly breeding with 7-10 pairs present, which	National
population	would equate to between 0.99% and 1.4% of the	
	national breeding population.	



Important Ecological Feature	Key Attributes	Nature Conservation
		Value
Bat assemblage	Assemblage of at least eight species, including one	District
	Kent BAP species. However, the activity is	
	predominantly of common pipistrelle bats.	
	Two buildings on the Kent Project Site with	
	confirmed bat roosts, including a common	
	pipistrelle summer day roost within building B32	
	and a soprano pipistrelle summer day roost within	
	building B67. Other buildings with	
	high/moderate/low potential to support roosting	
	bats are also present, as well as buildings that	
	could not be surveyed due to access restrictions.	
	No tree roosts confirmed but nine trees with high	
	bat roost potential, five with moderate bat roost	
	potential and six with low bat roost potential are	
	present.	
	Three tunnels (TU/011, TU/013A and TU/014A)	
	returned low numbers of recordings of common	
	pipistrelle and soprano pipistrelle bats during	
	monitoring for winter hibernation but considered	
	unlikely to be hibernation roosts. One tunnel	
	(TU/018) with low hibernation potential was not	
	surveyed due to access restrictions.	
Bat assemblage	Tunnel TU/07 confirmed as a winter hibernation	County
Tunnel TU/07 and	roost for Myotis sp and potentially low numbers of	
TU/016	pipistrelle species.	
hibernation roost		
	Tunnel TU/016 possible winter hibernation roost	
	for low numbers of common -pipistrelle and	
	soprano pipistrelle bats.	



Important Ecological Feature	Key Attributes	Nature Conservation Value
Dormouse	Confirmed present within the Kent Project Site. Activity greatest within the former landfill, with the majority of nests, adults and juveniles found there. Confirmed breeding/ considered very likely breeding within the southern half of the Lent-Kent Project Site in Station Quarter South, former landfill, Bamber Pit and the Sports ground. Individuals dispersing across to the Swanscombe peninsula (taken to be the land north of Tiltman Avenue, London Road and Galley Hill Road) for summer foraging, with no evidence of breeding on the peninsula. Likely to be a meta population with that close to the Bluewater shopping centre.	District
Otter	Confirmed present within Black Duck Marsh, and assumed present in low numbers on the suitable habitat throughout the ditch network, reedbeds, marshes and on the River Ebbsfleet.	Local
Water vole	Latrines and feeding signs found in Botany Marsh East and West and NE tip – likely breeding and therefore meets LWS selection criteria.	Local to district
Harvest mouse	Confirmed present on Swanscombe Peninsula especially in Broadness grassland and on Botany Marsh.	Local
Amphibian assemblage	Likely to support four species and meets criteria for LWS selection.	Local to district
Reptile assemblage	Three species supported, with several populations of varying sizes across the Kent Project Site. Many parts of the Kent Project site meet criteria for LWS selection.	District



		Value
species, comprising sampling method A total of 142 species status in the UK was survey, including Principal importation species under Section well as 33 species post-2001 IUCN range RDB3 or 'Near That species currently UK. Four distinct invector on the Kent Project invertebrates chies chalk; open short mineral sediment brackish marsh. The Kent Project State Swanscombe assessment of pothe assemblages in the Swanscombe assessment of pothe session and the UK species comprision in the UK was species under Section in the UK was spec	mblage comprising a total of 1446 ng 1,304 derived from terrestrial is and 142 from aquatic sampling. It cies of recognised conservation were recorded from the 2020 10 species listed as 'Species of ince' and two 'research only' it ion 41 of the NERC Act (2006); as a listed in one of the pre-1994 or ied data book categories as being reatened', or rarer and 159 classed as Nationally Scarce in the interest assemblages are present in the interest assemblages of interest assemblages on the invertebrate assemblages on the invertebrate assemblages on the invertebrate assemblages on it is and saltmarsh and transitional the invertebrate assemblages on it is are reasons for notification of it is are reasons for notification of it is included within the assessment included within the assessment included SSSI.	National

- <u>12.20012.225</u> In accordance with the assessment methodology described at paragraph 12.40, all other habitats and species/species groups are deemed to be of only Site level nature conservation importance or less, and will not be taken forward for detailed assessment, since effects upon such features are unlikely to be 'significant' in EIA terms.
- <u>12.201</u>12.226 This is with the exception of a freshwater fish assemblage which are considered to inform a 'no deterioration assessment' of the River Ebbsfleet, as described above, and provided in Appendix 12.8 (Document Reference 6.2.12.8).

Future Baseline Conditions

12.20212.227 It is anticipated that, in the absence of development and cessation of industrial practices, the Kent Project Site would continue to be neglected resulting in is likely to be subject to further scrub encroachment and natural succession through to woodland, although this process would be relatively slow owing to the nutrient poor skeletal soils and highly alkaline conditions which would supress rates of vegetation colonisation and growth. The notification of the majority of the Kent Project Site as Swanscombe Peninsula



SSSI places a responsibility on the landowners to maintain the SSSI in favourable condition, however, it is likely to be challenging to replicate the previously unplanned disturbances and deposition of industrial waste products etc. which have created the current site conditions.

12.20312.228 Material changes in the habitat composition and the species populations they support are unlikely to occur during the relatively short time period between the recent surveys and construction activities starting on site. However, in the medium to long-term, it is anticipated that the existing species populations/assemblages which are dependent on woody/scrub habitats, such as for example the terrestrial breeding bird assemblage and dormice would flourish, whilst others, reliant on open grassland/marsh habitats and an 'open mosaic', including for example waders, waterfowl, reptiles and invertebrates, would diminish.

12.20412.229 There is also evidence of fly tipping, littering, illegal camping, and fires across the Kent Project Site, which has a negative impact on the existing biodiversity, and can reasonably be expected to continue if the site remains undeveloped, albeit the SSSI notification may assist in discouraging some of these damaging activities.

POTENTIAL SIGNIFICANT ENVIRONMENTAL EFFECTS OF THE PROPOSALS

Mitigation Hierarchy

12.20512.230 The mitigation hierarchy requires that the design of a development follows what is known as the mitigation hierarchy to reduce impacts upon Important Ecological Features (IEFs). The Chartered Institute for Ecology and Environmental Management (CIEEM) guidelines for Ecological Impact Assessment (EcIA)¹⁸ outline the principles as follows:

'Avoidance: Seek options that avoid harm to ecological features (for example, by locating on an alternative site).

Mitigation: Negative effects should be avoided or minimised through mitigation measures, either through the design of the project or subsequent measures that can be guaranteed – for example, through a condition or planning obligation.

Compensation: Where there are significant residual negative ecological effects despite the mitigation proposed, these should be offset by appropriate compensatory measures.

Enhancement: Seek to provide net benefits for biodiversity over and above requirements for avoidance, mitigation or compensation.'

¹⁸ CIEEM (2018) Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, Freshwater, Coastal and Marine version 1.1. Chartered Institute of Ecology and Environmental Management, Winchester.



_

- <u>12.20612.231</u> The Project Site was chosen by the project team after careful consideration of a number of sites from a wide search area which included land within 100km of central London. The full consideration of alternative sites and site selection process is detailed in full in Chapter 4: *Project development and alternatives* (Document Reference 6.1.4).
- 12.20712.232 The Project Site was selected as the preferred location following review of a number of criteria including (in no order of priority) land availability, land use, planning and environmental constraints, proximity to and connectivity with central London, transport and accessibility, environmental constraints, planning constraints, regeneration and economic benefit and micro-climate. The Swanscombe Peninsula was subsequently chosen as it offers a unique combination of advantages, as detailed in Chapter 4 of the ES (Document Reference 6.1.4):

'It centres upon a large and generally unused brownfield site with a broadly level terrain, large enough to accommodate a full resort development. It is close to the edge of London but outside of the metropolitan green belt. It lies only 1 km north of Ebbsfleet International Station, which offers high speed train connections to London St Pancras International station with a journey time as low as 17 minutes and services to and from continental Europe.

Strategic highway routes in the locality include the A2(T), which passes 3 km to the south of the peninsula and provides a connection to Junction 2 of the M25 motorway to the west and onwards into London. The Dartford Tunnels and Queen Elizabeth II Bridge crossings of the River Thames lie approximately 3 km to the west of the site.

The Swanscombe Peninsula does not contain any international or national wildlife or heritage designations, and it offers the potential to dovetail the resort development with significant local economic regeneration initiatives.'

- 12.20812.233 Having determined the best location of the entertainment resort, the development layout for the site was appraised with consideration of a number of variables including (but not limited to) existing land use, land ownership, ground conditions, drainage, land contamination, local terrain, and transport requirements. From an ecological perspective, the development layout has been situated to avoid impacts on the most sensitive habitats where possible, and retains areas of habitat within Black Duck Marsh, Botany Marsh Local Wildlife Site and Broadness grassland, as well as areas of seminatural habitat within Bamber Pit and through the Ebbsfleet Valley, and the majority of the existing coastal saltmarsh and inter-tidal mudflats around the edge of Swanscombe Peninsula. Furthermore, the land take from the Proposed Development mostly includes existing development or formerly developed land and landfill sites. In order to deliver a viable entertainment resort, the site needs to be large enough to accommodate the entertainment resort, including two theme park gates, attendant visitor attractions and amenities, hotels and transport facilities. Some habitat loss is therefore unavoidable.
- 12.20912.234 Positioning the resort further west or north within the Swanscombe Peninsula, whilst avoiding impacts upon Botany Marsh West, the CTRL wetlands, and coastal



floodplain grazing marsh (Priority habitat), and areas of Open Mosaic Habitat on Previously Developed Land (OMHPDL), would necessitate loss of further wetland habitats including the reedbed (Priority habitat) within Black Duck Marsh and bring the development closer to the estuary front and coastal saltmarsh (Priority habitat) and is also constrained by the ultra high voltage connectors to the super pylon. Black Duck Marsh is considered to be 'functionally linked' to nearby statutory designated sites, including the Thames Estuary and Marshes Ramsar/Special Protection Area (SPA)/SSSI and the Medway Estuary and Marshes Ramsar/SPA/SSSI, as it supports various waterfowl species over winter. Furthermore, it also supports a diverse breeding bird assemblage not associated with the aforementioned designated sites, including small populations of breeding pochard, bearded tit and Cetti's warbler. On balance it is considered that situating the resort further west on the peninsula would have greater ecological impact.

12.21012.235 The ecology mitigation strategy has been developed to mitigate and compensate for negative impacts remaining after application of the first 'avoid' stage of the mitigation hierarchy. This includes mitigation for impacts to the mosaic of habitats within the Kent Project Site, and associated fauna, as discussed further in this Chapter. Mitigation is not required at the Essex Project Site based on the absence of any impacts on terrestrial and freshwater ecological features.

Inherent Mitigation Measures

12.21112.236 The Proposed Development includes a range of inherent mitigation measures which are illustrated within various design documents submitted along with the DCO application including the Illustrative Masterplan (Document Reference 6.3.3.1) and Landscape Masterplan enclosed within the Landscape Strategy (Document Reference 6.2.11.7), with details of management/enhancement provided within the Ecological Mitigation and Management Framework (Document Reference 6.2.12.3).

<u>12.212</u>12.237 The inherent mitigation measures are provided below in respect of measures that will benefit biodiversity generally, and specifically in relation to key species/taxa:

General:

- Retention of areas of ecological value including Black Duck Marsh, Botany Marsh East LWS, Broadness grasslands and saltmarsh, the majority of Bamber Pit and areas of semi-natural habitat throughout the Ebbsfleet Valley;
- Retention of large areas of saltmarsh and inter-tidal mudflats around the edge of the Swanscombe Peninsula; and
- Provision of a large area managed primarily for biodiversity including Black Duck Marsh, Broadness grassland and Botany Marsh East LWS, comprising a mixture of reedbed, open grassland, scrub, ditches, ponds and wetland/SuDS features.



Bats:

• Retention and management of foraging and commuting habitats, as described by the general points above.

Dormouse:

- The retention of areas of suitable habitat within Broadness grassland, including existing woodland, tree and scrub habitats, and where possible within Bamber Pit;
- The retention and management of dense bramble and low growing scrub and trees along the north western boundaries of Black Duck Marsh, to maintain potential dispersal routes from Swanscombe Peninsula to the south-west;
- The retention and enhancement of a continuous belt of woodland habitat along the southern boundaries of Black Duck Marsh, connecting to additional green corridors proposed along the southern boundary adjacent to Tiltman Avenue, to ensure the continued functioning of existing dispersal routes to valuable off-site habitats to the south-west, including the woodland at the Swanscombe Heritage Park; and
- The retention, enhancement and creation of additional woodland habitat alongside the sensitive design of new landscaping around the peripheries of Botany Marsh, necessary to further promote habitat connectivity between Swanscombe Peninsula and habitats across the wider landscape to the south.

Water vole:

• The maintenance of habitat connectivity between Botany Marsh East, Broadness saltmarsh and Black Duck marsh through the inclusion of a chain of water courses and water bodies wrapping around the side of the Proposed Development.

Invertebrates:

Retention (and enhancement where possible) of key areas of invertebrate habitat
including within the eastern portion of Broadness Saltmarsh (including a specific zone
of habitats around the base of the central electricity pylon which previous surveys
have identified as being particularly important for thermophilic spider fauna open
mosaic habitats), saltmarsh on the north west and north east fringes of Swanscombe
peninsula, and Botany Marsh on the eastern side of Swanscombe peninsula.



Construction Impacts and Effects

- 12.213 In the absence of additional mitigation, the potential effects of the Proposed Development during the construction phase on the IEFs identified above are summarised below:
 - Direct habitat loss, damage or degradation;
 - Habitat fragmentation/loss of flight paths/dispersal routes;
 - Habitat disturbance;
 - Killing, injuring and disturbance of wild animals;

•

- Increased collision risk to birds and bats;
- Increased noise, vibration, visual and light disturbance;
- Changes in air quality including from dust, construction waste and pollutants, and exhaust emissions;
- Hydrological effects, including changes to water quality/quantity;
- Pollution/contamination incidents; and
- Spread of invasive species.
- <u>12.21412.239</u> Further details, including which IEFs these potential effects relate to, are provided within **Table 12.10**.
- <u>12.215</u>12.240 The following should be read in conjunction with the Shadow Habitat Regulations Assessment (Document Reference 6.2.12.4), which provides further information on the assessment on potential effects upon European Sites.
- 12.21612.241 In addition, an impact assessment specific to each species/species assemblage/taxa is also provided within the individual mitigation strategies enclosed as appendices to Appendix 12.3: Ecological Mitigation and Management Framework (Document Reference 6.2.12.3).
- 12.21712.242 The assessment of potential effects during construction has been undertaken using information derived from other chapters within the Environmental Statement, notably Chapter 15: Noise and Vibration (Document reference 6.1.15), Chapter 16: Air Quality (Document reference 6.1.16), and Chapter 17: Water Resources and flood risk (Document reference 6.1.17).



Table 12.10: Construction effects on important ecological features without mitigation.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
Designations				
Thames	International	Potential impacts within the European/Ramsar S	ite itself:	
Estuary and		Damage to habitats within designated area	Temporary, reversible, minor	Not significant ²⁰
Marshes		used by cited bird species from changes to	magnitude and extent.	
SPA/Ramsar,		water and/or sediment quality (either from		
and the		surface or groundwater discharges from the		
component		Project Site including construction waste and		
SSSIs		pollutants; or from disruption of contaminated		
including		Thames sediments), with potential associated		
South		knock-on risk of bioaccumulation.		
Thames		Damage to habitats within the designated area	Temporary, reversible, minor	Not significant
Estuary and		used by cited bird species from changes in air	magnitude and extent.	
Marshes SSSI		quality, including from dust, construction		
and Mucking		waste and pollutants, exhaust emissions and		
Flats and		increased river traffic.		
Marshes		Damage to habitats used by cited bird species	Temporary, reversible, minor	Not significant ²¹
SSSI ¹⁹		within the designated area from introduction	magnitude and extent.	

South Thames Estuary and Marshes SSSI and Mucking Flats and Marshes SSSI are component SSSIs to the Thames Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar

²¹ Construction river traffic will be limited to that coming from Tilbury and from central London, and therefore the risk of INNS introductions in ballast water discharged by vessels is unlikely. Furthermore, the distance form regular shipping lanes to the SPA/Ramsar site is considered a limiting factor in the potential for INNS introductions. See Shadow Habitat Regulations Assessment (ES Appendix 12.3), Stage 1 Screening Matrix (Annex 4)



²⁰ By virtue of the distance between the Project Site and either the SPA/Ramsar/SSSI site, no significant effects are anticipated. Changes to sediment circulation patterns are likely to be minor given the open structure of the proposed new jetty at the Kent Project Site and of the extension of facilities at Tilbury. At this point no dredging is anticipated, Standard avoidance and management measures will reduce the likelihood of pollution events, and it is considered that any minor incidents that occur will be sufficiently dilute by the time they reach the TE and M SPA/Ramsar 2.8km downstream that they will not have the potential to cause LSE on the SPA/Ramsar.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
(see		or proliferation of Invasive Non-Native Species		
Appendix		(INNS), as a result of construction works (and in		
12.5)		particular shipping/transport of materials).		
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Not significant ²²
		giving rise to displacement, behavioural	magnitude and extent.	
		changes or physiological stress to cited bird		
		species within the designated area.		
		Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ²³²⁴
		with increased shipping traffic and construction	magnitude and extent.	
		activities, giving rise to displacement,		
		behavioural changes or physiological stress to		
		cited bird species within the designated area.		
		Potential impacts solely on functionally linked res	sources (outside of the European/I	Ramsar Site):
		Direct loss of/damage to functionally linked	Permanent, moderate	Significant negative
		land on the Kent Project Site, potentially used	magnitude and extent.	at district
		by cited bird species, totalling a net loss of		level Significant

It is anticipated that no additional shipping movements would occur along the Thames past either site during construction or operation. Whilst construction phase movements will include additional barge movements between the Kent and Essex Project Sites, this is not anticipated to increase shipping traffic within or nearby to either the Thames Estuary and Marshes or Medway Estuary and Marshes SPA/Ramsar Sites. These additional barge movements will be accommodated within the normal and ongoing delivery pattern and will not represent an uplift on disturbance due to the combined and absolute limitations of berthing capacity and tidal restrictions at that site. See Shadow Habitat Regulations Assessment (ES Appendix 12.3), Stage 1 Screening Matrix

In regards to noise, peak or mean (i.e. 24hr) noise in excess of 55dB is not predicted to be experienced outside of the DCO Order Limits for most construction or operational activities, with the exception of construction-phase jetty piling and dredging and pavement construction, for which values are not yet known. However, based on the information provided for more extensive works in association with Tilbury2, the foremost of these could see noise levels of 63dB at 300m from source with the latter having the potential to slightly exceed the 55dB level at 300m (Tilbury2 ES Chapter 17 Table 17.30 [APP-031]). These data indicate that noise levels during construction would not be sufficient to elicit any behavioural responses in birds at over 2.8km (the nearest point of the Thames Estuary and Marshes SPA/Ramsar Site). See Shadow Habitat Regulations Assessment (ES Appendix 12.3), Stage 1 Screening Matrix



The distance between the Project Site and the nearest part of the Thames Estuary and Marshes SPA (foreshore adjoining Eastcourt/Shorne Marshes) is approximately 3.3km and the nearest part of the Ramsar approximately 2.8km. At these distances significant impacts from lighting are not anticipated.

Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		15.49ha, including 14.55ha of coastal and		negative at District
		floodplain grazing marsh (Botany Marsh West)		level
		and 0.94ha of reedbeds (Black Duck Marsh) to		
		accommodate construction of the Leisure Core.		
		Damage to functionally linked habitats	Temporary, moderate	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at district
		changes to hydrology (as a result of drainage		level Significant
		and infill of the CTRL wetland, swamp/wetland		negative at District
		habitat along the east/south east of Black Duck		<u>level</u>
		Marsh and other low-lying seasonally		
		inundated areas); and changes to water and		
		/or sediment quality (either from surface or		
		groundwater discharges from Project Site		
		including construction waste and pollutants;		
		or from disruption of contaminated Thames		
		sediments), with potential associated knock-on		
		risk of bioaccumulation.		
		Damage to functionally linked habitats	Temporary, minor magnitude	Significant negative
		potentially used by cited bird species from	and extent.	at district
		changes in air quality including from dust,		levelSignificant
		construction waste and pollutants ²⁵ .		negative at District
				<u>level</u> ²⁶
		Damage to functionally linked habitats	Temporary, reversible, minor	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at district

No retained functionally linked habitats lie within 200m of an existing road. Effects due to increased construction traffic have therefore been screened out of this assessment.

Given the proximity of construction work to retained functionally linked habitats (<50m in some cases, especially around Black Duck Marsh and along the estuary front), significant dust deposition is anticipated.



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
	посерье:	introduction or proliferation of Invasive Non- Native Species (INNS)INNS.		levelSignificant negative at District
		Disturbance (from shipping/ferry movements) giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	Temporary, reversible, minor magnitude and extent.	Significant negative at district level Significant negative at District level
		Disturbance (noise and lighting) associated with increased shipping traffic and construction activities, giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	Temporary, reversible, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level ²⁷
		Disturbance (human movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during construction.	Temporary, reversible, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
Medway Estuary and Marshes	International	Potential impacts within the European/Ramsar S Damage to habitats within designated area used by the cited bird species from changes to water and/or sediment quality (either from	ite itself: Temporary, reversible, minor magnitude and extent.	Not significant ²⁹

Although baseline ambient noise levels are high surrounding the site (50db-71db) and average within (44-48db), general construction activities at Gate 1 are expected to increase noise levels by 16-20db at NSR 22, although the effect is attenuated further from the source, with negligible increases at the SW edge of Black Duck Marsh and along the estuary foreshore.

Given the distance between the Project Site and the mouth of the Medway estuary (33.4km) (i.e. the receptor pathway for pollutants), no likely significant effects are anticipated on the Medway Estuary and Marshes SPA/Ramsar/SSSI. See Shadow Habitat Regulations Assessment (ES Appendix 12.3), Stage 1 Screening Matrix (Annex 4)



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
SPA/Ramsar/		surface or groundwater discharges from the		
SSSI ²⁸		Project Site including construction waste and		
		pollutants; or from disruption of contaminated		
		Thames sediments), with potential associated		
		knock-on risk of bioaccumulation.		
		Damage to habitats within the designated area	Temporary, reversible, minor	Not significant
		used by cited bird species from changes in air	magnitude and extent.	
		quality, including from dust, construction		
		waste and pollutants, and exhaust emissions.		
		Damage to habitats used by cited bird species	Temporary, reversible, minor	Not significant
		within the designated area from introduction	magnitude and extent.	
		or proliferation of Invasive Non-Native Species		
		(INNS) INNS, as a result of construction works		
		(and in particular shipping/transport of		
		materials).		
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Not significant
		giving rise to displacement, behavioural	magnitude and extent.	
		changes or physiological stress to cited bird		
		species within the designated area.		
		Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ^{30 31}
		with increased shipping traffic and construction	magnitude and extent.	

Medway Estuary and Marshes SSSI is a component SSSI to the Medway Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar

In regards to noise, peak or mean (i.e. 24hr) noise in excess of 55dB is not predicted to be experienced outside of the DCO Order Limits for most construction or operational activities, with the exception of construction-phase jetty piling and dredging and pavement construction, for which values are not yet known. However,



The distance between the Project Site and the nearest part of the Medway Estuary and Marshes SPA is approximately 13.1km. At these distances significant impacts from lighting are not anticipated.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		activities, giving rise to displacement,		
		behavioural changes or physiological stress to		
		cited bird species within the designated area.		
		Potential impacts solely on functionally linked res	sources (outside of the European/I	Ramsar Site):
		Direct loss of/damage to functionally linked	Permanent, moderate	Significant negative
		land on the Kent Project Site, potentially used	magnitude and extent.	at district
		by cited bird species, totalling a net loss of		levelSignificant
		15.49ha, including 14.55ha of coastal and		negative at District
		floodplain grazing marsh (Botany Marsh West)		level
		and 0.94ha of reedbeds (Black Duck Marsh) to		
		accommodate construction of the Leisure Core.		
		Damage to functionally linked habitats	Temporary, moderate	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at district
		changes to hydrology (as a result of drainage		levelSignificant
		and infill of the CTRL wetland, swamp/wetland		negative at District
		habitat along the east/south east of Black Duck		<u>level</u>
		Marsh and other low-lying seasonally		
		inundated areas); and changes to water and		
		/or sediment quality (either from surface or		
		groundwater discharges from Project Site		
		including construction waste and pollutants;		
		or from disruption of contaminated Thames		
		sediments), with potential associated knock-on		
		risk of bioaccumulation.		

based on the degree of geographical separation between the Project Site and the Medway Estuary and Marshes SPA/Ramsar/SSSI significant effects are not anticipated at this distance



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Damage to functionally linked habitats	Temporary, minor magnitude	Significant negative
		potentially used by cited bird species from	and extent.	at district
		changes in air quality including from dust,		levelSignificant
		construction waste and pollutants ³² .		negative at District
				<u>level</u> ³³
		Damage to functionally linked habitats	Temporary, reversible, minor	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at district
		introduction or proliferation of Invasive Non-		level Significant
		Native Species (INNS)INNS.		negative at District
				<u>level</u>
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Significant negative
		giving rise to displacement, behavioural	magnitude and extent.	at district
		changes or physiological stress to cited bird		levelSignificant
		species potentially using functionally linked		negative at District
		habitats.		<u>level</u>
		Disturbance (noise and lighting) associated	Temporary, reversible,	Significant negative
		with increased shipping traffic and construction	moderate magnitude and	at district
		activities, giving rise to displacement,	extent.	levelSignificant
		behavioural changes or physiological stress to		negative at District
		cited bird species potentially using functionally		<u>level</u> ³⁴
		linked habitats.		

No retained functionally linked habitats lie within 200m of an existing road. Effects due to increased construction traffic have therefore been screened out of this assessment.

Although baseline ambient noise levels are high surrounding the site (50db-71db) and average within (44-48db), general construction activities at Gate 1 are expected to increase noise levels by 16-20db at NSR 22, although the effect is attenuated further from the source, with negligible increases at the SW edge of Black Duck Marsh and along the estuary foreshore.



³³ Given the proximity of construction work to retained functionally linked habitats (<50m in some cases, especially around Black Duck Marsh and along the estuary front), significant dust deposition is anticipated.

Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
reature	Receptor	Disturbance (human movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during construction.	Temporary, reversible, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
North Downs Woodland SAC, and the component SSSI Wouldham to Detling Escarpment SSSI ³⁵	International	Construction traffic generated air quality impacts – due to potential increases in vehicle movements along the A229 and A259 trunk roads which lie within 200m of areas of North Downs Woodland SAC and link the M2/A2 corridor and the M20.	Temporary, minor magnitude and extent.	Not significant ³⁶
Swanscombe Peninsula SSSI	<u>National</u>	Overall reduction in SSSI area, and fragmentation by the Resort (109.6ha lost, i.e. 42.2%) Damage to habitats within the designated area from changes in air quality, including from dust, construction waste and pollutants, and exhaust emissions.	Permanent, major magnitude and extent Temporary, reversible, major magnitude and extent. Likely to be major magnitude due to proximity of construction	Significant negative at the national level Significant negative at National level

As set out with Air Quality Assessment (ES Chapter 16 Air Quality; Document Reference 6.1.16), for construction the SAC is not within 350m of the site boundary and/or within 50m of the kerb of a road used by construction traffic and any impacts can be ruled as insignificant.



Wouldham to Detling Escarpment SSSI is a component SSSI to the North Downs Woodland SAC, with overlapping reasons for designation. Potential impacts to the SAC as described are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar

Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Special feature of interest: pPopulations of rai	re plants ³⁷ s:	
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality	Temporary, moderate magnitude and extent.	Significant negative at national levelSignificant negative at National level
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, major magnitude and extent.	Significant negative at national levelSignificant negative at National level
		<u>Damage to supporting habitats from</u> introduction or proliferation of <u>Invasive Non-Native Species (INNS)INNS.</u>	Temporary, reversible, minor magnitude and extent.	Significant negative at national levelSignificant negative at National level
		Habitat fragmentation.	Permanent, moderate magnitude and extent.	Significant negative at national levelSignificant negative at National level

Includes only those rare plant populations listed as a reason for notification of the Swanscombe Peninsula SSSI, including divided sedge (*Carex divisa*), yellow vetchling (*Lathyrus aphaca*), slender hare's-ear (*Bupleurum tenuissimum*), Bithynian vetch (*Vicia bithynica*) and round-leaved wintergreen (*Pyrola rotundifolia subsp. maritima*). A separate assessment of potentially significant effects upon 'non-SSSI' rare plants is provided under the 'rare plants' IEF.



Ecological Feature	Importance of	Nature of Impact	Impact Magnitude	Significance
reature	Receptor	Disturbance (lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at national levelSignificant negative at National level
Darenth Woods SSSI	National	Direct loss of habitat – not applicable. The Proposed Development will result in no direct land take from within the SSSI. Whilst the DCO boundary previously included the ancient woodland parcels between the A2(T) and A296 slip roads, these have since been removed and will not be affected by The London Resort proposals, which will be limited to minor highway works in this area such as road markings and signage. As a point of clarity, a small number of individual trees will be lost on the edge of 'The Thrift' Ancient Woodland on the southern boundary of the A2(T) and the slip road as a result of the separate A2(T) Bean and Ebbsfleet Junction improvement works, which were permitted in May of this year 2020. Damage to habitats within the designated area from changes in air quality, including from	Temporary, reversible, minor magnitude and extent.	Significant at the district level
		dust, construction waste and pollutants, and exhaust emissions.		<u>level</u>



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ³⁸
		with increased construction traffic along A2(T).	magnitude and extent.	
Shorne and	National	Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ³⁹
Ashenbank		with increased construction traffic along A2(T).	magnitude and extent.	
Woods SSSI				
nner Thames	National	Potential impacts within the SSSI itself:		
Marshes SSSI		Damage to habitats within designated area	Temporary, reversible, minor	Not significant
		used by cited bird species from changes to	magnitude and extent.	
		water and/or sediment quality (either from		
		surface or groundwater discharges from the		
		Project Site including construction waste and		
		pollutants; or from disruption of contaminated		
		Thames sediments), with potential associated		
		knock-on risk of bioaccumulation.		
		Damage to habitats within the designated area	Temporary, reversible, minor	Not significant
		used by cited bird species from changes in air	magnitude and extent.	
		quality, including from dust, construction		
		waste and pollutants, and exhaust emissions.		
		Damage to habitats used by cited bird species	Temporary, reversible, minor	Not significant
		within the designated area from introduction	magnitude and extent.	
		or proliferation of INNS, as a result of		

Construction traffic is predicted to create an additional 0-0.1db of road noise along the A2(T), as modelled and shown within the Construction Noise Assessment (Document Reference Part 6.2, Chapter 15.3). General construction noise levels are not expected to change significantly at NSR 5, which is significantly closer to construction activities (<500m) and situated between the noise source and SSSI.



Construction traffic is predicted to create an additional 0-0.1db of road noise along the A2(T), as modelled and shown within the Construction Noise Assessment (Document Reference Part 6.2, Chapter 15.3). General construction noise levels are not expected to change significantly at NSR 5, which is significantly closer to construction activities (<500m) and situated between the noise source and SSSI.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		construction works (and in particular		
		shipping/transport of materials).		
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Not significant
		giving rise to displacement, behavioural	magnitude and extent.	
		changes or physiological stress to cited bird		
		species within the designated area.		
		Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant
		with increased shipping traffic and construction	magnitude and extent.	
		activities, giving rise to displacement,		
		behavioural changes or physiological stress to		
		cited bird species within the designated area.		
		Potential impacts solely on functionally linked res	sources (outside of the SSSI):	
		Direct loss of/damage to functionally linked	Permanent, moderate	Significant negative
		habitats on the Kent Project Site, potentially	magnitude and extent.	at District level
		used by bird species, totalling a net loss of		
		15.49ha, including 14.55ha of coastal and		
		floodplain grazing marsh (Botany Marsh West)		
		and 0.94ha of reedbeds (Black Duck Marsh) to		
		accommodate construction of the Leisure Core.		
		Damage to functionally linked habitats	Temporary, moderate	Significant negative
		potentially used by bird species from changes	magnitude and extent	at District level
		to hydrology (as a result of drainage and infill		
		of the CTRL wetland, swamp/wetland habitat		
		along the east/south east of Black Duck Marsh		
		and other low-lying seasonally inundated		
		areas); and changes to water and /or sediment		
		quality (either from surface or groundwater		
		discharges from Project Site including		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		construction waste and pollutants; or from		
		disruption of contaminated Thames		
		sediments), with potential associated knock-on		
		risk of bioaccumulation.		
		Damage to functionally linked habitats	Temporary, minor magnitude	Significant negative
		potentially used by cited bird species from	and extent.	at District level ⁴¹
		changes in air quality including from dust,		
		construction waste and pollutants ⁴⁰ .		
		Damage to functionally linked habitats	Temporary, reversible, minor	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at District level
		introduction or proliferation of INNS.		
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Significant negative
		giving rise to displacement, behavioural	magnitude and extent.	at District level
		changes or physiological stress to cited bird		
		species potentially using functionally linked		
		habitats.		
		Disturbance (noise and lighting) associated	Temporary, reversible,	Significant negative
		with increased shipping traffic and construction	moderate magnitude and	at District level ⁴²
		activities, giving rise to displacement,	extent.	
		behavioural changes or physiological stress to		

It is likely that the Project Site is functionally linked to the Inner Thames Marshes SSSI. Although baseline ambient noise levels are high surrounding the site (50db-71db) and average within (44-48db), general construction activities at Gate 1 are expected to increase noise levels by 16-20db at NSR 22, although the effect is attenuated further from the source, with negligible increases at the SW edge of Black Duck Marsh and along the estuary foreshore.



No retained functionally linked habitats lie within 200m of an existing road. Effects due to increased construction traffic have therefore been screened out of this assessment.

⁴¹ Given the proximity of construction work to retained functionally linked habitats (<50m in some cases, especially around Black Duck Marsh and along the estuary front), significant dust deposition is anticipated.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		cited species potentially using functionally		
		linked habitats.		
		Disturbance (human movement or activity), to	Temporary, reversible, minor	Significant negative
		cited bird species potentially using functionally	magnitude and extent.	at District level
		linked habitats, as a result of increased human		
		presence on the Project Site during		
		construction.		
West	<u>National</u>	Potential impacts within the SSSI itself:		
<u>Thurrock</u>		Damage to habitats within designated area	Temporary, reversible, minor	Not significant
Lagoon and		used by cited bird species from changes to	magnitude and extent.	
Marshes SSSI		water and/or sediment quality (either from		
		surface or groundwater discharges from the		
		Project Site including construction waste and		
		pollutants; or from disruption of contaminated		
		Thames sediments), with potential associated		
		knock-on risk of bioaccumulation.		
		Damage to habitats within the designated area	Temporary, reversible, minor	Not significant
		used by cited bird species from changes in air	magnitude and extent.	
		quality, including from dust, construction		
		waste and pollutants, and exhaust emissions		
		Damage to habitats used by cited bird species	Temporary, reversible, minor	Not significant
		within the designated area from introduction	magnitude and extent.	
		or proliferation of INNS, as a result of		
		construction works (and in particular		
		shipping/transport of materials).		
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Not significant
		giving rise to displacement, behavioural	magnitude and extent.	



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		changes or physiological stress to cited bird		
		species within the designated area.		
		Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ⁴³
		with increased shipping traffic and construction	magnitude and extent.	
		activities, giving rise to displacement,		
		behavioural changes or physiological stress to		
		cited bird species within the designated area.		
		Potential impacts solely on functionally linked res	sources (outside of the SSSI):	
		Direct loss of/damage to functionally linked	Permanent, moderate	Significant negative
		habitats on the Kent Project Site, potentially	magnitude and extent.	at District level
		used by cited bird species, totalling a net loss		
		of 15.49ha, including 14.55ha of coastal and		
		floodplain grazing marsh (Botany Marsh West)		
		and 0.94ha of reedbeds (Black Duck Marsh) to		
		accommodate construction of the Leisure Core.		
		Damage to functionally linked habitats	Temporary, moderate	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at District level
		changes to hydrology (as a result of drainage		
		and infill of the CTRL wetland, swamp/wetland		
		habitat along the east/southeast of Black		
		Duck Marsh and other low-lying seasonally		
		inundated areas); and changes to water and		
		/or sediment quality (either from surface or		
		groundwater discharges from Project Site		
		including construction waste and pollutants;		

Noise levels further from the source into the Thames Estuary and across to Thurrock Lagoon and Marshes SSSI will attenuate are expected to be as low as 35-40db (worst case 45db) on the north bank of the Thames.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		or from disruption of contaminated Thames		
		sediments), with potential associated knock-on		
		risk of bioaccumulation.		
		Damage to functionally linked habitats	Temporary, minor magnitude	Significant negative
		potentially used by cited bird species from	and extent.	at District level ⁴⁵
		changes in air quality including from dust,		
		construction waste and pollutants ⁴⁴ .		
		Damage to functionally linked habitats	Temporary, reversible, minor	Significant negative
		potentially used by cited bird species from	magnitude and extent.	at District level
		introduction or proliferation of INNS.		
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Significant negative
		giving rise to displacement, behavioural	magnitude and extent.	at District level
		changes or physiological stress to cited bird		
		species potentially using functionally linked		
		habitats.		
		Disturbance (noise and lighting) associated	Temporary, reversible,	Significant negative
		with increased shipping traffic and construction	moderate magnitude and	at District level ⁴⁶
		activities, giving rise to displacement,	extent.	
		behavioural changes or physiological stress to		
		cited species potentially using functionally		
		linked habitats.		

No retained functionally linked habitats lie within 200m of an existing road. Effects due to increased construction traffic have therefore been screened out of this assessment.

It is likely that the Project Site is functionally linked to the Inner Thames Marshes SSSI. Although baseline ambient noise levels are high surrounding the site (50db-71db) and average within (44-48db), general construction activities at Gate 1 are expected to increase noise levels by 16-20db at NSR 22, although the effect is attenuated further from the source, with negligible increases at the SW edge of Black Duck Marsh and along the estuary foreshore



⁴⁵ Given the proximity of construction work to retained functionally linked habitats (<50m in some cases, especially around Black Duck Marsh and along the estuary front), significant dust deposition is anticipated.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (human movement or activity), to	Temporary, reversible, minor	Significant negative
		cited bird species potentially using functionally	magnitude and extent.	at District level
		linked habitats, as a result of increased human		
		presence on the Project Site during		
		construction.		
Cobham	<u>National</u>	Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ⁴⁷
Woods SSSI		with increased construction traffic along A2(T).	magnitude and extent.	
<u>Great</u>	<u>National</u>	Disturbance (noise and lighting) associated	Temporary, reversible, minor	Not significant ⁴⁸
<u>Crabbles</u>		with increased construction traffic along A2(T).	magnitude and extent.	
Wood SSSI				
Botany	County	Damage to habitats from changes in air	Temporary, major magnitude	Significant at local
Marshes LWS		quality, including from dust, construction	and extent.	levelSignificant at
		waste and pollutants.		Local level
		Damage to habitats from changes to	Permanent, major magnitude	Significant at local
		hydrological regime (through destruction of	and extent.	levelSignificant at
		adjoining wetland/grazing marsh), and changes		Local level
		in water quality/quantity.		
		Damage to habitats from adjacent construction	Temporary, moderate	Significant at local
		works and deposition of construction materials.	magnitude and extent.	levelSignificant at
				Local level

⁴⁸ Construction traffic is predicted to create an additional 0-0.1db of road noise along the A2(T), as modelled and shown within the Construction Noise Assessment (Document Reference Part 6.2, Chapter 15.3). General construction noise levels are not expected to change significantly at NSR 5, which is significantly closer to construction activities (<500m) and situated between the noise source and SSSI.



⁴⁷ Construction traffic is predicted to create an additional 0-0.1db of road noise along the A2(T), as modelled and shown within the Construction Noise Assessment (Document Reference Part 6.2, Chapter 15.3). General construction noise levels are not expected to change significantly at NSR 5, which is significantly closer to construction activities (<500m) and situated between the noise source and SSSI.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise and lighting) associated	Temporary, minor magnitude	Significant at local
		with construction activities, giving rise to	and extent.	levelSignificant at
		displacement, behavioural changes or		Local level ⁴⁹
		physiological stress to wildlife species within		
		the designated area.		
Ebbsfleet	County	Direct loss of/damage to habitat through	Permanent, moderate	Significant at district
Marshes,		Station Quarter South .	magnitude and extent.	level District level
Northfleet		Damage to habitats from changes in air	Temporary, moderate	Significant at local
LWS		quality, including from dust, construction	magnitude and extent.	levelSignificant at
		waste and pollutants.		Local level
		Damage to habitats from changes to	Temporary, moderate	Significant at local
		hydrological regime (through destruction of	magnitude and extent.	levelSignificant at
		habitat within flood zone of River Ebbsfleet),		Local level
		and changes in water quality/quantity.		
		Damage to habitats from adjacent construction	Temporary, moderate	Significant at local
		works and deposition of construction materials.	magnitude and extent.	levelSignificant at
				Local level
		Disturbance (noise and lighting) associated	Temporary, moderate	Significant at local
		with construction activities, giving rise to	magnitude and extent.	levelSignificant at
		displacement, behavioural changes or		Local level ⁵⁰
		physiological stress to wildlife species within		
		the designated area.		

⁵⁰ Noise levels not expected to exceed 59db outside of construction area (within retained habitat), which is not considered significant over the baseline noise levels, which are already significant.



⁴⁹ Noise levels are expected to reach between 54db and 64db at Botany Marshes LWS, however, minor adverse effects are anticipated because baseline noise and vibration levels from HGV and industrial operations within the Manor Way business park and CEMEX concrete works facility are already significant.

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Alkerden Lane Pit LWS	Receptor County	Direct loss of habitat – not applicable. The Proposed Development will result in no direct land take from within the LWS.	N/A	N/A
		Loss of part of plant and invertebrate meta- population through loss of habitat in the Kent Project Site.	Permanent, minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from changes in air quality, including from dust, construction waste and pollutants.	Temporary, major magnitude and extent.	Significant at local levelSignificant at Local level
Disused Hospital Grounds, Mabledon LWS	County	Damage to habitats from changes in air quality, including from dust, construction waste and pollutants.	Temporary, major magnitude and extent.	Not significant ⁵¹
Tilbury Marshes LWS	County	Direct loss of habitat – not applicable. The Proposed Development will result in no direct land take from within the LWS.	N/A	N/A
		Direct loss of/damage to functionally linked habitats on the Kent Project Site, potentially used by bird species, totalling a net loss of 15.49ha, including 14.55ha of coastal and floodplain grazing marsh (Botany Marsh West) and 0.94ha of reedbeds (Black Duck Marsh) to accommodate construction of the Leisure Core.	Permanent, moderate magnitude and extent.	

In accordance with IAQM guidance, an assessment of construction dust will normally be required where there is an 'ecological receptor' within 50m of the boundary of the site; or 50m of the route(s) used by construction vehicles on the public highway, up to 500m from the site entrance. The Disused Hospital Grounds LWS is located 1.3km south-west of the Kent Project Site. Chapter 16: Air Quality (Document reference: 6.1.16) confirms that "...80-90% of construction materials will be transported by river." The chapter confirms that "The impact from construction traffic emissions is predicted to be negligible at all modelled receptor locations"



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (from shipping/ferry movements)	Temporary, reversible, minor	Significant negative
		giving rise to displacement, behavioural	magnitude and extent.	at district
		changes or physiological stress to cited bird		levelSignificant
		species potentially using functionally linked		negative at District
		habitats.		<u>level</u>
		Disturbance (noise and lighting) associated	Temporary, reversible,	Significant negative
		with increased shipping traffic and construction	moderate magnitude and	at district
		activities, giving rise to displacement,	extent.	level Significant
		behavioural changes or physiological stress to		negative at District
		bird species potentially using functionally		<u>level</u> ⁵²
		linked habitats.		
		Disturbance (human movement or activity), to	Temporary, reversible, minor	Significant negative
		bird species potentially using functionally	magnitude and extent.	at district
		linked habitats, as a result of increased human		level Significant
		presence on the Project Site during		negative at District
		construction.		<u>level</u>
Habitats/Flora	1			
Populations	National	N.B. Assessment of potential effects upon	Permanent, major magnitude	Significant at
of nationally		scarce plants included under Swanscombe	and extent.	<mark>regional level</mark>
scarce plants		Peninsula SSSI Direct loss of individual plants,		
		and part of plant meta-population through		
		loss of habitat in the Kent Project Site.		
Broad leaved	Local	Direct loss of habitat from woodland south of	Permanent, major magnitude	Significant at local
Semi Natural		Black Duck Marsh – 3.84ha lost from a total of	and extent.	levelSignificant at
Woodland		21.37ha (18% of the existing total).		Local level

Noise levels further from the source into the Thames Estuary and across to Tilbury during general construction will attenuate are expected to be as low as 35-40db (worst case 45db) on the north bank of the Thames. Additional noise from construction work at Tilbury is expected to be negligible as the existing car park and buildings will be used, with only the addition of a floating jetty.



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Damage to habitat from construction dust,	Temporary, minor magnitude	Significant at local
		waste and pollutants.	and extent.	levelSignificant at
				<u>Local level</u>
Ancient	County	Damage to habitats within the designated area	Temporary, reversible, minor	Significant at the
woodland		from changes in air quality, including from	magnitude and extent.	district District level
		dust, construction waste and pollutants, and		
		exhaust emissions.		
Scrub	Local	Direct loss of habitat, a total of 4534.65ha	Permanent, major magnitude	Significant at local
		79ha lost (4641.9971% of the existing total)	and extent.	levelSignificant at
		including: 38.70ha of dense/continuous scrub		Local level
		(46% of the existing total), and 7.17ha of		
		scattered scrub (taken as 50% of the total area		
		measured as grassland/scrub mosaic, 57.08% of		
		the existing total). 4.10ha of lost scrub is also		
		included under the OMHPDL grouping and		
		0.57ha under Floodplain Wetland Mosaic,		
		within the biodiversity net gain calculations		
		(Document reference 6.2, Appendix 12.2).		
		Damage to habitats from adjacent construction	Temporary, minor magnitude	Significant at local
		works and deposition of construction materials	and extent.	levelSignificant at
		in Root Protection Area (RPA).		Local level
		Damage to habitat from construction dust,	Temporary, minor magnitude	Significant at local
		waste and pollutants.	and extent.	levelSignificant at
				Local level
		Damage to habitats from introduction or	Temporary, reversible, minor	Significant negative
		proliferation of Invasive Non Native Species	magnitude and extent.	at district
		(INNS)INNS		level Significant



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
				negative at District level
Semi- improved grassland	Local to District	Direct loss of habitat, 4236.19ha lost (4844.4279% of the existing total)-28.23ha of lost semi-improved grassland is also included under the OMHPDL grouping and 1.54ha under Floodplain Wetland Mosaic, within the biodiversity net gain calculations (Document reference 6.2, Appendix 12.2).	Permanent, major magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from adjacent construction works and deposition of construction materials.	Temporary, minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitat from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
Coastal/flood plain grazing marsh	District	Direct loss of habitat, 1113.99ha-74ha lost ((95.08100)% of the existing total). All CFGM is also included under the Floodplain Wetland Mosaic grouping, within the biodiversity net gain calculations biodiversity impact calculations (Document reference 6.2, Appendix 12.2).	Permanent, major magnitude and extent.	Significant at district level District level



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
Open mosaic on previously developed land	District	Direct loss of habitat, 5539.13ha-83ha lost (7663.72% of the existing total), includes a range of individual habitat components including areas of bare ground, ephemeral vegetation, open grassland and patches of scrub.	Permanent, major magnitude and extent.	Significant at district level District level
		Damage to habitats from adjacent construction works and deposition of construction materials.	Temporary, minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitat from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant at local level Significant at Local level
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
Waterbodies (ponds, standing water and ditch network)	District	Direct loss of habitat, 2.30ha of waterbodies lost (50.79% of the existing total) and 4.7km of ditches (50.98% of the existing total). 0.22ha of lost waterbodies are also included under the Floodplain Wetland Mosaic grouping, within the biodiversity net gain calculations biodiversity impact calculations (Document reference 6.2, Appendix 12.2).	Permanent, major magnitude and extent.	Significant at district level District level
		Damage to habitats from changes to	Temporary-	Significant at local
		hydrological regime (through destruction of	Temporary permanent, reversible-irreversible,	level Significant at Local level



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		surrounding habitats), and changes in water quality/quantity.	moderate magnitude and extent.	
		Damage to habitat from construction dust, waste and pollutants.	Temporary, <u>reversible</u> , minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
Swamp (reedbed)	County	Direct loss of habitat from 13.02ha-62ha lost (4241.9975% of total area). 2.84ha of reedbed is also included under the OMHPDL grouping and 0.88ha under Floodplain Wetland Mosaic, within the biodiversity net gain calculations biodiversity impact calculations (Document reference 6.2, Appendix 12.2).	Permanent, major magnitude and extent.	Significant at county level
		Damage to habitats from adjacent construction works and deposition of construction materials.	Temporary, minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from changes to hydrological regime (through destruction of surrounding habitats), and changes in water quality/quantity.	Temporary, moderate magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitat from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant at local levelSignificant at Local level



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District
River	Local	Direct loss of connected floodplain habitat	Permanent, moderate	level Significant at local
Ebbsfleet	Local	within Ebbsfleet Valley, such as waterbodies and semi-improved grassland.	magnitude and extent.	levelSignificant at Local level
		Damage to habitat from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from changes to hydrological regime (through destruction of surrounding habitats), and changes in water quality/quantity.	Temporary, moderate magnitude and extent.	Significant at local levelSignificant at Local level
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
Species				1 <u></u>
Wintering Waterfowl and Wading Bird Assemblage	International	N.B. Assessment of potential habitat loss and disturbance effects upon wintering waterfowl and wading bird assemblage included under SPA/Ramsar SitesDirect killing, injury or harm to individuals — during vegetation clearance.	Permanent, major magnitude and extent	Significant negative at district level
		<u>Direct killing, injury or harm to individuals –</u> <u>during vegetation clearance.</u>	Permanent, major magnitude and extent.	Significant negative at District level



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
Wintering terrestrial bird assemblage	County	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Direct habitat loss – loss of suitable habitat such as scrub, grassland, reedbed and woodland, as detailed above.	Permanent, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant negative at district level Significant negative at District level
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS) — as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, minor magnitude and extent.	Significant negative at district level Significant negative at District level
		Habitat fragmentation, loss of flight paths and dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at district levelSignificant



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
				negative at District level
		Disturbance (from shipping/ferry movements) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Disturbance (noise and lighting) – as described above in relation to impacts on habitats functionally linked to SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Disturbance (human movement or activity) – as described above in relation to impacts on habitats functionally linked to SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
Breeding Bird Assemblage	Regional Natio nal	N.B. Assessment of potential effects upon breeding bird assemblage included under Swanscombe Peninsula SSSIDirect killing, injury or harm to individuals — during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at district level
Bat Assemblage	District_ County	Direct killing, injury or harm to individuals – during demolition of buildings with bat roost potential and trees with bat roost potential.	Permanent, major magnitude and extent.	Significant negative at local levelSignificant negative at Local level



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Direct loss of potential roosting habitat –	Permanent, major magnitude	Significant negative
		including trees, and buildings and structures	and extent.	at local
		with bat roost potential. <u>Tunnels TU/016</u>		level Significant
		(negligible/low potential with possible		negative at Local-
		pipistrelle hibernation roost) and TU/018 (low		<u>District level</u>
		potential and winter bat roost uncertain) will		
		be used for access between the transport		
		interchange and staff accommodation.		
		Direct loss of foraging, commuting and	Permanent, moderate	Significant negative
		dispersal habitats - loss of suitable foraging	magnitude and extent.	at local
		habitat such as scrub, grassland, reedbed and		levelSignificant
		woodland, as detailed above, including		negative at Local
		potential winter foraging habitat.		<u>level</u>
		Damage to habitats from changes to	Temporary, moderate	Significant negative
		hydrological regime and changes to water	magnitude and extent.	at local
		and/or sediment quality – as described above		levelSignificant
		in relation to impacts on habitats.		negative at Local
				<u>level</u>
		Damage to supporting habitats from	Temporary, minor magnitude	Significant negative
		construction dust, waste and pollutants.	and extent.	at local
				levelSignificant
				negative at Local
				<u>level</u>
		Damage to supporting habitats from	Temporary, reversible, minor	Significant negative
		introduction or proliferation of Invasive Non-	magnitude and extent.	at local
		Native Species (INNS)INNS.		levelSignificant
				negative at Local
				<u>level</u>



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
reature	песерие	Habitat fragmentation, loss of flight paths and dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at local
				levelSignificant negative at Local level
		Disturbance (noise and lighting) – including on retained roosts, foraging areas and commuting routes.	Temporary, reversible, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
Dormouse	District	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Direct habitat loss – loss of suitable habitat such as scrub and woodland, as detailed above, and within the Dormouse Mitigation Strategy within the Ecological Mitigation and Management Framework (Document Reference 6.2.12.3).	Permanent, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	Temporary, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant negative at district



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
				levelSignificant negative at District
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	level Significant negative at district level Significant negative at District level
		Habitat fragmentation and loss of dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at district levelSignificant negative at District level
Otter	Local	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Direct habitat loss – loss of suitable habitat such as reedbed and waterbodies, as detailed above.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	Temporary, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to supporting habitats from construction dust, waste and pollutants	Temporary, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Habitat fragmentation and loss of dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
Water Vole	Local to district	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at local levelSignificant



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
				negative at Local level
		Direct habitat loss – including 6.1km of suitable ditches and bankside habitat and reedbed (as detailed above).	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	Temporary, moderate magnitude and extent.	level Significant negative at local level Significant negative at Local
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	level Significant negative at local level Significant
		Damage to supporting habitats from introduction or proliferation of Invasive Non-	Temporary, reversible, minor magnitude and extent.	negative at Local level Significant negative at local
		Native Species (INNS)INNS.		levelSignificant negative at Local level
		Habitat fragmentation and loss of dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
Harvest Mouse	Local	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Direct habitat loss – loss of suitable habitat such as reedbed, semi-improved grassland, scrub and tall ruderal, as detailed above.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	Temporary, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
				negative at Local level
		Habitat fragmentation and loss of dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at local
				levelSignificant negative at Local level
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at local levelSignificant
				negative at Local level
Amphibian assemblage	Local to district	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Direct habitat loss – loss of suitable habitat such as reedbed and waterbodies, as detailed above.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	Temporary, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Habitat fragmentation and loss of dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
Reptile assemblage	District	Direct killing, injury or harm to individuals – during vegetation clearance.	Permanent, major magnitude and extent	Significant negative at local levelSignificant negative at Local level
		Direct habitat loss – loss of suitable habitat such as semi-improved grassland, reedbed and tall ruderal, as detailed above.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
				negative at Local level
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above	Temporary, moderate magnitude and extent.	Significant negative at local levelSignificant
		in relation to impacts on habitats.		negative at Local level
		Damage to supporting habitats from construction dust, waste and pollutants.	Temporary, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Habitat fragmentation and loss of dispersal routes.	Permanent, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	Temporary, reversible, moderate magnitude and extent.	Significant negative at local levelSignificant negative at Local level



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
Invertebrate	National	N.B. Assessment of potential effects upon	Permanent, major magnitude	Significant negative
assemblage		invertebrate assemblages included under	and extent.	<mark>at regional level</mark>
		Swanscombe Peninsula SSSI Direct habitat loss		
		- loss of suitable habitat, particularly Open		
		Mosaic Habitat on Previously Developed Land,		
		as detailed above.		



Operational Impacts and Effects

- <u>12.21812.243</u> Without mitigation, the potential effects of the proposed development on the IEFs identified above during the operational phase could include:
 - Habitat fragmentation/loss of flight paths/dispersal routes;
 - Increased lighting, noise and traffic leading to disturbance of species within retained and newly created habitats;
 - Increased collision risk to birds and bats;
 - Hydrological effects, including changes to water quality/quantity;
 - Damage or degradation to habitats and disturbance of wildlife through increased recreational pressure and trampling; and
 - Potential positive effects/benefits through provision of habitats with greater biodiversity value than those currently present, and implementation of appropriate management of the retained and created habitats to maximise their biodiversity potential.
- <u>12.219</u>12.244 Further details, including which IEFs these potential effects relate to, are provided within **Table 12.11**.
- <u>12.22012.245</u> In addition, further information on potential effects upon European Sites is provided within Appendix 12.4: *Shadow Habitat Regulations Assessment* (Document Reference 6.2.12.4).
- <u>12.221</u>12.246 An impact assessment specific to each species/species assemblage/taxa is also provided within the individual mitigation strategies enclosed within Appendix 12.3: *Ecological Mitigation and Management Framework* (Document Reference 6.2.12.3).
- 12.22212.247 The assessment of potential effects during operation has been undertaken using information derived from other chapters within the Environmental Statement, notably Chapter 15: Noise and Vibration (Document reference 6.1.15), Chapter 16: Air Quality (Document reference 6.1.16), and Chapter 17: Water Resources and flood risk (Document reference 6.1.17). An assessment of the potential effects of climate change on ecological receptors, using Chapter 20: Greenhouse gas and climate change (Document reference: 6.1.20) is provided towards the end of this chapter.

Air Quality

<u>12.248</u> To inform the assessment of potential ecological effects arising from changes in air quality, as predicted within Chapter 16: Air Quality (Document reference 6.1.16), transect data provided in Figures 16.20 – 16.24 have been used to calculate the extent of nitrogen



deposition at those ecological receptors included in the air quality assessment based on the area predicted to exceed 1% of the minimum nutrient nitrogen critical load and nitrous oxide critical level). The results in respect of Darenth Wood SSSI are presented in Tables_-12.12_and 12.13, and in respect of local sites (ancient woodland and LWSs) in Tables_-12.14_and Table-12.15-3.



Table 12.11: Operational Effects on Important Ecological Features without Mitigation-

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor	•		
Designations	•			
Thames	International	Potential impacts within the European/Ramsar S	ite itself:	
Estuary and		Damage to habitats within designated area	Temporary <u>-permanent</u> ,	Not significant
Marshes		used by cited bird species from changes to	reversible <u>-irreversible</u> , minor	
SPA/Ramsar,		water and/or sediment quality (from surface	magnitude and extent.	
and the		or groundwater discharges from the Project		
component		Site including operational waste and		
SSSIs		pollutants).		
including		Damage to habitats within the designated area	Temporary Permanent,	Not significant ⁵⁴
South		used by cited bird species and/or changes in	<u>irreversible</u> reversible, minor	
Thames		the food supply of cited species from changes	magnitude and extent.	
Estuary and		in air quality, including from exhaust		
Marshes SSSI		emissions.		
and Mucking		Disturbance (from increased ferry	Permanent Temporary,	Not significant
Flats and		movements) associated with the new Thames	<u>ir</u> reversible, minor magnitude	
Marshes		Clipper river bus service and operational	and extent.	
SSSI ⁵³		deliveries, potentially giving rise to		
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		

⁵³ South Thames Estuary and Marshes SSSI and Mucking Flats and Marshes SSSI are component SSSIs to the Thames Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also 54 With reference to the Air Quality Assessment in Chapter 16: Air Quality (Document reference 6.2.16), which is based on a traffic model which incorporates traffic data from relevant cumulative schemes, impacts upon the Thames Estuary and Marshes SPA/Ramsar/SSSI can be ruled insignificant. The designated site is located >200m from the roadside of roads predicted to experience an increase of >1000 AADT. The designated site is within 10km of the energy centre point source, however, the predicted



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor		 	
		Disturbance (noise and lighting) from the	<u>Permanent</u> Temporary ,	Not significant 55
		operation of the resort rides and attractions	<u>irreversible</u> reversible, minor	
		and entertainment facilities, giving rise to	magnitude and extent.	
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		
		Recreational disturbance of flora and fauna	Temporary Permanent,	Significant negative
		within designated area as a result of increased	reversible-	<u>at National level</u> No
		residential population (due to residential	<u>irreversible</u> reversible, minor	<mark>significant</mark>
		accommodation for staff).	magnitude and extent.	
		Potential impacts solely on functionally linked res	sources (outside of the European/I	Ramsar Site):
		Damage to functionally linked habitats	Temporary-	Significant negative
		potentially used by cited bird species from	permanent Temporary,	at district
		changes to hydrology; and changes to water	reversible-irreversible,	levelSignificant
		and /or sediment quality (from surface or	moderate magnitude and	negative at District
		groundwater discharges from the Project Site	extent.	<u>level</u>
		including operational waste and pollutants).		
		Damage to functionally linked habitats	Permanent Temporary,	Not significant
		potentially used by cited bird species from	<u>irreversible</u> , minor magnitude	
		changes in air quality, including from exhaust	and extent.	
		emissions.		

impact is less than 1% of the relevant critical loads and critical level and can therefore be ruled insignificant. Furthermore, the average N disposition at the SPA/SSSI remains below the critical level for supporting saltmarsh habitats (APIS, 2021) with atmospheric N of low importance for these systems.

Noise levels generated within the site during operation by traffic and visitors/rides are not anticipated to exceed 49dB at any offshore location. Peaks associated with the resort's operation can therefore be ruled out as having the potential to give rise to significant effect on either the SPA/Ramsar/SSSI



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (from increased ferry	Permanent Temporary,	Significant negative
		movements) associated with the new Thames	<u>ir</u> reversible, minor magnitude	at district
		Clipper river bus service and operational	and extent.	level Significant
		deliveries, potentially giving rise to		negative at District
		displacement, behavioural changes or		<u>level</u>
		physiological stress to cited bird species		
		potentially using functionally linked habitats.		
		Disturbance (noise and lighting) from the	Permanent Temporary,	Significant negative
		operation of the resort rides and attractions	<u>irreversible</u> reversible, minor	at district
		and entertainment facilities, giving rise to	magnitude and extent.	level Significant
		displacement, behavioural changes or		negative at District
		physiological stress to cited bird species within		level ⁵⁶
		the designated area.		
		Disturbance (human movement or activity), to	Permanent Temporary,	Significant negative
		cited bird species potentially using functionally	<u>irreversible</u> ,	at district
		linked habitats, as a result of increased human	moderate magnitude and	level Significant
		presence on the Project Site during operation	extent.	negative at District
		of the resort rides and attractions and		level
		entertainment facilities, including increased		
		recreational use of retained habitats.		

The most significant noise source during operation will be loudspeaker announcements, which are expected to produce up to 80db at the estuary front, dropping to c.50-59db throughout Black Duck Marsh. However, these announcements will be sporadic and ambient noise levels are expected to be around 40-49db within Black Duck Marsh and along the estuary foreshore.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance		
Feature	Receptor					
Medway	International	Potential impacts within the European/Ramsar Site itself:				
Estuary and		Damage to habitats within designated area	Temporary-	Not significant		
Marshes		used by cited bird species from changes to	permanent Temporary,			
SPA/Ramsar/		water and/or sediment quality (from surface	reversible-irreversible,			
SSSI ⁵⁷		or groundwater discharges from the Project	reversible, minor magnitude			
		Site including operational waste and	and extent.			
		pollutants).				
		Damage to habitats within the designated area	Permanent Temporary,	Not significant ⁵⁸		
		used by cited bird species and/or changes in	<u>irreversible</u> reversible, minor			
		the food supply of cited species from changes	magnitude and extent.			
		in air quality, including from exhaust				
		emissions.				
		Disturbance (from increased ferry	Permanent Temporary,	Not significant		
		movements) associated with the new Thames	<u>irreversible</u> reversible, minor			
		Clipper river bus service and operational	magnitude and extent.			
		deliveries, potentially giving rise to				
		displacement, behavioural changes or				
		physiological stress to cited bird species within				
		the designated area.				

With reference to the Air Quality Assessment in Chapter 16: Air Quality (Document reference 6.2.16), which is based on a traffic model which incorporates traffic data from relevant cumulative schemes, impacts upon the Medway Estuary and Marshes SPA/Ramsar/SSSI can be ruled insignificant. The designated site is located >200m from the roadside of roads predicted to experience an increase of >1000 AADT. The designated site is not within 10km of the energy centre point source. Average N disposition remains below the critical level for supporting saltmarsh habitats (APIS, 2021) with atmospheric N of low importance for these systems.



⁵⁷ _Medway Estuary and Marshes SSSI is a component SSSI to the Medway Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise and lighting) from the	<u>Permanent</u> Temporary,	Not significant 59
		operation of the resort rides and attractions	<u>ir</u> reversible, minor magnitude	
		and entertainment facilities, giving rise to	and extent.	
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		
		Recreational disturbance of flora and fauna	Permanent, reversible-	Not significant
		within designated area as a result of increased	<u>irreversible</u> Temporary,	
		residential population (due to residential	reversible, minor magnitude	
		accommodation for staff).	and extent.	
		Potential impacts solely on functionally linked re-	sources (outside of the European/	Ramsar Site):
		Damage to functionally linked habitats	Temporary-permanent,	Significant negative
		potentially used by cited bird species from	reversible-	at district
		changes to hydrology; and changes to water	<u>irreversible</u> Temporary,	level Significant
		and /or sediment quality (from surface or	moderate magnitude and	negative at District
		groundwater discharges from the Project Site	extent.	level
		including operational waste and pollutants).		
		Damage to functionally linked habitats	Permanent,	Not significant
		potentially used by cited bird species and/or	irreversible Temporary, minor	
		changes in the food supply of cited species	magnitude and extent.	
		from changes in air quality, including from		
		exhaust emissions.		

⁵⁹ _Noise levels generated within the site during operation by traffic and visitors/rides are not anticipated to exceed 49dB at any offshore location. Peaks associated with the resort's operation can therefore be ruled out as having the potential to give rise to significant effect on either the SPA/Ramsar/SSSI



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (from increased ferry	Permanent,	Significant negative
		movements) associated with the new Thames	<u>irreversible</u> Temporary,	at district
		Clipper river bus service and operational	reversible, minor magnitude	level Significant
		deliveries, potentially giving rise to	and extent.	negative at District
		displacement, behavioural changes or		level
		physiological stress to cited bird species		
		potentially using functionally linked habitats.		
		Disturbance (noise and lighting) from the	Permanent,	Significant negative
		operation of the resort rides and attractions	irreversible Temporary,	at district
		and entertainment facilities, giving rise to	reversible, minor magnitude	levelSignificant
		displacement, behavioural changes or	and extent.	negative at District
		physiological stress to cited bird species within		level ⁶⁰
		the designated area.		
		Disturbance (human movement or activity), to	Permanent,	Significant negative
		cited bird species potentially using functionally	<u>irreversible, Temporary,</u>	at district
		linked habitats, as a result of increased human	reversible, moderate	level Significant
		presence on the Project Site during operation	magnitude and extent.	negative at District
		of the resort rides and attractions and		level
		entertainment facilities, including increased		
		recreational use of retained habitats.		

⁶⁰ _The most significant noise source during operation will be loudspeaker announcements, which are expected to produce up to 80db at the estuary front, dropping to c.50-59db throughout Black Duck Marsh. However, these announcements will be sporadic and ambient noise levels are expected to be around 40-49db within Black Duck Marsh and along the estuary foreshore.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			_
North Downs	International	Traffic generated air quality impacts, resulting	Permanent,	Not significant ⁶²
Woodland		in nutrient enrichment and acidification of	<u>irreversible, Temporary</u> , minor	
SAC, and the		qualifying woodland habitat with subsequent	magnitude and extent.	
component		changes in structure and diversity.		
SSSI				
Wouldham to		For the ambient NoxNOx critical level, the		
Detling		critical level is predicted to be exceeded at		
Escarpment		North Downs Woodland, both with and		
SSSI ⁶¹		without the development in place.		
		Recreational disturbance of flora and fauna	Permanent,	Not significant
		within designated area as a result of increased	irreversible, Temporary,	
		residential population (due to residential	reversible, minor magnitude	
		accommodation for staff).	and extent.	
<u>Swanscombe</u>	<u>National</u>	Damage to habitats within the designated area	Permanent, irreversible, major	Significant negative
<u>Peninsula</u>		from traffic generated air quality impacts.	magnitude and extent.	at National level
<u>SSSI</u>			Likely to be major due to	
			proximity of SSSI to affected	
			road network and energy point	
			<u>source</u>	

⁶¹ _Wouldham to Detling Escarpment SSSI is a component SSSI to the North Downs Woodland SAC, with overlapping reasons for designation. Potential impacts to the SAC are therefore considered to adequately cover the designated interest features of the SSSI also

Circa 0.007% of the SAC land area lies within 200m of the A229, whilst 1.2% of SAC land area lies within 200m of the A249. Given the small extent of habitat located with the potential Zol of either main road, the potential for significant negative effects upon the favourable conservation status of the SAC is considered unlikely, whilst those SSSI units (units 26 and 15) in proximity to the A229 and A429 remain in favourable condition. As such, potential impacts arising are considered to be imperceptible and ruled insignificant. With reference to the Air Quality Assessment in Chapter 16: Air Quality (Document reference 6.2.16), which is based on a traffic model which incorporates traffic data from relevant cumulative schemes, the SAC is over 10km from the energy centre point source and over 200m from the roadside of any roads predicted to experience an increase of >1000 AADT, and the impact can be ruled insignificant.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Special feature of interest - populations of rare p	lants ⁶³ :	
		Damage to individual plants/supporting	Temporary-permanent,	Significant negative
		habitat, and part of plant meta-population,	reversible-irreversible,	at National level
		through increased recreational use, potentially	moderate magnitude and	
		resulting in trampling, littering, dog fouling,	extent.	
		unauthorised camping/campfires/barbecues.		
		Damage to individual plants Damage to	Temporary-permanent,	Significant negative
		individual plants/supporting habitat, and part	<u>reversible-</u>	at national
		of plant meta-population, through	irreversible Temporary,	level Significant
		inappropriate land management practices.	reversible, moderate	negative at National
			magnitude and extent.	level
		Damage to retained plants/population from	Temporary-permanent,	Significant negative
		changes to hydrological regime and changes in	reversible-	at national
		water quality/quantity.	irreversible Temporary,	level Significant
			reversible, moderate	negative at National
			magnitude and extent.	<u>level</u>
		Special feature of interest - breeding bird assemb	olages:	

Includes only those rare plant populations listed as a reason for notification of the Swanscombe Peninsula SSSI, including divided sedge (*Carex divisa*), yellow vetchling (*Lathyrus aphaca*), slender hare's-ear (*Bupleurum tenuissimum*), Bithynian vetch (*Vicia bithynica*) and round-leaved wintergreen (*Pyrola rotundifolia subsp. maritima*). A separate assessment of potentially significant effects upon 'non-SSSI' rare plants is provided under the 'rare plants' IEF.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor	·		
		Damage to on-site habitats potentially used by	Temporary-permanent,	Significant negative
		birds from changes to hydrology; and changes	<u>reversible-</u>	at national
		to water and /or sediment quality (from	irreversible Temporary,	level Significant
		surface or groundwater discharges from the	reversible, moderate	negative at National
		Project Site including operational waste and	magnitude and extent.	<u>level</u>
		pollutants).		
		Disturbance (noise ⁶⁴ and lighting) from the	Permanent,	Significant negative
		operation of the resort rides and attractions	irreversible Temporary,	at national
		and entertainment facilities, giving rise to	reversible, moderate	level Significant
		displacement, behavioural changes or	magnitude and extent.	negative at National
		physiological stress to bird species using the		<u>level</u>
		on-site habitats.		
		Disturbance (human movement or activity), to	Permanent,	Significant negative
		bird species, as a result of increased human	irreversible Temporary,	at national
		presence on the Project Site during operation	reversible, moderate	level Significant
		of the resort rides and attractions and	magnitude and extent.	negative at National
		entertainment facilities, including increased		<u>level</u>
		recreational use of retained habitats.		
		Increased collision risk, with potential for	<u>Permanent,</u>	Not significant
		killing/injuring birds, as a result of new tall	<u>irreversible</u> Permanent, minor	
		structures in the local environment.	magnitude and extent.	
		Special feature of interest - invertebrate assembl	<u>'ages:</u>	

Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Damage to on-site habitats potentially used by	Temporary-permanent,	Significant negative
		invertebrates from changes to hydrology; and	reversible-	at national
		changes to water and /or sediment quality	irreversible Temporary,	level Significant
		(from surface or groundwater discharges from	reversible, moderate	negative at National
		the Project Site including operational waste	magnitude and extent.	<u>level</u>
		and pollutants).		
		Disturbance (lighting) from the operation of	Permanent,	Significant negative
		the resort rides and attractions and	irreversible Temporary,	at national
		entertainment facilities.	reversible, minor magnitude	level Significant
			and extent.	negative at National
				level



Darenth	National	Damage to habitats within the designated area	Temporary-permanent,	Significant at the
Woods SSSI		from traffic generated air quality impacts.	<u>reversible-</u>	local level (on a
		resulting in nutrient enrichment and	irreversible Temporary,	precautionary basis,
		acidification of qualifying woodland habitat	reversible, minor magnitude	subject to further
		with subsequent changes in structure and	and extent.	investigation)
		diversity.		
		Chapter 16: Air Quality (Document reference		
		6.1.16) confirms that the impact of acid		
		deposition can be ruled insignificant.		
		In terms of the critical load for nutrient		
		nitrogen deposition, the contribution from the		
		Proposed Development is predicted to exceed		
		1% of the minimum critical load at Darenth		
		Wood SSSI. Transect data provided in Figure		
		16.23 have been used to calculate the extent of		
		nitrogen deposition based on the area		
		predicted to exceed 1% of the minimum		
		nutrient nitrogen critical load. A breakdown of		
		the areas to be affected, with reference to the		
		SSSI units, is provided in full in Table 12.22 . In		
		summary, the area predicted to exceed 1% of		
		the minimum nutrient nitrogen critical load is		
		8.01ha (6.52% of total site area). Three of the		
		nine SSSI units will be unaffected. At this level,		
		and based on the nature of the habitats		
		present, a significant ecological effect on the		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		site's integrity is considered unlikely.		
		For the ambient NOx critical level, the critical level		
		is predicted to be exceeded at Darenth Wood SSSI,		
		both with and without the development in place.		
		The contribution from the proposed		
		development is predicted to exceed 1% of the		
		critical level across 26.5ha; of the SSSI		
		accounting 21.72% of total site area.		
		Further investigation is needed to determine		
		the area predicted to exceed the 1% of the		
		critical level, using transect data, and if this will		
		result in a significant effect in ecology/ site		
		integrity, thereby requiring mitigation. On a		
		precautionary basis, based on the information		
		available, and subject to further investigation,		
		the effect is considered significant at the local		
		level.		
		Disturbance (noise and lighting) associated	Permanent,	Not significant ⁶⁵
		with traffic, operation of the resort rides and	irreversible Temporary,	
		attractions, and entertainment facilities.	reversible, minor magnitude	
			and extent.	

⁶⁵ Ambient noise level increases along the A2(t) are predicted to be <1db. Noise generated by rides and other facilities will be sub-significant within 300m of the DCO Order Limits.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Recreational disturbance of flora and fauna	<u>Permanent,</u>	Not significant
		within designated area as a result of increased	<u>irreversible</u> Temporary,	
		residential population (due to residential	reversible, minor magnitude	
		accommodation for staff).	and extent.	
Inner Thames	<u>National</u>	Potential impacts within the SSSI itself:		
Marshes SSSI		Damage to habitats within designated area	Temporary-permanent,	Not significant
		used by cited bird species from changes to	reversible-irreversible, minor	
		water and/or sediment quality (from surface	magnitude and extent.	
		or groundwater discharges from the Project		
		Site including operational waste and		
		pollutants).		
		Damage to habitats within the designated area	Permanent, irreversible, minor	Not significant ⁶⁶
		used by cited bird species and/or changes in	magnitude and extent.	
		the food supply of cited species from changes		
		in air quality, including from exhaust		
		emissions.		
		Disturbance (from increased ferry	Permanent, irreversible, minor	Not significant
		movements) associated with the new Thames	magnitude and extent.	
		Clipper river bus service and operational		
		deliveries, potentially giving rise to		
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		

The SSSI is over 2km from the energy centre point source and over 200m from the roadside of any roads predicted to experience an increase of >1000 AADT, and the impact can be ruled insignificant.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise and lighting) from the	Permanent, irreversible, minor	Not significant ⁶⁷
		operation of the resort rides and attractions	magnitude and extent.	
		and entertainment facilities, giving rise to		
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		
		Recreational disturbance of flora and fauna	Permanent, irreversible, minor	Not significant
		within designated area as a result of increased	magnitude and extent.	
		residential population (due to residential		
		accommodation for staff).		
		Potential impacts solely on functionally linked res	sources (outside of the SSSI):	
		Damage to functionally linked habitats	Temporary-permanent,	Significant negative
		potentially used by cited bird species from	reversible-irreversible,	at District level
		changes to hydrology; and changes to water	moderate magnitude and	
		and /or sediment quality (from surface or	<u>extent.</u>	
		groundwater discharges from the Project Site		
		including operational waste and pollutants).		
		Damage to functionally linked habitats	Permanent, irreversible, minor	Not significant
		potentially used by cited bird species and/or	magnitude and extent.	
		changes in the food supply of cited species		
		from changes in air quality, including from		
		exhaust emissions.		

Noise levels generated within the site during operation by traffic and visitors/rides are not anticipated to exceed 49dB at any offshore location. Peaks associated with the resort's operation can therefore be ruled out as having the potential to give rise to significant effect on the SSSI



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (from increased ferry	Permanent, irreversible, minor	Significant negative
		movements) associated with the new Thames	magnitude and extent.	at District level
		Clipper river bus service and operational		
		deliveries, potentially giving rise to		
		displacement, behavioural changes or		
		physiological stress to cited bird species		
		potentially using functionally linked habitats.		
		Disturbance (noise and lighting) from the	Permanent, irreversible, minor	Significant negative
		operation of the resort rides and attractions	magnitude and extent.	at District level
		and entertainment facilities, giving rise to		
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		
		Disturbance (human movement or activity), to	Permanent, irreversible,	Significant negative
		cited bird species potentially using functionally	moderate magnitude and	at District level
		linked habitats, as a result of increased human	extent.	
		presence on the Project Site during operation		
		of the resort rides and attractions, and		
		entertainment facilities.		
<u>West</u>	<u>National</u>	<u>Potential impacts within the SSSI itself:</u>		_
<u>Thurrock</u>		Damage to habitats within designated area	Temporary-permanent,	Not significant
Lagoon and		used by cited bird species from changes to	<u>reversible-</u>	
Marshes SSSI		water and/or sediment quality (from surface	irreversible Temporary,	
		or groundwater discharges from the Project	reversible, minor magnitude	
		Site including operational waste and	and extent.	
		pollutants).		



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Damage to habitats within the designated area	Permanent,	Not significant ⁶⁸
		used by cited bird species and/or changes in	irreversible Temporary,	
		the food supply of cited species from changes	reversible, minor magnitude	
		in air quality, including from exhaust	and extent.	
		emissions.		
		Disturbance (from increased ferry	Permanent,	Not significant
		movements) associated with the new Thames	<u>irreversible</u> Temporary,	
		Clipper river bus service and operational	reversible, minor magnitude	
		deliveries, potentially giving rise to	and extent.	
		displacement, behavioural changes or		
		physiological stress to cited bird species within		
		the designated area.		
		Disturbance (noise and lighting) from the	Permanent,	Not significant ⁶⁹
		operation of the resort rides and attractions	irreversible Temporary,	_
		and entertainment facilities, giving rise to	reversible, minor magnitude	
		displacement, behavioural changes or	and extent.	
		physiological stress to cited bird species within		
		the designated area.		
		Recreational disturbance of flora and fauna	Permanent,	Not significant
		within designated area as a result of increased	irreversible Temporary,	_
		residential population (due to residential	reversible, minor magnitude	
		accommodation for staff).	and extent.	

⁶⁸ _The SSSI is over 2km from the energy centre point source and over 200m from the roadside of any roads predicted to experience an increase of >1000 AADT, and the impact can be ruled insignificant. <u>Average N disposition remains below the critical level for supporting saltmarsh habitats (APIS, 2021) with atmospheric N of low importance for these systems.</u>

⁶⁹ Noise levels generated within the site during operation by traffic and visitors/rides are not anticipated to exceed 49dB at any offshore location. Peaks associated with the resort's operation can therefore be ruled out as having the potential to give rise to significant effect on the SSSI



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Potential impacts solely on functionally linked re	sources (outside of the SSSI):	
		Damage to functionally linked habitats	Temporary-permanent,	Significant negative
		potentially used by cited bird species from	reversible-	at district
		changes to hydrology; and changes to water	<u>irreversible</u> Temporary,	levelSignificant
		and /or sediment quality (from surface or	moderate magnitude and	negative at District
		groundwater discharges from the Project Site	extent.	level
		including operational waste and pollutants).		
		Damage to functionally linked habitats	Permanent,	Not significant
		potentially used by cited bird species and/or	<u>irreversible</u> Temporary , minor	
		changes in the food supply of cited species	magnitude and extent.	
		from changes in air quality, including from		
		exhaust emissions.		
		Disturbance (from increased ferry	Permanent,	Significant negative
		movements) associated with the new Thames	irreversible Temporary,	at district
		Clipper river bus service and operational	reversible, minor magnitude	levelSignificant
		deliveries, potentially giving rise to	and extent.	negative at District
		displacement, behavioural changes or		<u>level</u>
		physiological stress to cited bird species		
		potentially using functionally linked habitats.		
		Disturbance (noise and lighting) from the	Permanent,	Significant negative
		operation of the resort rides and attractions	irreversible Temporary,	at district
		and entertainment facilities, giving rise to	reversible, minor magnitude	level Significant
		displacement, behavioural changes or	and extent.	negative at District
		physiological stress to cited bird species within		level
		the designated area.		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (human movement or activity), to	Permanent,	Significant negative
		cited bird species potentially using functionally	<u>irreversible</u> Temporary,	at district
		linked habitats, as a result of increased human	reversible, moderate	levelSignificant
		presence on the Project Site during operation	magnitude and extent.	negative at District
		of the resort rides and attractions, and		level
		entertainment facilities.		
Shorne and	National	Damage to habitats within the designated area	Permanent,	Not significant ⁷⁰
Ashenbank		from traffic generated air quality impacts.	irreversible Temporary,	
Woods SSSI		Increased disposition of N likely to result in	reversible, minor magnitude	
		nutrient enrichment and acidification of	and extent.	
		qualifying woodland habitat with subsequent		
		changes in structure and diversity with such		
		impacts likely occurring across circa 0.2% of		
		total SSSI area is located within 200m of the A2		
		whilst the woodland is reported to be in		
		favourable or unfavourable recovering		
		condition.		

Given the relatively small area of qualifying woodland habitat within 200m of the A2, combined with the predominantly favourable condition of woodland habitat, significant negative effects upon the integrity of this SSSI and its qualifying features are considered unlikely.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
<u>Cobham</u>	<u>National</u>	Damage to habitats within the designated area	<u>Permanent,</u>	Not significant ⁷¹
Woods SSSI		from traffic generated air quality impacts.	irreversible Temporary,	
		Increased disposition of N likely to result in	reversible, minor magnitude	
		nutrient enrichment and acidification of	and extent.	
		qualifying woodland habitat with subsequent		
		changes in structure and diversity with such		
		impacts likely occurring across circa 5.7% of		
		total SSSI area is located within 200m of the		
		M2 motorway. Total area of impact is,		
		however, reduced given N sensitive woodland		
		habitat- is only present across a proportion of		
		this area.		

Given the relatively small area of qualifying woodland habitat within 200m of the M2 motorway, combined with its favourable condition, significant negative effects upon the integrity of this SSSI and its qualifying features are considered unlikely.



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
Great	<u>National</u>	Damage to habitats within the designated area	Permanent,	Significant at the
<u>Crabbles</u>		from traffic generated air quality impacts.	irreversible Temporary,	local level
Wood SSSI		Increased disposition of N likely to result in	reversible, minor magnitude	
		nutrient enrichment and acidification of	and extent	
		qualifying woodland habitat with subsequent		
		changes in structure and diversity with such		
		impacts likely occurring across circa 13.6% of		
		total SSSI area is located within 200m of the		
		A289 whilst the woodland is reported to be in		
		unfavourable recovering condition. Total area		
		of impact is, however, reduced given N		
		sensitive woodland habitat is only present		
		across a proportion of this area.		
Botany	County	Damage to habitats from changes in air	Permanent,	Not significant
Marshes LWS		quality, including from exhaust emissions.	irreversible Temporary,	
			reversible, minor magnitude	
			and extent.	
		Damage to habitats from changes to	Temporary-permanent,	Significant at local
		hydrological regime, and changes in water	<u>reversible-</u>	levelSignificant at
		quality/quantity.	<u>irreversible</u> Temporary,	Local level
			moderate magnitude and	
			extent.	



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise and lighting) from the	Permanent,	Significant at local
		operation of the resort rides and attractions	<u>irreversible</u> Temporary ,	levelSignificant at
		and entertainment facilities, potentially giving	moderate magnitude and	<u>Local level</u>
		rise to displacement, behavioural changes or	extent.	
		physiological stress to wildlife species within		
		the designated area.		
		Recreational disturbance of flora and fauna	Permanent,	Significant at local
		within designated area as a result of increased	irreversible Temporary,	levelSignificant at
		residential population (due to residential	reversible, minor magnitude	Local level
		accommodation for staff) and potential	and extent.	
		increased recreational use of Swanscombe		
		Peninsula from members of the general public.		



Ebbsfleet	County	Damage to habitats within the designated area	Permanent,	Significant at the
Marshes,		from traffic generated air quality impacts.	<u>irreversible <mark>Temporary,</mark></u>	local level (on a
Northfleet		Increased disposition of N likely to result in	reversible, minor magnitude	precautionary basis,
LWS		nutrient enrichment and acidification of	and extent.	subject to further
		woodland habitat with subsequent changes in		investigation)
		structure and diversity. Open water and		
		eutrophic swamp are, however, less sensitive		
		to atmospheric N inputs such that significant		
		effects are unlikely.		
		Chapter 16: Air Quality (Document reference		
		6.1.16) confirms that the impact of acid		
		deposition can be ruled insignificant.		
		In terms of the critical load for nutrient		
		nitrogen deposition, the contribution from the		
		Proposed Development is predicted to exceed		
		1% of the minimum critical load at Ebbsfleet		
		Marshes, Northfleet LWS. Transect data		
		provided in Figure 16.21 have been used to		
		calculate the extent of nitrogen deposition		
		based on the area predicted to exceed 1% of		
		the minimum nutrient nitrogen critical load.		
		The area predicted to exceed 1% of the		
		minimum nutrient nitrogen critical load is		
		3.93ha (8.45% of total site area) (see in Table		
		12.22). At this level, a significant ecological		
		effect on the site's integrity is considered		
		unlikely.		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		For the ambient NOx critical level, the critical		
		level is predicted to be exceeded at Ebbsfleet		
		Marshes, Northfleet LWS, both with and		
		without the development in place. Further		
		investigation is needed to determine the area		
		predicted to exceed the 1% of the critical level,		
		using transect data, and if this will result in a		
		significant effect in ecology/ site integrity,		
		thereby requiring mitigation. On a		
		precautionary basis, based on the information		
		available, and subject to further investigation,		
		the effect is considered significant at the local		
		level.		
		Damage to habitats from changes to	Temporary-permanent,	Significant at local
		hydrological regime and changes in water	reversible-	levelSignificant at
		quality/quantity.	<u>irreversible</u> Temporary,	Local level
			moderate magnitude and	
			extent.	
		Disturbance (noise and lighting) from the	Permanent,	Significant at local
		operation of the resort rides and attractions	irreversible Temporary,	levelSignificant at
		and entertainment facilities, potentially giving	moderate magnitude and	Local level
		rise to displacement, behavioural changes or	extent.	
		physiological stress to wildlife species within		
		the designated area.		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Recreational disturbance of flora and fauna	<u>Permanent,</u>	Significant at local
		within designated area as a result of increased	irreversible Temporary,	levelSignificant at
		residential population (due to residential	reversible, minor magnitude	Local level
		accommodation for staff) and potential	and extent.	
		increased recreational use of Swanscombe		
		Peninsula from members of the general public.		
Alkerden	County	Damage to habitats from changes in air	Temporary, reversible, minor	Not significant
Lane Pit LWS		quality, including from exhaust emissions.	magnitude and extent.	_



Disused	County	Damage to habitats within the designated area	Permanent,	Significant at the
Hospital		from traffic generated air quality impacts.	<u>irreversible</u> Temporary,	local level (on a
Grounds,			reversible, minor magnitude	precautionary basis,
Mabledon		Chapter 16: Air Quality (Document reference	and extent.	subject to further
LWS		6.1.16) confirms that the impact of acid		investigation)
		deposition, and nutrient nitrogen deposition, can be ruled insignificant.		
		For the ambient NoOx critical level, the critical		
		level is predicted to be exceeded at Disused		
		Hospital Grounds, Mabledon LWS, both with		
		and without the development in place. The		
		contribution from the proposed development is		
		predicted to exceed 1% of the critical level		
		across 1.76ha of the LWS accounting for 21.81% of total site area.		
		21.81% of total site area.		
		For the ambient NOx critical level, the critical		
		level is predicted to be exceeded at Disused		
		Hospital Grounds, Mabledon LWS, both with		
		and without the development in place. Further		
		investigation is needed to determine the area		
		predicted to exceed the 1% of the critical level,		
		using transect data, and if this will result in a		
		significant effect in ecology/ site integrity,		
		thereby requiring mitigation. On a		
		precautionary basis, based on the information		
		available, and subject to further investigation,		



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		the effect is considered significant at the local level.		
Tilbury	County	Damage to habitats from changes to	Temporary-permanent,	Significant at local
Marshes LWS		hydrological regime and changes in water	<u>reversible-</u>	levelSignificant at
		quality/quantity.	<u>irreversible</u> Temporary,	<u>Local level</u>
			moderate magnitude and	
			extent.	
		Disturbance (from increased ferry	Permanent,	Significant at local
		movements) associated with the new Thames	irreversible Temporary,	levelSignificant at
		Clipper river bus service and operational	reversible, minor magnitude	Local level
		deliveries, potentially giving rise to	and extent.	
		displacement, behavioural changes or		
		physiological stress to bird species potentially		
		using functionally linked habitats.		
		Disturbance (noise and lighting) from the	Permanent,	Significant negative
		operation of the resort rides and attractions	irreversible Temporary,	at local
		and entertainment facilities, giving rise to	reversible, minor magnitude	level Significant
		displacement, behavioural changes or	and extent.	negative at Local
		physiological stress to cited bird species within		<u>level</u>
		the designated area.		
		Disturbance (human movement or activity), to	Permanent,	Significant negative
		cited bird species potentially using functionally	irreversible Temporary,	at local
		linked habitats, as a result of increased human	reversible, moderate	level Significant
		presence on the Project Site during operation	magnitude and extent.	negative at Local
		of the resort rides and attractions, and		<u>level</u>
		entertainment facilities.		
Habitats/Flora				



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
Populations	National	N.B. Assessment of potential effects upon	Temporary, reversible,	Significant at
of nationally		scarce plants included under Swanscombe	moderate magnitude and	regional level
scarce plants		Peninsula SSSI. Damage to individual plants,	extent.	
		and part of plant meta-population, through		
		increased recreational use, potentially		
		resulting in trampling, littering, dog fouling,		
		unauthorised camping/campfires/barbecues.		
Retained	Local to	Damage to retained habitats, through	Permanent,	Significant at local-
habitats ⁷²	County (see	increased recreational use, potentially	irreversible Temporary,	Ceounty -level
(including	Baseline	resulting in trampling, littering, dog fouling,	reversible , moderate	
woodland,	section)	unauthorised camping/campfires/barbecues.	magnitude and extent.	
scrub,		Damage to retained habitats through	Temporary-permanent,	Significant at Local-
grassland,		inappropriate land management practices.	<u>reversible-</u>	County
open mosaic			<u>irreversible</u> Temporary,	<u>level</u> Significant at
on previously			reversible, moderate	local level
developed			magnitude and extent.	
land,		Damage to retained habitats from changes to	Temporary-permanent,	Significant at Local-
wetlands,		hydrological regime and changes in water	<u>reversible-</u>	County
waterbodies		quality/quantity.	<u>irreversible</u> Temporary,	<u>level</u> Significant at
and the River			moderate magnitude and	local level
Ebbsfleet).			extent.	

⁷² Predicted impacts on retained habitats during the operation of the Proposed Development are considered applicable across the full suite of habitats retained, and are therefore considered collectively



Ancient	County	Damage to habitats within the designated area	Permanent,	Significant at the
woodland		from traffic generated air quality impacts.	<u>irreversible Temporary,</u>	local level (on a
		Increased disposition of N likely to result in	reversible, minor magnitude	precautionary basis,
		nutrient enrichment and acidification of	and extent.	subject to further
		woodland habitat with subsequent changes in		investigation)
		structure and diversity.		
		Chapter 16: Air Quality (Document reference		
		6.1.16) confirms that the impact of acid		
		deposition can be ruled insignificant.		
		In terms of the critical load for nutrient		
		nitrogen deposition, the contribution from the		
		Proposed Development is predicted to exceed		
		1% of the minimum critical load at the		
		following ecological sites. Transect data		
		provided in Figures 16.20 – 16.24 have been		
		used to calculate the extent of nitrogen		
		deposition based on the area predicted to		
		exceed 1% of the minimum nutrient nitrogen		
		critical load). These areas are provided in		
		brackets in the list below, and in full in		
		Table 12.22.		
		• Coombegreen Wood (0.75ha; 4.01% of total site area);		
		 Darenth Wood (8.01ha; 6.52% of total site area); 		



Parkhill Wood (1.03ha; 21.59% of total site area); and	
• The Thrift (1.84ha; 20.74% of total site area).	
For the ambient NeOx critical level, the critical level is predicted to be exceeded at all sites for all scenarios, both with and without the development in place.	
The contribution from the proposed development is predicted to exceed 1% of the critical level at the following ecological sites:	
 Darenth Wood (26.5ha; 21.72% of total site area); 	
• The Thrift (4.16ha; 50.30% of total site area);	
 Coombegreen Wood (3.77ha; 22.77% of total site area); 	
 Parkhill Wood; 2.87ha; 60.17% of total site area); 	
 Rams Wood (3.13ha; 86.23% of total site 	



area); • Disused Hospital; 1.76ha (21.81% of total site area);-• Cobham Hall Wood; 2.54ha; 49.61% of total site area); • Hobbs Hole (1.21ha; 75.16% of total site area); and • Jackson Wood (0.59ha; 13.50% of total site area). For the ambient NOx critical level, the critical level is predicted to be exceeded at all sites for all scenarios, both with and without the development in place. The contribution from the proposed development is predicted to exceed 1% of the critical level at the following ecological sites: Darenth Wood: • The Thrift: Coombegreen Wood; Parkhill Wood:



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor	-		
		• Rams Wood;		
		• Disused Hospital;		
		• Cobham Hall Wood;		
		• Hobbs Hole; and		
		• Jackson Wood.		
		Further investigation is needed to determine		
		the area predicted to exceed the 1% of the		
		critical level, and if this will result in a		
		significant effect in ecology/ site integrity,		
		thereby requiring mitigation. On a		
		precautionary basis, based on the information		
		available, and subject to further investigation,		
		the effect is considered significant at the local		
		level.		
Species	1	,		1
Wintering	International	N.B. Assessment of potential habitat loss and dis	turbance effects upon wintering v	waterfowl and wading
Waterfowl		bird assemblage included under SPA/Ramsar Site		
and Wading		used by birds from changes to hydrology; and cl	-	The state of the s
Bird		or groundwater discharges from the Project Site	_	
Assemblage		Temporary, reversible, minor magnitude and ext		• ,
J -		Significant negative at district level		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Increased collision risk, with potential for	Permanent, minor magnitude	Not significant ⁷³
		killing/injuring birds, as a result of new tall	and extent.	
		structures in the local environment.		
Wintering	County	Damage to on-site habitats potentially used by	Temporary-permanent,	Significant negative
terrestrial		birds from changes to hydrology; and changes	<u>reversible-</u>	at district
bird		to water and /or sediment quality (from	<u>irreversible</u> Temporary,	level Significant
assemblage		surface or groundwater discharges from the	reversible, minor magnitude	negative at District
		Project Site including operational waste and	and extent.	level
		pollutants).		
		Disturbance (noise ⁷⁴ and lighting) from the	Permanent,	Significant negative
		operation of the resort rides and attractions	irreversible Temporary,	at district
		and entertainment facilities, giving rise to	reversible, minor magnitude	level Significant
		displacement, behavioural changes or	and extent.	negative at District
		physiological stress to bird species using the		level
		on-site habitats.		_

⁷⁴ Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



⁷³ Unrestricted flight paths into core habitat area within Black Duck Marsh retained. Botany Marsh West lost resulting in birds less likely to fly over the Swanscombe Peninsula

Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (human movement or activity), to	Permanent,	Significant negative
		cited bird species potentially using functionally	<u>irreversible</u> Temporary,	at district
		linked habitats, as a result of increased human	reversible , moderate	level Significant
		presence on the Project Site during operation	magnitude and extent.	negative at District
		of the resort rides and attractions and		level
		entertainment facilities, including increased		
		recreational use of retained habitats.		
		Increased collision risk, with potential for	Permanent, minor magnitude	Not significant
		killing/injuring birds, as a result of new tall	and extent.	
		structures in the local environment.		
Breeding Bird	Regional Natio	N.B. Assessment of potential effects upon breedi	ing bird assemblage included und	er Swanscombe
Assemblage	<u>nal</u>	Peninsula SSSI. Damage to on-site habitats poten	tially used by birds from changes	to hydrology; and
		changes to water and /or sediment quality (from	n surface or groundwater dischar	ges from the Project
		Site including operational waste and pollutants)	.	
		Temporary, reversible, minor magnitude and ext	ent.	
		Significant negative at district level		
Bat	District_	Damage to on-site habitats potentially used by	Temporary-permanent,	Significant negative
Assemblage	<u>County</u>	bats from changes to hydrology; and changes	<u>reversible-</u>	at district
		to water and /or sediment quality (from	<u>irreversible</u> Temporary,	levelSignificant
		surface or groundwater discharges from the	reversible, minor magnitude	negative at District
		Project Site including operational waste and	and extent.	<u>level</u>
		pollutants).		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (lighting) from the operation of	Permanent,	Significant negative
		the resort rides and attractions and	irreversible Temporary,	at district
		entertainment facilities, giving rise to	reversible, minor magnitude	level Significant
		displacement, behavioural changes or	and extent.	negative at District
		physiological stress to bat species using the on-		<u>level</u>
		site habitats, including light pollution of		
		foraging, commuting and roosting habitats.		
		Increased collision risk, with potential for	Permanent, minor magnitude	Not significant
		killing/injuring bats, as a result of new tall	and extent.	
		structures in the local environment.		
Dormouse	District	Damage to on-site habitats potentially used by	Temporary-permanent,	Significant negative
		dormice from changes to hydrology; and	reversible-	at local
		changes to water and /or sediment quality	irreversible Temporary,	level Significant
		(from surface or groundwater discharges from	reversible, minor magnitude	negative at Local
		the Project Site including operational waste	and extent.	<u>level</u>
		and pollutants).		



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise ⁷⁵ and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to dormice using the onsite habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	Permanent, irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
Otter	Local	Damage to on-site habitats potentially used by otters from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	Temporary-permanent, reversible- irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level

⁷⁵ Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise ⁷⁶ and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to otters using the on-site habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	Permanent, irreversibleTemporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
Water Vole	Local to district	Damage to on-site habitats potentially used by water voles from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	Temporary-permanent, reversible- irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level

⁷⁶ Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance
Feature	Receptor			
		Disturbance (noise ⁷⁷ and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to water voles using the onsite habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	Permanent, irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
Harvest Mouse	Local	Damage to on-site habitats potentially used by harvest mice from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	Temporary-permanent, reversible- irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level

⁷⁷ Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological Feature	Importance of Receptor	Nature of Impact	Impact Magnitude	Significance
		Disturbance (noise ⁷⁸ and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to harvest mice using the on-site habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	Permanent, irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at local levelSignificant negative at Local level
•	Local to district	Damage to on-site habitats potentially used by amphibians from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	Temporary-permanent, reversible- irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level
		Disturbance (lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to amphibians using the onsite habitats, including light pollution of breeding, foraging and hibernation/shelter habitats.	Permanent, irreversible Temporary, reversible, minor magnitude and extent.	Significant negative at district levelSignificant negative at District level

⁷⁸ Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological	Importance of	Nature of Impact	Impact Magnitude	Significance		
Feature	Receptor					
Reptile	District	Damage to on-site habitats potentially used by	Temporary-permanent,	Significant negative		
assemblage		reptiles from changes to hydrology; and	<u>reversible-</u>	at district		
		changes to water and /or sediment quality	<u>irreversible</u> Temporary,	level Significant		
		(from surface or groundwater discharges from	reversible, minor magnitude	negative at District		
		the Project Site including operational waste	and extent.	<u>level</u>		
		and pollutants).				
		Disturbance (lighting) from the operation of	Permanent,	Significant negative		
		the resort rides and attractions and	<u>irreversible</u> Temporary,	at district		
		entertainment facilities, giving rise to	reversible, minor magnitude	level Significant		
		displacement, behavioural changes or	and extent.	negative at District		
		physiological stress to reptiles using the on-site		<u>level</u>		
		habitats, including light pollution of potential				
		shelter and hibernation habitats.				
Invertebrate	National	N.B. Assessment of potential effects upon inverte	ebrate assemblages included und	er Swanscombe		
assemblage		Peninsula SSSI. Damage to on-site habitats poter	ntially used by invertebrates from	changes to hydrology;		
		and changes to water and /or sediment quality (from surface or groundwater discharg				
		Temporary, reversible, minor magnitude and extent.				
		Significant negative at district level				

Table 12.12 Extent of nitrogen deposition: Darenth Wood SSSI (based on area predicted to exceed 1% of the minimum nutrient nitrogen critical load).



Receptor	Unit	Unit Name	Condition	Area (ha)	Impact area (ha) (within designation)	Impact area (% of total)
Darenth Wood (SSSI/Ancient woodland)	1	Darenth Country Park	Favourable	4.19	0.32	7.63
ŀ	2	North west	Favourable	17.03	1.25	7.34
	3	North east	Unfavourable - recovering	11.16	1.16	10.40
	4	East	Favourable	8.18	1.40	17.12
	5	Central (badgers mount)	Favourable	18.39	1.68	9.14
	6	West (including former hospital ground)	Favourable	15.41	2.20	14.28
	7	Ladies wood	Favourable	13.49	0.00	0.00
	8	Lords wood	Favourable	32.55	0.00	0.00
	9	Adjacent old hospital site	Favourable	2.52	0.00	0.00
				122.91	8.01	6.52



Table 12.13 Extent of NOx deposition: Darenth Wood SSSI (based on area predicted to exceed 1% of the minimum NOx critical load).

Receptor	<u>Unit</u>	<u>Unit Name</u>	Condition	<u>Area (ha)</u>	Impact area (ha) (within designation)	Impact area (% of total)
Darenth Wood (SSSI/Ancient woodland)	1	Darenth Country Park	<u>Favourable</u>	4.19	2.18	52.03
	<u>2</u>	North west	<u>Favourable</u>	<u>17.03</u>	<u>4.8</u>	<u>28.19</u>
	<u>3</u>	North east	<u>Unfavourable -</u> <u>recovering</u>	<u>11.16</u>	4.84	43.37
	<u>4</u>	<u>East</u>	<u>Favourable</u>	<u>8.18</u>	<u>6.17</u>	<u>75.43</u>
	<u>5</u>	Central (badgers mount)	<u>Favourable</u>	<u>18.39</u>	<u>4.86</u>	<u>26.43</u>
	<u>6</u>	West (including former hospital ground)	<u>Favourable</u>	<u>15.41</u>	<u>3.65</u>	<u>23.69</u>
	7	<u>Ladies wood</u>	<u>Favourable</u>	<u>13.49</u>	0.00	0.00



Receptor	<u>Unit</u>	Unit Name	Condition	<u>Area (ha)</u>	Impact area (ha) (within designation)	Impact area (% of total)
	<u>8</u>	<u>Lords wood</u>	<u>Favourable</u>	<u>32.55</u>	0.00	0.00
	9	Adjacent old hospital site	<u>Favourable</u>	2.52	1.01	40.08
		-		<u>122.91</u>	<u>26.5</u>	<u>21.72</u>

Table 12.143 Extent of nitrogen deposition: Ancient Woodlands and Local Wildlife Sites (based on area predicted to exceed 1% of the minimum nutrient nitrogen critical load).

Receptor	Status	Site Area (ha)	Area (ha) within designation exceeding 1% of the critical load)	Impact area (% of total)
Rams Wood	Ancient woodland	3.63	0.21	5.79
The Thrift	Ancient woodland	8.87	1.84	20.74
Parkhill Wood	Ancient woodland	4.77	1.03	21.59
Jackson Wood	Ancient woodland	N/A	0	N/A



Receptor	Status	Site Area (ha)	Area (ha) within designation exceeding 1% of the critical load)	Impact area (% of total)
Coombegreen	Ancient woodland	18.71	0.75	4.01
Ebbsfleet Marshes	Local Wildlife Site	46.49	3.93	8.45
Disused Hospital	Local Wildlife Site	8.07	0.26	3.22



Table 12.15 Extent of nitrogen deposition: Ancient Woodlands and Local Wildlife Sites (based on area predicted to exceed 1% of the minimum nutrient nitrogen critical load).

Receptor	<u>Status</u>	Site Area (ha)	Area (ha) within designation exceeding 1% of the critical load)	Impact area (% of total)
Rams Wood	Ancient woodland	3.63	<u>3.13</u>	86.23
The Thrift	Ancient woodland	8.87	<u>4.16</u>	50.30
Parkhill Wood	Ancient woodland	4.77	<u>2.87</u>	60.17
Jackson Wood	Ancient woodland	<u>N/A</u>	0.59	13.50
Coombegreen	Ancient woodland	<u>18.71</u>	3.77	22.77
Codham Hall Woods	Ancient woodland	5.12	2.54	<u>49.61</u>
Hobbs Hole	Ancient woodland	<u>1.61</u>	1.21	<u>75.16</u>
Disused Hospital	Local Wildlife Site	8.07	<u>1.76</u>	21.81





Biodiversity Net Gain Impact Assessment

- 12.249 Within the current draft of the Environment Bill, as submitted to Parliament, Under the Environment Act 2021 Nationally Significant Infrastructure Projects (NSIPs) are exempt from the requirement to deliverit will become mandatory for Nationally Significant Infrastructure Projects (NSIPs) to achieve 10% net gain. However, mandatory net gain for NSIPs will not come into effect until secondary legislation has been passed, which is not expected before November 2024.
- 12.250 Nonetheless, the Applicant is submitting the biodiversity metrichas commissioned a Biodiversity Impact Assessment (BIA) on a voluntary basis to demonstrate a commitment to delivering net gain in accordance with the NPPFof the Proposed Development. These biodiversity net gain The BIA assessment calculations, which are provided in Appendix 12.2: Biodiversity Net Gain Impact Assessment (Document reference 6.2.12.2), have been used to objectively provide an overall biodiversity score for the project—using a standardised metric.
- 12.251 The extent of important ecological—habitats within the Project Site and the quantities retained, lost and proposed have been assessed using the 'beta test' version of the Biodiversity Metric 2.0, developed by Natural England with reference to the biodiversity net gain requirement specified within the NPPF (paragraph 170.d). Within the current draft of the Environment Bill, as submitted to Parliament, Nationally Significant Infrastructure Projects (NSIPs) are exempt from the requirement to deliver 10% net gain. The Applicant is submitting the biodiversity metric on a voluntary basis to demonstrate a commitment to delivering net gain in accordance with the NPPF. These biodiversity net gain assessment calculations, which are provided in Appendix 12.2: Biodiversity Net Gain Assessment (Document reference 6.2.12.2), have been used to objectively provide an overall biodiversity score for the project. As shown in detail in Appendix 12.2 (Document Reference 6.2.12.2), and summarised below, overall, the Proposed Development is considered likely to result in a net loss in habitat units:
 - Net biodiversity units (habitat units) = -8291,211.908 units (-2436.7815% net loss).

12.1

12.212.252 It should be noted that the calculations are based on the London Resort Masterplan and additional detail regarding landscape treatment as shown on the Landscape Masterplan included within the Landscape Strategy (Document Reference 6.2.11.7), and therefore may be subject to a degree of variance as the detailed design of the resort progresses. However, subject to the delivery of appropriate habitat provision and management, this is considered to be sufficient for assessing the overall biodiversity impact of the Proposed Development at this stage.



⁷⁹ _http://publications.naturalengland.org.uk/publication/5850908674228224

- 12.312.1 As shown in detail in Appendix 12.2 (Document Reference 6.2.12.2), and summarised below, overall, the Proposed Development is considered likely to result in a net loss in habitat units:
 - Net biodiversity units (habitat units) = -829.98 units (-24.78% net loss).
- 12.412.253 To achieve a net gain, tThe Applicant is committed to the funding and delivery of sufficient an offsite compensation scheme involving habitat creation and enhancement sufficient to deliver the necessary credits to achieve a net gain, in line with the current requirements of the NPPF (paragraph 170.densure that the post development biodiversity value (as measured in habitat units) exceeds the pre-development baseline.
- 12.5 The applicant is working with local landowners, consultees, stakeholders and delivery partners to devise an appropriate biodiversity offsetting scheme. However, at the time of the submission of the DCO application, details of the offsetting scheme are not yet fixed, however a commitment to delivering off-site mitigation of an appropriate scale and type is set out with Appendix 12.10: General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10), which will secured as a requirement of the DCO.
- 12.254 As set out in Appendix 12.2: Biodiversity Net Gain Assessment (Document Reference 6.2.12.2), a range of theoretical mitigation land scenarios, involving habitat creation and enhancement, have been tested in order to illustrate how a net gain to biodiversity using the Defra Biodiversity Metric 2.0 might be achieved, and to demonstrate that it is technically achievable. The calculations are detailed in full in Appendix 12.2 (Document Reference 6.2.12.2), and confirm, in summary, that the Proposed Development is capable of delivering a net gain to biodiversity subject to the delivering of off-site mitigation on between 160 and 210 hectares of land It is important to note, however, that the standard metric-based approach to BIA does not adequately account for impacts on protected sites such as SSSIs. This is made clear in the Biodiversity Metric 2.0 User Guide (Beta Version, July 2019)⁸⁰, which includes the following principles and rules for when using the metric:
 - "Principle 1: The metric does not change the protection afforded to biodiversity. Existing levels of protection afforded to protected species and to habitats are not changed by use of this or any other metric. Statutory obligations will still need to be satisfied.
 - Principle 2: Biodiversity metric calculations can inform decision-making where application of the mitigation hierarchy and good practice principles⁸¹ conclude that compensation for habitat losses is justified.

⁸¹ CIEEM, CIRIA, IEMA. 2016 Biodiversity Net Gain – Good Practice Principles for Development. https://www.cieem.net/data/files/Publications/Biodiversity Net Gain Principles.pdf



⁸⁰ IAN CROSHER A, SUSANNAH GOLD B, MAX HEAVER D, MATT HEYDON A, LAUREN MOORE D, STEPHEN PANKS A, SARAH SCOTT C, DAVE STONE A & NICK WHITE A. 2019. The Biodiversity Metric 2.0: auditing and accounting for biodiversity value. User guide (Beta Version, July 2019). Natural England A – Natural England, B – Imperial College, University of London, C – Environment Agency, D – Department for Environment, Food and Rural Affairs

- Principle 3: The metric's biodiversity units are only a proxy for biodiversity. While it is underpinned by ecological evidence the metric is only a proxy for biodiversity and to be of practical use has been kept deliberately simple. The numerical values generated by the metric represent relative, not absolute, values.
- Principle 4: The metric focuses on widespread species and typical habitats. Area based habitats are considered a suitable proxy for widespread species found in typical examples of different habitat types.
 - Protected and locally important species needs are not considered through the metric,
 - Impacts on protected (e.g. SSSIs) and irreplaceable habitats are not adequately measured by this metric, and will likely require separate consideration."
- 12.255 Furthermore, in considering whether biodiversity net gain can also be achieved when compensation of impacts is required, planning policy guidance82 associated with the NPPF states:

"Biodiversity net gain complements and works with the biodiversity mitigation hierarchy set out in NPPF paragraph 175a. It does not override the protection for designated sites, protected or priority species and irreplaceable or priority habitats set out in the NPPF. Local planning authorities need to ensure that habitat improvement will be a genuine additional benefit, and go further than measures already required to implement a compensation strategy"

- 12.256 Thus, ecological enhancement measures can only be deemed to achieve biodiversity net gain where these provide a benefit that goes beyond that required to compensate for impacts which cannot be avoided or mitigated. As described above with respect to construction effects (and below with regard to operational effects) the Proposed Development will unavoidably result in direct negative effects on Swanscombe Peninsula SSSI and its notified features. Such effects are to be addressed through avoidance and mitigation where possible, but inevitably compensatory measures will also be required.
- 12.257 It is unlikely that the benefits of the off-site enhancement measures proposed will exceed that required to compensate for the on-site impacts on Swanscombe Peninsula SSSI.

 However, it is anticipated the proposed off-site enhancement measures will result in a net gain in biodiversity units when measured using a standard Biodiversity Metric.

12.6

PROPOSED MITIGATION

⁸² https://www.gov.uk/guidance/natural-environment; Paragraph: 024 Reference ID: 8-024-20190721, Revision date: 21 07 2019



12.712.258 The preceding section of this chapter describes the potential significant environmental effects of the Proposed Development and provides an overview of the inherent mitigation measures incorporated into the Proposed Development. However, not all likely significant negative effects identified can be avoided or reduced in severity through inherent mitigation alone. This section therefore identifies those additional mitigation measures required proposed to avoid, or reduce or offset the likely significant negative impacts.

12.812.259 The key mechanisms to deliver mitigation will include measures to:

- Conform with relevant and pertinent legislative requirements, particular those associated with legally protected species; and
- Deliver and, where possible, maximise opportunities for biodiversity enhancement and gain through the Proposed Development.
- 12.912.260 The mitigation strategy strategies in respect of certain statutory designations, some of the legally protected species and species of conservation concern present has ve been developed in consultation with Natural England through their Discretionary Advice Service and a series of meetings as set out in **Table 12.2**.

Construction Measures

<u>12.261</u> The key mechanisms to deliver avoidance or mitigation of potential significant construction effects are summarised below. Further details on these measures, including which IEFs <u>and effects</u> they relate to, are provided within **Table 12.14**.

12.223 [Mechanisms for delivering mitigation as set out below to be included within an overall 'SSSI Mitigation Strategy', to be agreed with Natural England]

Construction Environmental Management Plan (CEMP)

- 12.1012.262 The Principal Contractor for the Proposed Development will produce a Construction Environmental Management Plan (CEMP) to include mitigation measures to protect the environment during both the demolition and construction phases. The CEMP will be secured by a requirement in the DCO. An Outline CEMP has been prepared in support of the DCO application (Document Reference 6.2.3.2). With regard to the avoidance and mitigation of effects on ecological features, the following measures are to be included within the CEMP:
 - Restricted working hours and a Sensitive Lighting Strategy (SLS) to minimise impacts on nocturnal wildlife;
 - Suppression of construction noise and dust;



- Use of site screening/hoarding to minimise visual, aural and light disturbance;
- Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy; and
- Adherence to pollution prevention guidelines.

Ecological Construction Method Statements (ECMSs)

- <u>12.1112.263</u> A series of ECMSs are contained within an overarching Ecological Mitigation and Management Framework (EMMF) (Document Reference 6.2.12.3), which will be secured as a requirement of the DCO. The measures contained within the ECMSs include (but are not limited to):
 - Provision of an Ecological Clerk of Works (ECoW) to supervise construction activities in sensitive areas for habitats and species;
 - Provision of 'toolbox talks' to relevant site contractors;
 - Method statements for the delivery of works in sensitive areas, including controls over timing/duration of works to avoid sensitive times of year (e.g. the bird breeding season);
 - Precautionary methods of working and sensitive vegetation/site clearance (under a European Protected Species where necessary), e.g. 'soft-felling' trees with bat roost potential or removal of suitable dormouse habitat;
 - Translocation/displacement of protected species prior to construction works commencing (under a European Protected Species where necessary);
 - Sensitive lighting strategy (SLS) to minimise impacts on nocturnal wildlife, where possible;
 - Establishment of Ecological Protection Zones (EPZs), through use of protective fencing, to prevent construction activities damaging retained habitats;
 - Creation of EPZs coordinated with standard arboricultural management and tree protection measures;
 - Bird Monitoring Response Strategy (BMRS) monitoring of bird use on retained intertidal habitats and functionally linked land during construction with additional control/mitigation measures enforced if the numbers of birds within the monitored area fall below a certain threshold; and
 - Update ecology surveys prior to commencement of works.



Species-Mitigation Strategies and Licensees

- <u>12.1212.264</u> The EMMF also contains <u>individual feature-specific</u> Mitigation <u>and/or</u> <u>Management</u> Strategies, including measures to address construction impacts, for the following <u>species or species groupsfeatures</u>:
 - Breeding SPAs and wintering birds;
 - Swanscombe Peninsula SSSI (including rare plants, breeding birds and invertebrates)
 - Bats;
 - Dormouse;
 - Water vole;
 - Otters;
 - Harvest mouse;
 - Amphibians;
 - Reptiles;
 - Invertebrates;
 - OtherR rare plants; and
 - Invasive plants.
- 12.1312.265 In the case of certain legally protected species confirmed present on the Kent Project Site where mitigation licensees will be required before development can proceed, including bats, dormice, water vole, and otter, these Mitigation Strategies effectively provide details of the impact avoidance, mitigation and compensation measures which, when taken together with off-site compensatory measures where relevant, necessary towill ensure the Proposed Development is capable of achieving the relevant EPS/development -licensees, and will be used to inform applications for such licenses.

Creation/Enhancement of Functionally Linked Habitat

12.266 Potential indirect effects on several designated sites have been identified which relate to the net loss (and disturbance) of habitat within the Project Site, which is likely to be functionally linked in terms of cited bird species using on-site wetland and intertidal areas within the Project Site for feeding and/or refuge. The designated sites in question are as follows:



- Thames Estuary and Marshes SPA/Ramsar (including South Thames Estuary and Marshes SSSI and Mucking Flats and Marshes SSSI);
- Medway Estuary and Marshes SPA/Ramsar/SSSI;
- West Thurrock Lagoon and Marshes SSSI;
- Inner Thames Marshes SSSI; and
- Tilbury Marshes LWS
- 12.267 Potential negative effects within these designated sites as a consequence of habitat loss within the Project Site will be avoided or mitigated by the creation or enhancement of new habitat outside of the Project Site but within the functional range of the relevant bird species.
- 12.268 Owing to the nature of the potential effects, these off-site measures represent 'avoidance and mitigation' with reference to the mitigation hierarchy sequence, rather than being compensatory measures. This is an important distinction with respect to European Sites, since avoidance and mitigation measures are taken into consideration at the Appropriate Assessment Stage (Stage 2) of the Habitats Regulations Assessment (HRA) process (see Appendix 12.4; Document Reference 6.2.12.4). Similarly, these measures are taken into consideration in this ES Chapter when identifying any residual effects.
- 12.269 Whilst being deemed avoidance and mitigation for the reasons explained above, these measures will be delivered off-site and are therefore detailed within the Ecological Compensation Framework (Appendix 12.10; Document Reference 6.2.12.10) together with a range of other measures proposed to compensate for impacts on on-site features.
- 12.270 The extent of new functionally linked habitat (259.33ha) significantly exceeds the area to be lost directly as a result of the Proposed Development. This will therefore avoid or mitigate the effects of any further reduction in the habitat area which is used by cited bird species for feeding and/or refuge as a result of unavoidable disturbance from construction activities.

General Principles for Offsite Ecological Mitigation

- 12.14 An off-site mitigation strategy will be required to mitigate construction stage effects on some ecological receptors. Off-site ecological mitigation can take two main forms, comprising off-site mitigation land managed to improve biodiversity; or off-site mitigation payments towards an existing or developing ecological enhancement project.
- The final off-site mitigation 'package' may take the form of a combination of both payments and land acquisition and thus the quantum of land areas and payments would be adjusted accordingly.



12,224

12.15 As an off-site mitigation strategy has not yet been finalised, for the purposes of application for development consent, a set of general principles has been set out with regard to the type of land that might be acquired if that is the preferred off-site mitigation option, see Appendix 12.10: General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10).

12.225 Owing to the need to 'ready' the off-site land ahead of mitigation taking place, the associated habitat creation/enhancement works will need to be started or completed/established, during the construction phase. Off-site mitigation is therefore treated as a mitigation measure in respect of construction phase in this assessment.



Table 12.14: Avoidance and Mitigation of Potential Significant Construction Effects

Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures	
Feature	Receptor			
Thames	International	Potential impacts solely on functionally linked resources (outside of the European/Ramsar Site):		
Estuary and		Direct loss of/damage to functionally linked	Landscape Strategy - enhancement of retained wetland	
Marshes		land on the Kent Project Site, potentially used	habitat on Swanscombe Peninsula and creation of new	
SPA/Ramsar,		by cited bird species, totalling a net loss of	saltmarsh, wetland and reedbed habitats, managed via	
and the		15.49ha, including 14.55ha of coastal and	the Ecological Mitigation and Management Framework	
component		floodplain grazing marsh (Botany Marsh West)	(EMMF) (Document Reference 6.2.12.3).	
SSSIs		and 0.94ha of reedbeds (Black Duck Marsh) to		
including		accommodate construction of the Leisure Core.	Creation/Enhancement of 259.33ha Functionally	
South			<u>Linked Habitat Off-site Off-site Ecological Mitigation</u> —	
Thames			delivered in accordance with the General Principles for	
Estuary and			Offsite Ecological Mitigation ECF (Document Reference	
Marshes SSSI			6.2.12.10). , and comprising either off site mitigation	
and Mucking			land managed to improve biodiversity; or financial	
Flats and			contributions towards an existing or developing	
Marshes SSSI			ecological enhancement project (for example a wetland	
			establishment scheme); or a combination of both as part	
			of an overall off site mitigation 'package'.	
		Damage to functionally linked habitats	CEMP - Prevention of hydrological impacts through	
		potentially used by cited bird species from	adherence to an appropriate Surface Water	
		changes to hydrology (as a result of drainage	Management Strategy, as outlined within Chapter 17 –	
		and infill of the CTRL wetland, swamp/wetland	Water Resources and Flood Risk (Document Reference	
		habitat along the east/south east of Black Duck	6.1.17).	
		Marsh and other low-lying seasonally		
		inundated areas); and changes to water and		
		/or sediment quality (either from surface or		
		groundwater discharges from Project Site		



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		including construction waste and pollutants ; or from disruption of contaminated Thames sediments), with potential associated knock-on risk of bioaccumulation.	
		Damage to functionally linked habitats potentially used by cited bird species from changes in air quality including from dust, construction waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.
		Damage to functionally linked habitats potentially used by cited bird species from introduction or proliferation of Invasive Non-Native Species (INNS).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.
		Disturbance (from shipping/ferry movements) giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	CEMP – to include control measures to prevent spread of INNS and INNS impacted soils. Bird Monitoring Response Strategy (BMRS) - monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious construction activities then those disturbance activities will be temporally ceased.



Ecological Importance o Feature Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature Receptor	Disturbance (noise and lighting) associated with construction activities, giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats. Off-site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects. CEMP and Sensitive Lighting Strategy (SLS)Sensitive Lighting Strategy (SLS) - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP – sensitive timing of works causing in excess of 55dB of noise at the river frontage or functionally linked habitat, to take place in summer months, where possible, as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference 6.2.12.3). CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
		Disturbance (human movement or activity), to	CEMP – use of site screening/hoarding; sensitive timing
		cited bird species potentially using functionally	of works within 300m of the river frontage or
		linked habitats, as a result of increased human	functionally linked habitat and visible from that habitat
		presence on the Project Site during	to take place in summer months, where possible; and
		construction.	access by construction workers to estuary front
			restricted within 500m of visible wetland habitats,
			where possible; as detailed in the Bird Mitigation
			Strategy enclosed within EMMF (Document Reference 6.2.12.3).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
	International	Potential impacts solely on functionally linked res	sources (outside of the European/Ramsar Site):



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
Medway		Direct loss of/damage to functionally linked	Landscape Strategy - enhancement of retained wetland
Estuary and		land on the Kent Project Site, potentially used	habitat on Swanscombe Peninsula and creation of new
Marshes		by cited bird species, totalling a net loss of	saltmarsh, wetland and reedbed habitats, managed via
SPA/Ramsar/		15.49ha, including 14.55ha of coastal and	the EMMF (Document Reference 6.2.12.3).
SSSI ⁸³		floodplain grazing marsh (Botany Marsh West)	
		and 0.94ha of reedbeds (Black Duck Marsh) to	Creation/Enhancement of 259.33ha Functionally
		accommodate construction of the Leisure Core	<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10). Off-site
			Ecological Mitigation – delivered in accordance with the
			General Principles for Offsite Ecological Mitigation.
		Damage to functionally linked habitats	CEMP - Prevention of hydrological impacts through
		potentially used by cited bird species from	adherence to an appropriate Surface Water
		changes to hydrology (as a result of drainage	Management Strategy, as outlined within Chapter 17 –
		and infill of the CTRL wetland, swamp/wetland	Water Resources and Flood Risk (Document Reference
		habitat along the east/south east of Black Duck	6.1.17).
		Marsh and other low-lying seasonally	
		inundated areas); and changes to water and	
		/or sediment quality (either from surface or	
		groundwater discharges from Project Site	
		including construction waste and pollutants;	
		or from disruption of contaminated Thames	
		sediments), with potential associated knock-on	
		risk of bioaccumulation.	
		Damage to functionally linked habitats	CEMP – including details on safe/appropriate disposal of
		potentially used by cited bird species from	construction waste, and pollution prevention guidelines,

⁸³ Medway Estuary and Marshes SSSI is a component SSSI to the Medway Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		changes in air quality including from dust,	to be strictly adhered to. CEMP will also include a Dust
		construction waste and pollutants.	Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/ maintenance, site
			monitoring, operation of vehicles/ machinery/ power
			tools, and techniques for demolition works, earthworks
			and construction.
		Damage to functionally linked habitats	Non-native Invasive Plant Species Mitigation Strategy
		potentially used by cited bird species from	included within the EMMF (Document Reference
		introduction or proliferation of Invasive Non- Native Species (INNS) INNS.	6.2.12.3) – including details on control/eradication of
		Native species (INNS) INNS.	existing populations of non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Disturbance (from shipping/ferry movements)	Bird Monitoring Response Strategy (BMRS)BMRS - to
		giving rise to displacement, behavioural	monitor bird activity and inform the development of
		changes or physiological stress to cited bird species potentially using functionally linked	further control/mitigation measures as required.
		habitats.	Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats. Off-site
			Ecological Mitigation – delivered in accordance with the
			'General Principles for Offsite Ecological Mitigation'
			(Document Reference 6.2.12.10), to offset any residual
			effects.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	Receptor	Disturbance (noise and lighting) associated	CEMP and SLS - Restricted working hours and sensitive
		with increased shipping traffic and construction	design and location of new lighting to minimise impacts
		activities, giving rise to displacement,	on nocturnal/crepuscular wildlife.
		behavioural changes or physiological stress to	
		cited bird species potentially using functionally	CEMP – sensitive timing of works causing in excess of
		linked habitats.	55dB of noise at the river frontage or functionally linked
			habitat, to take place in summer months, where
			possible, as detailed in the Bird Mitigation Strategy
			enclosed within EMMF (Document Reference 6.2.12.3).
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as required.
			required.
			Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
		Disturbance (human movement or activity) , to	CEMP – use of site screening/hoarding; sensitive timing
		cited bird species potentially using functionally	of works within 300m of the the river frontage or
		linked habitats, as a result of increased human	functionally linked habitat and visible from that habitat
		presence on the Project Site during	to take place in summer months, where possible; and
		construction.	access by construction workers to estuary front



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			restricted within 500m of visible wetland habitats,
			where possible; as detailed in the Bird Mitigation
			Strategy enclosed within the EMMF (Document
			Reference 6.2.12.3).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
Swanscombe	<u>National</u>	Overall reduction in SSSI area, and	Landscape Strategy and Swanscombe Peninsula SSSI
<u>Peninsula</u>		fragmentation by the Resort.	Mitigation Strategy – maintenance of connectivity
<u>SSSI</u>			across the Peninsula through the inclusion of a chain of
			watercourses and wetland areas wrapping around the
			side of the Proposed Development footprint,
			maintenance of scrub habitats across the Peninsula to
			allow terrestrial bird species to disperse between
			habitats and maintenance of unrestricted flight path
			from River Thames into Black Duck Marsh via retaining
			and managing low-growing vegetation at northern edge
			of marsh. Creation of new wetland and saltmarsh
			habitats on Swanscombe Peninsula, delivered through
			the EMMF and Landscape Strategy.
		Damage to habitats within the designated area	CEMP – including details on safe/appropriate disposal of
		from changes in air quality, including from	construction waste, and pollution prevention guidelines,



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		dust, construction waste and pollutants, and	to be strictly adhered to. CEMP will also include a Dust
		exhaust emissions.	Management Plan (DMP) to provide measures to control
			dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/ maintenance, site
			monitoring, operation of vehicles/ machinery/ power
			tools, and techniques for demolition works, earthworks
			and construction.
		Special feature of interest - populations of rare p	lants ⁸⁴ :
		Direct loss of individual plants, and Direct loss	Landscape Strategy - enhancement of retained habitats
		of individual plants/supporting habitat, and	on Swanscombe Peninsula and creation of new habitats,
		part of plant meta-population through loss of	managed via the Swanscombe Peninsula SSSI
		habitat within SSSI.	Mitigation Strategy SSSI Mitigation Strategy.
			Swanscombe Peninsula SSSI Mitigation Strategy (Rare
			Plants)- Mitigation Strategy enclosed within the SSSI
			Mitigation Strategy - providing details and subsequent
			detailed strategies -for the translocation of nationally
			scarce plants/seed bank occurring on the Kent Project
			Site.
		Damage to retained plants from construction	CEMP – including details on safe/appropriate disposal of
		dust, waste and pollutants.	construction waste, and pollution prevention guidelines,

Includes only those rare plant populations listed as a reason for notification of the Swanscombe Peninsula SSSI, including divided sedge (*Carex divisa*), yellow vetchling (*Lathyrus aphaca*), slender hare's-ear (*Bupleurum tenuissimum*), Bithynian vetch (*Vicia bithynica*) and round-leaved wintergreen (*Pyrola rotundifolia subsp. maritima*). A separate assessment of potentially significant effects upon 'non-SSSI' rare plants is provided under the 'rare plants' IEF.



Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Receptor		
		to be strictly adhered to. CEMP will also include a Dust
		Management Plan (DMP) to provide measures to control
		dust and other emissions.
	Damage to individual plantsDamage to	CEMP and Swanscombe Peninsula SSSI Mitigation
	individual plants/supporting habitat from	Strategy (Rare Plants) SSSI Mitigation Strategy –
	adjacent construction works and deposition of	detailing sensitive construction methodology, including
	construction materials.	set up and maintenance of EPZs where no construction
		vehicles/works are permitted and where no materials or
		site facilities may be situated.
	Damage to plants/their habitats from	Swanscombe Peninsula SSSI Mitigation Strategy (Rare
	introduction or proliferation of Invasive Non-	Plants) Non-native Invasive Plant Species Mitigation
	Native Species (INNS)INNS.	Strategy included within SSSI Mitigation Strategy
		including details on control/eradication of existing
		populations of non-native species.
	Special feature of interest - breeding bird assemb	blages:
	Direct killing injury or harm to individuals –	CEMP, and Breeding and Winter Bird Mitigation
		Strategy included withinSwanscombe Peninsula SSSI
	during vegetation elearance.	Mitigation Strategy (Breeding Birds) – including
		sensitive timing and methodology for habitat clearance,
		establishment of EPZs to protect retained habitats.
	Direct habitat loss – loss of suitable habitat	Landscape Strategy and Swanscombe Peninsula SSSI
		Mitigation Strategy (Breeding Birds) Breeding Bird
	· ·	Mitigation Strategy included within SSSI Mitigation
	TO GRANGE	Strategy - enhancement of retained scrub habitat
		through rotational management to create age class and
		structural diversity, and maintenance of site wide open
	•	Damage to individual plantsDamage to individual plants/supporting habitat from adjacent construction works and deposition of construction materials. Damage to plants/their habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			mosaic habitats to provide variety of foraging resources,
			managed via the EMMF.
		Damage to supporting habitats from changes	CEMP - Prevention of hydrological impacts through
		to hydrological regime and changes to water	adherence to an appropriate Surface Water
		and/or sediment quality.	Management Strategy, as outlined within Chapter 17 –
			Water Resources and Flood Risk (Document Reference
			<u>6.1.17).</u>
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to supporting habitats from	Swanscombe Peninsula SSSI Mitigation StrategyNon-
		introduction or proliferation of Invasive Non-	native Invasive Plant Species Mitigation Strategy
		Native Species (INNS)INNS.	included within SSSI Mitigation Strategy – including
			details on control/eradication of existing populations of
			non-native species.
		Habitat fragmentation, loss of flight paths and	Landscape Strategy and Swanscombe Peninsula SSSI
		dispersal routes.	Mitigation Strategy (Breeding Birds) SSSI Mitigation
			Strategy - maintenance of connectivity across the
			Peninsula through the inclusion of a chain of
			watercourses and wetland areas wrapping around the
			side of the Proposed Development footprint,
			maintenance of scrub habitats across the Peninsula to
			allow terrestrial bird species to disperse between
			habitats and maintenance of unrestricted flight path
			from River Thames into Black Duck Marsh via retaining



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	•		and managing low-growing vegetation at northern edge of marsh. Creation of new wetland and saltmarsh
			habitats on Swanscombe Peninsula, delivered through
			the EMMF and Landscape Strategy.
		Disturbance (from shipping/ferry movements)	Bird Monitoring Response Strategy (BMRS)BMRS -
		<u>– as described above in relation to impacts on</u> <u>habitats functionally linked to-</u> SPA/Ramsar	monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds
		sites.	within the monitored area fall below a certain threshold
			in response to obvious construction activities then those disturbance activities will be temporally ceased.
			Off-site Ecological Mitigation — delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.
		Disturbance (noise and lighting) – as described	CEMP and Sensitive Lighting Strategy (SLS)SLS -
		above in relation to impacts on habitats	Restricted working hours and sensitive design and
		functionally linked to SPA/Ramsar sites.	location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.
			CEMP – sensitive timing of works causing in excess of
			55dB of noise at the river frontage or functionally linked
			habitat, to take place in summer months, where
			possible, as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference
			6.2.12.3).



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
		Disturbance (human movement or activity) –	CEMP – use of site screening/hoarding; sensitive timing
		as described above in relation to impacts on	of works within 300m of the river frontage or
		SPA/Ramsar sites.	functionally linked habitat and visible from that habitat
			to take place in summer months, where possible; and
			access by construction workers to river front restricted
			within 500m of visible wetland habitats, where possible;
			as detailed in the Bird Mitigation Strategy enclosed
			within the SSSI Mitigation Strategy
		Special feature of interest - invertebrate assembl	<u>aqes:</u>
		Direct habitat loss – loss of suitable habitat,	Landscape Strategy, and Swanscombe Peninsula SSSI
		particularly Open Mosaic Habitat on Previously	Mitigation Strategy (Invertebrates) Mitigation Strategy
		Developed Land.	included in the SSSI Mitigation Strategy – enhancement
			of existing retained open mosaic habitat (OMH) to
			better quality and creation of new brown roofs as
			described previously. Furthermore, enhancement of
			retained wetland habitat within Botany Marsh East and
			Black Duck Marsh through reduction of scrub



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	Receptor		encroachment, selective ditch re-profiling, and creation
			of additional scrapes and deep areas in Black Duck
			Marsh. Creation of additional saltmarsh habitat on the
			north eastern edge of Swanscombe Peninsula through
			'retiring' the flood defence. Creation of a number of
			habitats of value to invertebrates within the amenity
			spaces within the resort itself, including native tree and
			shrub planting, wildflower strips, green roofs and walls
			and 'bug hotels'.
			Off-site Ecological Mitigation – delivered in accordance
			with the General Principles for Offsite Ecological
			Mitigation (Document Reference 6.2.12.10) will also
			create additional valuable invertebrate habitat.
		Damage to habitats from changes to	CEMP - Prevention of hydrological impacts through
		hydrological regime and changes to water	adherence to an appropriate Surface Water
		and/or sediment quality.	Management Strategy, as outlined within Chapter 17 –
			Water Resources and Flood Risk (Document Reference
			<u>6.1.17).</u>
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to supporting habitats from	Swanscombe Peninsula SSSI Mitigation StrategyNon-
		introduction or proliferation of Invasive Non-	native Invasive Plant Species Mitigation Strategy
		Native Species (INNS)INNS.	included within the SSSI Mitigation Strategy – including



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
- Catal C	песерио		details on control/eradication of existing populations of
			non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Habitat fragmentation.	Landscape Strategy and Swanscombe Peninsula SSSI
			Mitigation Strategy (Invertebrates)SSSI Mitigation
			Strategy – retention and enhancement of large area of
			open mosaic habitat on the Kent Project Site, connected
			to existing semi-natural habitats.
		Disturbance (lighting) – as described above in	CEMP and SLS – Restricted working hours and sensitive
		relation to impacts on SPA/Ramsar sites.	design and location of new lighting to minimise impacts
			on nocturnal/crepuscular wildlife.
<u>Darenth</u>	<u>National</u>	Damage to habitats within the designated area	CEMP – including details on safe/appropriate disposal of
Woods SSSI		from changes in air quality, including from	construction waste, and pollution prevention guidelines,
		dust, construction waste and pollutants, and	to be strictly adhered to. CEMP will also include a Dust
		exhaust emissions.	Management Plan (DMP) to provide measures to control
			dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/-maintenance, site
			monitoring, operation of vehicles/-machinery/-power
			tools, and techniques for demolition works, earthworks
			and construction.
Inner Thames	<u>National</u>	Potential impacts solely on functionally linked res	sources (outside of the SSSI):
Marshes SSSI		Direct loss of/damage to functionally linked	Landscape Strategy - enhancement of retained wetland
		habitats on the Kent Project Site, potentially	habitat on Swanscombe Peninsula and creation of new
		used by cited bird species, totalling a net loss	



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor	of 45 40ha including 44 55ha of coastal and	
		of 15.49ha, including 14.55ha of coastal and	saltmarsh, wetland and reedbed habitats, managed via
		floodplain grazing marsh (Botany Marsh West)	the EMMF (Document Reference 6.2.12.3).
		and 0.94ha of reedbeds (Black Duck Marsh) to	0
		accommodate construction of the Leisure Core.	Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance with
			the ECF (Document Reference 6.2.12.10)
		Damage to functionally linked habitats	CEMP - Prevention of hydrological impacts through
		potentially used by cited bird species from	adherence to an appropriate Surface Water
		changes to hydrology (as a result of drainage	Management Strategy, as outlined within Chapter 17 –
		and infill of the CTRL wetland, swamp/wetland	Water Resources and Flood Risk (Document Reference
		habitat along the east/south east of Black Duck	<u>6.1.17).</u>
		Marsh and other low-lying seasonally	
		inundated areas); and changes to water and	
		/or sediment quality (either from surface or	
		groundwater discharges from Project Site	
		including construction waste and pollutants;	
		or from disruption of contaminated Thames	
		sediments), with potential associated knock-on	
		risk of bioaccumulation.	
		Damage to functionally linked habitats	CEMP – including details on safe/appropriate disposal of
		potentially used by cited bird species from	construction waste, and pollution prevention guidelines,
		changes in air quality including from dust,	to be strictly adhered to. CEMP will also include a Dust
		construction waste and pollutants.	Management Plan (DMP) to provide measures to control
			dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/-maintenance, site
			monitoring, operation of vehicles/-machinery/-power



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			tools, and techniques for demolition works, earthworks
			and construction.
		Damage to functionally linked habitats	Non-native Invasive Plant Species Mitigation Strategy
		potentially used by cited bird species from	included within the EMMF (Document Reference
		introduction or proliferation of INNS.	6.2.12.3) – including details on control/eradication of
			existing populations of non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Disturbance (from shipping/ferry movements)	BMRS - to monitor bird activity and inform the
		giving rise to displacement, behavioural	development of further control/mitigation measures as
		changes or physiological stress to cited bird	required.
		species potentially using functionally linked	
		habitats.	Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats
		Disturbance (noise and lighting) associated	CEMP and SLS - Restricted working hours and sensitive
		with increased shipping traffic and construction	design and location of new lighting to minimise impacts
		activities, giving rise to displacement,	on nocturnal/crepuscular wildlife.
		behavioural changes or physiological stress to	
		cited species potentially using functionally	CEMP – sensitive timing of works causing in excess of
		linked habitats.	55dB of noise at the river frontage or functionally linked
			habitat, to take place in summer months, where
			possible, as detailed in the Bird Mitigation Strategy
			enclosed within EMMF (Document Reference 6.2.12.3).



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats
		Disturbance (human movement or activity), to	CEMP – use of site screening/hoarding; sensitive timing
		cited bird species potentially using functionally	of works within 300m of the the river frontage or
		linked habitats, as a result of increased human	functionally linked habitat and visible from that habitat
		presence on the Project Site during	to take place in summer months, where possible; and
		construction.	access by construction workers to estuary front
		<u>construction.</u>	restricted within 500m of visible wetland habitats,
			where possible; as detailed in the Bird Mitigation
			Strategy enclosed within the EMMF (Document
			Reference 6.2.12.3).
			Meterence 0.2.12.3).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			_



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
West	National	Potential impacts solely on functionally linked res	sources (outside of the SSSI):
Thurrock		Direct loss of/damage to functionally linked	Landscape Strategy - enhancement of retained wetland
Lagoon and		habitats on the Kent Project Site, potentially	habitat on Swanscombe Peninsula and creation of new
Marshes SSSI		used by cited bird species, totalling a net loss	saltmarsh, wetland and reedbed habitats, managed via
		of 15.49ha, including 14.55ha of coastal and	the EMMF (Document Reference 6.2.12.3).
		floodplain grazing marsh (Botany Marsh West)	
		and 0.94ha of reedbeds (CTRL wetland and	Creation/Enhancement of 259.33ha Functionally
		Black Duck Marsh) to accommodate	<u>Linked Habitat Off-site</u> – delivered in accordance with
		construction of the Leisure Core.	the ECF (Document Reference 6.2.12.10). Off-site
			Ecological Mitigation – delivered in accordance with the
			General Principles for Offsite Ecological Mitigation.
		Damage to functionally linked habitats	CEMP — Prevention of hydrological impacts through
		potentially used by cited bird species from	adherence to an appropriate Surface Water
		changes to hydrology (as a result of drainage	Management Strategy, as outlined within Chapter 17 –
		and infill of the CTRL wetland, swamp/wetland	Water Resources and Flood Risk (Document Reference
		habitat along the east/south east of Black Duck	6.1.17).
		Marsh and other low-lying seasonally	
		inundated areas); and changes to water and	
		/or sediment quality (either from surface or	
		groundwater discharges from Project Site	
		including construction waste and pollutants;	
		or from disruption of contaminated Thames	
		sediments), with potential associated knock-on	
		risk of bioaccumulation.	



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	·	Damage to functionally linked habitats	CEMP – including details on safe/appropriate disposal of
		potentially used by cited bird species from	construction waste, and pollution prevention guidelines,
		changes in air quality including from dust,	to be strictly adhered to. CEMP will also include a Dust
		construction waste and pollutants.	Management Plan (DMP) to provide measures to control
			dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/ maintenance, site
			monitoring, operation of vehicles/ machinery/ power
			tools, and techniques for demolition works, earthworks
			and construction.
		Damage to functionally linked habitats	Non-native Invasive Plant Species Mitigation Strategy
		potentially used by cited bird species from	included within the EMMF (Document Reference
		introduction or proliferation of Invasive Non-	6.2.12.3) – including details on control/eradication of
		Native Species (INNS)INNS.	existing populations of non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Disturbance (from shipping/ferry movements)	Bird Monitoring Response Strategy (BMRS)BMRS — to
		giving rise to displacement, behavioural	monitor bird activity and inform the development of
		changes or physiological stress to cited bird	further control/mitigation measures as required.
		species potentially using functionally linked habitats.	
		nuoreacs.	Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats Off-site
			Ecological Mitigation – delivered in accordance with the



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			'General Principles for Offsite Ecological Mitigation'
			(Document Reference 6.2.12.10), to offset any residual
			effects.
		Disturbance (noise and lighting) associated	CEMP and SLS - Restricted working hours and sensitive
		with increased shipping traffic and construction	design and location of new lighting to minimise impacts
		activities, giving rise to displacement,	on nocturnal/crepuscular wildlife.
		behavioural changes or physiological stress to	
		cited species potentially using functionally	CEMP – sensitive timing of works causing in excess of
		linked habitats.	55dB of noise at the river frontage or functionally linked
			habitat, to take place in summer months, where
			possible, as detailed in the Bird Mitigation Strategy
			enclosed within EMMF (Document Reference 6.2.12.3).
			CENTR adoution of witingtion was a control
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
		Disturbance (human movement or activity) , to	CEMP – use of site screening/hoarding; sensitive timing
		cited bird species potentially using functionally	of works within 300m of the river frontage or



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		linked habitats, as a result of increased human presence on the Project Site during construction.	functionally linked habitat and visible from that habitat to take place in summer months, where possible; and access by construction workers to estuary front restricted within 500m of visible wetland habitats, where possible; as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference 6.2.12.3).
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
			Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats.
Botany Marshes LWS	County	Damage to habitats from changes in air quality, including from dust, construction waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Damage to habitats from changes to hydrological regime (through destruction of adjoining wetland/grazing marsh), and changes in water quality/quantity.	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Damage to habitats from adjacent construction works and deposition of construction materials.	CEMP and EMMF – detailing sensitive construction methodology, including set up and maintenance of Ecological Protection Zones (EPZs) where no construction vehicles/works are permitted and where no materials or site facilities may be situated.
		Disturbance (noise and lighting) associated with construction activities, giving rise to displacement, behavioural changes or physiological stress to wildlife species within	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.
		the designated area.	CEMP – sensitive timing of works, where possible, as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference 6.2.12.3).
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).
Ebbsfleet Marshes, Northfleet LWS	County	Direct loss of/damage to habitat through Station Quarter South.	Landscape Strategy – Retention of site wide open mosaic of habitats within retained parts of Swanscombe Peninsula, and creation of new constructed wetland, new ponds and ditches, and water vole receptor site, managed through the EMMF (Document Reference 6.2.12.3).



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	nesceptor	Damage to habitats from changes in air quality, including from dust, construction waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.
		Damage to habitats from changes to hydrological regime (through destruction of habitat within floodzone of River Ebbsfleet), and changes in water quality/quantity.	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Damage to habitats from adjacent construction works and deposition of construction materials.	CEMP and EMMF – detailing sensitive construction methodology, including set up and maintenance of EPZs where no construction vehicles/works are permitted and where no materials or site facilities may be situated.
		Disturbance (noise and lighting) associated with construction activities, giving rise to displacement, behavioural changes or physiological stress to wildlife species within the designated area.	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP – sensitive timing of works, where possible, as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference 6.2.12.3).



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
Alkerden	County	Loss of part of plant and invertebrate meta-	Landscape Strategy - enhancement of retained habitats
Lane Pit LWS		population through loss of habitat in the Kent	on Swanscombe Peninsula and creation of new habitats,
		Project Site.	managed via the EMMF (Document Reference 6.2.12.3).
			EMMF and subsequent detailed strategies Rare Plant
			Mitigation Strategy enclosed within the EMMF -
			providing details for the translocation of nationally
			scarce plants/seed bank occurring on the Kent Project
			Site.
		Damage to habitats from changes in air	CEMP – including details on safe/appropriate disposal of
		quality, including from dust, construction	construction waste, and pollution prevention guidelines,
		waste and pollutants.	to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/ maintenance, site
			monitoring, operation of vehicles/ machinery/ power
			tools, and techniques for demolition works, earthworks
			and construction.
Tilbury	County	Direct loss of/damage to functionally linked	<u>Landscape Strategy</u> - enhancement of retained wetland
Marshes LWS		habitats on the Kent Project Site, potentially	habitat on Swanscombe Peninsula and creation of new
		used by bird species, totalling a net loss of	saltmarsh, wetland and reedbed habitats, managed via
		15.49ha, including 14.55ha of coastal and	the EMMF (Document Reference 6.2.12.3).



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		floodplain grazing marsh (Botany Marsh West)	
		and 0.94ha of reedbeds (Black Duck Marsh) to	Creation/Enhancement of 259.33ha Functionally
		accommodate construction of the Leisure	<u>Linked Habitat Off-site</u> – delivered in accordance with
		Core. Disturbance (from shipping/ferry	the ECF (Document Reference 6.2.12.10). Bird
		movements) giving rise to displacement,	Monitoring Response Strategy (BMRS) to monitor bird
		behavioural changes or physiological stress to	activity and inform the development of further
		bird species potentially using functionally linked habitats.	control/mitigation measures as required.
			Off-site Ecological Mitigation – delivered in accordance
			with the 'General Principles for Offsite Ecological
			Mitigation' (Document Reference 6.2.12.10), to offset
			any residual effects.
		Disturbance (from shipping/ferry movements)	BMRS - to monitor bird activity and inform the
		giving rise to displacement, behavioural	development of further control/mitigation measures as
		changes or physiological stress to bird species	required.
		potentially using functionally linked habitats.	
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
		Disturbance (noise and lighting) associated	CEMP and SLS - Restricted working hours and sensitive
		with increased shipping traffic and construction	design and location of new lighting to minimise impacts
		activities, giving rise to displacement,	on nocturnal/crepuscular wildlife.
		behavioural changes or physiological stress to	
		bird species potentially using functionally	CEMP – sensitive timing of works causing in excess of
		linked habitats.	55dB of noise at the river frontage or functionally linked
			habitat, to take place in summer months, where



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			possible, as detailed in the Bird Mitigation Strategy
			enclosed within EMMF (Document Reference 6.2.12.3).
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measures as
			required.
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance with
			the ECF (Document Reference 6.2.12.10) to address any
			residual disturbance effects on retained habitats.
		Disturbance (human movement or activity) , to	CEMP – use of site screening/hoarding; sensitive timing
		cited bird species potentially using functionally	of works within 300m of the river frontage or
		linked habitats, as a result of increased human	functionally linked habitat and visible from that habitat
		presence on the Project Site during	to take place in summer months, where possible; and
		construction.	access by construction workers to estuary front
			restricted within 500m of visible wetland habitats,
			where possible; as detailed in the Bird Mitigation
			Strategy enclosed within the EMMF (Document
			Reference 6.2.12.3).



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
			Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats
Habitats/Flora	1		
Populations of nationally scarce plants	National	N.B. Mitigation of potential effects upon scarce plants included under Swanscombe Peninsula SSSI Direct loss of individual plants, and part of plant meta-population through loss of habitat	Landscape Strategy - enhancement of retained habitats on Swanscombe Peninsula and creation of new habitats, managed via the EMMF (Document Reference 6.2.12.3).
		in the Kent Project Site.	Rare Plant Mitigation Strategy enclosed within the EMMF – providing details for the translocation of nationally scarce plants/seed bank occurring on the Kent Project Site.
Broad leaved Semi Natural Woodland	Local	Direct loss of habitat from woodland south of Black Duck Marsh – 3.84ha lost from a total of 21.37ha (18% of the existing total).	Landscape Strategy - enhancement of retained habitats on Swanscombe Peninsula and creation of new habitats, managed via the EMMF (Document Reference 6.2.12.3). Off-site Ecological Mitigation — delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10).
		Damage to habitat from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
Ancient	County	Damage to habitats within the designated area	CEMP – including details on safe/appropriate disposal of
woodland		from changes in air quality, including from	construction waste, and pollution prevention guidelines,
		dust, construction waste and pollutants, and	to be strictly adhered to. CEMP will also include a Dust
		exhaust emissions.	Management Plan (DMP) to provide measures to control
			dust and other emissions. The CEMP will secure a suite
			of construction mitigation measures as detailed in
			Chapter 16: Air Quality (Document reference 6.1.16),
			relating to site management/ maintenance, site
			monitoring, operation of vehicles/ machinery/ power
			tools, and techniques for demolition works, earthworks
			and construction.
Scrub	Local	Direct loss of habitat, a total of 34.79ha lost	Landscape Strategy and EMMF –. enhancement of
		(41.71% of the existing total Direct loss of	retained scrub habitats and commencement of
		habitat, a total of 45.65ha lost (46.99% of the	rotational management to avoid excessive
		existing total), including: 38.70ha of	encroachment and to maintain varied structure and
		dense/continuous scrub (46% of the existing	open mosaic habitat.
		total), and 7.17ha of scattered scrub (taken as	
		50% of the total area measured as	Off-site Ecological Mitigation – delivered in accordance
		grassland/scrub mosaic, 57.08% of the existing	with the 'General Principles for Offsite Ecological
		total). 4.10ha of lost scrub is also included	Mitigation' (Document Reference 6.2.12.10), to offset
		under the OMHPDL grouping and 0.57ha under	any residual effects.
		Floodplain Wetland Mosaic, within the	
		biodiversity net gain calculations (Document	
		Reference 6.2.12.2).	



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Damage to habitats from adjacent construction works and deposition of construction materials in Root Protection Area (RPA). Damage to habitat from construction dust, waste and pollutants.	CEMP and EMMF – detailing sensitive construction methodology, including set up and maintenance of EPZs where no construction vehicles/works are permitted and where no materials or site facilities may be situated. CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
Semi- improved grassland	Local to District	Direct loss of habitat, a total of 34.79ha lost (41.71% of the existing total Direct loss of habitat, 42.19ha lost (48.42% of the existing total) 28.23ha of lost semi-improved grassland is also included under the OMHPDL grouping and 1.54ha under Floodplain Wetland Mosaic, within the biodiversity net gain calculations (Document reference 6.2, check (6.2.12.2)	Landscape Strategy and EMMF – management of existing, retained habitats on Swanscombe Peninsula through commencement of rotational management to avoid excessive scrub encroachment and to maintain varied structure and open mosaic habitat, and create new opportunities for the development of species-rich wildflower grassland.
		Appendix 12.2).	Off-site Ecological Mitigation — delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Damage to habitats from adjacent construction	CEMP and EMMF – detailing sensitive construction
		works and deposition of construction materials.	methodology, including set up and maintenance of EPZs
			where no construction vehicles/works are permitted
			and where no materials or site facilities may be situated.
		Damage to habitat from construction dust,	CEMP – including details on safe/appropriate disposal of
		waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to habitats from introduction or	Non-native Invasive Plant Species Mitigation Strategy
		proliferation of Invasive Non-Native Species	included within EMMF (Document Reference 6.2.12.3) –
		(INNS)INNS.	including details on control/eradication of existing
			populations of non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
Coastal/flood	District	Direct loss of habitat, 13.74ha lost (100% of	No on-site measures possible/proposed. Off-site
plain grazing	District	the existing 11.99ha lost (95.08% of the existing	Ecological Mitigation – delivered in accordance with the
marsh		total). All CFGM is also included under the	'General Principles for Offsite Ecological Mitigation'
		Floodplain Wetland Mosaic grouping, within	(Document Reference 6.2.12.10).
		the biodiversity net gain	, ,
		calculations biodiversity impact calculations	
		(Document reference 6.2, Appendix 12.2).	
Open mosaic	District	Direct loss of habitat, 39.83ha lost (63.7% of	Landscape Strategy, and Invertebrate Swanscombe
on previously		the existing total), includes a range of individual	Peninsula SSSI Mitigation Strategy Mitigation Strategy
developed		habitat components including areas of bare	included in the EMMF – enhancement of existing
land		ground, ephemeral vegetation, open grassland	retained open mosaic habitat (OMH) through
		and patches of scrub. Direct loss of habitat,	introduction of greater variety and complexity,



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		55.13ha lost (76.72% of the existing total), includes a range of individual habitat components including areas of bare ground, ephemeral vegetation, open grassland and patches of scrub.	maintaining a range of microhabitats, foodplants and nectar sources. Measures include creation of bare ground scrapes, creation of shallow pools of varying depth, creation of piles/mounds of mixed crushed and coarse concrete rubble, and creation of chalk mounds and low bunds. Furthermore, creation of brown roof habitat on new buildings using substrate (crushed concrete and chalk) originating from the Proposed Development footprint.
			Off-site Ecological Mitigation – delivered in accordance with the General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10), will also create additional valuable invertebrate habitat.
		Damage to habitats from adjacent construction	CEMP and EMMF – detailing sensitive construction
		works and deposition of construction materials.	methodology, including set up and maintenance of EPZs where no construction vehicles/works are permitted and where no materials or site facilities may be situated.
		Damage to habitat from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
Waterbodies	District	Direct loss of habitat , 2.30ha of waterbodies	Landscape Strategy - Creation of new
(ponds,		lost (50.79% of the existing total) and 4.7km of	waterbodies/wetlands on Swanscombe Peninsula,
standing		ditches (50.98% of the existing total). 0.22ha of	including new perimeter swaleand_new constructed
water and		lost waterbodies are also included under the	wetland,, new ponds and ditches, and water vole
ditch		Floodplain Wetland Mosaic grouping, within	receptor site, managed through the EMMF (Document
network)		the biodiversity net gain	Reference 6.2.12.3).
		calculations biodiversity impact calculations	
		(Document reference 6.2 <u>.12.2</u> , Appendix 12.2).	Off-site Ecological Mitigation – delivered in accordance
			with the General Principles for Offsite Ecological
			Mitigation (Document Reference 6.2.12.10).
		Damage to habitats from changes to	CEMP - Prevention of hydrological impacts through
		hydrological regime (through destruction of	adherence to an appropriate Surface Water
		surrounding habitats), and changes in water	Management Strategy, as outlined within Chapter 17 –
		quality/quantity.	Water Resources and Flood Risk (Document Reference
			6.1.17).
		Damage to habitat from construction dust,	CEMP – including details on safe/appropriate disposal of
		waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to habitats from introduction or	Non-native Invasive Plant Species Mitigation Strategy
		proliferation of Invasive Non-Native Species	included within the EMMF (Document Reference
		(INNS) INNS.	6.2.12.3) – including details on control/eradication of
			existing populations of non-native species.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
Swamp (reedbed)	County	Direct loss of habitat from 13.02ha lost (42.99% of total area). 2.84ha of reedbed is also included under the OMHPDL grouping and 0.88ha under Floodplain Wetland Mosaic, within the biodiversity net gain calculations biodiversity impact calculations (Document reference 6.2.12.2, Appendix 12.2).	Landscape Strategy - Creation of new waterbodies/wetlands on Swanscombe Peninsula, including new perimeter swaleand_ new constructed wetland, new ponds and ditches, and water vole receptor site, managed through the EMMF (Document Reference 6.2.12.3).
			Off-site Ecological Mitigation — delivered in accordance with the General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10).
		Damage to habitats from adjacent construction works and deposition of construction materials.	CEMP and EMMF – detailing sensitive construction methodology, including set up and maintenance of EPZs where no construction vehicles/works are permitted and where no materials or site facilities may be situated.
		Damage to habitats from changes to hydrological regime (through destruction of surrounding habitats), and changes in water quality/quantity.	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Damage to habitat from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
River Ebbsfleet	Local	Direct loss of connected floodplain habitat within Ebbsfleet Valley, such as waterbodies and semi-improved grassland.	Landscape Strategy - Creation of new waterbodies/wetlands on Swanscombe Peninsula, including new perimeter swale and new constructed wetland, new constructed wetland, new ponds and ditches, and water vole receptor site, managed through the EMMF (Document Reference 6.2.12.3). Off-site Ecological Mitigation — delivered in accordance with the General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10).
		Damage to habitat from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.
		Damage to habitats from changes to hydrological regime (through destruction of surrounding habitats), and changes in water quality/quantity.	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).



Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Non-native Invasive Plant Species Mitigation Strategy included within the EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.
		CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
International	N.B. Assessment of potential habitat loss and disturbance effects upon wintering waterfowl and wading bird assemblage included under SPA/Ramsar Sites. Direct killing, injury or harm to individuals — during vegetation clearance. Direct killing, injury or harm to individuals — during vegetation clearance.	CEMP, and Breeding and Winter Bird Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) — including sensitive timing and methodology for habitat clearance, establishment of EPZs to protect retained habitats. CEMP, and SPA and Winter Bird Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) — including sensitive timing and methodology for habitat clearance, establishment of EPZs to protect retained habitats.
County	Direct killing, injury or harm to individuals – during vegetation clearance. Direct habitat loss – loss of suitable habitat such as scrub, grassland, reedbed and woodland, as detailed above.	CEMP ₇ and SPA and Winter Bird Mitigation StrategyBreeding and Winter Bird Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including sensitive timing and methodology for habitat clearance, establishment of EPZs to protect retained habitats. Landscape Strategy and SPA and Winter Bird Mitigation StrategyBreeding and Winter Bird Mitigation Strategy included within EMMF - enhancement of retained scrub
	International	Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS. International N.B. Assessment of potential habitat loss and disturbance effects upon wintering waterfowl and wading bird assemblage included under SPA/Ramsar Sites. Direct killing, injury or harm to individuals — during vegetation clearance. Direct killing, injury or harm to individuals — during vegetation clearance. Direct killing, injury or harm to individuals — during vegetation clearance. Direct habitat loss — loss of suitable habitat such as scrub, grassland, reedbed and



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			class and structural diversity, and maintenance of site
			wide open mosaic habitats to provide variety of foraging
			resources, managed via the EMMF.
			Off-site Ecological Mitigation – delivered in accordance
			with the 'General Principles for Offsite Ecological
			Mitigation' (Document Reference 6.2.12.10), will also
			create/enhance woodland/scrub habitat, with potential
			benefit to a wintering terrestrial bird assemblage in the
			local area.
		Damage to functionally linked habitats from	CEMP - Prevention of hydrological impacts through
		changes to hydrological regime and changes to	adherence to an appropriate Surface Water
		water and/or sediment quality – as described	Management Strategy, as outlined within Chapter 17 –
		above in relation to impacts on SPA/Ramsar	Water Resources and Flood Risk (Document Reference
		sites.	6.1.17).
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to functionally linked habitats from	Non-native Invasive Plant Species Mitigation Strategy
		introduction or proliferation of Invasive Non-	included within EMMF (Document Reference 6.2.12.3) –
		Native Species (INNS)INNS – as described	including details on control/eradication of existing
		above in relation to impacts on SPA/Ramsar	populations of non-native species.
		sites.	CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Habitat fragmentation, loss of flight paths and	Landscape Strategy and EMMF (Document Reference
		dispersal routes.	6.2.12.3)— maintenance of scrub habitats across the



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			Peninsula to allow terrestrial bird species to disperse between habitats.
		Disturbance (from shipping/ferry movements) – as described above in relation to impacts on SPA/Ramsar sites.	Bird Monitoring Response Strategy (BMRS)BMRS - monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious construction activities then those disturbance activities will be temporally ceased.
			Off-site Ecological Mitigation — delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	CEMP and Sensitive Lighting Strategy (SLS)SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.
			CEMP – sensitive timing of works causing in excess of 55dB of noise at the river frontage or functionally linked habitat, to take place in summer months, where possible, as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference 6.2.12.3).
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
		Disturbance (human movement or activity) – as described above in relation to impacts on SPA/Ramsar sites.	CEMP – use of site screening/hoarding; sensitive timing of works within 300m of the river frontage or functionally linked habitat and visible from that habitat to take place in summer months, where possible; and access by construction workers to river front restricted within 500m of visible wetland habitats, where possible; as detailed in the Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3).
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
Breeding Bird Assemblage	National		semblage included under Swanscombe Peninsula SSSI. ding bird assemblage included under Swanscombe
<mark>Breeding Bird</mark> Assemblage	Regional	Direct killing, injury or harm to individuals—during vegetation clearance.	CEMP, and Breeding and Winter Bird Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) — including sensitive timing and methodology for habitat clearance, establishment of EPZs to protect retained habitats.
Bat Assemblage	District County	Direct killing, injury or harm to individuals – during demolition of buildings with bat roost potential and trees with bat roost potential.	CEMP, and Bat Mitigation Strategy included within EMMF (Document Reference 6.2.12.3)— including precommencement surveys of structures/trees to be



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		demolished/felled, and receipt of European Protected Species Mitigation Licence (EPSML), or site registration under the Bat Mitigation Class Licensing (BMCL) scheme, prior to works commencing, with precautionary method of working (as prescribed by the license), strictly adhered to. Establishment of EPZs to protect retained habitats.
		Direct loss of potential roosting habitat – including trees, and buildings and structures with bat roost potential.	Landscape Strategy and Bat Mitigation Strategy included in EMMF Document Reference 6.2.12.3) — creation of new bat roosting habitat, including bat boxes incorporated into new buildings, and into the two new bird watching towers, and enhancement of hibernation habitat within retained tunnels
		Direct loss of foraging, commuting and dispersal habitats - loss of suitable foraging habitat such as scrub, grassland, reedbed and woodland, as detailed above, including potential winter foraging habitat.	Landscape Strategy and EMMF (Document Reference 6.2.12.3) – creation/maintenance of a range of habitats including woodland, scrub, wetlands, waterbodies, reedbeds, saltmarsh and invertebrate rich habitats, collectively will maintain foraging and commuting bat population on site.
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Damage to supporting habitats from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to supporting habitats from	Non-native Invasive Plant Species Mitigation Strategy
		introduction or proliferation of Invasive Non	included within EMMF (Document Reference 6.2.12.3) –
		Native Species (INNS)INNS.	including details on control/eradication of existing
			populations of non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Habitat fragmentation, loss of flight paths and	Landscape Strategy and EMMF (Document Reference
		dispersal routes.	6.2.12.3) – maintenance of connectivity across the
			Peninsula through the inclusion of a chain of
			watercourses and wetland areas wrapping around the
			side of the Proposed Development footprint,
			maintenance of scrub habitats across the Peninsula, and
			retention of habitat link to the River Thames to allow
			bats to disperse between habitats on- and off-site.
		Disturbance (noise and lighting) – including on	CEMP and SLS - Restricted working hours and sensitive
		retained roosts, foraging areas and commuting	design and location of new lighting to minimise impacts
		routes.	on nocturnal/crepuscular wildlife.
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
Dormouse	District	Direct killing, injury or harm to individuals –	CEMP, and Dormouse Mitigation Strategy included
		during vegetation clearance.	within EMMF (Document Reference 6.2.12.3) – including
			pre-commencement surveys of suitable dormouse



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			habitat, and receipt of European Protected Species
			Mitigation Licence (EPSML) prior to habitat clearance
			commencing, with dormouse displacement methodologies and precautionary method of working
			(as prescribed by the license), strictly adhered to.
			Establishment of EPZs to protect retained habitats.
		Direct habitat loss – loss of suitable habitat	Landscape Strategy and Dormouse Mitigation Strategy
		such as scrub and woodland, as detailed above,	included EMMF (Document Reference 6.2.12.3) –
		and within the Dormouse Mitigation Strategy	enhancement of existing scrub habitats to ensure
		within the Ecological Mitigation and	habitats remain suitable for dormice, through rotational
		Management Framework (Document	scrub management to improve structural and age class
		Reference 6.2.12.3).	diversity and maintain variety of food sources
			throughout the dormouse active season.
			Off-site Ecological Mitigation – delivered in accordance
			with the 'General Principles for Offsite Ecological
			Mitigation' (Document Reference 6.2.12.10), to mitigate
			any residual effects.
		Damage to habitats from changes to	CEMP - Prevention of hydrological impacts through
		hydrological regime and changes to water	adherence to an appropriate Surface Water
		and/or sediment quality – as described above	Management Strategy, as outlined within Chapter 17 –
		in relation to impacts on habitats.	Water Resources and Flood Risk (Document Reference
		Daniel Lander and Mark habitate from	6.1.17).
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust
			•
			Management Plan (DMP) to provide measures to control



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species. CEMP – to include control measures to prevent spread
		Habitat fragmentation and loss of dispersal routes.	Landscape Strategy and Dormouse Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) — retention of areas of suitable habitat within Broadness grassland, including existing woodland, tree and scrub habitats; retention and management of dense bramble and low growing scrub and trees along the north western boundaries of Black Duck Marsh, to maintain potential dispersal routes from Swanscombe Peninsula to the south-west; retention and enhancement of a continuous belt of woodland habitat along the southern boundaries of Black Duck Marsh, connecting to additional green corridors proposed along the southern boundary adjacent to Tiltman Avenue, to ensure the continued functioning of existing dispersal routes to valuable off-site habitats to the south-west, including
			the woodland at the Swanscombe Heritage Park; and retention, enhancement and creation of additional woodland habitat alongside the sensitive design of new landscaping around the peripheries of Botany Marsh, necessary to further promote habitat connectivity



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		between Swanscombe Peninsula and habitats across the wider landscape to the south.
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	CEMP and SLS – Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).
Otter	Local	Direct killing, injury or harm to individuals – during vegetation clearance.	CEMP, and Otter Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including precommencement surveys of suitable habitat, sensitive habitat clearance methodology, and establishment of EPZs to protect retained habitats. In the event an otter holt is found within the Proposed Development footprint, a European Protected Species Mitigation Licence (EPSML) will be needed prior to habitat clearance commencing.
		Direct habitat loss – loss of suitable habitat such as reedbed and waterbodies, as detailed above.	Landscape Strategy - Creation of new wetlands on Swanscombe Peninsula, including new constructed wetland, and new ponds and ditches, managed through the EMMF (Document Reference 6.2.12.3). Creation of two new artificial otter holts as described in the Otter Mitigation Strategy included in the EMMF.
		Damage to habitats from changes to hydrological regime and changes to water	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 –



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		and/or sediment quality – as described above	Water Resources and Flood Risk (Document Reference
		in relation to impacts on habitats.	6.1.17).
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to supporting habitats from	Non-native Invasive Plant Species Mitigation Strategy
		introduction or proliferation of Invasive Non-	included within EMMF (Document Reference 6.2.12.3) –
		Native Species (INNS)INNS.	including details on control/eradication of existing
			populations of non-native species.
			CEMP – to include control measures to prevent spread
			of INNS and INNS impacted soils.
		Habitat fragmentation and loss of dispersal	Landscape Strategy and EMMF (Document Reference
		routes.	6.2.12.3) – maintenance of connectivity across the
			Peninsula through the inclusion of a chain of
			watercourses and wetland areas wrapping around the
			side of the Proposed Development footprint. Creation of
			new wetland and saltmarsh habitats on Swanscombe
			Peninsula.
		Disturbance (noise and lighting) – as described	CEMP and SLS – Restricted working hours and sensitive
		above in relation to impacts on SPA/Ramsar	design and location of new lighting to minimise impacts
		sites.	on nocturnal/crepuscular wildlife.
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
Water Vole	Local to	Direct killing, injury or harm to individuals –	CEMP, and Water Vole Mitigation Strategy included
	district	during vegetation clearance.	within EMMF (Document Reference 6.2.12.3) – including
			pre-commencement surveys of suitable habitat,
			establishment of suitable receptor site, and water vole
			relocation under licence from Natural England (granted
			prior to habitat clearance commencing). Establishment
			of EPZs to protect retained habitats.
		Direct habitat loss – including 6.1km of suitable	Landscape Strategy - Creation of new wetlands and
		ditches and bankside habitat and reedbed (as	ditches_ , ponds and waterbodies -on Swanscombe
		detailed above).	Peninsula, as documented within the Water Vole
			Mitigation Strategy included within EMMF (Document
			Reference 6.2.12.3).
		Damage to habitats from changes to	CEMP - Prevention of hydrological impacts through
		hydrological regime and changes to water	adherence to an appropriate Surface Water
		and/or sediment quality – as described above	Management Strategy, as outlined within Chapter 17 –
		in relation to impacts on habitats.	Water Resources and Flood Risk (Document Reference
			6.1.17).
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.
		Damage to supporting habitats from	Non-native Invasive Plant Species Mitigation Strategy
		introduction or proliferation of Invasive Non-	included within EMMF (Document Reference 6.2.12.3) –
		Native Species (INNS)INNS.	including details on control/eradication of existing
			populations of non-native species.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
		Habitat fragmentation and loss of dispersal routes.	Landscape Strategy and EMMF (Document Reference 6.2.12.3) – maintenance of connectivity across the Peninsula through the inclusion of a chain of watercourses and wetland areas wrapping around the side of the Proposed Development footprint. Creation of new wetland and saltmarsh habitats on Swanscombe Peninsula.
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration
Harvest Mouse	Local	Direct killing, injury or harm to individuals – during vegetation clearance.	Mitigation (Document reference 6.2.15.5). CEMP, and Harvest Mouse Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including pre-commencement check of suitable habitat, and precautionary method of working strictly adhered to. Establishment of EPZs to protect retained habitats.
		Direct habitat loss – loss of suitable habitat such as reedbed, semi-improved grassland, scrub and tall ruderal, as detailed above.	Landscape Strategy and Harvest Mouse Mitigation Strategy included within the EMMF (Document Reference 6.2.12.3) – maintenance of site wide open mosaic habitat including a range of habitats of value to harvest mice including rough, tussocky grassland, tall



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	Receptor		riparian/marginal/reedbed habitat with adjacent scrub
			to provide refuge, foraging and breeding opportunities.
		Damage to habitats from changes to	CEMP - Prevention of hydrological impacts through
		hydrological regime and changes to water	adherence to an appropriate Surface Water
		and/or sediment quality – as described above	Management Strategy, as outlined within Chapter 17 –
		in relation to impacts on habitats.	Water Resources and Flood Risk (Document Reference 6.1.17).
		Damage to supporting habitats from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.
		Damage to supporting habitats from	Non-native Invasive Plant Species Mitigation Strategy
		introduction or proliferation of Invasive Non-	included within EMMF (Document Reference 6.2.12.3) –
		Native Species (INNS)INNS.	including details on control/eradication of existing
			populations of non-native species.
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
		Habitat fragmentation and loss of dispersal	Landscape Strategy and EMMF (Document Reference
		routes.	6.2.12.3) – maintenance of connectivity across the
			Peninsula through the inclusion of a chain of
			watercourses and wetland areas wrapping around the
			side of the Proposed Development footprint, and
			maintenance of scrub and rough grassland habitats
			across the Peninsula to allow individuals to disperse
			between habitats.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).
Amphibian assemblage	Local to Delistrict	Direct killing, injury or harm to individuals – during vegetation clearance.	CEMP, and Amphibian Mitigation Strategy included within EMMF – including translocation of amphibians in tandem with reptile translocation, precautionary methods of working including sensitive drain down of any waterbodies to be removed, timing of works, and establishment of EPZs to protect retained habitats.
		Direct habitat loss – loss of suitable habitat such as reedbed and waterbodies, as detailed above.	Landscape Strategy and Amphibian Mitigation Strategy included within EMMF - Creation of new wetlands, ditches, ponds and waterbodies on Swanscombe Peninsula, as described above in relation to water voles and otters, will also mitigate habitat loss impacts to amphibians.
		Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Damage to supporting habitats from construction dust, waste and pollutants.	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	·		Management Plan (DMP) to provide measures to control dust and other emissions.
		Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
		Habitat fragmentation and loss of dispersal routes.	Landscape Strategy and EMMF — maintenance of connectivity across the Peninsula through the inclusion of a chain of watercourses and wetland areas wrapping around the side of the Proposed Development footprint. Creation of new wetland and saltmarsh habitats on Swanscombe Peninsula, delivered through the EMMF and Landscape Strategy.
		Disturbance (noise and lighting) – as described above in relation to impacts on SPA/Ramsar sites.	cemp and sls – Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. cemp – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration
Reptile assemblage	District	Direct killing, injury or harm to individuals – during vegetation clearance.	Mitigation (Document reference 6.2.15.5). CEMP, and Reptile Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including establishment of suitable receptor site and subsequent translocation of reptiles from construction footprint



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			prior to vegetation clearance commencing, and
			precautionary methods of working. Establishment of
			EPZs to protect retained habitats.
			Off-site Ecological Mitigation – delivered in
			accordance with the General Principles for Offsite
			Ecological Mitigation (Document Reference 6.2.12.10).
		Direct habitat loss – loss of suitable habitat	Landscape Strategy and Reptile Mitigation Strategy
		such as semi-improved grassland, reedbed and	included within EMMF (Document Reference 6.2.12.3) –
		tall ruderal, as detailed above.	maintenance of site wide open mosaic habitats on
			Swanscombe Peninsula to maintain a variety of foraging
			resources, breeding, basking, sheltering and hibernation
			habitats. Management of grassland areas to encourage
			development of structurally complex grassland sward
			with substantial 'litter layer' and areas of bare ground.
			Creation of log/brash piles and hibernaculum to provide
			sheltering and hibernating habitats, and grass snake
			breeding sites from piling arisings from grass cutting.
		Damage to habitats from changes to	CEMP - Prevention of hydrological impacts through
		hydrological regime and changes to water	adherence to an appropriate Surface Water
		and/or sediment quality – as described above	Management Strategy, as outlined within Chapter 17 –
		in relation to impacts on habitats.	Water Resources and Flood Risk (Document Reference
			6.1.17).
		Damage to supporting habitats from	CEMP – including details on safe/appropriate disposal of
		construction dust, waste and pollutants.	construction waste, and pollution prevention guidelines,
			to be strictly adhered to. CEMP will also include a Dust
			Management Plan (DMP) to provide measures to control
			dust and other emissions.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Damage to supporting habitats from	Non-native Invasive Plant Species Mitigation Strategy
		introduction or proliferation of Invasive Non-	included within EMMF (Document Reference 6.2.12.3) –
		Native Species (INNS)INNS.	including details on control/eradication of existing
			populations of non-native species.
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.
		Habitat fragmentation and loss of dispersal routes.	Landscape Strategy and EMMF (Document Reference 6.2.12.3) – maintenance of connectivity across the
			Peninsula through the inclusion of a chain of
			watercourses and wetland areas wrapping around the side of the Proposed Development footprint, and
			maintenance of scrub and rough grassland habitats
			across the Peninsula to allow individuals to disperse
			between habitats. Creation of new wetland and
			saltmarsh habitats on Swanscombe Peninsula.
		Disturbance (noise and lighting) – as described	CEMP and SLS - Restricted working hours and sensitive
		above in relation to impacts on SPA/Ramsar	design and location of new lighting to minimise impacts
		sites.	on nocturnal/crepuscular wildlife.
			CEMP – adoption of mitigation measures to control
			construction noise and vibration, as described in
			Appendix 15.5: Construction Noise and Vibration
			Mitigation (Document reference 6.2.15.5).
Invertebrate	National	N.B. Impact assessment for the invertebrate asse	emblage included under Swanscombe Peninsula
assemblage		SSSI. Direct habitat loss — loss of suitable habitat,	particularly Open Mosaic Habitat on Previously
		Developed Land, as detailed above.	



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		N.B. Proposed mitigation measures for the invertebrate assemblage included under Swanscombe	
		Peninsula SSSILandscape Strategy, and Invertebrate Mitigation Strategy included in the EMMF	
		(Document Reference 6.2.12.3) - enhancement	of existing retained open mosaic habitat (OMH) to better
		quality and creation of new brown roofs as described previously. Furthermore, enhancement of retained	
		wetland habitat within Botany Marsh East and Black Duck Marsh through reduction of scrub	
		encroachment, selective ditch re-profiling, and creation of additional scrapes and deep areas in Black Duck	
		Marsh. Creation of additional saltmarsh habitat on the north eastern edge of Swanscombe Peninsula	
		through 'retiring' the flood defence. Creation of a number of habitats of value to invertebrates within the	
		amenity spaces within the resort itself, including native tree and shrub planting, wildflower strips, green	
		roofs and walls and 'bug hotels'.	
		Off-site Ecological Mitigation — delivered in accordance with the General Principles for Offsite Ecological	
		Mitigation (Document Reference 6.2.12.10) will a	also create additional valuable invertebrate habitat.



Post-construction and Operational Measures

<u>12.16</u>12.271 The key mechanisms to <u>deliver avoidance or mitigation of avoid or mitigate</u> potential significant operational effects, <u>compensation where effects cannot be avoided or mitigated</u>, and to deliver ecology enhancements are summarised below. Further details on these measures, including which IEFs they relate to, are provided within **Table 12.15**.

A Landscape Management Plan (LMP)

- 12.1712.272 A Landscape Management Plan (LMP) (ES Appendix 11.8, Document Reference 6.2.11.8) has been prepared in support of the DCO application and should be read in conjunction with the overarching Ecological Mitigation and Management Framework (EMMF) provided as Appendix 12.3 (Document Reference 6.2.12.3). The measures contained within the LMP include (but are not limited to):
 - Prescriptions for the ongoing management, maintenance and monitoring of the IEFs and of those newly created habitats to maximise opportunities for biodiversity enhancement and gain;
 - Management of open spaces for biodiversity, with controlled/restricted recreational use within sensitive areas;
 - Objectives and principles for the long-term management of ecology interests; and
 - Compliance checks and monitoring to ensure the success of the implemented measures against the objectives and principles, with interventions carried out as necessary.

Species Mitigation Strategies and Licensees

<u>12.18</u>12.273 The EMMF also contains individual Mitigation Strategies, including measures to address operational impacts, for the following species or species groups:

- Breeding and wintering birds;
- Bats;
- Dormouse;
- Water vole;
- Otters;
- Harvest mouse;



- Amphibians;
 - •—
- Reptiles;
- Invertebrates; and
- Rare plants

12.1912.274 In the case of certain legally protected species confirmed present on the Kent Project Site where mitigation licensees will be required before development can proceed, including bats, dormice, water vole, and otter, these Mitigation Strategies effectively provide details of the impact avoidance, mitigation and compensation measures necessary to ensure the Proposed Development is capable of achieving the relevant licensees, and will be used to inform applications for such licensees.

Landscape Strategy

<u>12.2012.275</u> A Landscape Strategy has been prepared in support of the DCO application and is provided as Appendix 11.7 (Document Reference 6.2.11.7). This provides the overall landscape and green infrastructure design for the Proposed Development and forms the basis for a fully detailed Soft Landscaping Scheme (SLS) to be prepared post-consent. Together these will provide overarching design principles and full specifications of new habitat creation and the necessary infrastructure to provide controlled/restricted access for recreation in the open space.

General Principles for Off-site Ecological Mitigation

- 12.21 As described previously, an off-site mitigation strategy will be required to mitigate construction stage effects on some ecological receptors. This may include a mitigation 'package' of both off-site mitigation land and financial contributions to ecological enhancement projects, the full details of which are not yet finalised, see Appendix 12.10 General Principles for Off site Ecological Mitigation (Document Reference 6.2.12.10).
- 12.22 The long-term management of any off-site mitigation land is considered a mitigation measure to be delivered throughout the operational phase of the Proposed Development, and therefore discussed further in this section.

Sustainable Drainage System

12.23 12.276 Sustainable Drainage Systems (SuDS) will be incorporated across the Kent Project Site to manage surface water flows and minimise the risk of pollution to the water environment. Appendix 17.2: Surface Water Drainage Strategy (Document reference 6.2.17.1) has been prepared to both avoid or mitigate any potential effects on water quality and flow within sensitive habitats, and to provide suitable hydrological conditions for new wetland habitats of ecological value to become established. Future SuDS design



will adhere to Appendix 17.2: Surface Water Drainage Strategy (Document reference 6.2.17.1).

Sensitive Lighting Strategy (SLS)

12.2412.277 An Artificial Lighting Environmental Impact Assessment (Document reference 6.2.12.11) has been prepared for the Proposed Development. This document, along with the Lighting Statement (Document reference 7.9) sets out a lighting strategy and design principles to avoid/minimise light spill on habitats used by sensitive/nocturnal species, where possible. Collectively, these documents are considered part of the Sensitive Lighting Strategy (SLS)Sensitive Lighting Strategy, as referred to in Table 12.13 below, which provides mitigation for operational effects, and is to be secured as a requirement of the DCO.

<u>12.25</u>12.278 The SLS should be read in conjunction with Figure 12.45: Light Mitigation Strategy for Biodiversity (Document Reference 6.3.12.45).

Contribution to SAMMS

- <u>12.279 Potential indirect effects on effects upon Thames Estuary and Marshes SPA have been</u> <u>identified, which relate to increased recreational disturbance resulting from the new</u> <u>residential dwellings (staff accommodation) associated with the Proposed Development.</u>
- 12.280 As described within the Shadow HRA (Appendix 12.4; Document Reference 6.2.12.4), these effects will be avoided or mitigated through a financial contribution to the Strategic Access Management and Monitoring Strategy (SAMMS), which is in operation across the Thames, Medway and Swale estuaries. This strategy ensures that recreational pressures on the three SPA/Ramsar sites relating to those estuaries is managed sufficiently, through a combination of education, access restrictions, enhancement and ranger presence. This is implemented through financial contributions from developers.

Creation/Enhancement of Functionally Linked Habitat

- 12.281 As described above with respect to mitigation of construction effects, the creation or enhancement of new habitat, outside of the Project Site but within the functional range of relevant bird species, is proposed to mitigate potential effects on relevant statutory designated sites from the loss of functionally linked habitat within the Project Site.
- 12.282 Whilst being deemed avoidance and mitigation for the reasons explained above, these measures will be delivered off-site and are therefore detailed within the Ecological Compensation Framework (Appendix 12.10; Document Reference 6.2.12.10) together with a range of other measures proposed to compensate for impacts on on-site features.
- 12.283 The extent of new functionally linked habitat (259.33ha) significantly the area to be lost directly as a result of the Proposed Development. This will therefore also avoid or mitigate the effects of any further reduction in the habitat area which is used by cited bird species



for feeding and/or refuge as a result of unavoidable disturbance of retained habitats during the operational phase of the development. This is discussed in further detail in respect of European Sites within Appendix 12.4; Document Reference 6.2.12.4).



Table 12.15: Avoidance and Mitigation of Potential Significant Operational Effects-and Compensation/Enhancement Measures

Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
Thames	International	Potential impacts within the European/Ramsar S	<u>ite itself:<mark>Potential impacts solely on functionally</mark></u>
Estuary and		linked resources (outside of the European/Ramsa	<mark>ır Site):</mark>
Marshes		Recreational disturbance of flora and fauna	Contribution to SAMMS – proportionate financial
SPA/Ramsar,		within designated area as a result of increased	contribution to strategic access
and the		residential population (due to residential	management. Sustainable Drainage System (SuDS)
component		accommodation for staff). Damage to	- designed in accordance with best practice
SSSIs		functionally linked habitats potentially used by	guidance as outlined within Chapter 17 – Water
including		cited bird species from changes to hydrology;	Resources and Flood Risk (Document Reference
South		and changes to water and /or sediment	6.1.17).
Thames		quality (from surface or groundwater	
Estuary and		discharges from the Project Site including	Breeding and Winter Bird Mitigation Strategy
Marshes SSSI		operational waste and pollutants)	enclosed within EMMF (Document Reference
and Mucking			6.2.12.3) - updated bird surveys of retained and
Flats and			newly created wetlands and functionally linked
Marshes			habitats in years 3, 5 and 10 post-completion to
SSSI ⁸⁵			ensure habitats continue to support the intended
			habitats and desired flora/fauna.
		Potential impacts solely on functionally linked res	sources (outside of the European/Ramsar Site):
		Damage to functionally linked habitats	Sustainable Drainage System (SuDS) – designed in
		potentially used by cited bird species from	accordance with best practice guidance as outlined
		changes to hydrology; and changes to water	within Chapter 17 – Water Resources and Flood Risk
		and-/or sediment quality (from surface or	(Document Reference 6.1.17).

⁸⁵ South Thames Estuary and Marshes SSSI and Mucking Flats and Marshes SSSI are component SSSIs to the Thames Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	·	groundwater discharges from the Project Site	
		including operational waste and pollutants).	SPA and Winter Bird Mitigation Strategy enclosed
			within EMMF (Document Reference 6.2.12.3) -
			updated bird surveys of retained and newly created
			wetlands and functionally linked habitats in years 3,
			5 and 10 post-completion to ensure habitats
			continue to support the intended habitats and
			desired flora/fauna.
		Disturbance (from increased ferry	Bird Monitoring Response Strategy (BMRS)BMRS -
		movements) associated with the new Thames	monitoring of bird use on the intertidal habitats and
		Clipper river bus service and operational	other functionally linked land – if the numbers of
		deliveries, potentially giving rise to	birds within the monitored area fall below a certain
		displacement, behavioural changes or	threshold in response to obvious operational
		physiological stress to cited bird species	shipping then those disturbance activities will be
		potentially using functionally linked habitats.	temporally ceased.
			Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on
			retained habitats. Off-site Ecological Mitigation —
			delivered in accordance with the 'General Principles
			for Offsite Ecological Mitigation' (Document
			Reference 6.2.12.10), to offset any residual effects.
		Disturbance (noise and lighting) from the	Sensitive Lighting Strategy (SLS)SLS - Sensitive
		operation of the resort rides and attractions	design and location of new lighting to minimise
		and entertainment facilities, giving rise to	impacts on nocturnal/crepuscular wildlife. Retained



Ecological Importance Feature Receptor	of Nature of Impact	Summary of Avoidance/Mitigation Measures
	displacement, behavioural changes or physiological stress to cited bird species using functionally linked habitats within the designated area.	intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone — Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes — Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). BMRS — to monitor bird activity and inform the development of further control/mitigation measures as required. Disturbance control measures — final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15)
		Creation/Enhancement of 259.33ha Functionally



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on retained
			habitats.
		Disturbance (human movement or activity), to	Landscape Strategy, Landscape and Ecology
		cited bird species potentially using functionally	Management Plan (LMP) and EMMF (Document
		linked habitats, as a result of increased human	Reference 6.2.12.3) – Recreational use managed via
		presence on the Project Site during operation	trail network, signage and interpretation boards to
		of the resort rides and attractions and	mitigate against knock-on effects on on-site wildlife.
		entertainment facilities, including increased	Creation of new bird hides at key observation
		recreational use of retained habitats.	locations to reduce visual impact.
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on
			retained habitats.
Medway	International	Potential impacts solely on functionally linked res	
Estuary and		Damage to functionally linked habitats	SuDS – designed in accordance with best practice
Marshes		potentially used by cited bird species from	guidance as outlined within Chapter 17 – Water
SPA/Ramsar/		changes to hydrology; and changes to water	Resources and Flood Risk (Document Reference
SSSI ⁸⁶		and /or sediment quality (from surface or	6.1.17).
		groundwater discharges from the Project Site	
		including operational waste and pollutants)	

⁸⁶ Medway Estuary and Marshes SSSI is a component SSSI to the Medway Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			SPA and Winter Bird Mitigation StrategyBreeding
			and Winter Bird Mitigation Strategy enclosed
			within EMMF (Document Reference 6.2.12.3) -
			updated bird surveys of retained and newly created
			wetlands and functionally linked habitats in years 3,
			5 and 10 post-completion to ensure habitats
			continue to support the intended habitats and
			desired flora/fauna.
		Disturbance (from increased ferry	BMRS – monitoring of bird use on the intertidal
		movements) associated with the new Thames	habitats and other functionally linked land – if the
		Clipper river bus service and operational	numbers of birds within the monitored area fall
		deliveries, potentially giving rise to	below a certain threshold in response to obvious
		displacement, behavioural changes or	operational shipping then those disturbance
		physiological stress to cited bird species	activities will be temporally ceased.
		potentially using functionally linked habitats	
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on
			retained habitats. Off site Ecological Mitigation -
			delivered in accordance with the 'General Principles
			for Offsite Ecological Mitigation' (Document
			Reference 6.2.12.10), to offset any residual effects.
		Disturbance (noise and lighting) from the	SLS - Sensitive design and location of new lighting to
		operation of the resort rides and attractions	minimise impacts on nocturnal/crepuscular wildlife.
		and entertainment facilities, giving rise to	Retained intertidal and marsh habitats within the
		displacement, behavioural changes or	



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		physiological stress to cited bird species within	Kent Project Site to remain in the following
		the designated area.	Environmental Lighting Zones:
			 River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).
			<u>Creation/Enhancement of 259.33ha Functionally</u> <u>Linked Habitat Off-site – delivered in accordance</u>



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on retained
			<u>habitats</u>
		Disturbance (human movement or activity), to	Landscape Strategy, LMP and EMMF (Document
		cited bird species potentially using functionally	Reference 6.2.12.3) – Recreational use managed via
		linked habitats, as a result of increased human	trail network, signage and interpretation boards to
		presence on the Project Site during operation	mitigate against knock-on effects on on-site wildlife.
		of the resort rides and attractions and	Creation of new bird hides at key observation
		entertainment facilities, including increased	locations to reduce visual impact.
		recreational use of retained habitats.	
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on
			retained habitats.
<u>Swanscombe</u>	<u>National</u>	Damage to habitats within the designated area	Landscape Strategy, and LMP, included within and
<u>Peninsula</u>		from traffic generated air quality impacts	SSSI Mitigation Strategy – Recreational use
<u>SSSI</u>		resulting in nutrient enrichment and	managed via trail network, signage and
		acidification of qualifying woodland habitat	interpretation boards to mitigate against knock on
		with subsequent changes in structure and	effects on on site wildlife habitat management and
		diversity. Damage to individual plants, and part	monitoring.
		of plant meta-population, through increased	
		recreational use, potentially resulting in	
		trampling, littering, dog fouling, unauthorised	
		camping/campfires/barbecues.	



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		Special feature of interest - populations of rare p	lants ⁸⁷ :
		Damage to individual plants/supporting	Landscape Strategy, LMP and Swanscombe
		habitat, and part of plant meta-population,	Peninsula SSSI Mitigation Strategy (Rare Plants)
		through increased recreational use, potentially	Recreational use managed via trail network, signage
		resulting in trampling, littering, dog fouling,	and interpretation boards to mitigate against
		unauthorised camping/campfires/barbecues.	knock-on effects on on-site wildlife.
		Damage to individual plantsDamage to	Landscape Strategy, LMP and Landscape Strategy
		individual plants/supporting habitat, and part	and LMP, included within-Swanscombe Peninsula
		of plant meta-population, through	SSSI Mitigation Strategy (Rare Plants) SSSI
		inappropriate land management practices.	Mitigation Strategy – Appropriate management of
			all retained, enhanced or newly created habitats to
			ensure they continue to support the intended flora
			and fauna, and provide long term benefits to
			biodiversity.
		Damage to retained plants/population from	SuDS – designed in accordance with best practice
		changes to hydrological regime and changes in	guidance as outlined within Chapter 17 – Water
		water quality/quantity.	Resources and Flood Risk (Document Reference
			<u>6.1.17).</u>
		Special feature of interest - breeding bird assemb	plages:
		Damage to on-site habitats potentially used by	SuDS – designed in accordance with best practice
		birds from changes to hydrology; and changes	guidance as outlined within Chapter 17 – Water
		to water and-/or sediment quality (from	

Includes only those rare plant populations listed as a reason for notification of the Swanscombe Peninsula SSSI, including divided sedge (*Carex divisa*), yellow vetchling (*Lathyrus aphaca*), slender hare's-ear (*Bupleurum tenuissimum*), Bithynian vetch (*Vicia bithynica*) and round-leaved wintergreen (*Pyrola rotundifolia subsp. maritima*). A separate assessment of potentially significant effects upon 'non-SSSI' rare plants is provided under the 'rare plants' IEF.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	•	surface or groundwater discharges from the	Resources and Flood Risk (Document Reference
		Project Site including operational waste and	6.1.17).
		pollutants).	
			Swanscombe Peninsula SSSI Mitigation Strategy
			(Breeding Birds) Breeding and Winter Bird
			Mitigation Strategy enclosed within SSSI Mitigation
			Strategy – updated bird surveys of retained and
			newly created habitats in years 3, 5 and 10 post-
			completion to ensure habitats continue to support
			the intended habitats and desired flora/fauna.
		<u>Disturbance (noise⁸⁸ and lighting)</u> from the	SLS - Sensitive design and location of new lighting to
		operation of the resort rides and attractions	minimise impacts on nocturnal/crepuscular wildlife
		and entertainment facilities, giving rise to	Retained scrub habitats within the Kent Project Sit
		displacement, behavioural changes or	to remain in the following Environmental Lightin
		physiological stress to bird species using the	Zones:
		on-site habitats.	
			 Environmental Zone E2 (Typical of sparsely
			inhabited rural areas. No lighting sources visible
			from animal habitats. Protect the natural areas
			that are to be conserved and enhanced).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation measure
			as required.

Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			<u>Disturbance control measures</u> – final ride design to
			include deliberate placement of 'scream zones';
			strict noise specification to eliminate clanking;
			building placement to reduce noise spread; and
			controls on location, direction and sound pressure
			level of loudspeaker systems. (See Chapter 15,
			Document reference: 6.1.15)
		Disturbance (human movement or activity), to	Landscape Strategy, LMP and Landscape Strategy
		bird species, as a result of increased human	and LMP, included within-Swanscombe Peninsula
		presence on the Project Site during operation	SSSI Mitigation Strategy (Breeding Birds) SSSI
		of the resort rides and attractions and	Mitigation Strategy – Recreational use managed via
		entertainment facilities, including increased	trail network, signage and interpretation boards to
		recreational use of retained habitats.	mitigate against knock-on effects on on-site wildlife.
			Creation of new bird hides at key observation
			locations to reduce visual impact.
		Special feature of interest - invertebrate assemble	ages:
		Damage to on-site habitats potentially used by	SuDS – designed in accordance with best practice
		invertebrates from changes to hydrology; and	guidance as outlined within Chapter 17 – Water
		changes to water and /or sediment quality	Resources and Flood Risk (Document Reference
		(from surface or groundwater discharges from	<u>6.1.17).</u>
		the Project Site including operational waste	
		and pollutants).	
		Disturbance (lighting) from the operation of	SLS - Sensitive design and location of new lighting to
		the resort rides and attractions and	minimise impacts on wildlife. Retained intertidal
		entertainment facilities.	and marsh habitats within the Kent Project Site to



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			remain in the following Environmental Lighting
			Zones:
			 River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
Darenth	National	Potential damage to habitats within the	If significant effects are identified following further
Woods SSSI		designated area from traffic generated air	investigation, mitigation measures to reduce traffic
		quality impacts – subject to further	emissions will need to be developed and agreed
		investigation to determine the nature of any	with Natural England.
		significant effects.	
Inner Thames	National	Potential impacts solely on functionally linked res	sources (outside of the SSSI):
Marshes SSSI		Damage to functionally linked habitats	SuDS – designed in accordance with best practice
		potentially used by bird species from changes	guidance as outlined within Chapter 17 – Water
		to hydrology; and changes to water and /or	Resources and Flood Risk (Document Reference
		sediment quality (from surface or groundwater	6.1.17).
		discharges from the Project Site including	
		operational waste and pollutants).	SPA and Winter Bird Mitigation Strategy Breeding
			and Winter Bird Mitigation Strategy enclosed
			within EMMF (Document Reference 6.2.12.3) -



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor	Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	updated bird surveys of retained and newly created wetlands and functionally linked habitats in years 3, 5 and 10 post-completion to ensure habitats continue to support the intended habitats and desired flora/fauna. BMRS – monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased. Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitatsOff-site Ecological Mitigation—delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area.	Reference 6.2.12.10), to offset any residual effects. SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		 River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).
			<u>Creation/Enhancement of 259.33ha Functionally</u> <u>Linked Habitat Off-site – delivered in accordance</u> with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	кесерио	Disturbance (human movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during operation of the resort rides and attractions, and entertainment facilities.	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) – Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation locations to reduce visual impact. Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on
			retained habitats.
West	National	Potential impacts solely on functionally linked res	, , ,
Thurrock		Damage to functionally linked habitats	SuDS – designed in accordance with best practice
Lagoon and		potentially used by cited bird species from	guidance as outlined within Chapter 17 – Water
Marshes SSSI		changes to hydrology; and changes to water	Resources and Flood Risk (Document Reference
		and-/or sediment quality (from surface or	6.1.17).
		groundwater discharges from the Project Site	
		including operational waste and pollutants).	SPA and Winter Bird Mitigation StrategyBreeding and Winter Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3) - updated bird surveys of retained and newly created wetlands and functionally linked habitats in years 3, 5 and 10 post-completion to ensure habitats continue to support the intended habitats and desired flora/fauna.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	BMRS – monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased.
			Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats. Off-site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
			from animal habitats. Protect the natural areas
			that are to be conserved and enhanced).
			BMRS – to monitor bird activity and inform the
			development of further control/mitigation
			measures as required.
			Disturbance control measures – final ride design to
			include deliberate placement of 'scream zones';
			strict noise specification to eliminate clanking;
			building placement to reduce noise spread; and
			controls on location, direction and sound pressure
			level of loudspeaker systems. (See Chapter 15,
			Document reference: 6.1.15) <u>.</u>
			Creation/Enhancement of 259.33ha Functionally
			<u>Linked Habitat Off-site</u> – delivered in accordance
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on retained
			habitats.
		Disturbance (human movement or activity), to	Landscape Strategy, LMP and EMMF (Document
		cited bird species potentially using functionally	Reference 6.2.12.3) – Recreational use managed via
		linked habitats, as a result of increased human	trail network, signage and interpretation boards to
		presence on the Project Site during operation of the resort rides and attractions, and	mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation
		entertainment facilities.	locations to reduce visual impact.
		entertainment facilities.	locations to reduce visual impact.



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		Creation/Enhancement of 259.33ha Functionally
			Linked Habitat Off-site – delivered in accordance
			with the ECF (Document Reference 6.2.12.10) to
			address any residual disturbance effects on
			retained habitats.
Great	National	Damage to habitats within the designated area	Mitigation measures to reduce traffic emissions will
Crabbles		from traffic generated air quality impacts.	need to be developed and agreed with Natural
Wood SSSI		Increased disposition of N likely to result in	England.
		nutrient enrichment and acidification of	
		qualifying woodland habitat with subsequent	
		changes in structure and diversity with such	
		impacts likely occurring across circa 13.6% of	
		total SSSI area is located within 200m of the	
		A289 whilst the woodland is reported to be in	
		unfavourable recovering condition. Total area	
		of impact is, however, reduced given N	
		sensitive woodland habitat is only present	
		across a proportion of this area.	
Botany	County	Damage to habitats from changes to	SuDS – designed in accordance with best practice
Marshes LWS		hydrological regime, and changes in water	guidance as outlined within Chapter 17 – Water
		quality/quantity.	Resources and Flood Risk (Document Reference
			6.1.17).
		Disturbance (noise and lighting) from the	SLS - Sensitive design and location of new lighting to
		operation of the resort rides and attractions	minimise impacts on nocturnal/crepuscular wildlife.
		and entertainment facilities, potentially giving	Botany Marshes LWS to remain in the following
		rise to displacement, behavioural changes or	Environmental Lighting Zone:



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	·	physiological stress to wildlife species within the designated area.	Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15)
		Recreational disturbance of flora and fauna	Landscape Strategy, LMP and EMMF (Document
		within designated area as a result of increased	Reference 6.2.12.3) – Recreational use managed via
		residential population (due to residential	trail network, signage and interpretation boards to
		accommodation for staff) and potential	mitigate against knock-on effects on on-site wildlife.
		increased recreational use of Swanscombe	Creation of new bird hides at key observation
		Peninsula from members of the general public.	locations to reduce visual impact.
Ebbsfleet Marshes, Northfleet LWS	County	Potential damage to habitats within the designated area from traffic generated air quality impacts – subject to further investigation to determine the nature of any significant effects.	If significant effects are identified following further investigation, mitigation measures to reduce traffic emissions will need to be developed and agreed with Natural England.
		Damage to habitats from changes to	SuDS – designed in accordance with best practice
		hydrological regime and changes in water	guidance as outlined within Chapter 17 – Water
		quality/quantity <u>.</u>	Resources and Flood Risk (Document Reference 6.1.17).



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor	Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, potentially giving rise to displacement, behavioural changes or physiological stress to wildlife species within the designated area.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Where possible, parts of Ebbsfleet Marshes, Northfleet LWS to remain in the following Environmental Lighting Zone: Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). With remaining areas (which are already subject to high levels of lighting) within: Environmental Zone E3 (Typical of well inhabited rural and urban settlements). Necessary lighting to create a safe and comfortable environment without over illuminating and reducing light spill onto adjacent areas of the E2 zone.
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).
		Recreational disturbance of flora and fauna	Landscape Strategy, LMP and EMMF (Document
		within designated area as a result of increased	Reference 6.2.12.3) – Recreational use managed via
			trail network, signage and interpretation boards to



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		residential population (due to residential	mitigate against knock-on effects on on-site wildlife.
		accommodation for staff).	Creation of new bird hides at key observation
			locations to reduce visual impact.
Disused	County	Potential damage to habitats within the	If significant effects are identified following further
Hospital		designated area from traffic generated air	investigation, Mmitigation measures to reduce
Grounds,		quality impacts – subject to further	traffic emissions will need to be developed and
Mabledon		investigation to determine the nature of any	agreed with Natural England.
LWS		significant effects.	
Tilbury	County	Damage to habitats from changes to	SuDS – designed in accordance with best practice
Marshes LWS		hydrological regime and changes in water	guidance as outlined within Chapter 17 – Water
		quality/quantity.	Resources and Flood Risk (Document Reference
			6.1.17).
		Disturbance (from increased ferry	BMRS – monitoring of bird use on the intertidal
		movements) associated with the new Thames	habitats and other functionally linked land – if the
		Clipper river bus service and operational	numbers of birds within the monitored area fall
		deliveries, potentially giving rise to	below a certain threshold in response to obvious
		displacement, behavioural changes or	operational shipping then those disturbance
		physiological stress to bird species potentially	activities will be temporally ceased.
		using functionally linked habitats.	
			Off-site Ecological Mitigation – delivered in
			accordance with the 'General Principles for Offsite
			Ecological Mitigation' (Document Reference
			6.2.12.10), to offset any residual
			effects.Creation/Enhancement of 259.33ha
			Functionally Linked Habitat Off-site – delivered in
			accordance with the ECF (Document Reference



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	песериот		6.2.12.10) to address any residual disturbance
			effects on retained habitats.
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Entire stretch of the River Thames maintained in the following Environmental Lighting Zone: Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting.
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).
		Disturbance (human movement or activity) , to cited bird species potentially using functionally linked habitats, as a result of increased human	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) – Recreational use managed via trail network, signage and interpretation boards to
		presence on the Project Site during operation of the resort rides and attractions, and	mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation
		entertainment facilities.	locations to reduce visual impact.



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
Populations of nationally scarce plants	National	N.B. Mitigation of potential effects upon scarce plants included under Swanscombe Peninsula SSSI. Damage to individual plants, and part of plant meta-population, through increased recreational use, potentially resulting in trampling, littering, dog fouling, unauthorised camping/campfires/barbecues	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) — Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife.
Retained habitats ⁸⁹ (including woodland, scrub, grassland, open mosaic on previously developed	Local to County (see Baseline section)	Damage to retained habitats, through increased recreational use, potentially resulting in trampling, littering, dog fouling, unauthorised camping/campfires/barbecues. Damage to retained habitats through inappropriate land management practices.	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) – Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife. LMP and EMMF (Document Reference 6.2.12.3) – Appropriate management of all retained, enhanced or newly created habitats to ensure they continue to support the intended flora and fauna, and provide long term benefits to biodiversity.
land, wetlands, waterbodies and the River Ebbsfleet).		Damage to retained habitats from changes to hydrological regime and changes in water quality/quantity.	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
Ancient woodland	County	Potential damage to habitats within the designated area from traffic generated air	If significant effects are identified following further investigation, Mmitigation measures to reduce

⁸⁹ Predicted impacts on retained habitats during the operation of the Proposed Development are considered applicable across the full suite of habitats retained, and are therefore considered collectively



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures		
		quality impacts subject to further investigation to determine the nature of any significant effects	traffic emissions will need to be developed and agreed with Natural England.		
Species					
Wintering Waterfowl and Wading Bird	International	N.B. Assessment of potential habitat loss and disturbance effects upon wintering waterfowl and wading bird assemblage included under SPA/Ramsar Sites. Damage to functionally linked habitats potentially used by birds from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and			
Assemblage		Sustainable Drainage System (SuDS) — designed in accordance with best practice guidance as within Chapter 17 — Water Resources and Flood Risk (Document Reference 6.1.17). Breeding and Winter Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.1.17).			
		- updated bird surveys of retained and newly cre	eated wetlands and functionally linked habitats in itats continue to support the intended habitats and		
Wintering terrestrial bird assemblage	County	Damage to on-site habitats potentially used by birds from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	Sustainable Drainage System (SuDS) – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17). SPA and Winter Bird Mitigation Strategy Breeding and Winter Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3) – updated bird surveys of retained and newly created wetlands and functionally linked habitats in years 3, 5 and 10 post-completion to ensure habitats		



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	Receptor	Dicturbance (noise and lighting) from the	continue to support the intended habitats and desired flora/fauna.
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to bird species using the on-site habitats.	 SLS – Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes – Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	-		level of loudspeaker systems. (See Chapter 15,
			Document reference: 6.1.15)
		Disturbance (human movement or activity), to	Landscape Strategy, LMP and EMMF (Document
		cited bird species potentially using functionally	Reference 6.2.12.3) – Recreational use managed via
		linked habitats, as a result of increased human	trail network, signage and interpretation boards to
		presence on the Project Site during operation	mitigate against knock-on effects on on-site wildlife.
		of the resort rides and attractions and	Creation of new bird hides at key observation
		entertainment facilities, including increased	locations to reduce visual impact.
		recreational use of retained habitats.	
Breeding Bird	Regional Natio	N.B. Impact assessment for the breeding bird ass	emblage included under Swanscombe Peninsula
Assemblage	<u>nal</u>	SSSI. Damage to on-site habitats potentially used	by birds from changes to hydrology; and changes to
		water and/or sediment quality (from surface or	groundwater discharges from the Project Site
		including operational waste and pollutants)	
		N.B. Proposed mitigation measures for the breed	ling bird assemblage included under Swanscombe
		Peninsula SSSISuDS – designed in accordance wit	h best practice guidance as outlined within Chapter
		17 – Water Resources and Flood Risk (Document	Reference 6.1.17).
		Breeding and Winter Bird Mitigation Strategy er	nclosed within EMMF (Document Reference 6.2.12.3)
		- updated bird surveys of retained and newly cre	eated habitats in years 3, 5 and 10 post completion to
		ensure habitats continue to support the intended	d habitats and desired flora/fauna.
Bat	District	Damage to on-site habitats potentially used by	SuDS – designed in accordance with best practice
Assemblage	County	bats from changes to hydrology; and changes	guidance as outlined within Chapter 17 – Water
		to water and /or sediment quality (from	Resources and Flood Risk (Document Reference
		surface or groundwater discharges from the	6.1.17).
		Project Site including operational waste and	
		pollutants) <u>.</u>	



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		Disturbance (lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to bat species using the onsite habitats, including light pollution of foraging, commuting and roosting habitats.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
Dormouse	District	Damage to on-site habitats potentially used by dormice from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to dormice using the onsite habitats, including light and noise pollution	 SLS — Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained scrub habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
		of foraging, dispersal and potential breeding/hibernating habitats.	from animal habitats. Protect the natural areas that are to be conserved and enhanced).
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).
Otter	Local	Damage to on-site habitats potentially used by otters from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to otters using the on-site habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking;
			building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems.
Water Vole	Local to dDistrict	Damage to on-site habitats potentially used by water voles from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to water voles using the onsite habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures
	Посория		inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
Hanvest	Local	Damage to on-site habitats potentially used by	Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15). SuDS – designed in accordance with best practice
Harvest Mouse	Local	harvest mice from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).
		Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to harvest mice using the on-site habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and



Ecological Feature	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
reature	Receptor		Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).
Amphibian	Local to D	Damage to on-site habitats potentially used by	SuDS – designed in accordance with best practice
assemblage	district	amphibians from changes to hydrology; and	guidance as outlined within Chapter 17 – Water
		changes to water and /or sediment quality	Resources and Flood Risk (Document Reference
		(from surface or groundwater discharges from	6.1.17).
		the Project Site including operational waste and pollutants).	
		Disturbance (lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to amphibians using the onsite habitats, including light pollution of breeding, foraging and hibernation/shelter habitats.	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively



Ecological Feature	Importance of Receptor	Nature of Impact	Summary of Avoidance/Mitigation Measures	
reacure	песерио		uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).	
Reptile assemblage	District	Damage to on-site habitats potentially used by reptiles from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	
		Disturbance (lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to reptiles using the on-site habitats, including light pollution of potential shelter and hibernation habitats.	 SLS - Sensitive design and location of new lighting to minimise impacts on wildlife. Retained grassland/scrub habitats within the Kent Project Site to remain in the following Environmental Lighting Zone: Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). 	
Invertebrate	National	N.B. Impact assessment for the invertebrate asse		
assemblage		SSSI. Damage to on-site habitats potentially used by invertebrates from changes to hydrology; and		
		Site including operational waste and pollutants	m surface or groundwater discharges from the Project	



Ecological	Importance of	Nature of Impact	Summary of Avoidance/Mitigation Measures
Feature	Receptor		
		N.B. Proposed mitigation measures for the invert	tebrate assemblage included under Swanscombe
		Peninsula SSSISuDS - designed in accordance wit	h best practice guidance as outlined within Chapter
		17 – Water Resources and Flood Risk (Document	Reference 6.1.17).
		·	·



RESIDUAL ENVIRONMENTAL EFFECTS

- <u>12.284</u> The residual effects are the likely effects occurring following implementation of the design measures <u>and the</u>—construction phase and operational phase <u>avoidance and</u> mitigation measures described above.
- <u>12.26</u>12.285 Aside from those off-site measures which constitute avoidance and mitigation for indirect effects on off-site designated sites, namely effects relating to loss/disturbance of functionally linked land and those relating to recreational disturbance, all other off-site (compensatory) measures are not taken into account in determining residual effects.
- 12.27 The assessment of residual effects assumes that off site mitigation will be provided in line with the 'General Principles for Off site Mitigation' provided in Appendix 12.10 (Document Reference 6.2.12.10).
- <u>12.28</u>12.286 The residual effects predicted to arise during construction are detailed within **Table 12.16**.
- <u>12.29</u>12.287 The residual effects predicted to arise during the operational phase are detailed within **Table 12.17**.



Table 12.16: Summary of ecological impact assessment and residual effects during construction phase-

Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
Thames Estuary	Potential impacts solely on funct	ionally linked resources (o	utside of the European/Ramsar Site):	
and Marshes	Direct loss of/damage to	Significant at District	Landscape Strategy - enhancement	No significant effect
SPA/Ramsar, and	functionally linked land on the	<u>level (moderate</u>	of retained wetland habitat on	
the component	Kent Project Site, potentially	adverse) (permanent).	Swanscombe Peninsula and creation	(potential for significant
SSSIs including	used by cited bird species,		of new saltmarsh, wetland and	positive effect in the
South Thames	totalling a net loss of 15.49ha,		reedbed habitats, managed via the	long term).
Estuary and	including 14.55ha of coastal		Ecological Mitigation and	
Marshes SSSI and	and floodplain grazing marsh		Management Framework (EMMF)	
Mucking Flats and	(Botany Marsh West) and		(Document Reference 6.2.12.3).	
Marshes SSSI ⁹⁰	0.94ha of reedbeds (Black Duck			
	Marsh) to accommodate		Creation/Enhancement of 259.33ha	
	construction of the Leisure		Functionally Linked Habitat Off-site	
	<u>Core.</u>		 delivered in accordance with the 	
			ECF (Document Reference	
			6.2.12.10).	
	Damage to functionally linked	Significant at district	CEMP - Prevention of hydrological	No significant effect. No
	habitats potentially used by	level District level	impacts through adherence to an	significant effect
	cited bird species from	(moderate adverse)	appropriate Surface Water	Significant circut
	changes to hydrology (as a	(temporary).	Management Strategy, as outlined	
	result of drainage and infill of	(33, 63.3.7).	within Chapter 17 – Water	
	the CTRL wetland,		within Chapter 17 – water	
	swamp/wetland habitat along			

South Thames Estuary and Marshes SSSI and Mucking Flats and Marshes SSSI are component SSSIs to the Thames Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and	Residual Effect including Significance
		melaung Significance	Mitigation	including Significance
	the east/southeast of Black		Resources and Flood Risk (Document	
	Duck Marsh and other low-		Reference 6.1.17).	
	lying seasonally inundated		,	
	areas); and changes to water			
	and-/or sediment quality			
	(either from surface or			
	groundwater discharges from			
	Project Site including			
	construction waste and			
	pollutants; or from disruption			
	of contaminated Thames			
	sediments), with potential			
	associated knock-on risk of			
	bioaccumulation. Damage to			
	functionally linked habitats			
	potentially used by cited bird			
	species from changes to			
	hydrology (as a result of			
	drainage and infill of the CTRL			
	wetland, swamp/wetland			
	habitat along the east/south			
	east of Black Duck Marsh and			
	other low lying seasonally			
	inundated areas); and changes			
	to water and /or sediment			
	quality (either from surface or			
	groundwater discharges from			



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	Project Site including construction waste and pollutants; or from disruption of contaminated Thames sediments), with potential associated knock-on risk of			
	Damage to functionally linked habitats potentially used by cited bird species from changes in air quality including from dust, construction waste and pollutants. Damage to functionally linked habitats potentially used by cited bird species from changes in air quality including from dust, construction waste and pollutants.	Significant at district level [level District level] (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.	No significant effect. No significant effect
	Damage to functionally linked habitats potentially used by cited bird species from	Significant at district level District level	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference	No significant effect. No significant effect



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
	introduction or proliferation of	(minor adverse)	6.2.12.3) – including details on	
	INNS. Damage to functionally	(temporary, reversible).	control/eradication of existing	
	linked habitats potentially used		populations of non-native species.	
	by cited bird species from			
	introduction or proliferation of		CEMP – to include control measures	
	Invasive Non-Native Species		to prevent spread of INNS and INNS	
	(INNS).		impacted soils. Non-native Invasive	
			Plant Species Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) – including	
			details on control/eradication of	
			existing populations of non-native	
	!		species.	
			Species:	
			CEMP – to include control measures	
	!		to prevent spread of INNS and INNS	
			impacted soils.	
	Disturbance (from	Significant at district	BMRS - monitoring of bird use on	No significant effect. No
	shipping/ferry movements)	level District level	the intertidal habitats and other	significant effect
	giving rise to displacement,	(minor adverse)	functionally linked land – if the	
	behavioural changes or	(temporary, reversible).	numbers of birds within the	
	physiological stress to cited	, , , , , , , , , , , , , , , , , , , ,	monitored area fall below a certain	
	bird species potentially using		threshold in response to obvious	
	functionally linked		construction activities then those	
	habitats. Disturbance (from		disturbance activities will be	
	shipping/ferry movements)		temporally ceased.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	giving rise to displacement,			
	behavioural changes or		Creation/Enhancement of 259.33ha	
	physiological stress to cited		Functionally Linked Habitat Off-site	
	bird species potentially using		– delivered in accordance with the	
	functionally linked habitats.		ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats.Bird	
			Monitoring Response Strategy	
			(BMRS) monitoring of bird use on	
			the intertidal habitats and other	
			functionally linked land – if the	
			numbers of birds within the	
			monitored area fall below a certain	
			threshold in response to obvious	
			construction activities then those	
			disturbance activities will be	
			temporally ceased.	
			Off site Ecological Mitigation –	
			delivered in accordance with the	
			'General Principles for Offsite	
			Ecological Mitigation' (Document	
			Reference 6.2.12.10), to offset any	
			residual effects.	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Disturbance (noise and	Significant at district	CEMP and Sensitive Lighting	No significant effect: No
	lighting) associated with	level District level	Strategy (SLS)SLS - Restricted	significant effect
	increased shipping traffic and	(moderate adverse)	working hours and sensitive design	
	construction activities, giving	(temporary, reversible).	and location of new lighting to	
	rise to displacement,		minimise impacts on	
	behavioural changes or		nocturnal/crepuscular wildlife.	
	physiological stress to cited bird species potentially using			
	functionally linked habitats.		CEMP – sensitive timing of works	
	Turictionally liffked flabitats.		causing in excess of 55dB of noise at	
			the river frontage or functionally	
			linked habitat, to take place in	
			summer months, where possible, as	
			detailed in the Bird Mitigation	
			Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3).	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
			BMRS – to monitor bird activity and	
			inform the development of further	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			control/mitigation measures as required.	
			Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10)	
	Disturbance (human	Significant at district	to address any residual disturbance effects on retained habitats. CEMP – use of site	No significant effect. No
	movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during construction.	levelDistrict level (moderate adverse) (temporary, reversible).	screening/hoarding; sensitive timing of works within 300m of the river frontage or functionally linked habitat and visible from that habitat to take place in summer months, where possible; and access by construction workers to river front restricted within 500m of visible wetland habitats, where possible; as detailed in the Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3).	significant effect
			BMRS – to monitor bird activity and inform the development of further	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
			<u>Mitigation</u>	
			control/mitigation measures as	
			required.	
			Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			 delivered in accordance with the 	
			ECF (Document Reference	
			<u>6.2.12.10).</u>	
Medway Estuary	Potential impacts solely on funct	ionally linked resources (c	outside of the European/Ramsar Site):	
and Marshes	Direct loss of/damage to	Significant at district	Landscape Strategy - enhancement	No significant effect. No
SPA/Ramsar/SSSI ⁹¹	functionally linked land on the	level District level	of retained wetland habitat on	significant effect.
	Kent Project Site, potentially	(moderate adverse)	Swanscombe Peninsula and creation	
	used by cited bird species,	(permanent).	of new saltmarsh, wetland and	(potential for significant
	totalling a net loss of 15.49ha,		reedbed habitats, managed via the	positive effect in the
	including 14.55ha of coastal		EMMF (Document Reference	long term -subject to
	and floodplain grazing marsh		<u>6.2.12.3).</u>	final off-site mitigation
	(Botany Marsh West) and			package).
	0.94ha of reedbeds (Black Duck		Creation/Enhancement of 259.33ha	
	Marsh) to accommodate		Functionally Linked Habitat Off-site	
	construction of the Leisure		<u>– delivered in accordance with the</u>	
	Core.		ECF (Document Reference	
			6.2.12.10). Landscape Strategy	
			enhancement of retained wetland	

⁹¹ Medway Estuary and Marshes SSSI is a component SSSI to the Medway Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also. No potential impacts are identified to the SSSI that are additional or different in their nature to those identified to the SPA/Ramsar



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			habitat on Swanscombe Peninsula and creation of new saltmarsh, wetland and reedbed habitats, managed via the EMMF (Document Reference 6.2.12.3). Off-site Ecological Mitigation – delivered in accordance with the General Principles for Offsite	
	Damage to functionally linked habitats potentially used by cited bird species from changes to hydrology (as a result of drainage and infill of the CTRL wetland, swamp/wetland habitat along the east/south east of Black Duck Marsh and other low-lying seasonally inundated areas); and changes to water and /or sediment quality (either from surface or groundwater discharges from Project Site including construction waste and pollutants; or from disruption	Significant at district level District level (moderate adverse) (temporary).	Ecological Mitigation. CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	of contaminated Thames sediments), with potential associated knock-on risk of bioaccumulation.			
	Damage to functionally linked habitats potentially used by cited bird species from changes in air quality including from dust, construction waste and pollutants.	Significant at district levelDistrict level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.	No significant effect. No significant effect.
	Damage to functionally linked habitats potentially used by cited bird species from introduction or proliferation of	Significant at district level District level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within the EMMF (Document Reference 6.2.12.3) – including details on	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	Invasive Non-Native Species (INNS)INNS.		control/eradication of existing populations of non-native species.	
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	
	Disturbance (from	Significant at district	Bird Monitoring Response Strategy	No significant effect. No
	shipping/ferry movements)	level District level	(BMRS)BMRS - to monitor bird	significant effect
	giving rise to displacement,	(minor adverse)	activity and inform the development	
	behavioural changes or	(temporary, reversible).	of further control/mitigation	
	physiological stress to cited		measures as required.	
	bird species potentially using			
	functionally linked habitats.		Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			— delivered in accordance with the	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitatsOff-site	
			Ecological Mitigation – delivered in	
			accordance with the 'General	
			Principles for Offsite Ecological	
			Mitigation' (Document Reference	
			6.2.12.10), to offset any residual	
	Distruction of the income	Cignificant of district	CEMP and SIS Postricted working	No significant offeet No
	Disturbance (noise and	Significant at district	CEMP and SLS - Restricted working	No significant effect. No
	lighting) associated with	level District level	hours and sensitive design and	significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	increased shipping traffic and	(moderate adverse)	location of new lighting to minimise	
	construction activities, giving	(temporary, reversible).	impacts on nocturnal/crepuscular	
	rise to displacement,		wildlife.	
	behavioural changes or			
	physiological stress to cited		CEMP – sensitive timing of works	
	bird species potentially using		causing in excess of 55dB of noise at	
	functionally linked habitats.		the river frontage or functionally	
			linked habitat, to take place in	
			summer months, where possible, as	
			detailed in the Bird Mitigation	
			Strategy enclosed within EMMF	
			(Document Reference 6.2.12.3).	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			— delivered in accordance with the	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats.	
	Disturbance (human	Significant at district	CEMP – use of site	No significant effect. No
	movement or activity), to cited	level District level	screening/hoarding; sensitive timing	significant effect
	bird species potentially using	(moderate adverse)	of works within 300m of the river	
	functionally linked habitats, as	(temporary, reversible).	frontage or functionally linked	
	a result of increased human		habitat and visible from that habitat	
	presence on the Project Site		to take place in summer months,	
	during construction.		where possible; and access by	
			construction workers to river front	
			restricted within 500m of visible	
			wetland habitats, where possible; as	
			detailed in the Bird Mitigation	
			Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3).	
			,	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			<u>– delivered in accordance with the</u>	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats	
<u>Swanscombe</u>	Overall reduction in SSSI area,	Significant at national	Landscape Strategy and	Significant negative at
Peninsula SSSI	and fragmentation by the	levelNational level	Swanscombe Peninsula SSSI	County National level.
	Resort.	(major adverse)	Mitigation StrategySSSI Mitigation	
		(permanent).	Strategy – maintenance of	
			connectivity across the Peninsula	
			through the inclusion of a chain of	
			watercourses and wetland areas	
			wrapping around the side of the	
			Proposed Development footprint,	
			maintenance of scrub habitats	
			across the Peninsula to allow	
			terrestrial bird species to disperse	
			between habitats and maintenance	
			of unrestricted flight path from River	
			Thames into Black Duck Marsh via	
			retaining and managing low-growing	
			vegetation at northern edge of	
			marsh. Creation of new wetland and	
			saltmarsh habitats on Swanscombe	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
			Mitigation Peninsula, delivered through the	
			EMMF and Landscape Strategy.	
	Damage to habitats within the	Significant at National	CEMP – including details on	Not significant
	designated area from changes	level (major adverse)	safe/appropriate disposal of	
	in air quality, including from	(temporary).	construction waste, and pollution	
	dust, construction waste and		prevention guidelines, to be strictly	
	pollutants, and exhaust		adhered to. CEMP will also include a	
	emissions.		<u>Dust Management Plan (DMP) to</u>	
			provide measures to control dust	
			and other emissions. The CEMP will	
			secure a suite of construction	
			mitigation measures as detailed in	
			Chapter 16: Air Quality (Document	
			reference 6.1.16), relating to site	
			management/ maintenance, site	
			monitoring, operation of vehicles/	
			machinery/-power tools, and	
			techniques for demolition works,	
			earthworks and construction.	
	Special feature of interest - popu	lations of rare plants		
	Direct loss of individual plants,	Significant at national	Landscape Strategy - enhancement	Significant negative at
	and Direct loss of individual	levelNational level	of retained habitats on Swanscombe	National level. No
	plants/supporting habitat, and	(major adverse)	Peninsula and creation of new	significant effect.
	part of plant meta-population	(permanent).	habitats, managed via the	
	through loss of habitat within	11.	Swanscombe Peninsula SSSI	
	SSSI.			



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			Mitigation Strategy SSSI Mitigation	
			Strategy.	
			Swanscombe Peninsula SSSI	
			Mitigation Strategy (Rare Plants)	
			Rare Plant Mitigation Strategy	
			enclosed within the SSSI Mitigation	
			Strategy and subsequent detailed	
			strategies providing details for the	
			translocation of nationally scarce	
			plants/seed bank occurring on the	
			Kent Project Site.	
	Damage to retained plants	Significant at national	CEMP – including details on	No significant effect. No
	from construction dust, waste	levelNational level	safe/appropriate disposal of	significant effect.
	and pollutants.	(major adverse)	construction waste, and pollution	
		(temporary).	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions.	
	Damage to individual	Significant at national	CEMP and Swanscombe Peninsula	No significant effect. No
	plants Damage to individual	levelNational level	SSSI Mitigation Strategy (Rare	significant effect.
	plants/supporting habitat	(moderate adverse)	Plants) SSSI Mitigation Strategy -	
	from adjacent construction	(temporary).	detailing sensitive construction	
	works and deposition of		methodology, including set up and	
	construction materials.		maintenance of EPZs where no	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			construction vehicles/works are	
			permitted and where no materials	
			or site facilities may be situated.	
	Damage to plants/their	Significant at national	Swanscombe Peninsula SSSI	No significant effect. No
	habitats from introduction or	levelNational level	Mitigation Strategy (Rare Plants)	significant effect.
	proliferation of Invasive Non	(minor adverse)	Non-native Invasive Plant Species	
	Native Species (INNS)INNS.	(temporary).	Mitigation Strategy included within	
			SSSI Mitigation Strategy – including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
	Special feature of interest - breed	ding bird assemblages:		
	Direct killing, injury or harm to	Significant at national	CEMP, and Swanscombe Peninsula	No significant effect.No
	individuals – during vegetation	levelNational level	SSSI Mitigation Strategy (Breeding	significant effect.
	<u>clearance.</u>	(major adverse)	Birds) Breeding and Winter Bird	
		(permanent).	Mitigation Strategy included within	
			SSSI Mitigation Strategy – including	
			sensitive timing and methodology	
			for habitat clearance, establishment	
			of EPZs to protect retained habitats.	
	Direct habitat loss – loss of	Significant at national	Landscape Strategy and	Significant negative at
	suitable habitat such as scrub,	levelNational level	Swanscombe Peninsula SSSI	National level. No
	grassland, reedbed and	(moderate adverse)	Mitigation Strategy (Breeding Birds)	significant effect.
	woodland.	(permanent).	Breeding Bird Mitigation Strategy	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			included within SSSI Mitigation	
			Strategy - enhancement of retained	
			scrub habitat through rotational	
			management to create age class and	
			structural diversity, and	
			maintenance of site wide open	
			mosaic habitats to provide variety of	
			foraging resources, managed via the	
			EMMF.	
	Damage to supporting	Significant at national	CEMP - Prevention of hydrological	No significant effect. No
	habitats from changes to	levelNational level	impacts through adherence to an	significant effect.
	hydrological regime and	(moderate adverse)	appropriate Surface Water	
	changes to water and/or	(temporary).	Management Strategy, as outlined	
	sediment quality.		within Chapter 17 – Water	
			Resources and Flood Risk (Document	
			Reference 6.1.17).	
	Damage to supporting	Significant at national	CEMP – including details on	No significant effect. No
	habitats from construction	levelNational level	safe/appropriate disposal of	significant effect.
	dust, waste and pollutants.	(major adverse)	construction waste, and pollution	
		(temporary).	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
			Mitigation	
	Damage to supporting	Significant at national	Swanscombe Peninsula SSSI	No significant effect. No
	habitats from introduction or	levelNational level	Mitigation StrategyNon-native	significant effect.
	proliferation of Invasive Non-	(minor adverse)	Invasive Plant Species Mitigation	
	Native Species (INNS)INNS.	(temporary).	Strategy included within SSSI	
			Mitigation Strategy - including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
	Habitat fragmentation, loss of	Significant at national	Landscape Strategy and	Significant negative at
	flight paths and dispersal	levelNational level	Swanscombe Peninsula SSSI	National level. No
	routes.	(moderate adverse)	Mitigation Strategy (Breeding Birds)	significant effect.
		(permanent).	SSSI Mitigation Strategy -	
			maintenance of connectivity across	
			the Peninsula through the inclusion	
			of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint, maintenance of scrub	
			habitats across the Peninsula to	
			allow terrestrial bird species to	
			disperse between habitats and	
			maintenance of unrestricted flight	
			path from River Thames into Black	
			Duck Marsh via retaining and	
			managing low-growing vegetation at	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			northern edge of marsh. Creation of	
			new wetland and saltmarsh habitats	
			on Swanscombe Peninsula, delivered	
			through the EMMF and Landscape	
			Strategy.	
	Disturbance (from	Significant at national	Bird Monitoring Response Strategy	Significant negative at
	shipping/ferry movements) -	levelNational level	(BMRS)BMRS - monitoring of bird	National level. No
	as described above in relation	(minor adverse)	use on the intertidal habitats and	significant effect.
	to impacts on habitats	(temporary).	other functionally linked land – if the	
	functionally linked to		numbers of birds within the	
	SPA/Ramsar sites.		monitored area fall below a certain	
			threshold in response to obvious	
			construction activities then those	
			disturbance activities will be	
			temporally ceased.	
			Off-site Ecological Mitigation -	
			delivered in accordance with the	
			'General Principles for Offsite	
			Ecological Mitigation' (Document	
			Reference 6.2.12.10), to offset any	
			residual effects.	
	Disturbance (noise and	Significant at national	CEMP and Sensitive Lighting	No significant effect.
	lighting) – as described above	levelNational level	Strategy (SLS)SLS - Restricted	significant effect.
	in relation to impacts on		working hours and sensitive design	
	habitats-SPA/Ramsar sites.		and location of new lighting to	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	functionally linked to SPA/Ramsar sites.	(moderate adverse) (temporary).	minimise impacts on nocturnal/crepuscular wildlife. CEMP – sensitive timing of works causing in excess of 55dB of noise at	
			the river frontage or functionally linked habitat, to take place in summer months, where possible, as detailed in the Bird Mitigation Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3). CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document	
			reference 6.2.15.5). BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.	
	Disturbance (human movement or activity) – as described above in relation to impacts on SPA/Ramsar sites.	Significant at national levelNational level	CEMP – use of site screening/hoarding; sensitive timing of works within 300m of the river	Significant negative at National level. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	Special feature of interest - inver	(moderate adverse) (temporary).	frontage or functionally linked habitat and visible from that habitat to take place in summer months, where possible; and access by construction workers to river front restricted within 500m of visible wetland habitats, where possible; as detailed in the Swanscombe Peninsula SSSI Mitigation Strategy (Breeding Birds).	
	Direct habitat loss – loss of suitable habitat, particularly Open Mosaic Habitat on Previously Developed Land, Iprovide further detail, including assessment of impacts upon all four assemblages and& provide quantities of habitat loss].	Significant at national levelNational level (major adverse) (permanent).	Landscape Strategy, and Swanscombe Peninsula SSSI Mitigation Strategy (Invertebrates) – enhancement of existing retained open mosaic habitat (OMH) to better quality and creation of new brown roofs as described previously. Furthermore, enhancement of retained wetland habitat within Botany Marsh East and Black Duck Marsh through reduction of scrub encroachment, selective ditch re- profiling, and creation of additional scrapes and deep areas in Black	Significant negative at National level. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			<u>Duck Marsh. Creation of additional</u>	
			saltmarsh habitat on the north-	
			eastern edge of Swanscombe	
			Peninsula through 'retiring' the flood	
			defence. Creation of a number of	
			habitats of value to invertebrates	
			within the amenity spaces within the	
			resort itself, including native tree	
			and shrub planting, wildflower	
			strips, green roofs and walls and	
			<u>'bug hotels'.</u>	
			Off-site Ecological Mitigation -	
			delivered in accordance with the	
			General Principles for Offsite	
			Ecological Mitigation (Document	
			Reference 6.2.12.10) will also create	
			additional valuable invertebrate	
			habitat.	
	Damage to habitats from	Significant at national	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelNational level	impacts through adherence to an	significant effect.
	regime and changes to water	(moderate adverse)	appropriate Surface Water	
	and/or sediment quality.	(temporary).	Management Strategy, as outlined	
			within Chapter 17 – Water	
			Resources and Flood Risk (Document	
			Reference 6.1.17).	
			MCICICINCE U.I.I/J.	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Damage to supporting	Significant at national	CEMP – including details on	No significant effect. No
	habitats from construction	levelNational level	safe/appropriate disposal of	significant effect.
	dust, waste and pollutants.	(major adverse)	construction waste, and pollution	
		(temporary).	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions.	
	Damage to supporting	Significant at national	Swanscombe Peninsula SSSI	No significant effect. No
	habitats from introduction or	levelNational level	Mitigation Strategy – including	significant effect.
	proliferation of Invasive Non-	(minor adverse)	details on control/eradication of	
	Native Species (INNS)INNS.	(temporary).	existing populations of non-native	
			species.	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation.	Significant at national	Landscape Strategy and	Significant negative at
		levelNational level	Swanscombe Peninsula SSSI	National level. No
		(moderate adverse)	Mitigation Strategy (Invertebrates)	significant effect.
		(permanent).	- retention and enhancement of	
			large area of open mosaic habitat on	
			the Kent Project Site, connected to	
			existing semi-natural habitats.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	<u>Disturbance (lighting)</u> – as described above in relation to impacts on SPA/Ramsar sites.	Significant at national levelNational level (moderate adverse) (temporary).	CEMP and SLS – Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.	No significant effect. No significant effect.
Darenth Woods SSSI	Damage to habitats within the designated area from changes in air quality, including from dust, construction waste and pollutants, and exhaust emissions.	Significant at National level (moderate adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/-maintenance, site monitoring, operation of vehicles/machinery/-power tools, and techniques for demolition works, earthworks and construction.	No significant effect.
	Potential impacts solely on funct	ionally linked resources (o	utside of the SSSI):	1



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
Inner Thames Marshes SSSI	Direct loss of/damage to functionally linked habitats on the Kent Project Site, potentially used by bird species, totalling a net loss of 15.49ha, including 14.55ha of coastal and floodplain grazing marsh (Botany Marsh West) and 0.94ha of reedbeds (Black Duck Marsh) to accommodate construction of the Leisure Core.	Significant at district level District level (moderate adverse) (permanent).	Landscape Strategy - enhancement of retained wetland habitat on Swanscombe Peninsula and creation of new saltmarsh, wetland and reedbed habitats, managed via the EMMF (Document Reference 6.2.12.3). Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10).Off site Ecological Mitigation — delivered in accordance with the General Principles for Offsite Ecological Mitigation.	No significant effect. No significant effect. (potential for significant positive effect in the long term subject to final off-site mitigation package).
	habitats potentially used by bird species from changes to hydrology (as a result of drainage and infill of the CTRL wetland, swamp/wetland habitat along the east/south east of Black Duck Marsh and other low-lying seasonally inundated areas); and changes to water and-/or sediment	Significant at district level District level (moderate adverse) (temporary).	cemp - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect. No significant effect



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	quality (either from surface or groundwater discharges from Project Site including construction waste and pollutants; or from disruption of contaminated Thames sediments), with potential associated knock-on risk of bioaccumulation.			
	Damage to functionally linked habitats potentially used by cited bird species from changes in air quality including from dust, construction waste and pollutants.	Significant at district level District level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
	Damage to functionally linked	Significant at district	Non-native Invasive Plant Species	No significant effect. No
	habitats potentially used by	level <u>District level</u>	Mitigation Strategy included within	significant effect.
	cited bird species from	(minor adverse)	the EMMF (Document Reference	
	introduction or proliferation of	(temporary, reversible).	6.2.12.3) – including details on	
	Invasive Non-Native Species		control/eradication of existing	
	(INNS) INNS.		populations of non-native species.	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Disturbance (from	Significant at district	Bird Monitoring Response Strategy	No significant effect. No
	shipping/ferry movements)	level <u>District level</u>	(BMRS)BMRS - to monitor bird	significant effect.
	giving rise to displacement,	(minor adverse)	activity and inform the development	
	behavioural changes or	(temporary, reversible).	of further control/mitigation	
	physiological stress to cited		measures as required.	
	bird species potentially using			
	functionally linked habitats.		Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			— delivered in accordance with the	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitatsOff site	
			Ecological Mitigation – delivered in	
			accordance with the 'General	
			Principles for Offsite Ecological	
			Mitigation' (Document Reference	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
			Mitigation 6.2.12.10), to offset any residual effects.	
	Disturbance (noise and lighting) associated with increased shipping traffic and construction activities, giving rise to displacement, behavioural changes or physiological stress to cited species potentially using functionally linked habitats.	Significant at district level District level (moderate adverse) (temporary, reversible).	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP — sensitive timing of works causing in excess of 55dB of noise at the river frontage or functionally linked habitat, to take place in summer months, where possible, as detailed in the Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3). CEMP — adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document	No significant effect. No significant effect.
			reference 6.2.15.5). Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			— delivered in accordance with the	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats.	
	Disturbance (human	Significant at district	CEMP – use of site	No significant effect. No
	movement or activity), to cited	level District level	screening/hoarding; sensitive timing	significant effect.
	bird species potentially using	(minor adverse)	of works within 300m of the river	
	functionally linked habitats, as	(temporary, reversible).	frontage or functionally linked	
	a result of increased human		habitat and visible from that habitat	
	presence on the Project Site		to take place in summer months,	
	during construction.		where possible; and access by	
			construction workers to river front	
			restricted within 500m of visible	
			wetland habitats, where possible; as	
			detailed in the Bird Mitigation	
			Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3).	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
			Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			– delivered in accordance with the	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats	Residual Effect including Significance
West Thurrock Lagoon and Marshes SSSI	Potential impacts solely on functionally linked habitats on the Kent Project Site, potentially used by cited bird species, totalling a net loss of 15.49ha, including 14.55ha of coastal and floodplain grazing marsh (Botany Marsh West) and 0.94ha of reedbeds (Black Duck Marsh) to accommodate construction of the Leisure Core.	Significant at district levelDistrict level (moderate adverse) (permanent).	Landscape Strategy - enhancement of retained wetland habitat on Swanscombe Peninsula and creation of new saltmarsh, wetland and reedbed habitats, managed via the EMMF (Document Reference 6.2.12.3). Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10). Off-site Ecological Mitigation — delivered in accordance	No significant effect. No significant effect (potential for significant positive effect in the long term subject to final off-site mitigation package).
	Damage to functionally linked habitats potentially used by cited bird species from changes to hydrology (as a result of drainage and infill of the CTRL wetland,	Significant at district level [evelDistrict level] (moderate adverse) (temporary).	with the General Principles for Offsite Ecological Mitigation. CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and	Residual Effect including Significance
			Mitigation	
	swamp/wetland habitat along		Resources and Flood Risk (Document	
	the east/south east of Black		Reference 6.1.17).	
	Duck Marsh and other low-		,	
	lying seasonally inundated			
	areas); and changes to water			
	and /or sediment quality			
	(either from surface or			
	groundwater discharges from			
	Project Site including			
	construction waste and			
	pollutants; or from disruption			
	of contaminated Thames			
	sediments), with potential			
	associated knock-on risk of			
	bioaccumulation.			
	Damage to functionally linked	Significant at district	CEMP – including details on	No significant effect. No
	habitats potentially used by	level District level	safe/appropriate disposal of	significant effect.
	cited bird species from	(minor adverse)	construction waste, and pollution	
	changes in air quality including	(temporary).	prevention guidelines, to be strictly	
	from dust, construction waste		adhered to. CEMP will also include a	
	and pollutants.		Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions. The CEMP will	
			secure a suite of construction	
			mitigation measures as detailed in	
			Chapter 16: Air Quality (Document	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			reference 6.1.16), relating to site	
			management/ maintenance, site	
			monitoring, operation of vehicles/	
			machinery/ power tools, and	
			techniques for demolition works,	
			earthworks and construction.	
	Damage to functionally linked	Significant at district	Non-native Invasive Plant Species	No significant effect. No
	habitats potentially used by	level District level	Mitigation Strategy included within	significant effect.
	cited bird species from	(minor adverse)	the EMMF (Document Reference	
	introduction or proliferation of	(temporary, reversible).	6.2.12.3) – including details on	
	Invasive Non-Native Species		control/eradication of existing	
	(INNS) INNS.		populations of non-native species.	
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	
	Disturbance (from	Significant at district	Bird Monitoring Response Strategy	No significant effect. No
	shipping/ferry movements)	levelDistrict level	(BMRS)BMRS - to monitor bird	significant effect.
	giving rise to displacement,	(minor adverse)	activity and inform the development	
	behavioural changes or	(temporary,	of further control/mitigation	
	physiological stress to cited	reversible) <u>.</u> -	measures as required.	
	bird species potentially using			
	functionally linked habitats.		Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			— delivered in accordance with the	
			ECF (Document Reference 6.2.12.10)	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			to address any residual disturbance effects on retained habitatsOff site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.	
	Disturbance (noise and lighting) associated with increased shipping traffic and construction activities, giving rise to displacement, behavioural changes or physiological stress to cited species potentially using functionally linked habitats.	Significant at district levelDistrict level (moderate adverse) (temporary, reversible).	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP – sensitive timing of works causing in excess of 55dB of noise at the river frontage or functionally linked habitat, to take place in summer months, where possible, as detailed in the Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3). CEMP – adoption of mitigation	No significant effect. No significant effect.
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).	
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.	
			Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats.	
	Disturbance (human movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during construction.	Significant at district level District level (minor adverse) (temporary, reversible).	cemp – use of site screening/hoarding; sensitive timing of works within 300m of the river frontage or functionally linked habitat and visible from that habitat to take place in summer months, where possible; and access by construction workers to river front restricted within 500m of visible wetland habitats, where possible; as	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			detailed in the Bird Mitigation	
			Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3).	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
			Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			 delivered in accordance with the 	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats.	
Botany Marshes	Damage to habitats from	Significant at local	CEMP – including details on	No significant effect. No
LWS	changes in air quality,	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	including from dust,	level (major adverse)	construction waste, and pollution	
	construction waste and	(temporary).	prevention guidelines, to be strictly	
	pollutants.	, , , , , , , , , , , , , , , , , , , ,	adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions. The CEMP will	
			secure a suite of construction	
			mitigation measures as detailed in	
			Chapter 16: Air Quality (Document	
			reference 6.1.16), relating to site	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			management/ maintenance, site monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.	
	Damage to habitats from changes to hydrological regime (through destruction of adjoining wetland/grazing marsh), and changes in water quality/quantity.	Significant at local levelSignificant at Local level (major adverse) (permanent).	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect. No significant effect.
	Damage to habitats from adjacent construction works and deposition of construction materials.	Significant at local levelSignificant at Local level (moderate adverse) (temporary).	CEMP and EMMF – detailing sensitive construction methodology, including set up and maintenance of Ecological Protection Zones (EPZs) where no construction vehicles/works are permitted and where no materials or site facilities may be situated.	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
	Disturbance (noise and	Significant at local	CEMP and SLS - Restricted working	No significant effect. No
	lighting) associated with	levelSignificant at Local	hours and sensitive design and	significant effect.
	construction activities, giving	<u>level</u> (moderate	location of new lighting to minimise	
	rise to displacement,	adverse) (temporary).	impacts on nocturnal/crepuscular	
	behavioural changes or		wildlife.	
	physiological stress to wildlife			
	species within the designated		CEMP – sensitive timing of works,	
	area.		where possible, as detailed in the	
			Bird Mitigation Strategy enclosed	
			within the EMMF (Document	
			Reference 6.2.12.3).	
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).	
Ebbsfleet	Direct loss of/damage to	Significant at district	Landscape Strategy – Retention of	No significant effect. No
Marshes,	habitat through Station	level District level	site wide open mosaic of habitats	significant effect
Northfleet LWS	Quarter South.	(moderate adverse)	within retained parts of	
		(permanent).	Swanscombe Peninsula, and	
			creation of new constructed	
			wetland , new ponds and ditches,	
			and water vole receptor site,	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			managed through the EMMF (Document Reference 6.2.12.3).	
	Damage to habitats from changes in air quality, including from dust, construction waste and pollutants.	Significant at local levelSignificant at Local level (moderate adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/-maintenance, site monitoring, operation of vehicles/ machinery/-power tools, and techniques for demolition works,	No significant effect. No significant effect.
	Damage to habitats from	Significant at local	earthworks and construction. CEMP - Prevention of hydrological	No significant effect.No
	changes to hydrological regime (through destruction of habitat within floodzoneflood zone of River Ebbsfleet), and changes in water quality/quantity.	levelSignificant at Local level (moderate adverse) (temporary).	impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Damage to habitats from	Significant at local	CEMP and EMMF – detailing	No significant effect. No
	adjacent construction works	levelSignificant at Local	sensitive construction methodology,	significant effect.
	and deposition of construction	<u>level</u> (moderate	including set up and maintenance of	
	materials.	adverse) (temporary).	EPZs where no construction	
			vehicles/works are permitted and	
			where no materials or site facilities	
			may be situated.	
	Disturbance (noise and	Significant at local	CEMP and SLS - Restricted working	No significant effect. No
	lighting) associated with	levelSignificant at Local	hours and sensitive design and	significant effect.
	construction activities, giving	<u>level</u> (moderate	location of new lighting to minimise	
	rise to displacement,	adverse) (temporary).	impacts on nocturnal/crepuscular	
	behavioural changes or		wildlife.	
	physiological stress to wildlife			
	species within the designated		CEMP – sensitive timing of works,	
	area.		where possible, as detailed in the	
			Bird Mitigation Strategy enclosed	
			within the EMMF (Document	
			Reference 6.2.12.3).	
			,	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
Alkerden Lane Pit LWS	Loss of part of plant and invertebrate meta-population through loss of habitat in the Kent Project Site.	Significant at local levelSignificant at Local level (minor adverse) (permanent).	Landscape Strategy - enhancement of retained habitats on Swanscombe Peninsula, and creation of new habitats, managed via the EMMF (Document Reference 6.2.12.3).	Significant negative at Local level. Significant positive effect (local level).
			EMMF and subsequent detailed strategies Rare Plant Mitigation Strategy enclosed within the EMMF - providing details for the translocation of nationally scarce plants/seed bank occurring on the Kent Project Site.	
	Damage to habitats from changes in air quality, including from dust, construction waste and pollutants.	Significant at local levelSignificant at Local level (major adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. The CEMP will secure a suite of construction mitigation measures as detailed in Chapter 16: Air Quality (Document reference 6.1.16), relating to site management/ maintenance, site	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			monitoring, operation of vehicles/ machinery/ power tools, and techniques for demolition works, earthworks and construction.	
Tilbury Marshes LWS	Direct loss of/damage to functionally linked habitats on the Kent Project Site, potentially used by cited bird species, totalling a net loss of 15.49ha, including 14.55ha of coastal and floodplain grazing marsh (Botany Marsh West) and 0.94ha of reedbeds (Black Duck Marsh) to accommodate construction of the Leisure Core. Disturbance (from shipping/ferry movements) giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats.	Significant at District level (moderate adverse) (permanent). Significant at district level (minor adverse) (temporary, reversible).	Landscape Strategy - enhancement of retained wetland habitat on Swanscombe Peninsula and creation of new saltmarsh, wetland and reedbed habitats, managed via the EMMF (Document Reference 6.2.12.3). Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) Bird Monitoring Response Strategy (BMRS) — to monitor bird activity and inform the development of further control/mitigation measures as required. Off-site Ecological Mitigation — delivered in accordance with the	No significant effect. No significant effect.
			'General Principles for Offsite Ecological Mitigation' (Document	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement	Residual Effect including Significance
			Mitigation	
			Reference 6.2.12.10), to offset any	
			residual effects.	
	Disturbance (from	Significant at District	BMRS - to monitor bird activity and	No significant effect.
	shipping/ferry movements)	<u>level (minor adverse)</u>	inform the development of further	
	giving rise to displacement,	(temporary, reversible).	control/mitigation measures as	
	<u>behavioural changes or</u>		required.	
	physiological stress to cited			
	bird species potentially using		Creation/Enhancement of 259.33ha	
	functionally linked habitats.		Functionally Linked Habitat Off-site	
			<u>– delivered in accordance with the</u>	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats	
	Disturbance (noise and	Significant at district	CEMP and SLS - Restricted working	No significant effect. No
	lighting) associated with	level <u>District level</u>	hours and sensitive design and	significant effect.
	increased shipping traffic and	(moderate adverse)	location of new lighting to minimise	
	construction activities, giving	(temporary, reversible).	impacts on nocturnal/crepuscular	
	rise to displacement,		wildlife.	
	behavioural changes or			
	physiological stress to bird species potentially using		CEMP – sensitive timing of works	
	functionally linked habitats.		causing in excess of 55dB of noise at	
	ranctionally linked habitats.		the river frontage or functionally	
			linked habitat, to take place in	
			summer months, where possible, as	
			detailed in the Bird Mitigation	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			Strategy enclosed within EMMF	
			(Document Reference 6.2.12.3).	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
			Creation/Enhancement of 259.33ha	
			Functionally Linked Habitat Off-site	
			 delivered in accordance with the 	
			ECF (Document Reference 6.2.12.10)	
			to address any residual disturbance	
			effects on retained habitats	
	Disturbance (human	Significant at district	CEMP – use of site	No significant effect.No
	movement or activity), to bird	level District level	screening/hoarding; sensitive timing	significant effect.
	species potentially using	(minor adverse)	of works within 300m of the river	
	functionally linked habitats, as	(temporary, reversible).	frontage or functionally linked	
	a result of increased human		habitat and visible from that habitat	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	presence on the Project Site during construction.		to take place in summer months, where possible; and access by construction workers to river front restricted within 500m of visible wetland habitats, where possible; as detailed in the Bird Mitigation Strategy enclosed within the EMMF (Document Reference 6.2.12.3). BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.	
Deputations of	N.D. Mitigation of actorial	Cionificant et regional	Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site - delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats	Cincificant acciding
Populations of nationally scarce plants	N.B. Mitigation of potential effects upon scarce plants included under Swanscombe Peninsula SSSI. Direct loss of individual plants, and part of plant meta-population through	Significant at regional level (major adverse) (permanent).	Landscape Strategy - enhancement of retained habitats on Swanscombe Peninsula and creation of new habitats, managed via the EMMF (Document Reference 6.2.12.3).	Significant positive effect (local level).



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	loss of habitat in the Kent		Rare Plant Mitigation Strategy	
	Project Site.		enclosed within the EMMF -	
			providing details for the	
			translocation of nationally scarce	
			plants/seed bank occurring on the	
			Kent Project Site.	
Broad leaved Semi	Direct loss of habitat from	Significant at local	Landscape Strategy - enhancement	Significant negative at
Natural Woodland	woodland south of Black Duck	levelSignificant at Local	of retained habitats on Swanscombe	Local level. No
	Marsh – 3.84ha lost from a	<u>level</u> (major adverse)	Peninsula and creation of new	significant effect.
	total of 21.37ha (18% of the	(permanent).	habitats, managed via the EMMF	
	existing total). Direct loss of		(Document Reference 6.2.12.3).	(potential for significant
	habitat.			positive effect in the
			Off-site Ecological Mitigation -	long term subject to
			delivered in accordance with the	final off-site mitigation
			'General Principles for Offsite	package).
			Ecological Mitigation' (Document	
			Reference 6.2.12.10).	
	Damage to habitat from	Significant at local	CEMP – including details on	No significant effect. No
	changes from construction	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	dust, waste and pollutants.	<u>level</u> (minor adverse)	construction waste, and pollution	
		(temporary).	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions.	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
Ancient woodland	Damage to habitats within the	Significant at District	CEMP – including details on	No significant effect.
	designated area from changes	level (minor adverse)	safe/appropriate disposal of	
	in air quality, including from	(temporary).	construction waste, and pollution	
	dust, construction waste and		prevention guidelines, to be strictly	
	pollutants, and exhaust		adhered to. CEMP will also include a	
	emissions.		Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions. The CEMP will	
			secure a suite of construction	
			mitigation measures as detailed in	
			Chapter 16: Air Quality (Document	
			reference 6.1.16), relating to site	
			management/-maintenance, site	
			monitoring, operation of vehicles/	
			machinery/-power tools, and	
			techniques for demolition works,	
			earthworks and construction.	
Scrub	Direct loss of habitat, a total of	Significant at local	Landscape Strategy and EMMF	Significant negative at
	34.79ha lost (41.71% of the	levelSignificant at Local	enhancement of retained scrub	Local level. No
	existing total) Direct loss of	<u>level</u> (major adverse)	habitats and commencement of	significant effect.
	habitat.	(permanent).	rotational management to avoid	
			excessive encroachment and to	(potential for significant
			maintain varied structure and open	positive effect in the
			mosaic habitat. Landscape Strategy	long term subject to
			& FMMF — enhancement of	final off-site mitigation
			retained scrub habitats and	package).



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			commencement of rotational	
			management to avoid excessive	
			encroachment and to maintain	
			varied structure and open mosaic	
			habitat.	
			Off site Ecological Mitigation –	
			delivered in accordance with the	
			'General Principles for Offsite	
			Ecological Mitigation' (Document	
			Reference 6.2.12.10), to offset any	
			residual effects.	
	Damage to habitats from	Significant at local	CEMP and EMMF – detailing	No significant effect. No
	adjacent construction works	levelSignificant at Local	sensitive construction methodology,	significant effect.
	and deposition of construction	<u>level</u> (minor adverse)	including set up and maintenance of	
	materials in Root Protection	(temporary).	EPZs where no construction	
	Area (RPA).		vehicles/works are permitted and	
			where no materials or site facilities	
		Temporary, minor	may be situated.	
		magnitude and extent.		
	Damage to habitat from	Significant at local	CEMP – including details on	No significant effect. No
	construction dust, waste and	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	pollutants.	<u>level</u> (minor adverse)	construction waste, and pollution	
		(temporary).	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	



	including Significance	Enhancement Avoidance and	including Significance
			, 5 5
		Mitigation	
· · · · · · · · · · · · · · · · · · ·		provide measures to control dust	
		and other emissions.	
Damage to habitats from	Significant at district	Non-native Invasive Plant Species	No significant effect. No
introduction or proliferation of	level District level	Mitigation Strategy included within	significant effect.
Invasive Non-Native Species	(minor adverse)	EMMF (Document Reference	
(INNS) INNS.	(temporary, reversible).	6.2.12.3) – including details on	
		control/eradication of existing	
		populations of non-native species.	
		CEMP – to include control measures	
Direct loss of habitat, 36.19ha	Significant at local	Landscape Strategy and EMMF –	Significant negative at
lost (44.79% of the existing	levelSignificant at Local	management of existing, retained	Local level. No
total)Direct loss of habitat.	<u>level</u> (major adverse)	habitats on Swanscombe Peninsula	significant effect.
	(permanent).	through commencement of	
			(potential for significant
		_	positive effect in the
			long term subject to
			final off-site mitigation
			package).
		wildilower grassiand.	
i 4 (ntroduction or proliferation of nvasive Non-Native Species (INNS) INNS. Direct loss of habitat, 36.19ha ost (44.79% of the existing	Introduction or proliferation of Invasive Non-Native Species (minor adverse) (temporary, reversible). Direct loss of habitat, 36.19ha ost (44.79% of the existing total) Direct loss of habitat. Significant at local level (major adverse)	INNS) INNS. Level District level (minor adverse) (temporary, reversible). Level District level (minor adverse) (temporary, reversible).



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation Off site Ecological Mitigation— delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any	Residual Effect including Significance
	Damage to habitats from adjacent construction works and deposition of construction materials.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	residual effects. CEMP and EMMF – detailing sensitive construction methodology, including set up and maintenance of EPZs where no construction vehicles/works are permitted and where no materials or site facilities may be situated.	No significant effect.No significant effect.
	Damage to habitat from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect.No significant effect.
	Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Significant at district level District level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on	No significant effect.No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			control/eradication of existing	
			populations of non-native species.	
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	
Coastal/floodplain	Direct loss of habitat, 13.74ha	Significant at district	Off site Ecological Mitigation –	Significant negative at
grazing marsh	lost (100% of the existing	level District level	delivered in accordance with the	District level. No
	total). All CFGM is also included	(major adverse)	'General Principles for Offsite	significant effect.
	under the Floodplain Wetland	(permanent).	Ecological Mitigation' (Document	
	Mosaic grouping, within the		Reference 6.2.12.10).No on-site	(potential for significant
	biodiversity impact calculations		measures possible/proposed.	positive effect in the
	(Document reference 6.2,			long term subject to
	Appendix 12.2). Direct loss of			final off-site mitigation
	habitat.			package).
Open mosaic on	Direct loss of habitat, 39.83ha	Significant at district	Landscape Strategy, and	Significant negative at
previously	lost (63.7% of the existing	level District level	Swanscombe Peninsula SSSI	District level. No
developed land	total), includes a range of	(major adverse)	Mitigation StrategyInvertebrate	significant effect.
	individual habitat components including areas of bare ground,	(permanent).	Mitigation Strategy included in the	
	ephemeral vegetation, open		EMMF – enhancement of existing	
	grassland and patches of		retained open mosaic habitat (OMH)	
	scrub. Direct loss of habitat.		through introduction of greater	
	Sold Sold Prince 1005 Of Habitati		variety and complexity, maintaining	
			a range of microhabitats, foodplants	
			and nectar sources. Measures	
			include creation of bare ground	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and	Residual Effect including Significance
		including Significance	Mitigation	including Significance
			scrapes, creation of shallow pools of	
			varying depth, creation of	
			piles/mounds of mixed crushed and	
			coarse concrete rubble, and creation	
			of chalk mounds and low bunds.	
			Furthermore, creation of brown roof	
			habitat on new buildings using	
			substrate (crushed concrete and	
			chalk) originating from the Proposed	
			Development footprint.	
			Development tootprint.	
			Off-site Ecological Mitigation -	
			delivered in accordance with the	
			General Principles for Offsite Ecological Mitigation (Document	
			(
			Reference 6.2.12.10), will also create	
			additional valuable invertebrate	
			habitat.	
	Damage to habitats from	Significant at local	CEMP and EMMF – detailing	No significant effect. N
	adjacent construction works	levelSignificant at Local	sensitive construction methodology,	significant effect
	and deposition of construction	<u>level</u> (minor adverse)	including set up and maintenance of	
	materials.	(temporary).	EPZs where no construction	
			vehicles/works are permitted and	
			where no materials or site facilities	
			may be situated.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	Damage to habitat from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect.No significant effect
	Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Significant at district levelDistrict level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species. CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	No significant effect.No significant effect
Waterbodies (ponds, standing water and ditch network)	Direct loss of habitat, 2.30ha of waterbodies lost (50.79% of the existing total) and 4.7km of ditches (50.98% of the existing total). 0.22ha of lost waterbodies are also included under the Floodplain Wetland Mosaic grouping, within the	Significant at district level District level (major adverse) (permanent).	Landscape Strategy - Creation of new waterbodies/wetlands on Swanscombe Peninsula, including new perimeter swale, new constructed wetland, new ponds and ditches, and water vole receptor site, managed through the EMMF (Document Reference 6.2.12.3).	Significant negative at District level. Significant positive effect (local level).



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation	Residual Effect including Significance
	biodiversity impact calculations (Document reference 6.2.12.2, Appendix 12.2). Direct loss of habitat. Damage to habitats from changes to hydrological	Significant at local levelSignificant at Local	Off-site Ecological Mitigation – delivered in accordance with the General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10). CEMP - Prevention of hydrological impacts through adherence to an	No significant effect. No significant effect.
	regime (through destruction of surrounding habitats), and changes in water quality/quantity.	level (moderate adverse) (temporarypermanent).	appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	<u></u>
	Damage to habitat from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect. No significant effect.
	Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS)INNS.	Significant at district level of the level of the level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within the EMMF (Document Reference 6.2.12.3) – including details on	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation control/eradication of existing populations of non-native species. CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	Residual Effect including Significance
Swamp (reedbed)	Direct loss of habitat from 13.62ha lost (41.75% of total area). 2.84ha of reedbed is also included under the OMHPDL grouping and 0.88ha under Floodplain Wetland Mosaic, within the biodiversity impact calculations (Document reference 6.2, Appendix 12.2). Direct loss of habitat.	Significant at <u>Ceounty</u> level (major adverse) (permanent).	Landscape Strategy - Creation of new waterbodies/wetlands on Swanscombe Peninsula, including new perimeter swale, and new constructed wetland, new ponds and ditches, and water vole receptor site, managed through the EMMF (Document Reference 6.2.12.3). Off site Ecological Mitigation delivered in accordance with the General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10).	Significant negative at County level. No significant effect. (potential for significant positive effect in the long term subject to final off-site mitigation package).
	Damage to habitats from adjacent construction works and deposition of construction materials.	Significant at local level (minor adverse) (temporary).	cemp and emmf – detailing sensitive construction methodology, including set up and maintenance of EPZs where no construction vehicles/works are permitted and	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			where no materials or site facilities	
			may be situated.	
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.
	regime (through destruction of	<u>level</u> (moderate	appropriate Surface Water	
	surrounding habitats), and	adverse) (temporary).	Management Strategy, as outlined	
	changes in water		within Chapter 17 – Water	
	quality/quantity.		Resources and Flood Risk (Document	
			Reference 6.1.17).	
			,	
	Damage to habitat from	Significant at local	CEMP – including details on	No significant effect. No
	construction dust, waste and	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	pollutants.	level (minor adverse)	construction waste, and pollution	
		(temporary)	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions.	
	Damage to habitats from	Significant at district	Non-native Invasive Plant Species	No significant effect. No
	introduction or proliferation of	level District level	Mitigation Strategy included within	significant effect.
	Invasive Non-Native Species	(minor adverse)	EMMF (Document Reference	
	(INNS) INNS.	(temporary, reversible).	6.2.12.3) – including details on	
			control/eradication of existing	
			populations of non-native species.	
			·	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
River Ebbsfleet	Direct loss of connected	Significant at local	Landscape Strategy - Creation of	No significant effect. No
	floodplain habitat within	levelSignificant at Local	new waterbodies/wetlands on	significant effect.
	Ebbsfleet Valley, such as	<u>level</u> (moderate	Swanscombe Peninsula, including	
	waterbodies and semi-	adverse) (permanent).	new perimeter swale and new	
	improved grassland. Direct loss		constructed wetland, new	
	of connected floodplain		constructed wetland, new ponds	
	habitat.		and ditches, and water vole receptor	
			site, managed through the EMMF	
			(Document Reference 6.2.12.3).	
			Off-site Ecological Mitigation -	
			delivered in accordance with the	
			General Principles for Offsite	
			Ecological Mitigation (Document	
			Reference 6.2.12.10).	
	Damage to habitat from	Significant at local	CEMP – including details on	No significant effect. No
	construction dust, waste and	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	pollutants.	<u>level</u> (minor adverse)	construction waste, and pollution	
		(temporary).	prevention guidelines, to be strictly	
			adhered to. CEMP will also include a	
			Dust Management Plan (DMP) to	
			provide measures to control dust	
			and other emissions.	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect		
		including Significance	Enhancement Avoidance and	including Significance		
			Mitigation			
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No		
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.		
	regime (through destruction of	<u>level</u> (moderate	appropriate Surface Water			
	surrounding habitats), and	adverse) (temporary).	Management Strategy, as outlined			
	changes in water		within Chapter 17 – Water			
	quality/quantity.		Resources and Flood Risk (Document			
			Reference 6.1.17).			
	Damage to habitats from	Significant at district	Non-native Invasive Plant Species	No significant effect. No		
	introduction or proliferation of	level District level	Mitigation Strategy included within	significant effect.		
	Invasive Non-Native Species	(minor adverse)	the EMMF (Document Reference			
	(INNS) INNS.	(temporary, reversible).	6.2.12.3) – including details on			
			control/eradication of existing			
			populations of non-native species.			
			CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.			
Wintering	N.B. Assessment of potential hab	pitat loss and disturbance (effects upon wintering waterfowl and w	vading bird assemblage		
Waterfowl and	included under SPA/Ramsar Sites	.Direct killing, injury or ha	irm to individuals – during vegetation c l	earance.		
Wading Bird	Significant at district level (major	adverse) (permanent).				
Assemblage	CEMP, and Breeding and Winter Bird Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including					
	sensitive timing and methodology for habitat clearance, establishment of EPZs to protect retained habitats.					
	No significant effect.					
	(potential for significant positive	effect in the long term su	bject to final off site mitigation package	2).		



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Direct killing, injury or harm to	Significant at District	CEMP , and SPA and Winter Bird	No significant effect.
	individuals – during vegetation	level (major adverse)	Mitigation Strategy included within	
	<u>clearance.</u>	(permanent).	EMMF (Document Reference	
			6.2.12.3) – including sensitive timing	
			and methodology for habitat	
			clearance, establishment of EPZs to	
			protect retained habitats.	
Wintering	Direct killing, injury or harm to	Significant at district	CEMP, and SPA and Winter Bird	No significant effect. No
terrestrial bird	individuals – during vegetation	level District level	Mitigation Strategy Breeding and	significant effect.
assemblage	clearance.	(major adverse)	Winter Bird Mitigation Strategy	
		(permanent).	included within EMMF (Document	
			Reference 6.2.12.3) – including	
			sensitive timing and methodology	
			for habitat clearance, establishment	
			of EPZs to protect retained habitats.	
	<u>Direct habitat loss</u> – loss of	Significant at district	Landscape Strategy and SPA and	Significant negative at
	suitable habitat such as scrub,	level District level	Winter Bird Mitigation	<u>District level.</u> No
	grassland, reedbed and	(moderate adverse)	Strategy Breeding and Winter Bird	significant effect.
	woodland, as detailed	(permanent).	Mitigation Strategy included within	
	above. Direct habitat loss.		EMMF - enhancement of retained	(potential for significant
			scrub habitat through rotational	positive effect in the
			management to create age class and	long term subject to
			structural diversity, and	final off-site mitigation
			maintenance of site wide open	package).
			mosaic habitats to provide variety of	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
Lcological Feature	Damage to functionally linked habitats from changes to hydrological regime and changes to water and/or sediment quality — as described above in relation to impacts on SPA/Ramsar sites. Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality — as described	_	. 0	
	above in relation to impacts on SPA/Ramsar sites. Damage to supporting habitats from construction	Significant at district level	CEMP – including details on safe/appropriate disposal of	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	dust, waste and pollutants. Damage to supporting habitats from construction dust, waste and pollutants.	(minor adverse) (temporary).	construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	
	Damage to functionally linked habitats from introduction or proliferation of INNS – as described above in relation to impacts on SPA/Ramsar sites.Damage to habitats from introduction or proliferation of Invasive Non-Native Species (INNS) – as described above in relation to impacts on SPA/Ramsar sites.	Significant at district level level District level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species. CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	No significant effect. No significant effect.
	Habitat fragmentation, loss of flight paths and dispersal routes.	Significant at district level District level (moderate adverse) (permanent).	Landscape Strategy and EMMF (Document Reference 6.2.12.3)— maintenance of scrub habitats across the Peninsula to allow terrestrial bird species to disperse between habitats.	Significant negative at District level. No significant effect.
	Disturbance (from shipping/ferry movements) – as described above in relation	Significant at district level District level (minor adverse) (temporary, reversible).	Bird Monitoring Response Strategy (BMRS)BMRS - monitoring of bird use on the intertidal habitats and other functionally linked land – if the	Significant negative at District level. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	to impacts on SPA/Ramsar		numbers of birds within the	
	sites.		monitored area fall below a certain	
			threshold in response to obvious	
			construction activities then those	
			disturbance activities will be	
			temporally ceased.	
			Off-site Ecological Mitigation –	
			delivered in accordance with the	
			'General Principles for Offsite	
			Ecological Mitigation' (Document	
			Reference 6.2.12.10), to offset any	
			residual effects.	
	Disturbance (noise and	Significant at district	CEMP and SLS - Restricted working	Significant negative at
	lighting) – as described above	level <u>District level</u>	hours and sensitive design and	<u>District level.</u> No
	in relation to impacts on	(moderate adverse)	location of new lighting to minimise	significant effect.
	SPA/Ramsar sites.	(temporary, reversible).	impacts on nocturnal/crepuscular	
			wildlife.	
			CEMP – sensitive timing of works	
			causing in excess of 55dB of noise at	
			the river frontage or functionally	
			linked habitat, to take place in	
			summer months, where possible, as	
			detailed in the Bird Mitigation	
			Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3).	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation CEAR adoption of mitigation	
			<u>CEMP – adoption of mitigation</u> measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
			<u></u>	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
			CEMP and Sensitive Lighting Strategy	
			(SLS) Restricted working hours and	
			sensitive design and location of new	
			lighting to minimise impacts on	
			nocturnal/crepuscular wildlife.	
			, , , , , , , , , , , , , , , , , , , ,	
			CEMP – sensitive timing of works	
			causing in excess of 55dB of noise at	
			the river frontage or functionally	
			linked habitat, to take place in	
			summer months, where possible, as	
			detailed in the Bird Mitigation	
			Strategy enclosed within the EMMF	
			(Document Reference 6.2.12.3).	
			(Seeding Reference 0.2.12.3).	
		<u> </u>		<u> </u>



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
	Disturbance (human	Significant at district	CEMP – use of site	Significant negative a
	movement or activity) – as	level District level	screening/hoarding; sensitive timing	<u>District level</u> .No
	described above in relation to	(moderate adverse)	of works within 300m of the the	significant effect.
	impacts on SPA/Ramsar sites.	(temporary, reversible).	river frontage or functionally linked	
			habitat and visible from that habitat	
			to take place in summer months,	
			where possible; and access by	
			construction workers to the river	
			front restricted within 500m of	
			visible wetland habitats, where	
			possible; as detailed in the Bird	
			Mitigation Strategy enclosed within	
			EMMF (Document Reference	
			6.2.12.3).	
			0.2.12.5).	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect	
		including Significance	Enhancement Avoidance and	including Significance	
			Mitigation		
			BMRS – to monitor bird activity and		
			inform the development of further		
			control/mitigation measures as		
			required.		
Breeding Bird	N.B. Impact assessment for the b	preeding bird assemblage	included under Swanscombe Peninsula	SSSI. Direct killing, injury	
Assemblage	or harm to individuals – during v	egetation clearance.			
	Significant at district level (major	, , ,			
			ncluded within EMMF (Document Refer		
	sensitive timing and methodology for habitat clearance, establishment of EPZs to protect retained habitats.				
	No significant effect. Included within Swanscombe Peninsula SSSI				
Bat Assemblage	Direct killing, injury or harm to	Significant at local	CEMP, and Bat Mitigation Strategy	No significant effect. No	
	individuals – during demolition	levelSignificant at Local	included within EMMF (Document	significant effect.	
	of buildings with bat roost	<u>level</u> (major adverse)	Reference 6.2.12.3)— including pre-		
	potential and trees with bat	(permanent).	commencement surveys of		
	roost potential.		structures/trees to be		
			demolished/felled, and receipt of		
			European Protected Species		
			Mitigation Licence (EPSML), or site		
			registration under the Bat Mitigation		
			Class Licensing (BMCL) scheme, prior		
			to works commencing, with		
			precautionary method of working		
			(as prescribed by the license), strictly		
			adhered to. Establishment of EPZs to		
			protect retained habitats.		



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Direct loss of potential	Significant at local	Landscape Strategy and Bat	No significant effect. No
	roosting habitat – including	levelSignificant at Local	Mitigation Strategy included in	significant effect.
	trees <u>,</u> and buildings <u>and</u>	-District level (major	EMMF Document Reference	
	structures with bat roost	adverse) (permanent).	6.2.12.3)— creation of new bat	
	potential.		roosting habitat, including bat boxes	
			incorporated into new buildings, and	
			into the two new bird watching	
			towers-, and enhancement of	
			hibernation habitat within retained	
			tunnels	
	Direct loss of foraging,	Significant at local	Landscape Strategy and EMMF	Significant negative at
	commuting and dispersal	levelSignificant at Local	(Document Reference 6.2.12.3) –	Local level. No
	habitats - loss of suitable	<u>level</u> (moderate	creation/maintenance of a range of	significant effect.
	foraging habitat such as scrub,	adverse) (permanent).	habitats including woodland, scrub,	
	grassland, reedbed and		wetlands, waterbodies, reedbeds,	
	woodland, as detailed above,		saltmarsh and invertebrate rich	
	including potential winter		habitats, collectively will maintain	
	foraging habitat. Direct loss of		foraging and commuting bat	
	foraging, commuting and		population on site.	
	dispersal habitats.			
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.
	regime and changes to water	<u>level</u> (moderate	appropriate Surface Water	
	and/or sediment quality – as	adverse) (temporary).	Management Strategy, as outlined	
	described above in relation to		within Chapter 17 – Water	
	impacts on habitats. Damage to		Resources and Flood Risk (Document	
	habitats from changes to		Reference 6.1.17).	
			NEIGIGING 0.1.1/J.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	hydrological regime and changes to water and/or sediment quality — as described above in relation to impacts on habitats. Damage to supporting habitats from construction dust, waste and pollutants. Damage to supporting habitats from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect. No significant effect.
	Damage to supporting habitats from introduction or proliferation of INNS. Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS).	Significant at local levelSignificant at Local level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species. CEMP – to include control measures to prevent spread of INNS and INNS impacted soils.	No significant effect. No significant effect.
	Habitat fragmentation, loss of flight paths and dispersal routes.	Significant at local levelSignificant at Local	Landscape Strategy and EMMF (Document Reference 6.2.12.3) — maintenance of connectivity across	Significant negative at Local levelNo significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
		<u>level</u> (moderate	the Peninsula through the inclusion	
		adverse) (permanent).	of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint, maintenance of scrub	
			habitats across the Peninsula, and	
			retention of habitat link to the River	
			Thames to allow bats to disperse	
			between habitats on- and off-site.	
	Disturbance (noise and	Significant at local	CEMP and SLS - Restricted working	No significant effect. No
	lighting) – including on	levelSignificant at Local	hours and sensitive design and	significant effect.
	retained roosts, foraging areas	<u>level</u> (moderate	location of new lighting to minimise	
	and commuting routes.	adverse) (temporary,	impacts on nocturnal/crepuscular	
		reversible).	wildlife.	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
Dormouse	Direct killing, injury or harm to	Significant at district	CEMP, and Dormouse Mitigation	No significant effect. No
	individuals – during vegetation	level District level	Strategy included within EMMF	significant effect.
	clearance.	(major adverse)	(Document Reference 6.2.12.3) –	
		(permanent).	including pre-commencement	
			surveys of suitable dormouse	
			habitat, and receipt of European	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			Protected Species Mitigation Licence	
			(EPSML) prior to habitat clearance	
			commencing, with dormouse	
			displacement methodologies and	
			precautionary method of working	
			(as prescribed by the license), strictly	
			adhered to. Establishment of EPZs to	
			protect retained habitats.	
	<u>Direct habitat loss – loss of</u>	Significant at district	Landscape Strategy and Dormouse	Significant negative at
	suitable habitat such as scrub	level District level	Mitigation Strategy included EMMF	District level. No
	and woodland, as detailed	(moderate adverse)	(Document Reference 6.2.12.3) –	significant effect.
	above. Direct habitat loss.	(permanent).	enhancement of existing scrub	
			habitats to ensure habitats remain	(potential for significant
			suitable for dormice, through	positive effect in the
			rotational scrub management to	long term subject to
			improve structural and age class	final off-site mitigation
			diversity and maintain variety of	package).
			food sources throughout the	
			dormouse active season.	
			Off-site Ecological Mitigation -	
			delivered in accordance with the	
			'General Principles for Offsite	
			Ecological Mitigation' (Document	
			Reference 6.2.12.10), to mitigate	
			any residual effects.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats. Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats.	Significant at district level District level (moderate adverse) (temporary).	CEMP - Prevention of hydrological impacts through adherence to an appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect. No significant effect.
	Damage to supporting habitats from construction dust, waste and pollutants. Damage to supporting habitats from construction dust, waste and pollutants.	Significant at district levelDistrict level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect. No significant effect.
	Damage to supporting habitats from introduction or proliferation of INNS. Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS).	Significant at district level District level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and	Residual Effect including Significance
		including Significance	Mitigation	including Significance
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation and	Significant at district	Landscape Strategy and Dormouse	Significant negative at
	loss of dispersal routes. Habitat	level District level	Mitigation Strategy included within	District level. No
	fragmentation and loss of	(moderate adverse)	EMMF (Document Reference	significant effect.
	dispersal routes.	(permanent).	6.2.12.3) –retention of areas of	
			suitable habitat within Broadness	
			grassland, including existing	
			woodland, tree and scrub habitats;	
			retention and management of dense	
			bramble and low growing scrub and	
			trees along the north western	
			boundaries of Black Duck Marsh, to	
			maintain potential dispersal routes	
			from Swanscombe Peninsula to the	
			south-west; retention and	
			enhancement of a continuous belt of	
			woodland habitat along the	
			southern boundaries of Black Duck	
			Marsh, connecting to additional	
			green corridors proposed along the	
			southern boundary adjacent to	
			Tiltman Avenue, to ensure the	
			continued functioning of existing	
			dispersal routes to valuable off-site	
			habitats to the south-west, including	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			the woodland at the Swanscombe	
			Heritage Park; and retention,	
			enhancement and creation of	
			additional woodland habitat	
			alongside the sensitive design of	
			new landscaping around the	
			peripheries of Botany Marsh,	
			necessary to further promote	
			habitat connectivity between	
			Swanscombe Peninsula and habitats	
			across the wider landscape to the	
			south.	
	Disturbance (noise and	Significant at district	CEMP and SLS - Restricted working	No significant effect. No
	<u>lighting</u>) – as described above	level <u>District level</u>	hours and sensitive design and	significant effect.
	in relation to impacts on	(moderate adverse)	location of new lighting to minimise	
	SPA/Ramsar sites. Disturbance	(temporary, reversible).	impacts on nocturnal/crepuscular	
	(noise and lighting) – as		wildlife.	
	described above in relation to			
	impacts on SPA/Ramsar sites.		CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
Otter	Direct killing, injury or harm to	Significant at local	CEMP, and Otter Mitigation	No significant effect. No
	individuals – during vegetation	levelSignificant at Local	Strategy included within EMMF	significant effect.
	clearance.		(Document Reference 6.2.12.3) –	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation	Residual Effect including Significance
		level (major adverse) (permanent).	including pre-commencement surveys of suitable habitat, sensitive habitat clearance methodology, and establishment of EPZs to protect retained habitats. In the event an otter holt is found within the Proposed Development footprint, a European Protected Species Mitigation Licence (EPSML) will be needed prior to habitat clearance commencing.	
	Direct habitat loss – loss of suitable habitat such as reedbed and waterbodies, as detailed above. Direct habitat loss.	Significant at local levelSignificant at Local level (moderate adverse) (permanent).	Landscape Strategy - new constructed wetland, -managed through the EMMF (Document Reference 6.2.12.3). Creation of two new artificial otter holts as described in the Otter Mitigation Strategy included in the EMMF. Landscape Strategy - Creation of new wetlands on Swanscombe Peninsula, including new constructed wetland, and new ponds and ditches, managed through the EMMF (Document Reference 6.2.12.3). Creation of two new artificial otter holts as described in the Otter Mitigation Strategy included in the EMMF.	Significant negative at Local level. Significant positive effect (local level).



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.
	regime and changes to water	<u>level</u> (moderate	appropriate Surface Water	
	and/or sediment quality – as	adverse) (temporary).	Management Strategy, as outlined	
	described above in relation to		within Chapter 17 – Water	
	impacts on habitats. Damage to		Resources and Flood Risk (Document	
	habitats from changes to		Reference 6.1.17).CEMP -	
	hydrological regime and		Prevention of hydrological impacts	
	changes to water and/or		through adherence to an	
	sediment quality – as described		appropriate Surface Water	
	above in relation to impacts on		Management Strategy, as outlined	
	habitats.		within Chapter 17 – Water	
			Resources and Flood Risk (Document	
			Reference 6.1.17).	
	Damage to supporting	Significant at local	CEMP – including details on	No significant effect. No
	habitats from construction	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	dust, waste and	<u>level</u> (minor adverse)	construction waste, and pollution	
	pollutants. Damage to	(temporary).	prevention guidelines, to be strictly	
	supporting habitats from		adhered to. CEMP will also include a	
	construction dust, waste and		Dust Management Plan (DMP) to	
	pollutants.		provide measures to control dust	
			and other emissions. CEMP -	
			including details on safe/appropriate	
			disposal of construction waste, and	
			pollution prevention guidelines, to	
			be strictly adhered to. CEMP will	
			also include a Dust Management	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			Plan (DMP) to provide measures to	
			control dust and other emissions.	
	Damage to supporting	Significant at local	Non-native Invasive Plant Species	No significant effect. No
	habitats from introduction or	levelSignificant at Local	Mitigation Strategy included within	significant effect.
	proliferation of INNS. Damage	<u>level</u> (minor adverse)	EMMF (Document Reference	
	to supporting habitats from	(temporary, reversible).	6.2.12.3) – including details on	
	introduction or proliferation of		control/eradication of existing	
	Invasive Non Native Species		populations of non-native species.	
	(INNS).			
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils. Non-native Invasive	
			Plant Species Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) - including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
			·	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation and	Significant at local	Landscape Strategy and EMMF	Significant negative at
	loss of dispersal routes. Habitat	levelSignificant at Local	(Document Reference 6.2.12.3) –	Local level. No
	fragmentation and loss of	level (moderate	maintenance of connectivity across	significant effect.
	dispersal routes.	adverse) (permanent).	the Peninsula through the inclusion	3



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint. Creation of new wetland	
			and saltmarsh habitats on	
			Swanscombe Peninsula. Landscape	
			Strategy and EMMF (Document	
			Reference 6.2.12.3) – maintenance	
			of connectivity across the Peninsula	
			through the inclusion of a chain of	
			watercourses and wetland areas	
			wrapping around the side of the	
			Proposed Development footprint.	
			Creation of new wetland and	
			saltmarsh habitats on Swanscombe	
			Peninsula.	
	Disturbance (noise and	Significant at local	CEMP and SLS – Restricted working	No significant effect. N
	lighting) – as described above	levelSignificant at Local	hours and sensitive design and	significant effect.
	in relation to impacts on	level (moderate	location of new lighting to minimise	
	SPA/Ramsar sites. Disturbance	adverse) (temporary,	impacts on nocturnal/crepuscular	
	(noise and lighting) — as	reversible).	wildlife.	
	described above in relation to	,		
	impacts on SPA/Ramsar sites.		CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			reference 6.2.15.5). CEMP and SLS	
			Restricted working hours and	
			sensitive design and location of new	
			lighting to minimise impacts on	
			nocturnal/crepuscular wildlife.	
Water Vole	Direct killing, injury or harm to	Significant at local	CEMP, and Water Vole Mitigation	No significant
	individuals – during vegetation	levelSignificant at Local	Strategy included within EMMF	effect. Significant
	clearance. Direct killing, injury	<u>level</u> (major adverse)	(Document Reference 6.2.12.3) –	positive effect (local
	or harm to individuals – during	(permanent).	including pre-commencement	level).
	vegetation clearance.		surveys of suitable habitat,	
			establishment of suitable receptor	
			site, and water vole relocation under	
			licence from Natural England	
			(granted prior to habitat clearance	
			commencing). Establishment of EPZs	
			to protect retained habitats. CEMP,	
			and Water Vole Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) – including pre-	
			commencement surveys of suitable	
			habitat, establishment of suitable	
			receptor site, and water vole	
			relocation under licence from	
			Natural England (granted prior to	
			habitat clearance commencing).	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and	Residual Effect including Significance
		merading oignificance	Mitigation	merading significance
			Establishment of EPZs to protect	
			retained habitats.	
	Direct habitat loss – including	Significant at local	Landscape Strategy - Creation of	Significant negative at
	6.1km of suitable ditches and	levelSignificant at Local	new wetlands on Swanscombe	Local level. No
	bankside habitat and reedbed	level (moderate	Peninsula, as documented within the	significant effect.
	(as detailed above). Direct	adverse) (permanent).	Water Vole Mitigation Strategy	
	habitat loss – including 6.1km		included within EMMF (Document	
	of suitable ditches and		Reference 6.2.12.3). Landscape	
	bankside habitat and reedbed.		Strategy Creation of new wetlands,	
			ditches, ponds and waterbodies on	
			Swanscombe Peninsula, as	
			documented within the Water Vole	
			Mitigation Strategy included within	
			EMMF (Document Reference	
			6.2.12.3).	
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.
	regime and changes to water	<u>level</u> (moderate	appropriate Surface Water	
	and/or sediment quality – as	adverse) (temporary).	Management Strategy, as outlined	
	described above in relation to		within Chapter 17 – Water	
	impacts on habitats. Damage to		Resources and Flood Risk (Document	
	habitats from changes to		Reference 6.1.17). CEMP	
	hydrological regime and		Prevention of hydrological impacts	
	changes to water and/or		through adherence to an	
	sediment quality – as described		8	
	above in relation to impacts on		appropriate Surface Water	
	habitats.		Management Strategy, as outlined	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			within Chapter 17 – Water	
			Resources and Flood Risk (Document	
			Reference 6.1.17).	
	Damage to supporting	Significant at local	CEMP – including details on	No significant effect. No
	habitats from construction	levelSignificant at Local	safe/appropriate disposal of	significant effect.
	dust, waste and	<u>level</u> (minor adverse)	construction waste, and pollution	
	pollutants. Damage to	(temporary).	prevention guidelines, to be strictly	
	supporting habitats from		adhered to. CEMP will also include a	
	construction dust, waste and		Dust Management Plan (DMP) to	
	pollutants.		provide measures to control dust	
			and other emissions. CEMP -	
			including details on safe/appropriate	
			disposal of construction waste, and	
			pollution prevention guidelines, to	
			be strictly adhered to. CEMP will	
			also include a Dust Management	
			Plan (DMP) to provide measures to	
			control dust and other emissions.	
	Damage to supporting	Significant at local	Non-native Invasive Plant Species	No significant effect. No
	habitats from introduction or	levelSignificant at Local	Mitigation Strategy included within	significant effect.
	proliferation of INNS. Damage	<u>level</u> (minor adverse)	EMMF (Document Reference	
	to supporting habitats from	(temporary, reversible).	6.2.12.3) – including details on	
	introduction or proliferation of		control/eradication of existing	
	Invasive Non-Native Species		populations of non-native species.	
	(INNS).			
			<u>CEMP – to include control measures</u>	
			to prevent spread of INNS and INNS	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
			<u>Mitigation</u>	
			impacted soils. Non-native Invasive	
			Plant Species Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) – including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation and	Significant at local	Landscape Strategy and EMMF	Significant negative at
	loss of dispersal routes. Habitat	levelSignificant at Local	(Document Reference 6.2.12.3) –	Local level. No
	fragmentation and loss of	<u>level</u> (moderate	maintenance of connectivity across	significant effect.
	dispersal routes.	adverse) (permanent).	the Peninsula through the inclusion	
			of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint. Creation of new wetland	
			and saltmarsh habitats on	
			Swanscombe Peninsula. Landscape	
			Strategy and EMMF (Document	
			Reference 6.2.12.3) – maintenance	
			of connectivity across the Peninsula	
			through the inclusion of a chain of	
			watercourses and wetland areas	
			wrapping around the side of the	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			Proposed Development footprint. Creation of new wetland and saltmarsh habitats on Swanscombe Peninsula.	
	Disturbance (noise and lighting) — as described above in relation to impacts on SPA/Ramsar sites. Disturbance (noise and lighting) — as described above in relation to impacts on SPA/Ramsar sites.	Significant at local level Significant at Local level (moderate adverse) (temporary, reversible).	CEMP and SLS - Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP — adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5). CEMP and SLS Restricted working hours and sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. CEMP — adoption of mitigation measures to control construction noise and vibration, as described in	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation and Vibration Mitigation (Document reference 6.2.15.5).	Residual Effect including Significance
Harvest Mouse	Direct killing, injury or harm to individuals – during vegetation clearance. Direct killing, injury or harm to individuals – during vegetation clearance.	Significant at local levelSignificant at Local level (major adverse) (permanent).	CEMP, and Harvest Mouse Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) — including pre- commencement check of suitable habitat, and precautionary method of working strictly adhered to. Establishment of EPZs to protect retained habitats. CEMP, and Harvest Mouse Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) — including pre- commencement check of suitable habitat, and precautionary method of working strictly adhered to. Establishment of EPZs to protect retained habitats.	No significant effect. No significant effect.
	<u>Direct habitat loss – loss of</u> <u>suitable habitat such as</u> <u>reedbed, semi-improved</u> <u>grassland, scrub and tall</u>	Significant at local levelSignificant at Local level (moderate adverse) (permanent).	Landscape Strategy and Harvest Mouse Mitigation Strategy included within the EMMF (Document Reference 6.2.12.3) – maintenance of site wide open mosaic habitat	Significant negative at Local level. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	ruderal, as detailed above. Direct habitat loss.		including a range of habitats of value to harvest mice including rough, tussocky grassland, tall riparian/marginal/reedbed habitat with adjacent scrub to provide refuge, foraging and breeding	
			opportunities. Landscape Strategy and Harvest Mouse Mitigation Strategy included within the EMMF (Document Reference 6.2.12.3) — maintenance of site wide open mosaic habitat including a range of habitats of value to harvest mice including rough, tussocky grassland,	
			tall riparian/marginal/reedbed habitat with adjacent scrub to provide refuge, foraging and breeding opportunities.	
	Damage to habitats from changes to hydrological regime and changes to water and/or sediment quality – as described above in relation to impacts on habitats. Damage to habitats from changes to hydrological regime and changes to water and/or	Significant at local levelSignificant at Local level (moderate adverse) (temporary).	<u>CEMP - Prevention of hydrological</u> <u>impacts through adherence to an</u> <u>appropriate Surface Water</u> <u>Management Strategy, as outlined</u> <u>within Chapter 17 – Water</u> <u>Resources and Flood Risk (Document</u> <u>Reference 6.1.17).CEMP –</u> <u>Prevention of hydrological impacts</u> <u>through adherence to an</u>	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation	Residual Effect including Significance
	sediment quality — as described above in relation to impacts on habitats.		appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	
	Damage to supporting habitats from construction dust, waste and pollutants. Damage to supporting habitats from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect. No significant effect.
	Damage to supporting habitats from introduction or proliferation of INNS. Damage to supporting habitats from introduction or proliferation of Invasive Non-Native Species (INNS).	Significant at local levelSignificant at Local level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and	Residual Effect including Significance
			<u>Mitigation</u>	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils. Non-native Invasive	
			Plant Species Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) – including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation and	Significant at local	Landscape Strategy and EMMF	Significant negative at
	loss of dispersal routes. Habitat	levelSignificant at Local	(Document Reference 6.2.12.3) –	Local level. No
	fragmentation and loss of	<u>level</u> (moderate	maintenance of connectivity across	significant effect.
	dispersal routes.	adverse) (permanent).	the Peninsula through the inclusion	
			of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint, and maintenance of scrub	
			and rough grassland habitats across	
			the Peninsula to allow individuals to	
			disperse between	
			habitats.Landscape Strategy and	
			EMMF (Document Reference	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			6.2.12.3) – maintenance of	
			connectivity across the Peninsula	
			through the inclusion of a chain of	
			watercourses and wetland areas	
			wrapping around the side of the	
			Proposed Development footprint,	
			and maintenance of scrub and rough	
			grassland habitats across the	
			Peninsula to allow individuals to	
			disperse between habitats.	
	Disturbance (noise and	Significant at local	CEMP and SLS - Restricted working	No significant effect. No
	lighting) – as described above	levelSignificant at Local	hours and sensitive design and	significant effect.
	in relation to impacts on	<u>level</u> (moderate	location of new lighting to minimise	
	SPA/Ramsar sites. Disturbance	adverse) (temporary,	impacts on nocturnal/crepuscular	
	(noise and lighting) – as	reversible).	wildlife.	
	described above in relation to			
	impacts on SPA/Ramsar sites.		CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5). CEMP and SLS	
			Restricted working hours and	
			sensitive design and location of new	
			lighting to minimise impacts on	
			nocturnal/crepuscular wildlife.	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
			CEMP – adoption of mitigation measures to control construction noise and vibration, as described in Appendix 15.5: Construction Noise and Vibration Mitigation (Document reference 6.2.15.5).	
Amphibian assemblage	Direct killing, injury or harm to individuals – during vegetation clearance. Direct killing, injury or harm to individuals – during vegetation clearance.	Significant at local levelSignificant at Local level (major adverse) (permanent).	CEMP, and Amphibian Mitigation Strategy included within EMMF— including translocation of amphibians in tandem with reptile translocation, precautionary methods of working including sensitive drain down of any waterbodies to be removed, timing of works, and establishment of EPZs to protect retained habitats. CEMP, and Amphibian Mitigation Strategy included within EMMF—including translocation of amphibians in tandem with reptile translocation, precautionary methods of working including sensitive drain down of any waterbodies to be removed, timing of works, and establishment	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
	Direct habitat loss – loss of	Significant at local	Landscape Strategy and Amphibian	Significant negative at
	suitable habitat such as	levelSignificant at Local	Mitigation Strategy included within	Local level. Significant
	reedbed and waterbodies, as	<u>level</u> (moderate	EMMF - Creation of new wetlands,	positive effect (local
	<u>detailed above</u> . Direct habitat	adverse) (permanent).	ditches, ponds and waterbodies on	level).
	loss.		Swanscombe Peninsula, as described	
			above in relation to water voles and	
			otters, will also mitigate habitat loss	
			impacts to amphibians. Landscape	
			Strategy and Amphibian Mitigation	
			Strategy included within EMMF	
			Creation of new wetlands, ditches,	
			ponds and waterbodies on	
			Swanscombe Peninsula, as described	
			above in relation to water voles and	
			otters, will also mitigate habitat loss	
			impacts to amphibians.	
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.
	regime and changes to water	<u>level</u> (moderate	appropriate Surface Water	
	and/or sediment quality – as	adverse) (temporary).	Management Strategy, as outlined	
	described above in relation to		within Chapter 17 – Water	
	impacts on habitats. Damage to		Resources and Flood Risk (Document	
	habitats from changes to		Reference 6.1.17).CEMP -	
	hydrological regime and		Prevention of hydrological impacts	
	changes to water and/or		through adherence to an	
	sediment quality – as described			



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and EnhancementAvoidance and Mitigation	Residual Effect including Significance
	above in relation to impacts on habitats.		appropriate Surface Water Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	
	Damage to supporting habitats from construction dust, waste and pollutants. Damage to supporting habitats from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. CEMP – including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to	No significant effect. No significant effect.
	Damage to supporting habitats from introduction or proliferation of INNS.Damage to supporting habitats from introduction or proliferation of	Significant at local levelSignificant at Local level (minor adverse) (temporary, reversible).	control dust and other emissions. Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
	Invasive Non Native Species			
	(INNS).		CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils. Non-native Invasive	
			Plant Species Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) – including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation and	Significant at local	Landscape Strategy and EMMF –	Significant negative at
	loss of dispersal routes. Habitat	levelSignificant at Local	maintenance of connectivity across	Local level. No
	fragmentation and loss of	<u>level</u> (moderate	the Peninsula through the inclusion	significant effect.
	dispersal routes.	adverse) (permanent).	of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint. Creation of new wetland	
			and saltmarsh habitats on	
			Swanscombe Peninsula, delivered	
			through the EMMF and Landscape	
			Strategy.Landscape Strategy and	
			EMMF – maintenance of	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			<u>Mitigation</u>	
			connectivity across the Peninsula	
			through the inclusion of a chain of	
			watercourses and wetland areas	
			wrapping around the side of the	
			Proposed Development footprint.	
			Creation of new wetland and	
			saltmarsh habitats on Swanscombe	
			Peninsula, delivered through the	
			EMMF and Landscape Strategy.	
	Disturbance (noise and	Significant at local	CEMP and SLS – Restricted working	No significant effect. No
	<u>lighting</u>) – as described above	levelSignificant at Local	hours and sensitive design and	significant effect.
	in relation to impacts on	<u>level</u> (moderate	location of new lighting to minimise	
	SPA/Ramsar sites. Disturbance	adverse) (temporary,	impacts on nocturnal/crepuscular	
	(noise and lighting) – as	reversible).	wildlife.	
	described above in relation to			
	impacts on SPA/Ramsar sites.		CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5). CEMP and SLS	
			Restricted working hours and	
			sensitive design and location of new	
			lighting to minimise impacts on	
			nocturnal/crepuscular wildlife.	
			Hoctarnal/crepascular whaline:	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5).	
Reptile	Direct killing, injury or harm to	Significant at local	CEMP, and Reptile Mitigation	No significant effect. No
assemblage	<u>individuals</u> – during vegetation	levelSignificant at Local	Strategy included within EMMF	significant effect.
	<u>clearance</u> . Direct killing, injury	<u>level</u> (major adverse)	(Document Reference 6.2.12.3) –	
	or harm to individuals — during	(permanent).	including establishment of suitable	(potential for significant
	vegetation clearance.		receptor site and subsequent	positive effect in the
			translocation of reptiles from	long term subject to
			construction footprint prior to	final off-site mitigation
			vegetation clearance commencing,	package).
			and precautionary methods of	
			working. Establishment of EPZs to	
			protect retained habitats. CEMP, and	
			Reptile Mitigation Strategy included	
			within EMMF (Document Reference	
			6.2.12.3) – including establishment	
			of suitable receptor site and	
			subsequent translocation of reptiles	
			from construction footprint prior to	
			vegetation clearance commencing,	
			2	
			and precautionary methods of	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			working. Establishment of EPZs to	
			protect retained habitats.	
			Off-site Ecological Mitigation -	
			delivered in accordance with the	
			General Principles for Offsite	
			Ecological Mitigation (Document	
			Reference 6.2.12.10).	
	Direct habitat loss – loss of	Significant at local	Landscape Strategy and Reptile	Significant negative at
	suitable habitat such as semi-	levelSignificant at Local	Mitigation Strategy included within	Local level. No
	improved grassland, reedbed	<u>level</u> (moderate	EMMF (Document Reference	significant effect.
	and tall ruderal, as detailed	adverse) (permanent).	6.2.12.3) – maintenance of site wide	
	above. Direct habitat loss.		open mosaic habitats on	
			Swanscombe Peninsula to maintain	
			a variety of foraging resources,	
			breeding, basking, sheltering and	
			hibernation habitats. Management	
			of grassland areas to encourage	
			development of structurally complex	
			grassland sward with substantial	
			'litter layer' and areas of bare	
			ground. Creation of log/brash piles	
			and hibernaculum to provide	
			sheltering and hibernating habitats,	
			and grass snake breeding sites from	
			piling arisings from grass cutting.	
			Landscape Strategy and Reptile	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			Mitigation Strategy included within	
			EMMF (Document Reference	
			6.2.12.3) – maintenance of site wide	
			open mosaic habitats on	
			Swanscombe Peninsula to maintain	
			a variety of foraging resources,	
			breeding, basking, sheltering and	
			hibernation habitats. Management	
			of grassland areas to encourage	
			development of structurally complex	
			grassland sward with substantial	
			'litter layer' and areas of bare	
			ground. Creation of log/brash piles	
			and hibernaculum to provide	
			sheltering and hibernating habitats,	
			and grass snake breeding sites from	
			piling arisings from grass cutting.	
	Damage to habitats from	Significant at local	CEMP - Prevention of hydrological	No significant effect. No
	changes to hydrological	levelSignificant at Local	impacts through adherence to an	significant effect.
	regime and changes to water	<u>level</u> (moderate	appropriate Surface Water	
	and/or sediment quality – as	adverse) (temporary).	Management Strategy, as outlined	
	described above in relation to		within Chapter 17 – Water	
	impacts on habitats. Damage to		Resources and Flood Risk (Document	
	habitats from changes to		Reference 6.1.17). CEMP	
	hydrological regime and		Prevention of hydrological impacts	
	changes to water and/or		through adherence to an	
	sediment quality – as described		appropriate Surface Water	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance
	above in relation to impacts on habitats.		Management Strategy, as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	
	Damage to supporting habitats from construction dust, waste and pollutants.Damage to supporting habitats from construction dust, waste and pollutants.	Significant at local levelSignificant at Local level (minor adverse) (temporary).	CEMP — including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions. CEMP—including details on safe/appropriate disposal of construction waste, and pollution prevention guidelines, to be strictly adhered to. CEMP will also include a Dust Management Plan (DMP) to provide measures to control dust and other emissions.	No significant effect. No significant effect.
	Damage to supporting habitats from introduction or proliferation of INNS. Damage to supporting habitats from introduction or proliferation of Invasive Non Native Species (INNS).	Significant at local levelSignificant at Local level (minor adverse) (temporary, reversible).	Non-native Invasive Plant Species Mitigation Strategy included within EMMF (Document Reference 6.2.12.3) – including details on control/eradication of existing populations of non-native species.	No significant effect. No significant effect.



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils. Non-native Invasive	
			Plant Species Mitigation Strategy	
			included within EMMF (Document	
			Reference 6.2.12.3) – including	
			details on control/eradication of	
			existing populations of non-native	
			species.	
			CEMP – to include control measures	
			to prevent spread of INNS and INNS	
			impacted soils.	
	Habitat fragmentation and	Significant at local	Landscape Strategy and EMMF	Significant negative at
	loss of dispersal routes. Habitat	levelSignificant at Local	(Document Reference 6.2.12.3) –	Local level. No
	fragmentation and loss of	<u>level</u> (moderate	maintenance of connectivity across	significant effect.
	dispersal routes.	adverse) (permanent).	the Peninsula through the inclusion	
			of a chain of watercourses and	
			wetland areas wrapping around the	
			side of the Proposed Development	
			footprint, and maintenance of scrub	
			and rough grassland habitats across	
			the Peninsula to allow individuals to	
			disperse between habitats. Creation	
			of new wetland and saltmarsh	
			habitats on Swanscombe	



Ecological Feature	Description of Effect	Potential Impact	Mitigation and	Residual Effect
		including Significance	Enhancement Avoidance and	including Significance
			Mitigation	
			Peninsula. Landscape Strategy and	
			EMMF (Document Reference	
			6.2.12.3) – maintenance of	
			connectivity across the Peninsula	
			through the inclusion of a chain of	
			watercourses and wetland areas	
			wrapping around the side of the	
			Proposed Development footprint,	
			and maintenance of scrub and rough	
			grassland habitats across the	
			Peninsula to allow individuals to	
			disperse between habitats. Creation	
			of new wetland and saltmarsh	
			habitats on Swanscombe Peninsula.	
	Disturbance (noise and	Significant at local	CEMP and SLS - Restricted working	No significant effect. No
	lighting) – as described above	levelSignificant at Local	hours and sensitive design and	significant effect.
	in relation to impacts on	<u>level</u> (moderate	location of new lighting to minimise	
	SPA/Ramsar sites. Disturbance	adverse) (temporary,	impacts on nocturnal/crepuscular	
	(noise and lighting) — as	reversible).	wildlife.	
	described above in relation to			
	impacts on SPA/Ramsar sites.		CEMP – adoption of mitigation	
			measures to control construction	
			noise and vibration, as described in	
			Appendix 15.5: Construction Noise	
			and Vibration Mitigation (Document	
			reference 6.2.15.5). CEMP and SLS	
			Restricted working hours and	



Ecological Feature	Description of Effect	Potential Impact including Significance	Mitigation and Enhancement Avoidance and Mitigation	Residual Effect including Significance	
			sensitive design and location of new		
			lighting to minimise impacts on		
			nocturnal/crepuscular wildlife.		
			CEMP – adoption of mitigation		
			measures to control construction		
			noise and vibration, as described in		
			Appendix 15.5: Construction Noise		
			and Vibration Mitigation (Document		
			reference 6.2.15.5).		
Invertebrate	N.B. Impact assessment for the invertebrate assemblage included under Swanscombe Peninsula SSSI. Included within				
assemblage	Swanscombe Peninsula SSSIDirect habitat loss.				
	Significant at regional level (major adverse) (permanent).				
	Landscape Strategy, and Invertebrate Mitigation Strategy included in the EMMF (Document Reference 6.2.12.3) –				
	enhancement of existing retained open mosaic habitat (OMH) to better quality and creation of new brown roofs as				
	described previously. Furthermore, enhancement of retained wetland habitat within Botany Marsh East and Black Duck				
	Marsh through reduction of scrub encroachment, selective ditch re-profiling, and creation of additional scrapes and deep				
	areas in Black Duck Marsh. Creation of additional saltmarsh habitat on the north eastern edge of Swanscombe Peninsula				
	through 'retiring' the flood defence. Creation of a number of habitats of value to invertebrates within the amenity spaces				
	within the resort itself, including native tree and shrub planting, wildflower strips, green roofs and walls and 'bug hotels'.				
	Off site Ecological Mitigation – delivered in accordance with the General Principles for Offsite Ecological Mitigation				
			l valuable invertebrate habitat.		
	No significant effect.	,			





Table 12.17: Summary of Ecological Impact Assessment and Residual Effects During Operational Phase

Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Designations				
Thames Estuary	Potential impacts within the Eur	opean/Ramsar Site itself: P	Potential impacts solely on functionally link	ed resources (outside
and Marshes	of the European/Ramsar Site):			
SPA/Ramsar, and	Recreational disturbance of	Significant at National	Contribution to SAMMS –	No significant
the component	flora and fauna within	level (minor adverse)	proportionate financial contribution to	effect. No significant
SSSIs including	designated area as a result of	(permanent).Significant	strategic access	effect
South Thames	increased residential	at the district level	management.Sustainable Drainage	
Estuary and	population (due to residential	(moderate adverse)	System (SuDS) – designed in	
Marshes SSSI and	accommodation for	(temporary)	accordance with best practice guidance	
Mucking Flats and	staff). Damage to functionally		as outlined within Chapter 17 – Water	
Marshes SSSI92	linked habitats potentially		Resources and Flood Risk (Document	
	used by cited bird species from		Reference 6.1.17).	
	changes to hydrology; and			
	changes to water and /or		Breeding and Winter Bird Mitigation	
	sediment quality (from surface		Strategy enclosed within EMMF	
	or groundwater discharges		(Document Reference 6.2.12.3)	
	from the Project Site including		updated bird surveys of retained and	
	operational waste and		newly created wetlands and	
	pollutants)		functionally linked habitats in years 3, 5	
			and 10 post-completion to ensure	
			habitats continue to support the	
			intended habitats and desired	
			flora/fauna.	
	Potential impacts solely on function	tionally linked resources (o	utside of the European/Ramsar Site):	1

⁹² South Thames Estuary and Marshes SSSI and Mucking Flats and Marshes SSSI are component SSSIs to the Thames Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Damage to functionally linked	Significant at the	Sustainable Drainage System (SuDS) –	No significant effect.
	habitats potentially used by	<u>District level (moderate</u>	designed in accordance with best	
	cited bird species from	adverse) (temporary-	practice guidance as outlined within	
	changes to hydrology; and	permanent).	Chapter 17 – Water Resources and	
	changes to water and /or		Flood Risk (Document Reference	
	sediment quality (from		<u>6.1.17).</u>	
	surface or groundwater			
	discharges from the Project		SPA and Winter Bird Mitigation	
	Site including operational		Strategy enclosed within EMMF	
	waste and pollutants).		(Document Reference 6.2.12.3) -	
			updated bird surveys of retained and	
			newly created wetlands and	
			functionally linked habitats in years 3, 5	
			and 10 post-completion to ensure	
			habitats continue to support the	
			intended habitats and desired	
			flora/fauna.	



Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats. Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats

Significant at the district level District level (minor adverse) (temporary, reversiblepermanent).

BMRS – monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased.

Creation/Enhancement of 259.33ha
Functionally Linked Habitat Off-site —
delivered in accordance with the ECF
(Document Reference 6.2.12.10) to
address any residual disturbance
effects on retained habitats. Bird
Monitoring Response Strategy (BMRS)—
monitoring of bird use on the intertidal
habitats and other functionally linked
land—if the numbers of birds within
the monitored area fall below a certain
threshold in response to obvious
operational shipping then those
disturbance activities will be temporally
ceased.

Off-site Ecological Mitigation –
delivered in accordance with the
'General Principles for Offsite
Ecological Mitigation' (Document



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
			Reference 6.2.12.10), to offset any	
			residual effects.	



Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species using functionally linked habitats. Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area

Significant at the district level District level (minor adverse) (temporary, reversiblepermanent).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.

Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and
- Black Duck, Botany and Broadness
 Marshes Environmental Zone E2
 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.

<u>placement of 'scream zones'; strict noise specification to eliminate</u>

No significant effect. No significant effect.



	noise spread; and controls on location,	
	direction and sound pressure level of	
	loudspeaker systems. (See Chapter 15,	
	Document reference: 6.1.15).	
	_	
	Creation/Enhancement of 259.33ha	
	Functionally Linked Habitat Off-site –	
	delivered in accordance with the ECF	
	(Document Reference 6.2.12.10) to	
	address any residual disturbance	
	effects on retained habitats. Sensitive	
	Lighting Strategy (SLS) - Sensitive	
	design and location of new lighting to	
	minimise impacts on	
	nocturnal/crepuscular wildlife.	
	Retained intertidal and marsh habitats	
	within the Kent Project Site to remain	
	in the following Environmental Lighting	
	Zones:	
	River Thames and Intertidal Zone –	
	Environmental Zone E1 (Typical of	
	relatively uninhabited rural areas. No	
	artificial lighting. Maintain the river	
	in its current condition); and	
	Black Duck, Botany and Broadness	
	Marshes Environmental Zone E2	
	(Typical of sparsely inhabited rural	
	areas. No lighting sources visible	
	from animal habitats. Protect the	



including significance	Mitigation and enhancement	Residual effect including significance
	natural areas that are to be conserved and enhanced).	
	BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.	
	Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate	
	clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15,	
		BMRS—to monitor bird activity and inform the development of further control/mitigation measures as required. Disturbance control measures — final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Disturbance (human	Significant at the	Landscape Strategy, LMP and EMMF	No significant
	movement or activity), to	district level District	(Document Reference 6.2.12.3) –	effect No significant
	cited bird species potentially	<u>level</u> (moderate	Recreational use managed via trail	effect.
	using functionally linked	adverse)	network, signage and interpretation	
	habitats, as a result of	(permanenttemporary,	boards to mitigate against knock-on	
	increased human presence on	reversible).	effects on on-site wildlife. Creation of	
	the Project Site during		new bird hides at key observation	
	operation of the resort rides		locations to reduce visual impact.	
	and attractions and			
	entertainment facilities,		Creation/Enhancement of 259.33ha	
	including increased		Functionally Linked Habitat Off-site –	
	recreational use of retained		delivered in accordance with the ECF	
	habitats. Disturbance (human		(Document Reference 6.2.12.10) to	
	movement or activity), to cited		address any residual disturbance	
	bird species potentially using		effects on retained habitats. Landscape	
	functionally linked habitats, as		Strategy, Landscape Management Plan	
	a result of increased human		(LMP) and EMMF (Document Reference	
	presence on the Project Site		6.2.12.3) – Recreational use managed	
	during operation of the resort		via trail network, signage and	
	rides and attractions and		interpretation boards to mitigate	
	entertainment facilities,		against knock on effects on on site	
	including increased		wildlife. Creation of new bird hides at	
	recreational use of retained		key observation locations to reduce	
	habitats		visual impact.	



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Potential impacts solely on functionally linked resources (outside of the European/Ramsar Site):			



Medway Estuary
and Marshes
SPA/Ramsar/SSSI ⁹³

Damage to functionally linked habitats potentially used by cited bird species from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants). Damage to functionally linked habitats potentially used by cited bird species from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants)

Significant at district levelDistrict level (moderate-minor adverse) (temporarypermanent temporary). <u>SuDS</u> – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).

SPA and Winter Bird Mitigation
Strategy enclosed within EMMF
(Document Reference 6.2.12.3) updated bird surveys of retained and
newly created wetlands and
functionally linked habitats in years 3, 5
and 10 post-completion to ensure
habitats continue to support the
intended habitats and desired
flora/fauna.SuDS — designed in
accordance with best practice guidance
as outlined within Chapter 17 — Water
Resources and Flood Risk (Document
Reference 6.1.17).

Breeding and Winter Bird Mitigation
Strategy enclosed within EMMF
(Document Reference 6.2.12.3) updated bird surveys of retained and
newly created wetlands and
functionally linked habitats in years 3, 5
and 10 post completion to ensure
habitats continue to support the



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
			intended habitats and desired flora/fauna.	

⁹³ Medway Estuary and Marshes SSSI is a component SSSI to the Medway Estuary and Marshes SPA/Ramsar, with overlapping reasons for designation. Potential impacts to the SPA/Ramsar Site are therefore considered to adequately cover the designated interest features of the SSSI also



Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats. Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats

Significant at district level District level (minor adverse) (permanent temporary, reversible).

BMRS – monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased.

Creation/Enhancement of 259.33ha
Functionally Linked Habitat Off-site —
delivered in accordance with the ECF
(Document Reference 6.2.12.10) to
address any residual disturbance
effects on retained habitats. BMRS—
monitoring of bird use on the intertidal
habitats and other functionally linked
land—if the numbers of birds within
the monitored area fall below a certain
threshold in response to obvious
operational shipping then those
disturbance activities will be temporally
ceased.

Off-site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.

No significant effect. No significant effect.



Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area. Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area

Significant at district level District level (minor adverse) (permanent temporary, reversible).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.

Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and
- Black Duck, Botany and Broadness
 Marshes Environmental Zone E2
 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.

<u>ride design to include deliberate</u> <u>placement of 'scream zones'; strict</u> <u>noise specification to eliminate</u> clanking; building placement to reduce No significant effect. No significant effect.



noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15). Creation/Enhancement of 259.33ha **Functionally Linked Habitat Off-site –** delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats.SLS-Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – **Environmental Zone E1 (Typical of** relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and • Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
			BMRS – to monitor bird activity and inform the development of further control/mitigation measures as	
			Disturbance control measures – final ride design to include deliberate	
			placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce	
			noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15)	



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Disturbance (human	Significant at district	Landscape Strategy, LMP and EMMF	No significant
	movement or activity), to	level District level	(Document Reference 6.2.12.3) –	effect No significant
	cited bird species potentially	(moderate adverse)	Recreational use managed via trail	effect.
	using functionally linked	(permanenttemporary,	network, signage and interpretation	
	habitats, as a result of	reversible).	boards to mitigate against knock-on	
	increased human presence on		effects on on-site wildlife. Creation of	
	the Project Site during		new bird hides at key observation	
	operation of the resort rides		locations to reduce visual impact.	
	and attractions and			
	entertainment facilities,		Creation/Enhancement of 259.33ha	
	including increased		Functionally Linked Habitat Off-site –	
	recreational use of retained		delivered in accordance with the ECF	
	habitats. Disturbance (human		(Document Reference 6.2.12.10) to	
	movement or activity), to cited		address any residual disturbance	
	bird species potentially using		effects on retained habitats. Landscape	
	functionally linked habitats, as		Strategy, LMP and EMMF (Document	
	a result of increased human		Reference 6.2.12.3) – Recreational use	
	presence on the Project Site		managed via trail network, signage and	
	during operation of the resort		interpretation boards to mitigate	
	rides and attractions and		against knock-on effects on on-site	
	entertainment facilities,		wildlife. Creation of new bird hides at	
	including increased		key observation locations to reduce	
	recreational use of retained		visual impact.	
	habitats			



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
<u>Swanscombe</u>	Damage to habitats within the	Significant at National	Landscape Strategy and LMP and SSSI	No significant effect.
Peninsula SSSI	designated area from traffic	level (major adverse)	Mitigation Strategy – habitat	
	generated air quality impacts	(permanent).	management and monitoring.	
	resulting in nutrient			
	enrichment and acidification of			
	qualifying woodland habitat			
	with subsequent changes in			
	structure and diversity.			
	Special feature of interest - popu	llations of rare plants		
	Damage to individual	Significant at national	Landscape Strategy, LMP and	No significant
	plantsDamage to individual	levelNational level	Landscape Strategy and LMP, included	effectNo significant
	plants/supporting habitat,	(moderate adverse)	within-Swanscombe Peninsula SSSI	effect.
	and part of plant meta-	(temporary-	Mitigation Strategy (Rare Plants) SSSI	
	population, through increased	permanent temporary,	Mitigation Strategy – Recreational use	
	recreational use, potentially	reversible).	managed via trail network, signage and	
	resulting in trampling, littering,		interpretation boards to mitigate	
	dog fouling, unauthorised		against knock-on effects on on-site	
	camping/campfires/barbecues.		<u>wildlife.</u>	
	Damage to individual	Significant at national	Landscape Strategy, LMP and	No significant
	plantsDamage to individual	levelNational level	Landscape Strategy and LMP, included	effectNo significant
	plants/supporting habitat,	(moderate adverse)	within-Swanscombe Peninsula SSSI	effect.
	and part of plant meta-	(temporary-	Mitigation Strategy (Rare Plants) SSSI	
	population, through	permanent temporary,	Mitigation Strategy – Appropriate	
	inappropriate land	reversible).	management of all retained, enhanced	
	management practices.		or newly created habitats to ensure	
			they continue to support the intended	
			flora and fauna, and provide long term	
			benefits to biodiversity.	



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Damage to retained	Significant at national	<u>SuDS</u> – designed in accordance with	No significant
	plants/population from	levelNational level	best practice guidance as outlined	effectNo significant
	changes to hydrological	(moderate adverse)	within Chapter 17 – Water Resources	effect.
	regime and changes in water	(temporary-	and Flood Risk (Document Reference	
	quality/quantity.	permanenttemporary,	<u>6.1.17).</u>	
		reversible).		
	Special feature of interest - bree	ding bird assemblages:		
	Damage to on-site habitats	Significant at national	<u>SuDS</u> – designed in accordance with	No significant
	potentially used by birds from	levelNational level	best practice guidance as outlined	effectNo significant
	changes to hydrology; and	(moderate adverse)	within Chapter 17 – Water Resources	effect.
	changes to water and /or	(temporary-	and Flood Risk (Document Reference	
	sediment quality (from	permanent temporary,	<u>6.1.17).</u>	
	surface or groundwater	reversible).		
	discharges from the Project		Swanscombe Peninsula SSSI Mitigation	
	Site including operational		Strategy (Breeding Birds) Breeding and	
	waste and pollutants).		Winter Bird Mitigation Strategy	
			enclosed within SSSI Mitigation	
			Strategy – updated bird surveys of	
			retained and newly created habitats in	
			years 3, 5 and 10 post-completion to	
			ensure habitats continue to support	
			the intended habitats and desired	
			flora/fauna.	



	cant negative
<u>lighting)</u> from the operation of <u>levelNational level</u> <u>new lighting to minimise impacts on</u> <u>at National level</u>	ional level. No
the resort rides and attractions (moderate adverse) nocturnal/crepuscular wildlife. significations	cant effect
and entertainment facilities, (permanenttemporary, Retained scrub habitats within the Kent	
giving rise to displacement, reversible). Project Site to remain in the following	
<u>behavioural changes or</u> <u>Environmental Lighting Zones:</u>	
physiological stress to bird	
<u>species using the on-site</u> • Environmental Zone E2 (Typical of	
<u>habitats.</u> <u>sparsely inhabited rural areas. No</u>	
lighting sources visible from animal	
habitats. Protect the natural areas	
that are to be conserved and	
<u>enhanced).</u>	
BMRS – to monitor bird activity and	
inform the development of further	
control/mitigation measures as	
required.	
Disturbance control measures – final	
ride design to include deliberate	
placement of 'scream zones'; strict	
noise specification to eliminate	
clanking; building placement to reduce	
noise spread; and controls on location,	
direction and sound pressure level of	



Document reference: 6.1.15)

loudspeaker systems. (See Chapter 15,

Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
_		including significance		including significance
	Disturbance (human	Significant at national	Landscape Strategy, LMP and	Significant negative
	movement or activity), to bird	levelNational level	Landscape Strategy and LMP, included	at National level. No
	species, as a result of	(moderate adverse)	within-Swanscombe Peninsula SSSI	significant effect
	increased human presence on	(permanent temporary,	Mitigation Strategy (Breeding Birds)	
	the Project Site during	reversible).	SSSI Mitigation Strategy – Recreational	
	operation of the resort rides		use managed via trail network, signage	
	and attractions and		and interpretation boards to mitigate	
	entertainment facilities,		against knock-on effects on on-site	
	including increased		wildlife. Creation of new bird hides at	
	recreational use of retained		key observation locations to reduce	
	<u>habitats.</u>		visual impact.	
	Special feature of interest - inver	<u>rtebrate assemblages:</u>		
	Damage to on-site habitats	Significant at national	SuDS – designed in accordance with	No significant
	potentially used by	levelNational level	best practice guidance as outlined	effect No significant
	invertebrates from changes to	(moderate adverse)	within Chapter 17 – Water Resources	effect.
	hydrology; and changes to	(temporary-	and Flood Risk (Document Reference	
	water and /or sediment	permanent temporary,	<u>6.1.17).</u>	
	quality (from surface or	reversible).		
	groundwater discharges from			
	the Project Site including			
	operational waste and			
	pollutants).			

⁹⁴ Operational noise impacts are limited to the operational hours of the resort. Negligible impacts are expected between 23:00-0700. See Chapter 15 – Noise and Vibration (Document reference 6.1.15)



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Disturbance (lighting) from	Significant at national	SLS - Sensitive design and location of	No significant
	the operation of the resort	levelNational level	new lighting to minimise impacts on	effectNo significant
	rides and attractions and	(minor adverse)	wildlife. Retained intertidal and marsh	effect.
	entertainment facilities.	(permanenttemporary,	habitats within the Kent Project Site to	
		reversible).	remain in the following Environmental	
			<u>Lighting Zones:</u>	
			 River Thames and Intertidal Zone – 	
			Environmental Zone E1 (Typical of	
			relatively uninhabited rural areas. No	
			artificial lighting. Maintain the river	
			in its current condition); and	
			 Black Duck, Botany and Broadness 	
			Marshes - Environmental Zone E2	
			(Typical of sparsely inhabited rural	
			areas. No lighting sources visible	
			from animal habitats. Protect the	
			natural areas that are to be	
			conserved and enhanced).	



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Darenth Woods	Potential damage to habitats	Significant at local	Mitigation measures to reduce traffic	No significant
SSSI	within the designated area	levelSignificant at Local	emissions will need to be developed	effect No significant
	from traffic generated air	<u>level</u> (minor adverse)	and agreed with Natural England,	effect, subject to
	quality impacts. Damage to	(temporary-	following completion of updated AQ	adoption of
	habitats within the designated	permanenttemporary,	assessment. If significant effects are	mitigation as
	area from traffic generated air	reversible) (on a	identified following further	required.
	quality impacts	precautionary basis,	investigation, mitigation measures to	
		subject to further	reduce traffic emissions will need to be	
		investigation)	developed and agreed with Natural	
			England.	
	Potential impacts solely on func	tionally linked resources (o	utside of the SSSI):	



Inner Thames	Damage to functionally linked	Significant at district	SuDS – designed in accordance with	No significant
Marshes SSSI	habitats potentially used by	level District level,	best practice guidance as outlined	effectNo significant
	bird species from changes to	(moderate adverse)	within Chapter 17 – Water Resources	effect.
	hydrology; and changes to	(temporary-	and Flood Risk (Document Reference	
	water and /or sediment	permanenttemporary).	<u>6.1.17).</u>	
	quality (from surface or			
	groundwater discharges from		SPA and Winter Bird Mitigation	
	the Project Site including		Strategy enclosed within EMMF	
	operational waste and		(Document Reference 6.2.12.3) -	
	pollutants). Damage to		updated bird surveys of retained and	
	functionally linked habitats		newly created wetlands and	
	potentially used by bird		functionally linked habitats in years 3, 5	
	species from changes to		and 10 post-completion to ensure	
	hydrology; and changes to		habitats continue to support the	
	water and /or sediment quality		intended habitats and desired	
	(from surface or groundwater		flora/fauna.SuDS – designed in	
	discharges from the Project		accordance with best practice guidance	
	Site including operational		as outlined within Chapter 17 – Water	
	waste and pollutants)		Resources and Flood Risk (Document	
			Reference 6.1.17).	
			Breeding and Winter Bird Mitigation	
			Strategy enclosed within EMMF	
			(Document Reference 6.2.12.3)	
			updated bird surveys of retained and	
			newly created wetlands and	
			functionally linked habitats in years 3, 5	
			and 10 post-completion to ensure	
			habitats continue to support the	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
		merading significance	intended habitats and desired	melading significance
			flora/fauna.	



Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats. Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats

Significant at district level, (minor adverse) (permanent temporary, reversible).

BMRS – monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased.

Creation/Enhancement of 259.33ha
Functionally Linked Habitat Off-site —
delivered in accordance with the ECF
(Document Reference 6.2.12.10) to
address any residual disturbance
effects on retained habitatsBMRS—
monitoring of bird use on the intertidal
habitats and other functionally linked
land—if the numbers of birds within
the monitored area fall below a certain
threshold in response to obvious
operational shipping then those
disturbance activities will be temporally
ceased.

Off-site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.



Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area. Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area

Significant at district level, (minor adverse) (permanent temporary, reversible).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.

Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and
- Black Duck, Botany and Broadness
 Marshes Environmental Zone E2
 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.

<u>placement of 'scream zones'; strict noise specification to eliminate</u> clanking; building placement to reduce



noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15) Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats.SLS Sensitive design and location of new lighting to minimise impacts nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and • Black Duck, Botany and Broadness Marshes Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
			Disturbance control measures – final	
			ride design to include deliberate	
			placement of 'scream zones'; strict	
			noise specification to eliminate	
			clanking; building placement to reduce	
			noise spread; and controls on location,	
			direction and sound pressure level of	
			loudspeaker systems. (See Chapter 15,	
			Document reference: 6.1.15)	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance		
	Disturbance (human movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during operation of the resort rides and attractions, and entertainment facilities. Disturbance (human movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during operation of the resort rides and attractions, and entertainment facilities	Significant at district level District level, (moderate adverse) (permanent temporary, reversible).	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) — Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation locations to reduce visual impact. Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats. Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) — Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation locations to reduce visual impact.	No significant effect No significant effect.		
	Potential impacts solely on functionally linked resources (outside of the SSSI):					



West Thurrock	Damage to functionally linked	Significant at district	SuDS – designed in accordance with	No significant
Lagoon and	habitats potentially used by	level District level	best practice guidance as outlined	effect No significant
Marshes SSSI	cited bird species from	(moderate adverse)	within Chapter 17 – Water Resources	effect.
	changes to hydrology; and	(temporary-	and Flood Risk (Document Reference	
	changes to water and /or	permanenttemporary).	<u>6.1.17).</u>	
	sediment quality (from			
	surface or groundwater		SPA and Winter Bird Mitigation	
	discharges from the Project		Strategy enclosed within EMMF	
	Site including operational		(Document Reference 6.2.12.3) -	
	waste and pollutants). Damage		updated bird surveys of retained and	
	to functionally linked habitats		newly created wetlands and	
	potentially used by cited bird		functionally linked habitats in years 3, 5	
	species from changes to		and 10 post-completion to ensure	
	hydrology; and changes to		habitats continue to support the	
	water and /or sediment quality		intended habitats and desired	
	(from surface or groundwater		flora/fauna.SuDS – designed in	
	discharges from the Project		accordance with best practice guidance	
	Site including operational		as outlined within Chapter 17 – Water	
	waste and pollutants)		Resources and Flood Risk (Document	
			Reference 6.1.17).	
			Breeding and Winter Bird Mitigation	
			Strategy enclosed within EMMF	
			(Document Reference 6.2.12.3) -	
			updated bird surveys of retained and	
			newly created wetlands and	
			functionally linked habitats in years 3, 5	
			and 10 post completion to ensure	
			habitats continue to support the	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
			intended habitats and desired flora/fauna.	



Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats. Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to cited bird species potentially using functionally linked habitats

Significant at district level District level (minor adverse) (permanent temporary, reversible).

BMRS – monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased.

Creation/Enhancement of 259.33ha
Functionally Linked Habitat Off-site —
delivered in accordance with the ECF
(Document Reference 6.2.12.10) to
address any residual disturbance
effects on retained habitats. BMRS—
monitoring of bird use on the intertidal
habitats and other functionally linked
land—if the numbers of birds within
the monitored area fall below a certain
threshold in response to obvious
operational shipping then those
disturbance activities will be temporally
ceased.

Off-site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual offsets.



Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area. Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area

Significant at district level District level (minor adverse) (permanent temporary, reversible).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife.

Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and
- Black Duck, Botany and Broadness
 Marshes Environmental Zone E2
 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

BMRS – to monitor bird activity and inform the development of further control/mitigation measures as required.

<u>placement of 'scream zones'; strict noise specification to eliminate</u>

No significant effect. No significant effect.



direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15) Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats.SLS – Sensitive	
Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects	
Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects	
Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects	
Functionally Linked Habitat Off-site — delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects	
delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects	
(Document Reference 6.2.12.10) to address any residual disturbance effects	
address any residual disturbance effects	
on retained habitats. SLS - Sensitive	
design and location of new lighting to	
minimise impacts on	
nocturnal/crepuscular wildlife.	
Retained intertidal and marsh habitats	
within the Kent Project Site to remain in	
the following Environmental Lighting	
Zones:	
River Thames and Intertidal Zone —	
Environmental Zone E1 (Typical of	
relatively uninhabited rural areas. No	
artificial lighting. Maintain the river	
in its current condition); and	
Black Duck, Botany and Broadness	
Marshes Environmental Zone E2	
(Typical of sparsely inhabited rural	
areas. No lighting sources visible	
from animal habitats. Protect the	
natural areas that are to be	
conserved and enhanced).	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
			BMRS – to monitor bird activity and	
			inform the development of further	
			control/mitigation measures as	
			required.	
			Disturbance control measures – final	
			ride design to include deliberate	
			placement of 'scream zones'; strict	
			noise specification to eliminate	
			clanking; building placement to reduce	
			noise spread; and controls on location,	
			direction and sound pressure level of	
			loudspeaker systems. (See Chapter 15,	
			Document reference: 6.1.15)	



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Disturbance (human	Significant at district	Landscape Strategy, LMP and EMMF	No significant
	movement or activity), to	level District level	(Document Reference 6.2.12.3) –	effect No significant
	cited bird species potentially	(moderate adverse)	Recreational use managed via trail	effect.
	using functionally linked	(permanenttemporary,	network, signage and interpretation	
	habitats, as a result of	reversible).	boards to mitigate against knock-on	
	increased human presence on		effects on on-site wildlife. Creation of	
	the Project Site during		new bird hides at key observation	
	operation of the resort rides		locations to reduce visual impact.	
	and attractions, and			
	<u>entertainment</u>		Creation/Enhancement of 259.33ha	
	<u>facilities</u> . Disturbance (human		Functionally Linked Habitat Off-site –	
	movement or activity), to cited		delivered in accordance with the ECF	
	bird species potentially using		(Document Reference 6.2.12.10) to	
	functionally linked habitats, as		address any residual disturbance	
	a result of increased human		effects on retained habitats. Landscape	
	presence on the Project Site		Strategy, LMP and EMMF (Document	
	during operation of the resort		Reference 6.2.12.3) – Recreational use	
	rides and attractions, and		managed via trail network, signage and	
	entertainment facilities		interpretation boards to mitigate	
			against knock-on effects on on-site	
			wildlife. Creation of new bird hides at	
			key observation locations to reduce	
			visual impact.	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
Great Crabbles	Damage to habitats within the	Significant at local	Mitigation measures to reduce traffic	No significant
Wood SSSI	designated area from traffic	levelSignificant at Local	emissions will need to be developed	effectNo significant
	generated air quality impacts.	level (minor adverse)	and agreed with Natural England,	effect, subject to
	Increased disposition of N	(permanent temporary,	following completion of updated AQ	adoption of
	likely to result in nutrient	reversible)	assessment.	mitigation as
	enrichment and acidification of		<u> </u>	required.
	qualifying woodland habitat			
	with subsequent changes in			
	structure and diversity with			
	such impacts likely occurring			
	across circa 13.6% of total SSSI			
	area is located within 200m of			
	the A289 whilst the woodland			
	is reported to be in			
	unfavourable recovering			
	condition. Total area of impact			
	is, however, reduced given N			
	sensitive woodland habitat is			
	only present across a			
	proportion of this			
	area. Damage to habitats			
	within the designated area			
	from traffic generated air			
	guality impacts			



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Botany Marshes	Damage to habitats from	Significant at local	SuDS – designed in accordance with	No significant
LWS	changes to hydrological	levelSignificant at Local	best practice guidance as outlined	effect No significant
	regime, and changes in water	<u>level</u> (moderate	within Chapter 17 – Water Resources	effect.
	quality/quantity.Damage to	adverse) (<u>temporary-</u>	and Flood Risk (Document Reference	
	habitats from changes to	permanenttemporary).	6.1.17). SuDS – designed in accordance	
	hydrological regime, and		with best practice guidance as outlined	
	changes in water		within Chapter 17 – Water Resources	
	quality/quantity		and Flood Risk (Document Reference	
			6.1.17).	



Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, potentially giving rise to displacement, behavioural changes or physiological stress to wildlife species within the designated area. Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities potentially giving rise to displacement, behavioural changes or physiological stress to wildlife species within the designated area

Significant at local levelSignificant at Local level (moderate adverse) (permanenttemporary). SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Botany Marshes LWS to remain in the following Environmental Lighting Zone:

 Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

Disturbance control measures — final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15). SLS—Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Botany Marshes LWS to remain in the following Environmental Lighting Zone:

 Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
			that are to be conserved and	
			enhanced).	
			Disturbance control measures – final	
			ride design to include deliberate	
			placement of 'scream zones'; strict	
			noise specification to eliminate	
			clanking; building placement to reduce	
			noise spread; and controls on location,	
			direction and sound pressure level of	
			loudspeaker systems. (See Chapter 15,	
			Document reference: 6.1.15)	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
	Recreational disturbance of	Significant at local	Landscape Strategy, LMP and EMMF	No significant
	flora and fauna within	levelSignificant at Local	(Document Reference 6.2.12.3) –	effectNo significant
	designated area as a result of	<u>level</u> (minor adverse)	Recreational use managed via trail	effect.
	increased residential	(permanenttemporary,	network, signage and interpretation	
	population (due to residential	reversible).	boards to mitigate against knock-on	
	accommodation for staff) and		effects on on-site wildlife. Creation of	
	potential increased		new bird hides at key observation	
	recreational use of		locations to reduce visual impact.	
	Swanscombe Peninsula from		Landscape Strategy, LMP and EMMF	
	members of the general		(Document Reference 6.2.12.3)	
	<u>public</u> .Recreational			
	disturbance of flora and fauna		Recreational use managed via trail	
	within designated area as a		network, signage and interpretation	
	result of increased residential		boards to mitigate against knock on	
	population (due to residential		effects on on-site wildlife. Creation of	
	accommodation for staff) and		new bird hides at key observation	
	potential increased		locations to reduce visual impact.	
	recreational use of			
	Swanscombe Peninsula from			
	members of the general public			



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Ebbsfleet	Potential damage to habitats	Significant at local	If significant effects are identified	No significant
Marshes,	within the designated area	levelSignificant at Local	following further investigation,	effect No significant
Northfleet LWS	from traffic generated air	<u>level</u> (minor adverse)	mitigation measures to reduce traffic	effect, subject to
	quality impacts – subject to	(permanenttemporary,	emissions will need to be developed	adoption of
	further investigation to	reversible) (on a	and agreed with Natural England.	mitigation as
	determine the nature of any	precautionary basis,		required.
	significant effects. Damage to	subject to further		
	habitats within the designated	investigation) <u>.</u>		
	area from traffic generated air			
	quality impacts			
	Damage to habitats from	Significant at local	SuDS – designed in accordance with	No significant effect.
	changes to hydrological	levelSignificant at Local	best practice guidance as outlined	
	regime and changes in water	<u>level</u> (moderate	within Chapter 17 – Water Resources	
	quality/quantity <u>.</u>	adverse) (<u>temporary-</u>	and Flood Risk (Document Reference	
		permanenttemporary).	6.1.17).	



Disturbance (noise and
lighting) from the operation of
the resort rides and attractions
and entertainment facilities,
potentially giving rise to
displacement, behavioural
changes or physiological stress
to wildlife species within the
designated area <u>.</u>

Significant at local levelSignificant at Local level (moderate adverse) (permanenttemporary). **SLS** - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Where possible, parts of Ebbsfleet Marshes, Northfleet LWS to remain in the following Environmental Lighting Zone:

 Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

With remaining areas (which are already subject to high levels of lighting) within:

 Environmental Zone E3 (Typical of well inhabited rural and urban settlements). Necessary lighting to create a safe and comfortable environment without over illuminating and reducing light spill onto adjacent areas of the E2 zone.

Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
			loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).	
	Recreational disturbance of flora and fauna within designated area as a result of increased residential population (due to residential accommodation for staff).	Significant at local levelSignificant at Local level (minor adverse) (permanenttemporary, reversible).	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) — Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation locations to reduce visual impact.	No significant effect.
Disused Hospital Grounds, Mabledon LWS	Potential damage to habitats within the designated area from traffic generated air quality impacts – subject to further investigation to determine the nature of any significant effects. Damage to habitats within the designated area from traffic generated air quality impacts	Significant at local levelSignificant at Local level (minor adverse) (permanenttemporary, reversible). (on a precautionary basis, subject to further investigation)	Mitigation measures to reduce traffic emissions will need to be developed and agreed with Natural England, following completion of updated AQ assessment. If significant effects are identified following further investigation, mitigation measures to reduce traffic emissions will need to be developed and agreed with Natural England.	No significant effect No significant effect, subject to adoption of mitigation as required.



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
Tilbury Marshes LWS	Damage to habitats from changes to hydrological regime and changes in water quality/quantity.Damage to habitats from changes to hydrological regime and changes in water quality/quantity	Significant at local levelSignificant at Local level (moderate adverse) (temporary-permanenttemporary).	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect.



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
	Disturbance (from increased ferry movements) associated with the new Thames Clipper river bus service and operational deliveries, potentially giving rise to displacement, behavioural changes or physiological stress to bird species potentially using functionally linked habitats.	Significant at local level (minor adverse) (permanent temporary, reversible).	BMRS - monitoring of bird use on the intertidal habitats and other functionally linked land – if the numbers of birds within the monitored area fall below a certain threshold in response to obvious operational shipping then those disturbance activities will be temporally ceased. Creation/Enhancement of 259.33ha Functionally Linked Habitat Off-site – delivered in accordance with the ECF (Document Reference 6.2.12.10) to address any residual disturbance effects on retained habitats Off-site Ecological Mitigation – delivered in accordance with the 'General Principles for Offsite Ecological Mitigation' (Document Reference 6.2.12.10), to offset any residual effects.	No significant effect.



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
	Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to cited bird species within the designated area.	Significant at local level Significant at Local level (minor adverse) (permanenttemporary, reversible).	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Entire stretch of the River Thames maintained in the following Environmental Lighting Zone: Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition). Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15) 	No significant effect.



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
	Disturbance (human	Significant at local	Landscape Strategy, LMP and EMMF	No significant effect.
	movement or activity), to	levelSignificant at Local	(Document Reference 6.2.12.3) –	
	cited bird species potentially	<u>level</u> (moderate	Recreational use managed via trail	
	using functionally linked	adverse)	network, signage and interpretation	
	habitats, as a result of	(<u>permanent</u> temporary,	boards to mitigate against knock-on	
	increased human presence on	reversible).	effects on on-site wildlife. Creation of	
	the Project Site during		new bird hides at key observation	
	operation of the resort rides		locations to reduce visual impact.	
	and attractions, and			
	entertainment facilities.			
Habitats/Flora				
Populations of	N.B. Mitigation of potential	Significant at regional	Landscape Strategy, LMP and EMMF	No significant effect
nationally scarce	effects upon scarce plants	level (moderate	(Document Reference 6.2.12.3) -	
plants	included under Swanscombe	adverse) (temporary,	Recreational use managed via trail	
	Peninsula SSSI. Damage to	reversible)	network, signage and interpretation	
	individual plants, and part of		boards to mitigate against knock-on	
	plant meta-population,		effects on on site wildlife.	
	through increased			
	recreational use, potentially			
	resulting in trampling, littering,			
	dog fouling, unauthorised			
	camping/campfires/barbecues			



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
Retained habitats ⁹⁵ (including woodland, scrub, grassland, open mosaic on	Damage to retained habitats, through increased recreational use, potentially resulting in trampling, littering, dog fouling, unauthorised	Significant at local levelLocal-County level (moderate adverse) (Permanent, irreversibletemporary,	Landscape Strategy, LMP and EMMF (Document Reference 6.2.12.3) — Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on	Significant effect at Local-District level. No significant effect
previously developed land, wetlands,	Damage to retained habitats through inappropriate land	Significant at local levelSignificant at Local	effects on on-site wildlife. LMP and EMMF (Document Reference 6.2.12.3) – Appropriate management of	No significant effect.
waterbodies and the River Ebbsfleet). See Appendix 12.8 for no deterioration assessment for	management practices.	level (moderate adverse) (temporary-permanenttemporary, reversible).	all retained, enhanced or newly created habitats to ensure they continue to support the intended flora and fauna, and provide long term benefits to biodiversity.	
the River Ebbsfleet	Damage to retained habitats from changes to hydrological regime and changes in water quality/quantity.	Significant at local levelSignificant at Local level (moderate adverse) (temporary- permanenttemporary)	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect.

⁹⁵ Predicted impacts on retained habitats during the operation of the Proposed Development are considered applicable across the full suite of habitats retained, and therefore are addressed collectively



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect		
		including significance		including significance		
Ancient woodland	Damage to habitats within the	Significant at the Local	Mitigation measures to reduce traffic	No significant effect,		
	designated area from traffic	level (moderate	emissions to be developed and agreed	subject to adoption		
	generated air quality impacts.	adverse) (permanent).	with Natural England.	of mitigation as		
	Increased disposition of N			required.		
	likely to result in nutrient					
	enrichment and acidification of					
	woodland habitat with					
	subsequent changes in					
	structure and diversity.					
Species						
Wintering	N.B. Assessment of potential habitat loss and disturbance effects upon wintering waterfowl and wading bird assemblage					
Waterfowl and	included under SPA/Ramsar Sites. Damage to functionally linked habitats potentially used by birds from changes to					
Wading Bird	hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site					
Assemblage	including operational waste and pollutants)					
	Significant at district level (minor adverse) (temporary, reversible)					
	Sustainable Drainage System (SuDS) – designed in accordance with best practice guidance as outlined within Chapter 17 –					
	Water Resources and Flood Risk (Document Reference 6.1.17).					
	Breeding and Winter Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3) - updated bird					
	surveys of retained and newly created wetlands and functionally linked habitats in years 3, 5 and 10 post-completion to					
	ensure habitats continue to supp			•		
	No significant effect		•			



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
Wintering terrestrial bird assemblage	Damage to on-site habitats potentially used by birds from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater discharges from the Project Site including operational waste and pollutants).	Significant at district level District level (minor adverse) (temporary-permanent temporary, reversible).	Sustainable Drainage System (SuDS) – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17). SPA and Winter Bird Mitigation StrategyBreeding and Winter Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3) - updated bird surveys of retained and newly created wetlands and functionally linked habitats in years 3, 5 and 10 post-completion to ensure habitats continue to support the intended habitats and desired flora/fauna.	No significant effect.



Disturbance (noise and	Significant at district	SLS – Sensitive design and location of	No sSignificant effect
lighting) from the operation of	level District level	new lighting to minimise impacts on	at Local level.
the resort rides and attractions	(minor adverse)	nocturnal/crepuscular wildlife.	
and entertainment facilities,	(permanenttemporary,	Retained intertidal and marsh habitats	
giving rise to displacement,	reversible).	within the Kent Project Site to remain	
behavioural changes or		in the following Environmental Lighting	
physiological stress to bird		Zones:	
species using the on-site		 River Thames and Intertidal Zone – 	
habitats.		Environmental Zone E1 (Typical of	
ilabitats <u>.</u>		relatively uninhabited rural areas. No	
		artificial lighting. Maintain the river	
		in its current condition); and	
		 Black Duck, Botany and Broadness 	
		Marshes - Environmental Zone E2	
		(Typical of sparsely inhabited rural	
		areas. No lighting sources visible	
		from animal habitats. Protect the	
		natural areas that are to be	
		conserved and enhanced).	
		BMRS – to monitor bird activity and	
		inform the development of further	
		control/mitigation measures as	
		required.	
		-4-	
		Disturbance control measures – final	
		ride design to include deliberate	
		placement of 'scream zones'; strict	
		noise specification to eliminate	
		noise specification to eliminate	



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
	Disturbance (human	Significant at district	clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15) Landscape Strategy, LMP and EMMF	Significant effect at
	movement or activity), to cited bird species potentially using functionally linked habitats, as a result of increased human presence on the Project Site during operation of the resort rides and attractions and entertainment facilities, including increased recreational use of retained habitats.	levelDistrict level (moderate adverse) (permanenttemporary, reversible).	(Document Reference 6.2.12.3) – Recreational use managed via trail network, signage and interpretation boards to mitigate against knock-on effects on on-site wildlife. Creation of new bird hides at key observation locations to reduce visual impact.	Local level. No significant effect



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect	
		including significance		including significance	
Breeding Bird	N.B. Impact assessment for the	breeding bird assemblage	included under Swanscombe Peninsula SSS	I. Included within	
Assemblage	Swanscombe Peninsula SSSI				
	Damage to on-site habitats pot	entially used by birds <mark>from</mark>	changes to hydrology; and changes to wa	ter and /or sediment	
	quality (from surface or ground	water discharges from the	Project Site including operational waste a	n d pollutants)	
	Significant at district level (mine	r adverse) (temporary, rev	versible)		
	SuDS – designed in accordance	with best practice guidance	e as outlined within Chapter 17 – Water Re	sources and Flood Risk	
	(Document Reference 6.1.17).				
	Breeding and Winter Bird Mitigation Strategy enclosed within EMMF (Document Reference 6.2.12.3) updated bird				
	surveys of retained and newly c	reated habitats in years 3,	5 and 10 post completion to ensure habita	ts continue to support	
	the intended habitats and desire	e d flora/fauna.			
	No significant effect	1	,		
Bat Assemblage	Damage to on-site habitats	Significant at district	SuDS – designed in accordance with	No significant effect.	
	potentially used by bats from	level District level	best practice guidance as outlined		
	changes to hydrology; and	(minor adverse)	within Chapter 17 – Water Resources		
	changes to water and /or	(temporary-	and Flood Risk (Document Reference		
	sediment quality (from	permanent temporary,	6.1.17).		
	surface or groundwater	reversible).			
	discharges from the Project				
	Site including operational				
	waste and pollutants).				



Ecological Feature Des	scription of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
light the and giving beht physisped hab noise com	sturbance (noise and hting) from the operation of e resort rides and attractions d entertainment facilities, ring rise to displacement, havioural changes or ysiological stress to bat ecies using the on-site bitats, including light and ise pollution of foraging, mmuting and roosting bitats.	Significant at district level District level (minor adverse) (permanent temporary, reversible).	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). 	No significant effect.



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Dormouse	Damage to on-site habitats	Significant at local	SuDS – designed in accordance with	No significant effect.
	potentially used by dormice	levelSignificant at Local	best practice guidance as outlined	
	from changes to hydrology;	<u>level</u> (minor adverse)	within Chapter 17 – Water Resources	
	and changes to water and /or	(temporary-	and Flood Risk (Document Reference	
	sediment quality (from	permanenttemporary,	6.1.17).	
	surface or groundwater	reversible).		
	discharges from the Project			
	Site including operational			
	waste and pollutants).			



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
	Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to dormice using the on-site habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.	Significant at local levelSignificant at Local level (minor adverse) (permanenttemporary, reversible).	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained scrub habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). 	No significant effect.
			Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).	



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Otter	Damage to on-site habitats	Significant at local	SuDS – designed in accordance with	No significant effect.
	potentially used by otters from	levelSignificant at Local	best practice guidance as outlined	
	changes to hydrology; and	<u>level</u> (minor adverse)	within Chapter 17 – Water Resources	
	changes to water and /or	(temporary-	and Flood Risk (Document Reference	
	sediment quality (from	permanent temporary,	6.1.17).	
	surface or groundwater	reversible).		
	discharges from the Project			
	Site including operational			
	waste and pollutants).			



Significant at local levelSignificant at Local level (minor adverse) (permanenttemporary, reversible).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of
 relatively uninhabited rural areas. No
 artificial lighting. Maintain the river
 in its current condition); and
- Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems.



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Water Vole	Damage to on-site habitats	Significant at district	SuDS – designed in accordance with	No significant effect.
	potentially used by otters from	level District level	best practice guidance as outlined	
	changes to hydrology; and	(minor adverse)	within Chapter 17 – Water Resources	
	changes to water and /or	(temporary- permanenttemporary, reversible).	and Flood Risk (Document Reference 6.1.17).	
	sediment quality (from			
	surface or groundwater			
	discharges from the Project			
	Site including operational			
	waste and pollutants).			



Disturbance (noise and
lighting) from the operation of
the resort rides and attractions
and entertainment facilities,
giving rise to displacement,
behavioural changes or
physiological stress to water
voles using the on-site
habitats, including light and
noise pollution of foraging,
dispersal and potential
breeding/hibernating habitats.

Significant at district level District level (minor adverse) (permanent temporary, reversible).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of
 relatively uninhabited rural areas. No
 artificial lighting. Maintain the river
 in its current condition); and
- Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Harvest Mouse	Damage to on-site habitats	Significant at local	SuDS – designed in accordance with	No significant effect.
	potentially used by harvest	levelSignificant at Local	best practice guidance as outlined	
	mice from changes to	<u>level</u> (minor adverse)	within Chapter 17 – Water Resources	
	hydrology; and changes to	(temporary-	and Flood Risk (Document Reference	
	water and /or sediment	permanent temporary,	6.1.17).	
	quality (from surface or	reversible).		
	groundwater discharges from			
	the Project Site including			
	operational waste and			
	pollutants) <u>.</u>			



Disturbance (noise and lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to harvest mice using the on-site habitats, including light and noise pollution of foraging, dispersal and potential breeding/hibernating habitats.

Significant at local levelSignificant at Local level (minor adverse) (permanenttemporary, reversible).

SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones:

- River Thames and Intertidal Zone –
 Environmental Zone E1 (Typical of
 relatively uninhabited rural areas. No
 artificial lighting. Maintain the river
 in its current condition); and
- Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced).

Disturbance control measures – final ride design to include deliberate placement of 'scream zones'; strict noise specification to eliminate clanking; building placement to reduce noise spread; and controls on location, direction and sound pressure level of loudspeaker systems. (See Chapter 15, Document reference: 6.1.15).



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Amphibian	Damage to on-site habitats	Significant at district	SuDS – designed in accordance with	No significant effect.
assemblage	potentially used by amphibians	level District level	best practice guidance as outlined	
	from changes to hydrology;	(minor adverse)	within Chapter 17 – Water Resources	
	and changes to water and /or	(temporary-	and Flood Risk (Document Reference	
	sediment quality (from	permanent temporary,	6.1.17).	
	surface or groundwater	reversible).		
	discharges from the Project			
	Site including operational			
	waste and pollutants) <u>.</u>			



Ecological Feature	Description of effect	Potential impact including significance	Mitigation and enhancement	Residual effect including significance
	Disturbance (lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to amphibians using the on-site habitats, including light pollution of breeding, foraging and hibernation/shelter habitats	Significant at district levelDistrict level (minor adverse) (permanenttemporary, reversible).	 SLS - Sensitive design and location of new lighting to minimise impacts on nocturnal/crepuscular wildlife. Retained intertidal and marsh habitats within the Kent Project Site to remain in the following Environmental Lighting Zones: River Thames and Intertidal Zone – Environmental Zone E1 (Typical of relatively uninhabited rural areas. No artificial lighting. Maintain the river in its current condition); and Black Duck, Botany and Broadness Marshes - Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). 	No significant effect.



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect
		including significance		including significance
Reptile assemblage	Damage to on-site habitats potentially used by reptiles from changes to hydrology; and changes to water and /or sediment quality (from surface or groundwater	Significant at district level District level (minor adverse) (temporary-permanent temporary, reversible).	SuDS – designed in accordance with best practice guidance as outlined within Chapter 17 – Water Resources and Flood Risk (Document Reference 6.1.17).	No significant effect.
	discharges from the Project Site including operational waste and pollutants)_			
	Disturbance (lighting) from the operation of the resort rides and attractions and entertainment facilities, giving rise to displacement, behavioural changes or physiological stress to reptiles using the on-site habitats, including light pollution of potential shelter and hibernation habitats.	Significant at district level District level (minor adverse) (permanent temporary, reversible).	 SLS - Sensitive design and location of new lighting to minimise impacts on wildlife. Retained grassland/scrub habitats within the Kent Project Site to remain in the following Environmental Lighting Zone: Environmental Zone E2 (Typical of sparsely inhabited rural areas. No lighting sources visible from animal habitats. Protect the natural areas that are to be conserved and enhanced). 	No significant effect <u>.</u>



Ecological Feature	Description of effect	Potential impact	Mitigation and enhancement	Residual effect					
		including significance		including significance					
Invertebrate	N.B. Impact assessment for the i	nvertebrate assemblage i	ncluded under Swanscombe Penir	nsula SSSI. Included within					
assemblage	Swanscombe Peninsula SSSI								
	Damage to on-site habitats potentially used by invertebrates from changes to hydrology; and changes to water and /or								
	sediment quality (from surface	or groundwater discharge	s from the Project Site including o	perational waste and					
	pollutants)								
	Significant at district level (mino	r adverse) (temporary, re v	rersible)						
	SuDS - designed in accordance \	with best practice guidanc	e as outlined within Chapter 17 —	Water Resources and Flood Risk					
	(Document Reference 6.1.17).								
	No significant effect								



COMPENSATION

- 12.288 As identified in the Tables above, the updated assessment has identified significant negative residual effects on a number of IEFs resulting from the construction and operational phases of the Proposed Development. In line with mitigation hierarchy, these effects should as a last resort be compensated for.
- 12.289 Table 12.18 summarises the IEFs which are anticipated to experience significant negative residual effects.

Table 12.18: Summary of significant negative residual effects requiring compensation

IEF(s)	Negative Effect	Level of
		<u>significance</u>
Swanscombe	Overall reduction in area and fragmentation	<u>National</u>
<u>Peninsula</u>	(construction)	
<u>SSSI</u>	Direct loss of notified rare plants/supporting habitat	<u>National</u>
	(construction)	
	Direct loss and fragmentation of habitat for notified	<u>National</u>
	breeding birds (construction)	
	Indirect/effective loss of habitat for notified breeding	<u>National</u>
	birds due to disturbance from shipping movements and	
	human activity (construction)	
	Indirect/effective loss of habitat for notified breeding	<u>National</u>
	birds due to disturbance from resort rides and	
	attractions, and increased recreational activity within	
	retained habitat (operation)	
	<u>Direct loss and fragmentation of habitat for notified</u>	<u>National</u>
	invertebrates (construction)	
<u>Alkerden</u>	Loss of part of plant and invertebrate meta-population	<u>Local</u>
Lane Pit LWS	through loss of habitat in the Kent Project Site	
<u>Habitats</u>	<u>Direct loss (construction) and damage through increase</u>	<u>Local -</u>
	recreational use (operation) of the following habitats:	<u>County</u>
	• Scrub;	
	Semi-improved grassland;	
	Coastal/floodplain grazing marsh;	
	 Open mosaic on previously developed land; 	
	 Waterbodies (ponds, standing water and ditch 	
	network); and	
	Swamp (reedbed).	
<u>Species</u>	<u>Direct loss and fragmentation of habitat for the</u>	<u>Local -</u>
	following species (construction):	<u>District</u>
	 Terrestrial wintering birds; 	



Bats (foraging);	
• Dormouse;	
• Otter;	
• Water Vole;	
Harvest Mouse;	
Amphibians; and	
• Reptiles.	
Indirect/effective loss of habitat for terrestrial wintering	<u>District</u>
birds due to disturbance from shipping movements and	
human activity (construction)	
Indirect/effective loss of habitat for terrestrial wintering	Local
birds due to from resort rides and attractions, and	
increased recreational activity within retained habitat	
(operation)	

- 12.290 Proposed compensatory measures in respect of the negative residual effects identified above are set out within the Ecological Compensation Framework (ECF) (Appendix 12.10; Document Reference 6.2.12.10). In essence, direct habitat loss and indirect/effective habitat loss for species due to predicted disturbance levels are to be compensated for by the creation of new habitat, or restoration of existing habitat in poor condition, off-site, which will fulfil the same function as the habitat being lost/disturbed/damaged.
- 12.291 As detailed in the ECF, a precautionary approach has been taken to off-site compensation, where the quantity of new/restored habitat exceeds the quantity to be lost/disturbed/damaged.
- 12.292 There is considerable overlap in the nature of these negative residual effects where they apply to both habitats and the species that the habitats support. Consequently, the same off-site habitat creation/restoration measures can often meet several compensation requirements.
- 12.293 Additional land is still being acquired to fulfil all of the compensation requirements and achieve ecological equivalence with those habitats being loss/disturbed on site, such that further details of specific off-site land parcels will be added to the ECF as these become available.
- <u>12.294</u> Based solely on the off-site land which has been secured to date, the predicted net change in habitat units (as measured using a standard Biodiversity Metric) is +456.10 units (+13.61%).

CUMULATIVE AND IN-COMBINATION EFFECTS

12.3012.295 The assessment of the Proposed Development concludes that, through the adoption of appropriate mitigation measures as detailed in the ES, the project would not



give rise to any significant negative residual effects on ecology. However, this does not exclude the potential for insignificant effects to become significant in combination with other consented/Proposed Developments and associated infrastructure within the surrounding landscape, on sensitive habitats, fauna and flora. A range of sites have been identified in the wider landscape which warrant further consideration in terms of their potential to give rise to in cumulative effects owing to their size or proximity to the Site.

12.3112.296 Table 12.18-19 below provides an assessment of those schemes identified within the 'short list' of cumulative sites set out in the Cumulative Assessment Chapter (ES Chapter 21; Document Reference 6.1.21) which were screened in as potentially capable of having cumulative impacts on ecological receptors. All other potential cumulative sites were screened out either on the basis of lying outside the ZOI of all pertinent receptors, or due to the temporal scope or scale/nature of the development proposed. The assessment has been completed following review of the ecological assessment submitted in support of the respective applications for each project as available from the relevant planning portal.

<u>12.32</u>12.297 In summary, the assessment concludes that there are considered to be no significant cumulative effects in combination with the other projects evaluated.



Table 12.18_19 Cumulative impact assessment on ecological receptors.

ID	Application Reference	Applicant and Brief Description	Distance from Project	Status	IEFs Potentially at Risk of Cumulative Impacts	Nature of Potential Impact	Assessment	Conclusion
1	e.planninginspector ate.gov.uk/projects/ south-east/tilbury2/	ctor Limited. cts/ DCO application for a new port facility acting cts/ DCO application for a new port facility acting cts/ DCO application for a new port facility acting cts/ DCO application for a new port facility acting cts/	c. 4.3km east of Kent site. c. 820 east of Essex site.	Secretary of State for Transport granted development consent for this application on 20/04/19.	Estuarine European Sites and SSSIs.	Temporary and permanent loss/disturbance of functionally linked habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
		Determined by: NSIP/Planning Inspectorate.			Brownfield Invertebrates (local metapopulation).	Permanent of loss of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
2	https://infrastructur e.planninginspector ate.gov.uk/projects/ south-east/thurrock- flexible-generation-	Thurrock Flexible Generation Plant, by Thurrock Plant Ltd. Determined by: NSIP/Planning Inspectorate.	c. 4.0km east of Kent site. c. 400m east of Essex site.	Pre-examination Stage (Part 2 of Preliminary Meeting postponed to February 2021).	Estuarine European Sites and SSSIs.	Temporary and permanent loss/disturbance of functionally linked habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
	plant/				Brownfield Invertebrates (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
3	https://infrastructur e.planninginspector ate.gov.uk/projects/ south-east/lower- thames-crossing/	Lower Thames Crossing, by Highways England. Determined by: NSIP/Planning Inspectorate.	c. 5.4km east of Kent site c. 2.6km east of Essex site.	Application withdrawn. The Applicant has not yet set a timetable for a revised application	Estuarine European Sites and SSSIs.	Temporary and permanent loss/disturbance of functionally linked habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
					Brownfield Invertebrates (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
4		olanninginspector e.gov.uk/projects/ uth-east/tilbury- NSIP/Planning Inspectorate. oo oo c.	c. 4.6km east of Kent site. c. 1.8km east of Essex site.	stage.	Estuarine European Sites and SSSIs.	Temporary and permanent loss/disturbance of functionally linked habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
					Brownfield Invertebrates (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.



ID	Application Reference	Applicant and Brief Description	Distance from Project	Status	IEFs Potentially at Risk of Cumulative Impacts	Nature of Potential Impact	Assessment	Conclusion
9	16/00201/EDCCON https://publicaccess. dartford.gov.uk/onli ne- applications/simpleS earchResults.do?acti on=firstPage Eastern Quarry, Swanscombe by Barton Wilmore. A mixed use mixed-use development of up to 6250 dwellings & in addition up to 231,000 square metres of built floorspace (in total), including: Use classes A1-5, B1, D1, D2, and C1. Development also includes open space provision, highways and public transport facilities, and facilities for mooring, launching and landing watercraft. Determined by: Dartford Borough Council.	c. 1.4km south of Kent site. c. 4.6 km south west of	Permission granted 23/01/13 A number of Reserved Matters and Discharge of	Estuarine European Sites and SSSIs	Recreational disturbance	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.	
		Essex site.	Conditions applications have been submitted since permission was granted. Most	Dormice (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.	
				recently, a discharge of conditions application was granted on 23/06/20.	Brownfield Invertebrates (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
10	https://highwayseng land.co.uk/projects/ a2-bean-and- ebbsfleet-junction- improvements/	A2 Bean and Ebbsfleet Junction Improvements by Highways England. Construction of 5 new slip roads and 1 modified roundabout to connect the A2 Trunk Road to the	c. 2.5km south west of Kent site. c. 6.0 km	On 2 nd June 2020 Highways England received confirmation from the Secretary of	Darenth Wood SSSI.	Temporary and permanent air quality impacts.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
	Ebbsfleet East Rou	B255 Bean North and Bean South Roundabouts and Ebbsfleet East Roundabout. Determined by: Secretary of State for Transport.		State for Transport that the scheme can proceed to construction.	Dormice (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
11	KCC/DA/0232/2019 https://www.kentpl anningapplications.c o.uk/Planning/Displa y/KCC/DA/0232/201 9	Bluewater Shopping Centre by Kent County Council. Full planning application for a new tunnel (the Bean Road Tunnel) and associated road works to include bus, cycling and pedestrian access to the east of Bluewater Shopping Centre to link to the Eastern Quarry development, including tree planting at land adjacent to lake 5 and tunnel infilling. Determined by: Kent County Council.	c. 2.4km south west of Kent site. c. 6.0km south west of Essex site.	Granted with conditions 24/06/2020.	Dormice (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
14	05/00221/OUT https://publicaccess. dartford.gov.uk/onlin e- applications/applicati	Stone Pit II, by Barton Wilmore Planning. Development of up to 870 dwellings nad up to 1,200sq metres of built floorspace for B1(a), (b), and (c), D1 and D2, and A1-A5 uses. Development	c. 3.0km south west of Kent site.	Permission granted 30/10/17. A discharge of conditions	Estuarine European Sites and SSSIs.	Recreational disturbance.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
		includes ancillary infrastructure and works to create ecological and natural reserves and refuge areas. Determined by: Dartford Borough Council.	west of Essex site.	application was submitted and approved on 04/05/20 relating to	Darenth Wood SSSI.	Temporary and permanent air quality impacts.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects	No significant cumulative effect.



ID	Application Reference	Applicant and Brief Description	Distance from Project	Status	IEFs Potentially at Risk of Cumulative Impacts	Nature of Potential Impact	Assessment	Conclusion															
				condition 16 of the original permission (Ecological Mitigation Strategy).			from Proposed Development.																
18	20150155 EDC http://applications.eb bsfleetdc.org.uk/onlin e- applications/applicati		Adjacent to southern boundary of Kent site.	Permission granted February 2016. A number of reserved matters	Darenth Wood SSSI.	Temporary and permanent air quality impacts.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.															
	onDetails.do?keyVal= NK267YHPFZZ00&acti veTab=summary	open space, roads and infrastructure. Determined by: Ebbsfleet Development Corporation	c. 3.0km south west of Essex site.	been submitted since this application was approved. The most recent RM	River Ebbsfleet.	Temporary and permanent hydrological impacts.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.																
		(Consulting with Gravesham Borough Council).		application was granted permission on 17/04/20 (EDC/19/0194). Note: this application is to be delivered in phases.	Dormice (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.															
19	EDC/18/0009 http://applications.eb bsfleetdc.org.uk/onlin e- applications/applicati	Land West of Springhead Road, by Countryside Properties (UK) Ltd. Reserved matters application pursuant to application 20150155 EDC relating to the erection of 172	Adjacent to southern boundary of Kent site.	Permission granted June 2018. A number of discharge of	Estuarine European Sites and SSSIs.	Recreational disturbance.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.															
	onDetails.do?keyVal= P2OU7PUAFG800∾ tiveTab=summary	residential dwellings in Phase 3 of Springhead Quarter. Determined by: Ebbsfleet Development Corporation (Consulting with Gravesham Borough Council).	c. 3.0km south west of Essex site.	west of Essex	west of Essex	west of Essex	west of Essex	west of Essex	west of Essex site.	west of Essex site. but the	west of Essex site.	west of Essex	west of Essex	west of Essex site.	west of Essex site. appl been this appl	west of Essex site.	west of Essex site.	west of Essex	•••	Darenth Wood SSSI.	Temporary and permanent air quality impacts.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
				recent discharge of conditions application was granted permission on 09/08/19. 7 such	River Ebbsfleet.	Temporary and permanent hydrological impacts.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.															
				applications are currently awaiting a decision. The most recent of these applications is a discharge of conditions	Dormice (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.															



ID	Application Reference	Applicant and Brief Description	Distance from Project	Status	IEFs Potentially at Risk of Cumulative Impacts	Nature of Potential Impact	Assessment	Conclusion
				application validated 24/03/20 (EC/20/0046).				
27	19/01058/OUT https://regs.thurrock. gov.uk/online- applications/caseDet ails.do?caseType=App	Outline application for 161 new dwellings and	c. 1.8km north east of Kent site. c. 2.3km north	Application validated 11/07/19. Determination deadline 10/10/19. Awaiting decision.	Estuarine European Sites and SSSIs.	Recreational disturbance.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
	lication&keyVal=PUFL M8QGINZ00		west of Essex site.	Application still receiving consultee comments.	Brownfield Invertebrates (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
31	17/01668/OUT https://regs.thurrock. gov.uk/online- applications/applicati onDetails.do?keyVal= P109FAQGKLX00&act iveTab=summary	matters reserved for subsequent approval, except for means of access, for mixed-use redevelopment involving the demolition of existing buildings and other structures, site preparation works, and the development of up to 2,850 dwelling houses (Use	site. c. 9.3km north west of Essex site.	Permission granted 20/12/19. A number of reserved matters applications have been submitted related to this application. The most recent of these applications were validated in March 2020 and are awaiting determination.	Estuarine SSSIs.	Temporary and permanent loss/disturbance of functionally linked habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.



ID	Application Reference	Applicant and Brief Description	Distance from Project	Status	IEFs Potentially at Risk of Cumulative Impacts	Nature of Potential Impact	Assessment	Conclusion
		rebuilding, repairing, replacing and upgrading of river wall and flood defence wall and associated works of repair and reinstatement of the former Yara Purfleet Terminal jetty and the former Cory's Wharf jetty to facilitate the river wall and flood defence works, the provision of four grade separated railway crossings including a new bridge as part of the re-profiling and realignment of London Road. Determined by: Thurrock Council.						
33	18/01404/OUT https://regs.thurrock. gov.uk/online- applications/caseDet ails.do?caseType=App lication&keyVal=PFPF EBQGHUX00	Outline planning permission with all matters (except for access) reserved for the demolition, phased remediation and redevelopment of 167 hectares of former Coryton Oil Refinery to provide up to 480,000 sq. m of commercial development including a Food	east of Essex site.	Awaiting decision. Most recent consultee comment submitted April 2020.	Estuarine European Sites and SSSIs.	Temporary and permanent loss/disturbance of functionally linked habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
39		Outline application for a mixed development and comprising: up to 532 Homes, up to 46,000 sq. m Employment Floorspace and a Mixed Use Neighbourhood Centre.	c. 600m east of Kent site. c. 1.9km south west of Essex	Approved subject to 106 08/06/18. A number of discharge of	Estuarine European Sites and SSSIs.	Recreational disturbance.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
	onDetails.do?keyVal= KH2EC1HP02600&act iveTab=summary	•	site.	conditions applications have been submitted, the most recent of these was submitted in	Breeding birds (local metapopulations).	Temporary and permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
				June 2020 and is	Brownfield	Temporary and	Planning policy requires	No significant



ID	Application Reference	Applicant and Brief Description	Distance from Project	Status	IEFs Potentially at Risk of Cumulative Impacts	Nature of Potential Impact	Assessment	Conclusion
				awaiting decision.	Invertebrates (local metapopulation).	permanent loss/disturbance of habitat.	scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	cumulative effect.
41	EDC/17/0038 http://applications.eb bsfleetdc.org.uk/onlin e- applications/applicati onDetails.do?keyVal= ON7TV7UA00K00∾ tiveTab=summary	Use Classes A1/A2/A3/B1(a)/C3/D1) and demolition of the boundary wall and rear portion of the WT	c. 2.2km south east of Kent site. c. 1.1 south west of Essex site.	Approved subject to 106 20/03/19. There are a number of linked applications. The most recent of these applications is the submission of the Rosherville Pier Management Plan required by the S106, which is awaiting determination (EDC/20/0063).	Estuarine European Sites and SSSIs.	Recreational disturbance.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.
42	EDC/17/0110 http://applications.eb bsfleetdc.org.uk/onlin e- applications/applicati onDetails.do?keyVal= OVFNEOUAKRW00&a ctiveTab=summary		Adjacent to southern site boundary of Kent site. c. 4.4km west of Essex site.	Approved subject to 106 18/12/18. There are a number of linked applications. The most recent of these application is an application to modify a planning obligation,	Estuarine European Sites and SSSIs. Breeding birds (local metapopulations).	permanent loss/disturbance of habitat.	Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development. Legislation and planning policy require scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect. No significant cumulative effect.
				which is awaiting decision (EDC/20/0090).	Brownfield Invertebrates (local metapopulation).	Temporary and permanent loss/disturbance of habitat.	Planning policy requires scheme to avoid or mitigate negative impacts. No residual effects from Proposed Development.	No significant cumulative effect.



CLIMATE CHANGE

- 12.3312.298 Potential climate change projections have been calculated for the Proposed Development and are detailed within Chapter 20: *Greenhouse gases and climate change* (Document Reference 6.1.20). In summary, the projections estimate an increase in winter and summer mean temperatures, an increase in winter mean precipitation, a decrease in summer mean precipitation, and rising sea levels.
- 42.3412.299 Given that the valued habitats and species within the Project Site are 'largely' widespread and the Project Site is largely not near the edge of their ranges; the projected change in temperatures is not anticipated to result in any significant impacts on the designated site, habitat and species IEFs. However, it can reasonably be expected that some species more typically associated with continental Europe, and warmer climates, may expand their range further north and/or, in the case of migratory species, arrive in the UK earlier.
- <u>12.3512.300</u> The projected changes in precipitation may have impacts on the aquatic habitats within the Kent Project Site, including the reedbed, grazing marsh, waterbodies and watercourse, and consequently the species associated with them, such as otter, water vole, amphibians and invertebrates. It is considered that the generous provision of greenspace within the Proposed Development, which will be managed to promote biodiversity, the large area of varied SuDs/attenuation features proposed throughout the scheme and the large buffer afforded to the watercourse corridor (with additional aquatic features incorporated within it), will provide sufficient resilience to any likely effects of future climate change.
- 12.3612.301 Furthermore, future monitoring of the new and retained habitats within the Project Site recommended to be detailed within the LMP at Appendix 11.8 (Document Reference 6.2.11.8), and as described above in the mitigation section, will allow an opportunity for management prescriptions to be reviewed and amended to reflect any impacts as a result of climate change. This will further safeguard the habitat and species interests at the Project Site over the long term.

SUMMARY AND CONCLUSIONS

- <u>12.3712.302</u> This chapter provides an assessment of the significance and consequences of likely ecological impacts upon identified Important Ecological Features (IEFs) arising from the Proposed Development for the London Resort. It has been prepared to accompany a DCO application for the Project Site.
- <u>12.3812.303</u> A suite of impact avoidance, mitigation and compensation measures have been prepared as part of a holistic ecology and landscape strategy for the Proposed Development, to address any likely significant effects that may arise during construction



and after completion (operation) of the Proposed Development. Additional measures to mitigate and compensate for potentially significant effects have been provided.

12.3912.304 Further baseline information in support of this chapter is included within Appendix 12.1: *Ecology Baseline Report* (Document Reference 6.2.12.1) and is referred to throughout the assessment. The approach taken in this assessment is made with reference to the guidelines published in 2018 by the Chartered Institute of Ecology and Environmental Management (CIEEM)⁹⁶.

<u>12.40</u>12.305 The baseline survey work, and consultation with statutory and non-statutory consultees, has identified the following IEFs pertinent to the Proposed Development:

Designated Sites

- Thames Estuary and Marshes SPA/Ramsar;
- Medway Estuary and Marshes SPA/Ramsar/SSSI;
- North Downs Woodland SAC;
- Darenth Woods SSSI;
- Inner Thames Marshes SSSI;
- South Thames Estuary and Marshes SSSI;
- West Thurrock Lagoon and Marshes SSSI;
- Mucking Flats and Marshes SSSI
- Wouldham to Detling Escarpment SSSI;
- Botany Marshes LWS;
- Ebbsfleet Marshes, Northfleet LWS;
- Alkerden Lane Pit LWS; and
- Tilbury Marshes LWS.

Habitats/Flora

Rare plants;

⁹⁶ CIEEM (2018) Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, Freshwater, Coastal and Marine version 1.1. Chartered Institute of Ecology and Environmental Management, Winchester.



- Broad leaved Semi Natural Woodland;
- Scrub;
- Semi-improved grassland;

Ancient woodland;

- Coastal/Floodplain Grazing Marsh;
- Open mosaic on previously developed land;
- Waterbodies (ponds, standing water and ditches);
- Swamp (reedbed); and
- River Ebbsfleet.

Species

- Wintering waterfowl and wading bird assemblage;
- Wintering terrestrial bird assemblage;
- Breeding bird Assemblage assemblage;
- Pochard breeding population;
- Bat assemblage;
- Dormouse;
- Otter;
- Water vole;
- Harvest mouse;
- Amphibian assemblage;
- Reptile assemblage; and
- Invertebrates.



- 12.306 The impact assessment has identified that certain actions could result in significant negative impacts on these IEFs without mitigation. Inherent avoidance and mitigation measures and the implementation of an *Ecological Mitigation and Management Framework* (Document Reference 6.2.12.3), which will be secured as a requirement of the DCO, including construction methods statements and long-term habitat management principles, are considered to ameliorate to some degree those significant impacts identified.
- 12.307 Despite the proposed avoidance and mitigation measures, the updated assessment has identified significant negative residual effects on a number of IEFs resulting from the construction and operational phases of the Proposed Development. In line with mitigation hierarchy, these effects should as a last resort be compensated for. Proposed compensatory measures in respect of these negative residual effects are set out within the Ecological Compensation Framework (ECF) (Appendix 12.10; Document Reference 6.2.12.10).
- 12.308 In essence, direct habitat loss and indirect/effective habitat loss for species due to predicted disturbance levels are to be compensated for by the creation of new habitat, or restoration of existing habitat in poor condition, off-site which will fulfil the same function as the habitat being lost/disturbed/damaged. As detailed in the ECF, a precautionary approach has been taken to off-site compensation, where the quantity of new/restored habitat exceeds the quantity to be lost/disturbed/damaged.
- 12.309 There is considerable overlap in the nature of these negative residual effects where they apply to both habitats and the species that the habitats support. Consequently, the same off-site habitat creation/restoration measures can often meet several compensation requirements.
- 12.310 Additional land is still being acquired to fulfil all of the compensation requirements and achieve ecological equivalence with those habitats being loss/disturbed on site, such that further details of specific off-site land parcels will be added to the ECF as these become available.
- <u>12.311</u> Based solely on the off-site land which has been secured to date, the predicted net change in habitat units (as measured using a standard Biodiversity Metric) is is +456.10 units (+13.61%).
- 12.41 Much of the mitigation measures proposed rely on the provision of off site mitigation to avoid significant residual effects, details of which have not yet been finalised. A set of General Principles for Offsite Ecological Mitigation (Document Reference 6.2.12.10), is provided, which will be a requirement of the DCO, and will be used to form the basis of any mitigation measures to be secured prior to the development consent being granted.
- <u>12.4212.312</u> Overall, it is considered that the Proposed Development is capable of <u>achieving a range of positive residual effects and delivering a net biodiversity gain subject to the adherence to impact avoidance and mitigation measures on-site, along with the</u>



enhancement and long term management of the mosaic of habitats as currently present, and the delivery of off site ecological mitigation compensating for those residual effects, which cannot be avoided or mitigated.

