



CLEVE HILL SOLAR PARK

RESPONSES TO THE EXA'S WRITTEN QUESTIONS - APPENDICES **Appendix 9 – A statement of English Nature's Views About The** **Management of The Swale Site of Special Scientific Interest**

June 2019
Revision A

Document Reference: 10.1.1
Submitted: Deadline 2

www.clevehillsolar.com



CLEVE HILL
SOLAR PARK

Views About Management



A statement of English Nature's views about the management of The Swale Site of Special Scientific Interest (SSSI).

This statement represents English Nature's views about the management of the SSSI for nature conservation. This statement sets out, in principle, our views on how the site's special conservation interest can be conserved and enhanced. English Nature has a duty to notify the owners and occupiers of the SSSI of its views about the management of the land.

Not all of the management principles will be equally appropriate to all parts of the SSSI. Also, there may be other management activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

The management views set out below do not constitute consent for any operation. English Nature's written consent is still required before carrying out any operation likely to damage the features of special interest (see your SSSI notification papers for a list of these operations). English Nature welcomes consultation with owners, occupiers and users of the SSSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

Wet grassland with breeding and wintering bird interest

Wet grasslands occur on land that is subject to periodic flooding or has a seasonally high water table and is waterlogged for much of the year. Wet grassland often supports a wide variety of plants and animals, in particular birds and invertebrates, and is an important habitat for breeding waders and wintering wildfowl.

Wet grassland requires active management if it is to retain its conservation interest. Generally, each year's growth of vegetation must be removed. Otherwise the sward becomes dominated by tall, vigorous grasses and rushes which, together with an associated build up of dead plant matter, suppress less vigorous species and lower the botanical richness of the sward. Traditionally, this management is achieved by grazing. Cattle are often the preferred stock, being relatively tolerant of wet conditions and able to control tall grasses and rank vegetation. Cattle also tend to produce a rather uneven, structurally diverse sward. However, ponies, or even hill sheep, can be used if necessary. Grazing usually takes place at times between late spring and early autumn, but the precise timing and intensity will depend on local conditions and requirements, such as the need to avoid trampling ground-nesting birds. Heavy poaching should be avoided but light trampling can be beneficial in breaking down leaf litter and providing areas for seed germination. Agricultural

operations in general should be avoided before mid-June to minimise disturbance to breeding birds or the destruction of nests. An element of managed scrub, both within and fringing a field can be of importance to birds and invertebrates, as can a surrounding hedge.

Partial winter flooding is important in maintaining suitable habitat conditions for wintering birds. A mosaic of winter flooded grassland and permanently un-flooded grassland is desirable, with both temporary and permanent pools present. The maintenance of a mosaic of shallow surface pools and un-flooded areas during the winter will provide roosting and feeding habitat for wintering wildfowl and waders. From April onwards, the area of standing surface water should be reduced to increase the area available for nesting waders and also by concentrating aquatic invertebrates in small pools to provide suitable feeding areas for their young. Some shallow areas of flooding should be maintained until late June to provide patches of bare muddy ground on which the birds and their young can feed as raised sward height makes feeding on the drier areas more difficult. The birds using these features are directly vulnerable to disturbance, which can cause them to lose time spent feeding or drive them to areas with a poorer supply of food. Management should seek to minimise any harmful disturbance, especially at times when bird populations are under stress, such as severely cold conditions. Predators, especially crows and related species, should be controlled and this may be best achieved by limiting their nesting sites.

Careful maintenance of existing ditches and drains is usually acceptable practice, but abandonment or deepening of ditches can be harmful.

Ditches

Ditches are artificial habitats created by land drainage, or occasionally by the channelisation of small streams. They may represent the only remaining freshwater habitat within former wetland areas, and often support a wide range of aquatic plant and animal (in particular invertebrate) species that would have previously been more widespread in ponds and wetlands.

If left unmanaged, silt accumulates in the bottom of the ditches, and emergent plants such as reeds are able to colonise across the width of the ditch, leading to a loss of aquatic plant diversity and a gradual drying out of the ditch. To prevent this, periodical removal of sediment and vegetation may be necessary to return the ditch to an early stage of the management cycle. Ideally, ditch management should be undertaken on a rotation, creating a series of different management stages across a site at any one time. All stages of the management cycle have wildlife interest; recently cleared ditches are good for plants and animals which favour newly created habitats and cannot tolerate competition with other species; middle-stage ditches support a rich aquatic plant flora; and late-stage ditches may be important for a variety of invertebrates. The removal of both sediment and vegetation is usually better than simply cutting the vegetation, which does not recreate the earliest stages of the ditch management cycle. Where possible, management should aim to create shallow shelving margins rather than steep ditch sides. Where water voles are known to be present, the relevant good practice guidelines for ditch management and conservation should be followed.

Most ditch systems are subject to water level control, which should be managed to ensure that there is a sufficient depth of water (0.3-0.5m) in ditches throughout the year. Rapid or extreme changes in water level should be avoided unless they are known to be important to plant or animal communities relying on such fluctuations.

Ditches are susceptible to increased levels of nutrients which can cause a loss of aquatic plants and increases in algal growth. Other activities that can lead to this include the control or removal of aquatic plants, or the introduction of species such as bottom feeding coarse fish which uproot plants and disturb ditch sediments. Ditches are also susceptible to invasion by non-native aquatic plants such as floating pennywort and water fern, which are able to grow rapidly taking up available habitat and smothering other plants. Some native plants including a number of duckweed species are also able to take over in this way (although such growths are usually exacerbated by increased nutrients in the water) and management may be necessary to control such invasions where they cause a problem.

An important feature of many coastal ditch systems relating to water quality is the transition between freshwater and brackish water conditions, which often results in interesting changes in plant and animal species along ditch lengths or across the site. Management should aim to maintain such freshwater-brackish transitions where they occur.

Littoral sediments (mud and sand flats)

Intertidal mud and sand flats include a range of generally muddy or sandy low-gradient shores that are exposed to air during low tide and submerged during the higher tides. High energy shores, such as those on open coasts, are generally sandy in nature whilst more sheltered, low energy flats are muddier. They support a wide variety of marine invertebrates that represent an important food source for many fish and bird species.

Good water quality and sediment quality should be maintained, and the sediment budget within the estuarine or coastal system should not be restricted by anthropogenic influences.

The birds that use mud and sandflats for feeding and roosting are vulnerable to disturbance from human activities, for example, bait digging, dog walking and wildfowling. These activities can lead to reduced time spent feeding, or individuals being restricted to areas with a poor food supply. Disturbance should therefore be minimised, especially at times when bird populations may be stressed, such as during severe winter weather.

The location and extent of mud or sandflats is dependent on the extent to which the estuary or coast where they occur is constrained from responding to sea level rise and changing sediment regimes. Management needs to create space to enable landward roll-back to take place in response to sea-level rise, and should also allow the system to be dynamic and retain the flexibility to respond to associated changes such as the movement of physical features within the system, e.g. migrating subtidal sandbanks.

Coastal saltmarsh

Saltmarshes form the upper vegetated portions of intertidal mudflats in sheltered coastal locations, such as estuaries, lagoons and beach plains. There is typically a zonation of vegetation, from plants adapted to regular immersion by the tides (halophytes), through to more widespread plant species in the areas less frequently covered by the sea. The halophyte plant species are confined to this type of habitat, and areas of structurally diverse vegetation provide good invertebrate habitat. Saltmarshes are also important nursery sites for several fish species, and important refuge, feeding and breeding grounds for wading birds and wildfowl.

Where saltmarshes require management this has traditionally been achieved by grazing, and previously used regimes should be continued. Grazing provides a variety of different habitats, particularly for wintering bird species, and if grazing were to cease there may be a loss of botanical diversity. The precise timing and intensity will vary according to local conditions and requirements, for example the type or availability of stock, or the need to avoid trampling ground nesting birds. However on many sites, the aim of will be to create a short turf that can be attractive to overwintering wildfowl, with a reduction in stock density in the early summer for the benefit of ground-nesting birds. Indeed, careful reduction of grazing can increase the number of breeding birds, without significantly altering the plant species composition. Care should be taken not to overgraze the site, as this may reduce the diversity of animal and plant species that the saltmarsh is able to support, as well as potentially impact the sediments supporting the saltmarsh.

Not all saltmarsh habitats require active management to retain their conservation interest. Where there has not been a history of grazing, the saltmarsh will be able to maintain itself and grazing-sensitive species are likely to be present, therefore grazing should not be introduced.

There are a number of factors that are contributing to saltmarsh change that management may need to take into consideration. These include coastal erosion as a result of coastal flood-defence works, rising sea-levels, variations in sediment deposition, and land claim for development.

Hog's fennel

In the UK, hog's fennel is restricted to a few coastal localities in north Kent, north-east Essex and Suffolk. A number of factors are important to the survival of this plant. Scrub encroachment can reduce its abundance and vigour, and a degree of scrub management may be required. Scrub can compete directly with adult plants, as well as reducing the area of open sward with bare ground suitable for the establishment of hog's fennel seedlings. In addition, the seeds of hog's fennel require high summer temperatures to fully ripen, and shading from nearby scrub may affect their viability. Hog's fennel should not be subject to extensive or frequent mowing, as this reduces the size of plants and the number of flowering stems, and can have a detrimental affect on associated invertebrates. However, where cutting proves necessary it should be undertaken only on a small proportion of the plants in any given population, on a 3-5 year rotation. The appropriate time to cut will vary between sites, dependent upon the sometimes conflicting requirements of seed

ripening and rare moths which depend on hog's fennel. Grazing may prove to be a useful management tool but further research is required.

Coastal defence and stabilisation works could prove damaging. They can cause direct loss of adult plants and habitat, and cliff stabilisation can create conditions suitable for the development of more mature vegetation communities which may smother hog's fennel in time. However, limited disturbance need not be harmful because it can create bare ground for seed germination and distribute sections of rootstock, from which adult plants can grow. Hog's fennel appears to be very sensitive to salt damage and requires protection from saline inundation.

All habitats

The habitats within this site are highly sensitive to inorganic fertilisers and pesticides, applications of which should be avoided both within the site itself and in adjacent surrounding areas. Herbicides may be useful in targeting certain invasive species, but should be used with extreme care. Access to this site, and any recreational activities within, may also need to be controlled.