FIVE ESTUARIES OFFSHORE WIND FARM

FIVE ESTUARIES OFFSHORE WIND FARM

VOLUME 5, REPORT 4.1: HABITATS REGULATIONS ASSESSMENTS SITE INTEGRITY MATRICES – REVISION B (CLEAN)

Application Reference Application Document Number Revision Pursuant to Ecodoc Number Date EN010115 5.4.1 B Deadline 2 005076719-03 October 2024 COPYRIGHT © Five Estuaries Wind Farm Ltd All pre-existing rights reserved.

In preparation of this document Five Estuaries Wind Farm Ltd has made reasonable efforts to ensure that the content is accurate, up to date and complete for the purpose.

Revision	Date	Status/Reason for Issue	Originator	Checked	Approved
А	Mar-24	ES	GoBe	GoBe	VE OWFL
В	Oct-24	Deadline 2	GoBe	GoBe	VE OWFL

$\bigvee \Xi$

CONTENTS

1	Matrix key	.5
2	Index to matrices	.6
3	References	54

MATRICES

HRA Integrity Matrix 1: Margate and Long Sands (SAC)	8
HRA Integrity Matrix 2: Essex Estuaries SAC	9
HRA Integrity Matrix 3: Berwickshire and North Northumberland Coast SAC	10
HRA Integrity Matrix 4: Humber Estuary SAC	11
HRA Integrity Matrix 5: Humber Estuary RAMSAR	12
HRA Integrity Matrix 6: Southern North Sea SAC	13
HRA Integrity Matrix 7: Wash and North Norfolk Coast SAC	14
HRA Integrity Matrix 8: Transboundary Sites for Seals	15
HRA Integrity Matrix 9: Outer Thames Estuary SPA	17
HRA Integrity Matrix 10: Alde-Ore Estuary SPA	18
End of Matrix 10 HRA Integrity Matrix 11: Alde-Ore Estuary RAMSAR	18
HRA Integrity Matrix 12: Minsmere-Walberswick SPA	20
HRA Integrity Matrix 13: Minsmere-Walberswick RAMSAR	21
HRA Integrity Matrix 14: Deben Estuary SPA	22
HRA Integrity Matrix 15: Deben Estuary RAMSAR	23
HRA Integrity Matrix 16: Hamford Water SPA	24
HRA Integrity Matrix 17: Hamford Water RAMSAR	25
HRA Integrity Matrix 18: Stour and Orwell Estuaries SPA	26
HRA Integrity Matrix 19: Stour and Orwell Estuaries RAMSAR	27
HRA Integrity Matrix 20: Colne Estuary (Mid-Essex Coast Phase 2) SPA	28
HRA Integrity Matrix 21: Colne Estuary (Mid-Essex Coast Phase 2) RAMSAR	29
HRA Integrity Matrix 22: Dengie (Mid-Essex Coast Phase 1) SPA	30
HRA Integrity Matrix 23: Dengie (Mid-Essex Coast Phase 1) RAMSAR	31
HRA Integrity Matrix 24: Blackwater Estuary (Mid-Essex Coast Phase 4) SPA	32
HRA Integrity Matrix 25: Blackwater Estuary (Mid-Essex Coast Phase 4) RAMSAR	33
HRA Integrity Matrix 26: Flamborough and Filey Coast SPA	34
End of Matrix 26HRA Integrity Matrix 27: Farne Islands SPA	35
HRA Integrity Matrix 28: Vlaamse Banken (Special Area of Conservation (SAC)	37
HRA Integrity Matrix 29: Hamford Water SAC	38
HRA Integrity Matrix 30: Hamford Water SPA	39
End of Matrix 30HRA Integrity Matrix 31: Hamford Water RAMSAR	40
HRA Integrity Matrix 32: Stour and Orwell Estuaries SPA	42
HRA Integrity Matrix 33: Stour and Orwell Estuaries RAMSAR	44
HRA Integrity Matrix 34: Colne Estuary (Mid-Essex Coast Phase 2) SPA	46
End of Matrix 34HRA Integrity Matrix 35: Colne Estuary (Mid-Essex Coast Phase 2)	
RAMSAR	46
HRA Integrity Matrix 36: Abberton Reservoir SPA	49
HRA Integrity Matrix 37: Abberton Reservoir RAMSAR	51
HRA Integrity Matrix 38: Blackwater Estuary SPA	52
HRA Integrity Matrix 39: Blackwater Estuary RAMSAR	53

$\bigvee \Xi$

DEFINITION OF ACRONYMS

Term	Definition
EMF	Electromagnetic Field
ECC	Export Cable Corridor
HRA	Habitats Regulations Assessment
INNS	Invasive Non-Native Species
LSE	Likely Significant Effect
OWF	Offshore Wind Farm
PEIR	Preliminary Environmental Impact Report
PINS	Planning Inspectorate
RIAA	Report to Inform Appropriate Assessment
SAC	Special Area of Conservation
SPA	Special Protected Area
VE	Five Estuaries
VEOWFL	Five Estuaries Offshore Windfarm Limited
WTG	Wind Turbine Generator
Zol	Zone of Influence

UNITS

Units	Definition
km	Kilometre
cm	Centimetre
m	Metre
ha	Hectare
kg	Kilogram

$\bigvee \Xi$

1 MATRIX KEY

 \checkmark = A potential for AEoI has been identified

X = No potential for AEoI has been identified

Evidence for, or against adverse effects on European site qualifying feature and Likely Significant Effect is detailed within the footnotes to the integrity matrices

- C = construction
- O = operation and maintenance

D = decommissioning

= Screened out as effect not relevant to feature (no pathway)



2 INDEX TO MATRICES

2.1.1 This appendix presents the Integrity matrices for Five Estuaries Offshore Wind Farm (OWF, hereafter 'VE') prompted by Five Estuaries Offshore Windfarm Limited (hereafter 'the Applicant') in accordance with the structure and format specified in PINS Advice Note 10 (version 8, from November 2022).

Table 2.1 Index to matrices

Matrix Number	European site included within the assessment
Benthic and Inter	tidal Ecology
1	Margate and Long Sands (SAC)
2	Essex Estuaries SAC
Marine Mammal	
3	Berwickshire and North Northumberland Coast SAC
4	Humber Estuary SAC
5	Humber Estuary RAMSAR
6	Southern North Sea SAC
7	Wash and North Norfolk Coast SAC
8	Transboundary Sites for Seals
Offshore and Inte	rtidal Ornithology
9	Outer Thames Estuary SPA
10	Alde-Ore Estuary SPA
11	Alde-Ore Estuary RAMSAR
12	Minsmere-Walberswick SPA
13	Minsmere-Walberswick RAMSAR
14	Deben Estuary SPA
15	Deben Estuary RAMSAR
16	Hamford Water SPA
17	Hamford Water RAMSAR
18	Stour and Orwell Estuaries SPA
19	Stour and Orwell Estuaries RAMSAR
20	Colne Estuary (Mid-Essex Coast Phase 2) SPA
21	Colne Estuary (Mid-Essex Coast Phase 2) RAMSAR
22	Dengie (Mid-Essex Coast Phase 1) SPA
23	Dengie (Mid-Essex Coast Phase 1) RAMSAR



Matrix Number	European site included within the assessment
24	Blackwater Estuary (Mid-Essex Coast Phase 4) SPA
25	Blackwater Estuary (Mid-Essex Coast Phase 4) RAMSAR
26	Flamborough and Filey Coast SPA
27	Farne Islands SPA
Migratory Fish	
28	Vlaamse Banken (Special Area of Conservation (SAC)
Onshore Ecology	
29	Hamford Water SAC
30	Hamford Water SPA
31	Hamford Water RAMSAR
32	Stour and Orwell Estuaries SPA
33	Stour and Orwell Estuaries RAMSAR
34	Colne Estuary (Mid-Essex Coast Phase 2) SPA
35	Colne Estuary (Mid-Essex Coast Phase 2) RAMSAR
36	Abberton Reservoir SPA
37	Abberton Reservoir RAMSAR
38	Blackwater Estuary SPA
39	Blackwater Estuary RAMSAR

BENTHIC AND INTERTIDAL ECOLOGY

HRA Integrity Matrix 1: Margate and Long Sands (SAC)

The fine grity matrix 1. margate and Long Sands (SAS)	/																	
Name of European site:	Marg	Margate and Long Sands (SAC)																
EU Code:	UK00	JK0030371																
Distance to Project:	23.61	23.61 km to array																
Likely Effects of Project																		
Effect	Physical habitat loss/ disturbance		itat loss/	Suspended sediment/deposition			Accid	Accidental pollution			Invasive Non-Native Species (INNS)					Changes to physical processes		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Sandbanks which are slightly covered by sea water all the time	Ха	Xb	Хс	Ха	Ха	Ха	Xd	Xd	Xd	Xe	Xe	Xe		Xf			Xb	

Evidence supporting conclusions:

- Ха Given the short-term nature of the disturbance, the existing tolerance of the benthic habitats to disturbance within this area, and the predicted medium to high recoverability of the biotopes, it is considered that the site's conservation objectives will be maintained in the long-term. As highlighted in paragraphs 11.2.24 and 11.2.25 of Volume 5, Report 4: Report to Inform Appropriate Assessment (RIAA), the biotopes within this area are typical of high energy environments and are therefore naturally subject to, and tolerant of, high levels of physical disturbance. The communities that predominantly characterise these biotopes include infaunal mobile species such as polychaetes and bivalves. The likely biotopes present within the Annex 1 habitat 'sandbanks which are slightly covered by seawater all the time' are deemed to be of low vulnerability, medium to high recoverability and of national value. There is, therefore, no potential for an AEol.
- Given the small area of the SAC which will undergo disturbance, the VE ECC overlaps with 1.36 km² of the SAC, and the total area expected to be disturbed by sandwave clearance is 0.63 Xb km² (see Volume 6, Part 2, Chapter 5: Benthic and Intertidal Ecology), which equates to 0.09 % of the total SAC, the change is therefore very small compared to total area of habitat available within the SAC and therefore the site's conservation objectives will be maintained in the long-term. There is, therefore, no potential for an AEoI.
- Effects are considered to be similar or less than the construction phase and therefore there is no potential for an AEoI. Xc
- The primary source of the pollution risk comes from vessel movements and construction activities. These activities will be managed through the PEMP, ensuring that there are no adverse Xd environmental effects from the works (see paragraph 11.2.50 of Volume 5, Report 4: RIAA and Volume 6, Part 2, Chapter 5: Benthic Ecology). Therefore, there is no potential for an AEoI.
- Through increased vessel movements during construction and decommissioning there is a risk that vessels could contribute to the potential introduction or spread of marine INNS through Xe ballast water discharge, however the movement of commercial vessels is common throughout the region (Volume 6, Part 2, Chapter 9: Shipping and Navigation) and this provides an existing and potentially more likely method of transport for Marine INNS (due to the higher variety of ports and passage routes). Furthermore, there is a lack of evidence of any adverse effect from other offshore wind farms within the North Sea of having any adverse effect on key species and habitats through increasing the spread of marine INNS. Additionally, project level commitments to mitigate the risk such as following best practice guidelines and standard operating practices (as managed through the PEMP and biosecurity plan) will ensure the site's conservation objectives will be maintained in the long-term. There is, therefore, no potential for an AEoI.
- Xf Impacts from changes in EMFs arising from cables, are not considered to result in a significant effect on benthic ecology and intertidal receptors. EMFs are likely to be generated by subsea cables and detectable above background levels in close proximity to the cables. Although burial does not mask EMFs it increases the distance between species that may be affected by EMFs and the source. As the cable will be buried or protected, any behavioural responses are likely to be mitigated (see paragraph 11.2.78 of Volume 5, Report 4: RIAA). There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 2: Essex Estuaries SAC

Name of European site:	Esse	Essex Estuaries SAC																	
EU Code:	UK00	UK0013690																	
Distance to Project:	64.38	4.38 km to array																	
Likely Effects of Project																			
Effect	Physical habitat loss/ disturbance			Susp sedir depo	Suspended sediment/ deposition			Accidental pollution			Invasive Non-Native Species (INNS)			EMF			Changes to physical processes		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	
Estuaries	Xa	Ха	Ха	Ха	Ха	Xa	Xb	Xb	Xb	Xc	Xc	Xc		Xd			Ха		
Mudflats and sandflats not covered by seawater at low tide	Ха	Ха	Ха	Ха	Ха	Ха	Xb	Xb	Xb	Xc	Хс	Xc		Xd			Xa		
Salicornia and other annuals colonizing mud and sand	Ха	Ха	Ха	Ха	Ха	Ха	Xb	Xb	Xb	Xc	Xc	Хс		Xd			Ха		
Spartina swards	Xa	Ха	Ха	Ха	Ха	Xa	Xb	Xb	Xb	Xc	Хс	Xc		Xd			Ха		
Atlantic salt meadows	Ха	Ха	Ха	Ха	Ха	Ха	Xb	Xb	Xb	Хс	Хс	Xc		Xd			Ха		
Mediterranean and thermo-Atlantic halophilous scrubs	Ха	Ха	Ха	Ха	Ха	Ха	Xb	Xb	Xb	Xc	Xc	Хс		Xd			Ха		
Sandbanks which are slightly covered by sea water all the time	Ха	Ха	Ха	Ха	Ха	Ха	Xb	Xb	Xb	Xc	Xc	Xc		Xd			Ха		

Evidence supporting conclusions:

- The Essex Estuaries SAC site sits outside the Order limits, however the benthic study area, secondary zone of influence does interact with the site. Nevertheless, given the distance of the Ха site to potential direct interaction with construction and decommissioning activities, the site's conservation objectives will be maintained in the long-term (see paragraph 11.2.97 and 11.2.101 of Volume 5, Report 4: RIAA and Volume 6, Part 2, Chapter 5: Benthic and Intertidal Ecology). There is, therefore, no potential for an AEoI.
- The primary source of the pollution risk comes from vessel movements and construction activities. These activities will be managed through the PEMP, ensuring that there are no adverse Xb environmental effects from the works (see paragraph 11.2.50 of Volume 5, Report 4: RIAA and Volume 6, Part 2, Chapter 5: Benthic Ecology). Therefore, there is no potential for an AEoI.
- Through increased vessel movements during construction and decommissioning there is a risk that vessels could contribute to the potential introduction or spread of marine INNS through Xc ballast water discharge, however the movement of commercial vessels is common throughout the region (Volume 6, Part 2, Chapter 9: Shipping and Navigation) and this provides an existing and potentially more likely method of transport for Marine INNS (due to the higher variety of ports and passage routes). Furthermore, there is a lack of evidence of any adverse effect from other offshore wind farms within the North Sea of having any adverse effect on key species and habitats through increasing the spread of marine INNS. Additionally, project level commitments to mitigate the risk such as following best practice guidelines and standard operating practices (as managed through the PEMP (Volume 9, Report 18) and biosecurity plan) will ensure the site's conservation objectives will be maintained in the long-term. There is, therefore, no potential for an AEoI.
- Considering the distance of the site from potential direct EMF exposure during O&M activities and ensuring the preservation of the site's conservation objectives over the long term, there is Xd consequently no anticipated occurrence for an AEoI.



MARINE MAMMAL

HRA Integrity Matrix 3: Berwickshire and North Northumberland Coast SAC

Name of European site:	Berwickshire and North Northumberland Coast SAC														
EU Code:	UK0017	.0017072													
Distance to Project:	445.9 kr	15.9 km to array													
Likely Effects of Project															
Effect	Underwater noise			Vessel collision risk			Changes to prey			Physical disturba	habitat lo nce	ss/	Disturbance at haul out		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Grey seal	Ха		Ха	Xb	Xb	Xb	Хс	Xc	Хс	Xd		Xd	Xb	Xb	Xb

Evidence supporting conclusions:

- Ха There are a number of sources of underwater noise associated with Five Estuaries during construction, operation and decommissioning. These are addressed for marine mammals, including Grey Seal, in Volume 6, Part 2, Chapter 7: Marine Mammals and Section 11.3 of Volume 5, Report 4: RIAA. Overall, the impact of underwater noise will be negligible due to the implementation of the SIP (Volume 9, Report 15: Outline SNS SAC Site Integrity Plan) and MMMP (Volume 9, Report 14.1 and Report 14.2: MMMP – Piling and MMMP – UXO, respectively) to bring disturbance levels down and reduce the risk of injury to negligible levels. As a result, there will be no adverse effects on marine species or ecosystems, and the conservation objectives related to underwater noise levels will not be compromised. Therefore, there is no potential for AEoI arising from underwater noise pollution.
- Volume 6, Part 2, Chapter 7: Marine Mammals and paragraphs 11.3.132 and 11.3.154 of Volume 5, Report 4: RIAA provides an assessment of vessel collision risk with marine mammals. Xb Volume 6, Part 2, Chapter 9: Shipping and Navigation also provides an assessment of the level of vessel traffic within the areas surrounding VE, which already experiences high levels of vessel traffic. Therefore, it is considered increased vessel traffic associated with activities is insufficient to result in an increase in the risk of mortality or injury to marine mammals through vessel collisions. Furthermore, the adoption of a Working in Proximity to Wildlife protocol (see Volume 9, Report 18.1) would minimise any risk of collision further. There is, therefore, no AEoI.
- Volume 6, Part, Chapter 6: Fish and Shellfish Ecology provides an assessment of the impacts on marine mammals as a result of changes to prey. Overall it is not predicted that there will be Xc any impacts on marine mammals as a result of changes to the populations or general distributions of fish species within the vicinity of VE. This, coupled with the fact that there may be certain fish species that comprise the main part of grey seal diet (i.e., grey seal are considered to be generalist feeders and are thus not reliant on a single prey species) means that there is low risk of changes in prey abundance and distribution affecting the distribution of the grey seal feature. There is, therefore, no AEoI.
- Given the highly mobile nature of the species, the low number of seals in the vicinity of VE, the widely available comparable habitat, the relatively small area of habitat loss/disturbed, and the Xd generalist/opportunist nature of grey seals (Volume 6, Part 2, Chapter 7: Marine Mammal Ecology) it is considered that there is no adverse effect from a loss of available supporting habitat on grey seals.



HRA Integrity Matrix 4: Humber Estuary SAC

Name of European site:	Humber	Estuary S	SAC													
EU Code:	UK00301	JK0030170														
Distance to Project:	203.32 km to array															
Likely Effects of Project																
Effect	Underwa	derwater noise			Collision risk			Changes to prey			Physical habitat loss/ disturbance			Disturbance at haul out		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	
Grey seal	Ха		Ха	Xb	Xb	Xb	Хс	Хс	Хс	Xd		Xd	Xb	Xb	Xb	

Evidence supporting conclusions:

- There are a number of sources of underwater noise associated with Five Estuaries during construction, operation and decommissioning. These are addressed for marine mammals, including Xa Grey Seal, in Volume 6, Part 2, Chapter 7: Marine Mammals and Section 11.3 of Volume 5, Report 4: RIAA. Overall, the impact of underwater noise will be negligible due to the implementation of the SIP (Volume 9, Report 15: Outline SNS SAC Site Integrity Plan) and MMMP (Volume 9, Report 14.1 and Report 14.2: MMMP – Piling and MMMP – UXO, respectively) to bring disturbance levels down and reduce the risk of injury to negligible levels. As a result, there will be no adverse effects on marine species or ecosystems, and the conservation objectives related to underwater noise levels will not be compromised. Therefore, there is no potential for AEoI arising from underwater noise pollution.
- Xb Volume 6, Part 2, Chapter 7: Marine Mammals and paragraphs 11.3.132 and 11.3.154 of Volume 5, Report 4: RIAA provides an assessment of vessel collision risk with marine mammals. Volume 6, Part 2, Chapter 9: Shipping and Navigation also provides an assessment of the level of vessel traffic within the areas surrounding VE, which already experiences high levels of vessel traffic. Therefore, it is considered increased vessel traffic associated with activities is insufficient to result in an increase in the risk of mortality or injury to marine mammals through vessel collisions. Furthermore, the adoption of a Working in Proximity to Wildlife protocol (see Volume 9, Report 18.1) would minimise any risk of collision further. There is, therefore, no AEoI.
- Volume 6, Part, Chapter 6: Fish and Shellfish Ecology provides an assessment of the impacts on marine mammals as a result of changes to prey. Overall, it is not predicted that there will be Xc any impacts on marine mammals as a result of changes to the populations or general distributions of fish species within the vicinity of VE. This, coupled with the fact that there may be certain fish species that comprise the main part of grey seals diet (i.e., grey seal are considered to be generalist feeders and are thus not reliant on a single prey species) means that there is low risk of changes in prev abundance and distribution affecting the distribution of the grev seal feature. There is, therefore, no AEoI.
- Given the highly mobile nature of the species, the low number of seals in the vicinity of VE, the widely available comparable habitat, the relatively small area of habitat loss/disturbed, and the Xd generalist/ opportunist nature of grey seals (Volume 6, Part 2, Chapter 7: Marine Mammal Ecology) it is considered that there is no adverse effect from a loss of available supporting habitat on grey seals.

HRA Integrity Matrix 5: Humber Estuary RAMSAR

Name of European site:	Humber	Humber Estuary RAMSAR													
EU Code:	663														
Distance to Project:	197.29 k	197.29 km to array													
Likely Effects of Project															
Effect	Underwater noise		Collision risk		Changes to prey			Physical disturbar	habitat los ice	ss/	Disturba	nce at hau	l out		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Grey seal	Ха		Xa	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb

Evidence supporting conclusions:

- Xa There are a number of sources of underwater noise associated with Five Estuaries during construction, operation and decommissioning. These are addressed for marine mammals, including Grey Seal, in Volume 6, Part 2, Chapter 7: Marine Mammals and Section 11.3 of Volume 5, Report 4: RIAA. Overall, the impact of underwater noise will be negligible due to the implementation of the SIP (Volume 9, Report 15: Outline SNS SAC Site Integrity Plan) and MMMP (Volume 9, Report 14.1 and Report 14.2: MMMP – Piling and MMMP – UXO, respectively) to bring disturbance levels down and reduce the risk of injury to negligible levels. As a result, there will be no adverse effects on marine species or ecosystems, and the conservation objectives related to underwater noise levels will not be compromised. Therefore, there is no potential for AEoI arising from underwater noise pollution.
- Volume 6, Part 2, Chapter 7: Marine Mammals and paragraphs 11.3.132 and 11.3.154 of Volume 5, Report 4: RIAA provides an assessment of vessel collision risk with marine mammals. Xb Volume 6, Part 2, Chapter 9: Shipping and Navigation also provides an assessment of the level of vessel traffic within the areas surrounding VE, which already experiences high levels of vessel traffic. Therefore, it is considered increased vessel traffic associated with activities is insufficient to result in an increase in the risk of mortality or injury to marine mammals through vessel collisions. Furthermore, the adoption of a Working in Proximity to Wildlife protocol (see Volume 9, Report 18.1) would minimise any risk of collision further. There is, therefore, no AEoI.
- Volume 6, Part, Chapter 6: Fish and Shellfish Ecology provides an assessment of the impacts on marine mammals as a result of changes to prey. Overall, it is not predicted that there will be Xc any impacts on marine mammals as a result of changes to the populations or general distributions of fish species within the vicinity of VE. This, coupled with the fact that there may be certain fish species that comprise the main part of grey seals diet (i.e., grey seal are considered to be generalist feeders and are thus not reliant on a single prey species) means that there is low risk of changes in prey abundance and distribution affecting the distribution of the grey seal feature. There is, therefore, no AEoI.
- Given the highly mobile nature of the species, the low number of seals in the vicinity of VE, the widely available comparable habitat, the relatively small area of habitat loss/disturbed, and the Xd generalist/opportunist nature of grey seals (Volume 6, Part 2, Chapter 7: Marine Mammal Ecology) it is considered that there is no adverse effect from a loss of available supporting habitat on grey seals.

\sim

HRA Integrity Matrix 6: Southern North Sea SAC

Name of European site:	Souther	Southern North Sea SAC													
EU Code:	UK0030	395													
Distance to Project:	0 km to	km to array													
Likely Effects of Project															
Effect	Underwater noise			Collision risk C			Change	Changes to prey			tal pollutio s in water	on and quality	Physical habitat loss/ disturbance		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Harbour porpoise	Xa	Ха	Ха	Xb	Xb	Xb	Хс	Хс	Хс	Xd		Xd	Xe	Xe	Xe

Evidence supporting conclusions:

- There are a number of sources of underwater noise associated with Five Estuaries during construction, operation and decommissioning. These are addressed for marine mammals in Volume Ха 6, Part 2, Chapter 7: Marine Mammals and Section 11.3 of Volume 5, Report 4: RIAA. Overall, the impact of underwater noise will be negligible due to the implementation of the SIP (Volume 9, Report 15: Outline SNS SAC Site Integrity Plan) and MMMP (Volume 9, Report 14.1 and Report 14.2: MMMP – Piling and MMMP – UXO, respectively) bring disturbance levels to below seasonal thresholds and reduce the risk of injury to negligible levels. As a result, there will be no adverse effects on marine species or ecosystems, and the conservation objectives related to underwater noise levels will not be compromised. Therefore, there is no potential for adverse effects on integrity (AEoI) arising from underwater noise pollution.
- Xb Volume 6, Part 2, Chapter 7: Marine Mammals and paragraphs 11.3.132 and 11.3.154 of Volume 5, Report 4: RIAA provides an assessment of vessel collision risk with marine mammals. Volume 6, Part 2, Chapter 9: Shipping and Navigation also provides an assessment of the level of vessel traffic within the areas surrounding VE, which already experiences high levels of vessel traffic. Therefore, it is considered increased vessel traffic associated with activities is insufficient to result in an increase in the risk of mortality or injury to marine mammals through vessel collisions. Furthermore, the adoption of a Working in Proximity to Wildlife protocol (see Volume 9, Report 18.1) would minimise any risk of collision further. There is, therefore, no AEoI.
- Volume 6, Part, Chapter 6: Fish and Shellfish Ecology provides an assessment of the impacts on marine mammals as a result of changes to prey. Overall, it is not predicted that there will be Xc any impacts on marine mammals as a result of changes to the populations or general distributions of fish species within the vicinity of VE. This, coupled with the fact that there may be certain fish species that comprise the main part of harbour porpoises' diet (i.e., harbour porpoise are considered to be generalist feeders and are thus not reliant on a single prey species) means that there is low risk of changes in prey abundance and distribution affecting the distribution of the harbour porpoise feature. There is, therefore, no AEoI.
- An Outline PEMP (Volume 9, Report 18) has been provided in the DCO application to ensure that the potential for contaminant release is strictly controlled. The PEMP will include a Marine Xd Pollution Contingency Plan (MPCP), and enables the conclusion that there is, therefore, no AEoI.
- Given the highly mobile nature of the species, the widely available comparable habitat, the relatively small area of habitat loss/ disturbed, and the generalist/ opportunist nature of harbour Xe porpoise (Volume 6, Part 2, Chapter 7: Marine Mammal Ecology) it is considered that there is no adverse effect from a loss of available supporting habitat on harbour porpoise.

HRA Integrity Matrix 7: Wash and North Norfolk Coast SAC

Name of European site:	Wash and	North Norf	olk Coast S	AC											
EU Code:	UK0017075														
Distance to Project:	126.45 km to array														
Likely Effects	Likely Effects of Project														
Effect	Underwate	r noise		Collision ris	sk		Changes to prey			Physical habitat loss/ disturbance			Disturbance at haul out		
Stage of Development	С	0	D	с	0	D	С	0	D	с	0	D	С	0	D
Harbour seal	Ха		Xa	Xb	Xb	Xb	Хс	Xc	Хс	Xd		Xd	Xb	Xb	Xb

Evidence supporting conclusions

- Ха There are a number of sources of underwater noise associated with Five Estuaries during construction, operation and decommissioning. These are addressed for marine mammals in Volume 6, Part 2, Chapter 7: Marine Mammals and Section 11.3 of Volume 5, Report 4: RIAA. Overall, the impact of underwater noise will be negligible due to the implementation of the SIP (Volume 9, Report 15: Outline SNS SAC Site Integrity Plan) and MMMP (Volume 9, Report 14.1 and Report 14.2: MMMP – Piling and MMMP – UXO, respectively) to bring disturbance levels down and reduce the risk of injury to negligible levels. As a result, there will be no adverse effects on marine species or ecosystems, and the conservation objectives related to underwater noise levels will not be compromised. Therefore, there is no potential for adverse effects on integrity (AEoI) arising from underwater noise pollution.
- Volume 6, Part 2, Chapter 7: Marine Mammals and paragraphs 11.3.132 and 11.3.154 of Volume 5, Report 4: RIAA provides an assessment of vessel collision risk with marine mammals. Xb Volume 6, Part 2, Chapter 9: Shipping and Navigation also provides an assessment of the level of vessel traffic within the areas surrounding VE, which already experiences high levels of vessel traffic. Therefore, it is considered increased vessel traffic associated with activities is insufficient to result in an increase in the risk of mortality or injury to marine mammals through vessel collisions. Furthermore, the adoption of a Working in Proximity to Wildlife protocol (see Volume 9, Report 18.1) would minimise any risk of collision further. There is, therefore, no AEoI. With regards to disturbance at haul out sites during construction, operation and decommissioning it is considered that the effect (in terms of disturbance) is of negligible significance for harbour seals, and there is therefore no AEoI.
- Volume 6, Part, Chapter 6: Fish and Shellfish Ecology provides an assessment of the impacts on marine mammals as a result of changes to prey. Overall it is not predicted that there will be Xc any impacts on marine mammals as a result of changes to the populations or general distributions of fish species within the vicinity of VE. This, coupled with the fact that there may be certain fish species that comprise the main part of harbour seals diet (i.e., harbour seal are considered to be generalist feeders and are thus not reliant on a single prey species) means that there is low risk of changes in prey abundance and distribution affecting the distribution of the harbour seal feature. There is, therefore, no AEoI.
- Given the highly mobile nature of the species, the low number of seals in the vicinity of VE, the widely available comparable habitat, the relatively small area of habitat loss/disturbed, and the Xd generalist/opportunist nature of harbour seals (Volume 6, Part 2, Chapter 7: Marine Mammal Ecology) it is considered that there is no adverse effect from a loss of available supporting habitat on harbour seals.

$\vee \Xi$

HRA Integrity Matrix 8: Transboundary Sites for Seals

Name of European site:	Transt	Transboundary sites for seals (Harbour seal; and Grey seal)													
EU Code:	Various	S													
Distance to Project:	Various	S													
Likely Effects of Project															
Effect	Underv	vater nois	e	Collis	Collision risk		Chang	jes to prey	/	Physical habitat loss/ disturbance			Disturbance at haul out		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Bancs des Flandres SCI	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
Vlaamse Banken SAC	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
Doggersbank (Netherlands) SAC	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
Klaverbank SCI	Ха		Xa	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
Noordzeekustone SCI	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Хс	Xd		Xd	Xb	Xb	Xb
SBZ 1 SCI	Ха		Xa	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
SBZ 2 SCI	Ха		Xa	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
SBZ 3 SCI	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Хс	Xd		Xd	Xb	Xb	Xb
Vlakte van de Raan SCI	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Хс	Xd		Xd	Xb	Xb	Xb
Voordelta SCI	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Хс	Xd		Xd	Xb	Xb	Xb
Waddenzee SCI	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
Westerschelde & Saeftinghe	Ха		Ха	Xb	Xb	Xb	Xc	Xc	Xc	Xd		Xd	Xb	Xb	Xb
*Note that some sites may be considered	d senarately fo	r other fo	aturo(s) n	otably so	ale										

Note that some sites may be considered separately for other feature(s), notably seals

Evidence supporting conclusions:

Ха There are a number of sources of underwater noise associated with Five Estuaries during construction and decommissioning. These are addressed for marine mammals in Volume 6, Part 2, Chapter 7: Marine Mammals and Section 11.3 of Volume 5, Report 4: RIAA. Overall, the impact of underwater noise will be negligible due to the implementation of the SIP (Volume 9, Report 15: Outline SNS SAC Site Integrity Plan) and MMMP (Volume 9, Report 14.1 and Report 14.2: MMMP – Piling and MMMP – UXO, respectively) to bring disturbance levels down and reduce the risk of injury to negligible levels. Furthermore, there is predicted to be a low number of seals to be impacted and the proportion of the population this represents. As a result, there will be



no adverse effects on marine species or ecosystems, and the conservation objectives related to underwater noise levels will not be compromised. Therefore, there is no potential for adverse effects on integrity (AEoI) arising from underwater noise pollution.

- Xb Volume 6, Part 2, Chapter 7: Marine Mammals and paragraphs 11.3.132 and 11.3.154 of Volume 5, Report 4: RIAA provides an assessment of vessel collision risk with marine mammals. Volume 6, Part 2, Chapter 9: Shipping and Navigation also provides an assessment of the level of vessel traffic within the areas surrounding VE, which already experiences high levels of vessel traffic. Therefore, it is considered increased vessel traffic associated with activities is insufficient to result in an increase in the risk of mortality or injury to marine mammals through vessel collisions. Furthermore, the adoption of a Working in Proximity to Wildlife protocol (see Volume 9, Report 18.1) would minimise any risk of collision further. There is, therefore, no AEoI. With regards to disturbance at haul out sites during construction, operation and decommissioning it is considered that the effect (in terms of disturbance) is of negligible significance for harbour seals, and there is therefore no AEoI.
- Volume 6, Part, Chapter 6: Fish and Shellfish Ecology provides an assessment of the impacts on marine mammals as a result of changes to prey. Overall, it is not predicted that there will be Xc any impacts on marine mammals as a result of changes to the populations or general distributions of fish species within the vicinity of VE. This, coupled with the fact that there may be certain fish species that comprise the main part of seals diet (i.e., seals are considered to be generalist feeders and are thus not reliant on a single prey species) means that there is low risk of changes in prey abundance and distribution affecting the distribution of any seal feature. There is, therefore, no AEoI.
- Given the highly mobile nature of seals, the low number of seals in the vicinity of VE, the widely available comparable habitat, the relatively small area of habitat loss/disturbed, and the Xd generalist/opportunist nature of harbour seals (ES Volume 6, Part 2, Chapter 7: Marine Mammal Ecology) it is considered that there is no adverse effect from a loss of available supporting habitat on seals.



OFFSHORE AND INTERTIDAL ORNITHOLOGY

HRA Integrity Matrix 9: Outer Thames Estuary SPA

Name of European site:	Outer Thames Estuary SPA						
EU Code:	UK9020309A						
Distance to Project:	17.24 km to array						
Likely Effects	of Project						
Effect	Disturbance and displacement due to work activity and ves	sel movements within the ECC only					
Stage of Development	C	0	D				
Red-throated diver	Ха		Ха				

Evidence supporting conclusions:

Volume 6, Part 2, Chapter 4: Offshore Ornithology and paragraph 11.4.50 onwards in Volume 5, Report 4: RIAA assess the potential impact upon the Outer Thames Estuary SPA and the Ха feature Red Throated Diver. Overall, based on available evidence regarding red-throated diver displacement by operational OWFs, it is suggested that there will be little or no impact on adult survival as a result of displacement, and that any impact would probably be undetectable at the population level. Furthermore, following Natural England's advice a best practice protocol to minimise disturbance on red-throated divers will be adopted and can be found in Volume 9, Report 18.1: Working in Proximity to Wildlife in the Marine Environment. Additionally, export cable installatiomn will not be carried out within the Outer Thames Estuary between 1 November to 31 March inclusive. As a result, there is, therefore no potential for an AEoI.





HRA Integrity Matrix 10: Alde-Ore Estuary SPA

Name of European site:	Alde-Ore Estuary SPA						
EU Code:	UK9009112						
Distance to Project:	37.44 km to array						
Likely Effects of Project							
Effect	Collision risk						
Stage of Development	C	0	D				
Lesser black- backed gull		√a					
Avocet		Xb					
Redshank		Xb					
Ruff		Xb					

Evidence supporting conclusions:

- Paragraphs 12.4.97 of Volume 5, Report 4: RIAA considers the impacts from collision of Lesser Black-Backed Gull (LBBG) as a feature of the Alde-Ore Estuary SPA, in-combination with √a other projects. The total in-combination number of lesser black-backed gulls from the Alde-Ore SPA predicted to be subject to collision resultant mortality from the assessed OWFs, including VE, is 57 (56.2) breeding adults. Considering the potential impact of this loss to the Alde-Ore SPA, with a citation population of 28,140 breeding adults and annual background mortality of 3,236 breeding adults per annum, the addition of 57 breeding adults suffering collision consequent mortality would represent a 1.736% increase in baseline mortality, of which VE contributes five (5.48) individuals, representing a 0.169% increase in baseline mortality. Taking into account the ongoing declines at this population, the potential for an AEoI on the conservation objectives for lesser black-backed gull at the Alde Ore Estuary SPA cannot be ruled out. As a result, a derogation case has been conceded for this site and the LBBG feature in-combination. Therefore, a LBBG Compensation – Evidence, Site Selection and Roadmap document (Volume 5, Report 5.3) and LBBG Implementation and Monitoring Plan (Volume 5, Report 5.6) have been submitted as part of the DCO application.
- Avocet, Redshank and Ruff were screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Xb Annex 14.4: Migratory Collision Risk Modelling, assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.





HRA Integrity Matrix 11: Alde-Ore Estuary RAMSAR

Name of European site:	Alde-Ore Estuary RAMSAR						
EU Code:	UK9009112						
Distance to Project:	37.44 km to array						
Likely Effects of Project							
Effect	Collision risk						
Stage of Development	С	0	D				
Lesser black- backed gull		√a					
Avocet		Xb					
Redshank		Xb					

Evidence supporting conclusions:

- Paragraphs 12.4.97 of Volume 5, Report 4: RIAA considers the impacts from collision of Lesser Black-Backed Gull (LBBG) as a feature of the Alde-Ore Estuary SPA and Ramsar, in-√a combination with other projects. The total in-combination number of lesser black-backed gulls from the Alde-Ore SPA and Ramsar predicted to be subject to collision resultant mortality from the assessed OWFs, including VE, is 57 (56.2) breeding adults. Considering the potential impact of this loss to the Alde-Ore SPA and Ramsar, with a citation population of 28,140 breeding adults and annual background mortality of 3,236 breeding adults per annum, the addition of 57 breeding adults suffering collision consequent mortality would represent a 1.736% increase in baseline mortality, of which VE contributes five (5.48) individuals, representing a 0.169% increase in baseline mortality. Taking into account the ongoing declines at this population, the potential for an AEoI on the conservation objectives for lesser black-backed gull at the Alde Ore Estuary SPA cannot be ruled out in-combination with other plans and projects. As a result, a derogation case has been conceded for this site and the LBBG feature in-combination. Therefore, a LBBG Compensation – Evidence, Site Selection and Roadmap document (Volume 5, Report 5.3) and LBBG Implementation and Monitoring Plan (Volume 5, Report 5.6) have been produced.
- Avocet and Redshank were screened due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Xb Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEol.





HRA Integrity Matrix 12: Minsmere-Walberswick SPA

Name of European site:	Minsmere-Walberswick SPA								
EU Code:	UK9009101	UK9009101							
Distance to Project:	41.88 km to array								
Likely Effects of Project									
Effect	Collision risk (migration)								
Stage of Development	С	0	D						
Avocet		Ха							
Bittern		Ха							
Gadwall		Ха							
Greater white-fronted goose		Ха							
Hen harrier		Ха							
Shoveler		Ха							
Teal		Ха							

Evidence supporting conclusions:

Avocet, Bittern, Gadwall, Greater white-fronted goose, Hen harrier, Shoveler and Teal were screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards Ха within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.

HRA Integrity Matrix 13: Minsmere-Walberswick RAMSAR

Name of European site:	Minsmere-Walberswick RAMSAR		
EU Code:	UK1044		
Distance to Project:	41.88 km to array		
Likely Effects	of Project		
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Avocet		Ха	
Bittern		Ха	
Gadwall		Ха	
Marsh harrier		Ха	
Shoveler		Ха	
Teal		Ха	
Bearded tit		Ха	

Evidence supporting conclusions:

Avocet, Bittern, Gadwall, Marsh harrier, Shoveler, Teal and Bearded tit were screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Ха Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 14: Deben Estuary SPA

Name of European site:	Deben Estuary SPA		
EU Code:	UK9009261		
Distance to Project:	48.45 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Dark-bellied brent goose		Ха	
Avocet		Ха	

Evidence supporting conclusions:

Dark-bellied brent goose and Avocet were screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part Ха 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.





HRA Integrity Matrix 15: Deben Estuary RAMSAR

Name of European site:	Deben Estuary RAMSAR		
EU Code:	UK9009261		
Distance to Project:	48.45 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Dark-bellied brent goose		Ха	

Evidence supporting conclusions:

Ха Dark-bellied brent goose is screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 16: Hamford Water SPA

Name of European site:	Hamford Water SPA		
EU Code:	UK0030377		
Distance to Project:	51.17 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Avocet		Ха	
Black-tailed godwit		Ха	
Dark-bellied brent goose		Ха	
Grey plover		Ха	
Redshank		Ха	
Ringed plover		Ха	
Shelduck		Ха	
Teal		Ха	

Evidence supporting conclusions:

Xa The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.

		<u> </u>

HRA Integrity Matrix 17: Hamford Water RAMSAR

Name of European site:	Hamford Water RAMSAR		
EU Code:	UK11028		
Distance to Project:	52.89 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Black-tailed godwit		Ха	
Dark-bellied brent goose		Ха	
Redshank		Ха	
Ringed plover		Ха	

Evidence supporting conclusions:

Xa The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 18: Stour and Orwell Estuaries SPA

Name of European site:	Stour and Orwell Estuaries SPA		
EU Code:	UK9009121		
Distance to Project:	54.81 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Black-tailed godwit		Ха	
Dark-bellied brent goose		Ха	
Dunlin		Ха	
Grey plover		Ха	
Knot		Ха	
Pintail		Ха	
Redshank		Ха	

Evidence supporting conclusions:

The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Ха Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.

	\bigvee	

HRA Integrity Matrix 19: Stour and Orwell Estuaries RAMSAR

Name of European site:	Stour and Orwell Estuaries RAMSAR		
EU Code:	UK9009121		
Distance to Project:	54.81 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Black-tailed godwit		Ха	
Dark-bellied brent goose		Ха	
Dunlin		Ха	
Grey plover		Ха	
Knot		Ха	
Pintail		Ха	
Redshank		Ха	

Evidence supporting conclusions:

The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Ха Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 20: Colne Estuary (Mid-Essex Coast Phase 2) SPA

Name of European site:	Colne Estuary (Mid-Essex Coast Phase 2) SPA	
EU Code:	UK9009243		
Distance to Project:	66.51 km to array		
Likely Effects of Project			
Effect	Collision risk (migration)		
Stage of Development	С	0	D
Dark-bellied brent goose		Ха	
Pochard		Ха	
Redshank		Ха	
Ringed Plover		Ха	

Evidence supporting conclusions:

Ха The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 21: Colne Estuary (Mid-Essex Coast Phase 2) RAMSAR

Name of European site:	Colne Estuary (Mid-Essex Coast Phase 2) RAMSAR										
EU Code:	JK9015022										
Distance to Project:	66.63 km to array										
Likely Effects of Project											
Effect	Collision risk (migration)										
Stage of Development	С	0	D								
Dark-bellied brent goose		Ха									
Redshank		Ха									

Evidence supporting conclusions:

The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Ха Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 22: Dengie (Mid-Essex Coast Phase 1) SPA

Name of European site:	Dengie (Mid-Essex Coast Phase 1) SPA										
EU Code:	UK9009242										
Distance to Project:	73.63 km to array area										
Likely Effects of Project											
Effect	Collision risk (migration)										
Stage of Development	С	0	D								
Dark-bellied brent goose		Ха									
Grey plover		Ха									
Knot		Ха									

Evidence supporting conclusions:

The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Ха Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 23: Dengie (Mid-Essex Coast Phase 1) RAMSAR

Name of European site:	Dengie (Mid-Essex Coast Phase 1) RAMSAR										
EU Code:	UK9009242										
Distance to Project:	73.63 km to array area										
Likely Effects of Project											
Effect	Collision risk (migration)										
Stage of Development	С	0	D								
Dark-bellied brent goose		Ха									
Grey plover		Ха									
Knot		Ха									

Evidence supporting conclusions:

The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Ха Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 24: Blackwater Estuary (Mid-Essex Coast Phase 4) SPA

Name of European site:	Blackwater Estuary (Mid-Essex Coast Phase 4) SPA											
EU Code:	UK9009245	UK9009245										
Distance to Project:	77.69 km to array											
Likely Effects of Project												
Effect	Collision risk (migration)											
Stage of Development	С	0	D									
Black-tailed godwit		Ха										
Dark-bellied Brent goose		Ха										
Dunlin		Ха										
Grey plover		Ха										

Evidence supporting conclusions:

Ха The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this SPA can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 25: Blackwater Estuary (Mid-Essex Coast Phase 4) RAMSAR

Name of European site:	Blackwater Estuary (Mid-Essex Coast Phase 4) RAMSAR												
EU Code:	UK9009245	UK9009245											
Distance to Project:	77.81 km to array												
Likely Effects of Project													
Effect	Collision risk (migration)												
Stage of Development	С	0	D										
Black-tailed godwit		Ха											
Dark-bellied Brent goose		Ха											
Dunlin		Ха											
Grey plover		Ха											

Evidence supporting conclusions:

Xa The above species are screened in due to the potential risk of collision during migration. Paragraph 11.4.226 onwards within Volume 5, Report 4: RIAA and Volume 6, Part 5, Annex 14.4: Migratory Collision Risk Modelling assesses the potential impact of collision upon these species, utilising MigroPath analyses. Overall and considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at this Ramsar can be considered to be minimal and make no material contribution to any changes in population or baseline mortality. There is, therefore, no potential for an AEoI.



HRA Integrity Matrix 26: Flamborough and Filey Coast SPA

Name of European site:	Flamborough and Filey	Flamborough and Filey Coast SPA												
EU Code:	UK9006101	K9006101												
Distance to Project:	275.50 km to array	.75.50 km to array												
Likely Effects of Project														
Effect	Collision risk			Direct disturbance and displacement										
Stage of Development	С	0	D	С	0	D								
Kittiwake		Ха												
Gannet		Ха		Xb		Xb								
Guillemot				Xb	Xb	Xb								
Razorbill				Xb	Xb	Xb								

Evidence supporting conclusions:

Ха As highlighted for Gannet in Paragraphs 11.4.174 onwards, of Volume 5, Report 4: RIAA for the effect of collision risk, the addition of less than two possible additional breeding adult mortality per annum equates to less than a 1% increase in baseline mortality, when considering either the citation or the latest colony count. This level of impact would be indistinguishable from natural fluctuations in the baseline mortality rate of breeding adults from this population per annum. Similarly, as highlighted for Kittiwake in Paragraphs 11.4.188 onwards, the addition of one additional adult mortality in the non-breeding equates to less than 1% (0.006%) increase in baseline mortality, when considering either the citation or the latest colony count. Considering the level of impact is <0.01% increase in baseline mortality it would be indistinguishable from natural fluctuations in the baseline mortality rate of breeding adults from this population per annum and is considered to be no material contribution to the natural baseline mortality rates of the colony. Therefore, for both species, there is no potential for an AEoI. Xb As highlighted in Paragraph 11.4.144 and 11.4.145 of Volume 5, Report 4: RIAA, across all bio-seasons the number of gannets estimated to occur in the array area and a 2 km buffer is 940 (939.8) individuals. The total predicted displacement consequent mortality from these birds is estimated at 7 (6.58) individuals per annum. The impact attributed to FFC SPA throughout the operational life of VE is under two (1.51) breeding adult from FFC SPA per annum across all bio-seasons. This prediction of this total consequential additional mortality represents an increase of 0.085% when considering the citation population or an increase of 0.047% when considering the recent colony count across all bio-seasons per annum. This level of impact would be indistinguishable from natural fluctuations in the population. As highlighted in Paragraph 11.4.149 onwards of Volume 5, Report 4: RIAA, in the non-breeding bio-season the number of quillemots estimated to occur in the array area and 2 km buffer is 3,698 (3,698.0) individuals. The total predicted consequent mortality of birds within the array from displacement (based on 50% displacement, 1% mortality) is estimated at less than 19 (18.49) individuals. On the assumption that 4.41% of these guillemots are deemed to be breeding adults from the FFC SPA during the non-breeding bio-season (presented in Volume 6, Part 5, Annex 4.15: Apportioning Note), then the consequent mortality from being displaced is estimated at less than one (0.82) breeding adult. Displacement consequent mortalities are based on the range advocated by Natural England (30% to 70% displacement, 1% to 2% mortality). Based on a citation population of 83,214 breeding adults and an annual background mortality of 5,076 breeding adults per annum, the addition of less than one predicted breeding adult mortality would represent an increase in baseline mortality of 0.016%. As the population of guillemot has increased significantly since the citation population count the potential impact on the population is more reasonably assessed against the latest population count undertaken in 2022, consisting of 149,980 individuals and an annual background mortality of 9,149 individuals. On this basis, this would represent a 0.009% increase in baseline mortality in the non-breeding bio-season. As highlighted in Paragraph 11.4.157 onwards of Volume 5, Report 4: RIAA, in the non-breeding bioseasons, the number of **razorbills** estimated to occur in the array area and 2 km buffer is 757 (756.5) individuals during the return migration, 284 (283.6) during the post-breeding migration, and 1,046 (1046.0) in the migration free winter bio-season. The total predicted consequent mortality of birds within the array area and 2 km buffer from displacement is four (3.8) individuals in the return migration bio-season, less than two (1.4) individuals in the post-breeding migration bio-season and five (5.2) individuals in the migration-free winter bio-season (based on 50% displacement, 1% mortality). On the assumption that 3.38% of the razorbills are deemed to be breeding adults from the FFC SPA during the return migration bio-season (presented in Volume 6. Part 5, Annex 4.15: Apportioning Note), then the consequent mortality from being displaced is estimated at less than one (0.13) breeding adult per annum. During the post-breeding migration bio-season, it is considered that 3.38% of the razorbills are breeding adults from the FFC SPA (presented in Volume 6, Part 5, Annex 4.15: Apportioning Note). Therefore, the consequent mortality of adult birds from FFC SPA from being displaced is estimated at <0.1 (0.05) breeding adult per annum. During the migration-free winter bio-season, it is considered that



0.91% of the razorbills are breeding adults from the FFC SPA (presented in Volume 6, Part 5, Annex 4.15: Apportioning Note). Therefore, the consequent mortality of adult birds from FFC SPA from being displaced is estimated at <0.1 (0.05) breeding adult per annum. This equates to a total consequent mortality from displacement across the entire non-breeding bio-season of less than one (0.22) breeding adults per annum. Based on the citation count of 21,140 breeding adults and a baseline mortality of 2,220 breeding adults per annum, the addition of less than one predicted breeding adult mortality would represent a 0.010% increase in baseline mortality during the non-breeding bio-season. As the population of razorbills has increased significantly since the citation population count the potential impact on the population is more reasonably assessed against the latest population count undertaken in 2022, consisting of 61,346 breeding adults and an annual background mortality of 6,441 breeding adults per annum. On this basis, this would represent a 0.004% increase in baseline mortality during the non-breeding bio-season. Overall, it is considered that there is no potential for an AEol to the conservation objectives of the gannet, guillemot and razorbill feature of the Flamborough and Filey Coast SPA.



HRA Integrity Matrix 27: Farne Islands SPA

Name of European site:	Farne Islands SPA										
EU Code:	UK9006021	IK9006021									
Distance to Project:	472.54 km to array										
Likely Effects of Project											
Effect	Direct disturbance and displacement										
Stage of Development	С	0	D								
Guillemot	Ха	Ха	Ха								

Evidence supporting conclusions:

As highlighted in Paragraph 11.4.168 onwards of Volume 5, Report 4: RIAA, in the non-breeding bio-season the number of guillemots estimated to occur in the array area and 2 km buffer is Ха 3,698 (3,697.98) individuals. The total predicted consequent mortality of birds within the array from displacement (based on 50% displacement, 1% mortality) is estimated at less than 19 (18.49) individuals. On the assumption that 3.73% of the guillemots are deemed to be breeding adults from the Farne Islands SPA during the non-breeding bio-season (presented in Volume 6, Part 5, Annex 4.15: Apportioning Note), then the consequent mortality from being displaced is estimated at less than one (0.69) breeding adult. Based on the citation population of 65,750 breeding adults and a baseline mortality of 4,011 breeding adults per annum, the addition of less than one mortality would represent a 0.017% increase in baseline mortality. As the population of guillemot has changed since the citation population count the potential impact on the population is more reasonably assessed against the latest population count undertaken in 2017, consisting of 64,042 breeding adults and an annual background mortality of 3,907 breeding adults per annum. On this basis, this would represent a 0.018% increase in baseline mortality in the nonbreeding bioseason. This level of impact would be indistinguishable from natural fluctuations in the population. Therefore, there is no potential for an AEoI.





MIGRATORY FISH

HRA Integrity Matrix 28: Vlaamse Banken (Special Area of Conservation (SAC)

Name of European site:	Vlaamse Banken SAC										
EU Code:	BEMNZ000										
Distance to Project:	34.75 km to array										
Likely Effects of Project											
Effect	Underwater noise										
Stage of Development	С	0	D								
Twaite shad	Ха		Ха								

Evidence supporting conclusions:

Although Group 3 fish species are considered to be the most sensitive to underwater noise, due to their mobile nature Twaite shad are considered a fleeing/ mobile receptor in the Ха assessment presented in Volume 6, Part 2, Chapter 6: Fish and Shellfish Ecology; as they are expected to transit an impacted area (unlike some spawning receptors that exhibit site fidelity such as herring and sandeel). Therefore, twaite shad are expected to recover quickly, returning to normal behaviours and recolonise areas shortly after an impact. Furthermore, Group 3 species are broadly distributed and present in abundance within the southern North Sea region with the small impact range potentially affecting only a small proportion of the regional population, according to Volume 6, Part 2, Chapter 6: Fish and Shellfish Ecology. Therefore, given that any impacts from underwater noise are expected to be of local scale and the intermittent nature of the noisy activities, the maximum magnitude of impact from mortality, potential mortal injury and recoverable injury is reported to be negligible, with an overall impact conclusion of minor adverse. If it is assumed that effects on a designated site generally reduce with increasing distance from an impact source, considering the distance of Vlaamse Banken SAC to VE (34.75 km to array area), the likelihood of exposure to lethal or injurious sounds levels (i.e., limited to within <100 m of the array for mortality, mortal injury and recoverable injury for both the temporal and spatial MDS) is expected to be low and limited to sporadic, low numbers of twaite shad associated with Vlaamse Banken SAC. As such, mortalities and or recoverable injuries due to exposure to underwater noise are not expected to manifest at levels that could compromise the maintenance of the twaite shad population. There is, therefore, no potential for an AEoI.





ONSHORE ECOLOGY

HRA Integrity Matrix 29: Hamford Water SAC

Name of European site:	Hamfo	Hamford Water SAC													
EU Code:	UK0030	0377													
Distance to Project:	0 km to	onshore	ECC												
Likely Effects of Project															
Effect	Impacts on supporting populations, food plant and potential habitat outside the SAC			Water of from site affection	quality: p e run-off g habitat	ollution quality	Decrea quantity	ses in wa y	ater	Decrease in air quality			Increas		
Stage of Development	C O D			С	0	D	С	0	D	С	0	D	С		
Fisher's estuarine moth	Xa Xa		Xb		Xb	Xb		Xb	Хс		Хс	Xd			

Evidence supporting conclusions:

- Xa The effects of construction and decommissioning activities on the site are expected to be minor as Fisher's estuarine moths are limited to the areas outside of the project site and therefore their habitat is expected to remain intact and undisturbed. As highlighted, within Paragraph 11.6.54 of Volume 5, Report 4: RIAA, there would be de minimis risk (for unscheduled maintenance only, no risk for scheduled maintenance) of undermining the conservation objectives for Fisher's estuarine moth. As with the outlined mitigation in Volume 5, Report 4: RIAA, hog's fennel plants would remain in place, available for the larval stage of the moth, leading to population outside of the SAC being maintained. Such populations will support the restoration of the Fisher's estuarine moth population within the SAC via immigration. There is, therefore, no potential for AEoI.
- Xb As highlighted in Paragraph 11.6.61 of Volume 5, Report 4: RIAA onwards, and with the actions outline in the Code of Construction Practice, there will be no impact on water quality or quantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI.
- Xc As highlighted in Paragraph 11.6.60 of Volume 5, Report 4: RIAA, as construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. There is, therefore, no potential for AEoI.
- Xd Whilst the period of illuminated construction partially overlaps with the flight period during a maximum of two years, there is little possibility for it to interact with the individuals that form part of the population of Fisher's Estuarine Moth for which the SAC is designated and limited interaction with a supporting population outside the SAC and therefore would not affect the population of moths. Therefore, lighting will not undermine the conservation objectives of Hamford Water SAC when considering the Project alone.
- Xe With the low likelihood of hog's fennel/ Fisher's estuarine moth being present in the vicinity of the VE onshore ECC, due to its rarity, despite the lack of information appertaining to North Falls, the situation is likely to be similar to that of VE. There is, therefore, no potential for AEoI, in-combination.



e in light	ing	In-com	bination	
0	D	С	0	D
	Xd	Xe		Xe

HRA Integrity Matrix 30: Hamford Water SPA

Name of European site:	HAMFOR	D WATER S	SPA												
EU Code:	UK900913	31													
Distance to Project:	51.04 km	to array													
Likely Effects of Project															
Effect	Habitat loss			Disturband SPA	sturbance of birds outside the			Pollution (air quality)			Decreases in water quantity			lity: pollution	n from site at quality
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Avocet	Ха		Ха	Xb	Xb	Xb	Хс		Xc	Xd		Xd	Xd		Xd
Black-tailed godwit	Ха		Ха	Xb	Xb	Xb	Хс		Xc				Xd		Xd
Dark-bellied brent goose	Ха		Ха	Xb	Xb	Xb	Хс		Хс				Xd		Xd
Grey plover	Ха		Ха	Xb	Xb	Xb	Хс		Xc				Xd		Xd
Redshank	Ха		Ха	Xb	Xb	Xb	Хс		Хс				Xd		Xd
Ringed plover	Ха		Ха	Xb	Xb	Xb	Хс		Хс				Xd		Xd
Shelduck	Xa		Ха	Xb	Xb	Xb	Xc		Xc				Xd		Xd
Teal	Xa		Ха	Xb	Xb	Xb	Xc		Xc				Xd		Xd
Little tern	Xa		Ха	Xb	Xb	Xb	Xc		Xc				Xd		Xd

Evidence supporting conclusions:

- Xa See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Xb Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity.
- Xc As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives species with similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.



Qualifying avian interest features will not be affected by any hydrological changes and there will be no adverse effect on Hamford Water SPA. With the actions outlined in the Code of Construction Practice (Volume 9, Report 21), there will be no impact on water quality or quantity, in relation to the construction or decommissioning of the Project. There is, therefore, no Xd potential for AEoI.



HRA Integrity Matrix 31: Hamford Water RAMSAR

Name of European site:	Hamfo	rd Water R	AMSAR															
EU Code:	UK11028																	
Distance to Project:	0.72 km to array																	
Likely Effects of Project																		
Effect	Disturbance of birds outside the Ramsar			Decrease in air quality			Habitat loss			Water quality: pollution from site run-off affecting prey availability			Decreases in water quantity			Loss of foraging and roosting habitat outside the Ramsar		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Black-tailed godwit	Ха	Ха	Ха	Xb		Xb	Xc		Хс	Xd		Xd	Xd		Xd	Xe		Xe
Dark-bellied brent goose	Ха	Ха	Ха	Xb		Xb	Xc		Хс	Xd		Xd	Xd		Xd	Xe		Xe
Redshank	Ха	Ха	Ха	Xb		Xb	Xc		Хс	Xd		Xd	Xd		Xd	Xe		Xe
Ringed plover	Ха	Ха	Ха	Xb		Xb	Xc		Хс	Xd		Xd	Xd		Xd	Xe		Xe

Evidence supporting conclusions:

- Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering Ха populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity.
- Xb As construction air guality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives species with similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Хс construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Qualifying avian interest features will not be affected by any hydrological changes and there will be no adverse effect on Hamford Water Ramsar. With the actions outlined in the Code of Xd Construction Practice (Volume 9, Report 21), there will be no impact on water guality or guantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI.
- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Xe construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.



HRA Integrity Matrix 32: Stour and Orwell Estuaries SPA

Name of European site:	te: Stour and Orwell Estuaries SPA																	
EU Code:	UK9009	9121																
Distance to Project:	54.81 k	m to array	/															
Likely Effects of Project																		
Effect	Disturb: outside	ance of bi the SPA	rds	Decreas quantity	ses in wat	er	Decrea	se in air q	uality	Habitat	loss		Pollution affecting	n from site g prey ava	e run-off ailability	Loss of foraging and roosting habitat outside the SPA		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Black-tailed godwit	Ха	Ха	Ха				Xb		Xb	Хс		Xc	Xd		Xd	Xe		Xe
Dark-bellied brent goose	Ха	Ха	Ха				Xb		Xb	Xc		Xc	Xd		Xd	Xe		Xe
Dunlin	Ха	Xa	Ха				Xb		Xb	Xc		Xc	Xd		Xd	Xe		Хе
Grey plover	Ха	Ха	Ха				Xb		Xb	Xc		Xc	Xd		Xd	Xe		Хе
Knot										Хс		Xc	Xd		Xd	Xe		Хе
Pintail	Ха	Xa	Ха				Xb		Xb	Xc		Xc	Xd		Xd	Xe		Хе
Redshank	Ха	Xa	Ха				Xb		Xb	Xc		Xc	Xd		Xd	Xe		Хе
Avocet	Ха	Xa	Ха	Xd	Xd	Xd	Xb		Xb	Xc		Xc	Xd		Xd	Xe		Xe
Waterbird assemblage	Ха	Ха	Ха	Xd	Xd	Xd	Xb		Xb	Хс		Xc	Xd		Xd	Xe		Хе

Evidence supporting conclusions:

- Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering Ха populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity.
- As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Xb similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Xc construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Qualifying avian interest features will not be affected by any hydrological changes and there will be no adverse effect on the Stour and Orwell Estuaries SPA. With the actions outlined in the Xd Code of Construction Practice (Volume 9, Report 21), there will be no impact on water quality or quantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI.



Xe Habitat loss will be limited and will not undermine conservation objectives and therefore will have no adverse effects on the integrity of the designated site identified. There is, therefore, no AEol.



HRA Integrity Matrix 33: Stour and Orwell Estuaries RAMSAR

Name of European site:	Stour and Orwell Estuaries RAMSAR																	
EU Code:	UK9009	9121																
Distance to Project:	54.80 km to array																	
Likely Effects of Proje																		
Effect	Disturbance / displacement of birds outside of Ramsar					ality	Loss of foraging and roosting habitat outside the SPA			Decreas quantity.	es in wate	er	Pollutior affective	n from site prey ava	run-off ilability	Collisior	Risk	
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Black-tailed godwit	Ха	Ха	Ха	Xb		Xb	Xc			Xd		Xd	Xe		Xe		Xf	
Dark-bellied brent goose	Xg	Xg	Xg	Xb		Xb	Хс			Xd		Xd	Xe		Xe		Xf	
Dunlin	Ха	Ха	Ха	Xb		Xb	Xc			Xd		Xd	Xe		Xe		Xf	
Grey plover	Ха	Ха	Ха	Xb		Xb	Xc			Xd		Xd	Xe		Xe		Xf	
Knot	Ха	Ха	Ха				Xc			Xd		Xd	Xe		Xe		Xf	
Pintail	Xh	Xh	Xh	Xb		Xb	Xc			Xd		Xd	Xe		Xe		Xf	
Redshank	Ха	Ха	Ха	Xb		Xb	Xc			Xd		Xd	Xe		Xe		Xf	
Waterbird assemblage	Ха	Ха	Ха	Xb		Xb	Xc			Xd		Xd	Xe		Xe		Xf	
Wetland invertebrate assemblage				Xb		Xb				Xd		Xd						
Wetland plant assemblage				Xb		Xb				Xd		Xd						

Evidence supporting conclusions:

- Ха Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity.
- As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Xb similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Xc construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Xd Qualifying avian interest features will not be affected by any hydrological changes and there will be no adverse effect on the Stour and Orwell Estuaries SPA. With the actions outlined in the Code of Construction Practice (Volume 9, Report 21), there will be no impact on water guality or guantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI.
- The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey Xe populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.



- Considering the highly precautionary nature of the outputs of the MigroPath analyses, impacts to migrating birds at the scoped in SPAs and RAMSARs can be considered minimal and make Xf no material contribution to any changes in population or baseline mortality. Therefore, there is no potential for an AEoI.
- With consideration of the mitigation being implemented (timing of works/maintenance, vibro-piling technology, fencing for visual and acoustic screening, suspending works during very cold Xg periods, construction lighting at HDD locations would be at the lowest, safest permissible level and with light spill minimised and on-site measures overseen by an ECoW), the predicted potential disturbance to the species is reduced to negligible levels, and therefore there is no potential for AEoI.
- With consideration of the distance from any construction this species was recorded and the infrequency of observations, we conclude that the conservation objectives will not be undermined Xh by this effect and there is no potential for AEoI.



HRA Integrity Matrix 34: Colne Estuary (Mid-Essex Coast Phase 2) SPA

Name of European site:	Colne Estuary (Mid-Essex Coast Phase 2) SPA															
EU Code:	UK90092	243														
Distance to Project:	66.51 km	to array														
Likely Effects of Project																
Effect	Habitat Ic	DSS		Direct dis displacer	sturbance a nent	nd	Pollution	(air quality)		Decrease	es in water o	quantity	Pollution from site run-off affecting prey availability			
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	
Dark-bellied brent goose	Xa		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	
Hen harrier	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	
Pochard	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	
Redshank	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	
Ringed plover	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	
Little tern	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	
Waterbird assemblage	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe	

Evidence supporting conclusions:

- Xa See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Xb Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on inte
- populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- Xd Qualifying avian interest features will not be affected by any hydrological changes and there will be no adverse effect on the Stour and Orwell Estuaries SPA. With the actions outlined in the Code of Construction Practice (Volume 9, Report 21), there will be no impact on water quality or quantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI.
- Xe The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.



Name of European site:	Colne I	Estuary	(Mid-Esse	ex Coast	Phase 2) I	RAMSAR									
EU Code:	UK901	5022													
Distance to Project:	66.63 k	m to arra	ıy												
Likely Effects of Project															
Effect	Loss of roosting the SP/	Loss of foraging and bosting habitat outside he SPA Disturbance/displacement of birds outside of Ramsar Pollution (air quality) Invasive Non-Native Species (INNS)												on from s ecting pre bility	ite run- y
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Redshank	Ха		Ха	Xb	Xb	Xb	Хс		Xc	Xd	Xd	Xd	Xe	Xe	Xe
Dark-bellied brent goose	Ха		Xa	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe
Waterbird assemblage	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd	Xe	Xe	Xe
Wetland invertebrate assemblage										Xd	Xd	Xd			
Wetland plant assemblage										Xd	Xd	Xd			
Saltmarsh										Xd	Xd	Xd			

Evidence supporting conclusions:

- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Ха construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering Xb populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity.
- As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Xc similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- Xd Through increased vessel movements during construction and decommissioning there is a risk that vessels could contribute to the potential introduction or spread of marine INNS through ballast water discharge, however the movement of commercial vessels is common throughout the region (Volume 6, Part 2, Chapter 9: Shipping and Navigation) and this provides an existing and potentially more likely method of transport for Marine INNS (due to the higher variety of ports and passage routes). Furthermore, there is a lack of evidence of any adverse effect from other offshore wind farms within the North Sea of having any adverse effect on key species and habitats through increasing the spread of marine INNS. Additionally, project level commitments to mitigate the risk such as following best practice guidelines and standard operating practices (as managed through the PEMP and biosecurity plan) will ensure the site's conservation objectives will be maintained in the long-term. There is, therefore, no potential for an AEoI.



The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey Xe populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.



HRA Integrity Matrix 36: Abberton Reservoir SPA

Name of European site:	Abberton R	eservoir SPA										
EU Code:	UK9009141											
Distance to Project:	11.4 km to o	nshore EEC										
Likely Effects of Project				1			1					
Effect	Habitat loss			Disturbance	of birds outsid	e the SPA	Decrease in	air quality		Water quality affecting hat	y: pollution from pitat quality	n site run-off
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D
Cormorant	Xa		Ха	Xb	Xb	Xb	Хс		Xc	Xd	Xd	Xd
Coot	Xa		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd
Gadwall	Xa		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd
Great crested grebe	Xa		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd
Mute swan	Xa		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd
Pochard	Xa		Ха	Xb	Xb	Xb	Хс		Xc	Xd	Xd	Xd
Shoveler	Ха		Ха	Xb	Xb	Xb	Хс		Xc	Xd	Xd	Xd
Widgeon	Ха		Ха	Xb	Xb	Xb	Хс		Xc	Xd	Xd	Xd
Teal	Ха		Ха	Xb	Xb	Xb	Хс		Хс	Xd	Xd	Xd
Tufted Duck	Ха		Ха	Xb	Xb	Xb	Хс		Хс	Xd	Xd	Xd
Waterbird assemblage	Ха		Ха	Xb	Xb	Xb	Xc		Хс	Xd	Xd	Xd

Evidence supporting conclusions:

- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Ха construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Xb The conservation objectives would not be undermined for the identified sites in relation to important wintering populations of the designated species during construction, operation and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Хс similar thresholds. Air guality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air guality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEol.



The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey Xd populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.



HRA Integrity Matrix 37: Abberton Reservoir RAMSAR

Name of European site:	Abberton R	eservoir RAM	ISAR									
EU Code:	UK9009141											
Distance to Project:	11.4 km to E	CC										
Likely Effects of Project										1		
Effect	Habitat loss			Disturbance Ramsar	e of birds outsi	de the	Decrease in	air quality		Water quality affecting habi	: pollution from tat quality	site run-off
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D
Gadwall	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd
Shoveler	Xa		Xa	Xb	Xb	Xb	Xc		Xc	Xd	Xd	Xd
Widgeon	Ха		Ха	Xb	Xb	Xb	Хс		Хс	Xd	Xd	Xd
Waterbird assemblage	Ха		Ха	Xb	Xb	Xb	Хс		Хс	Xd	Xd	Xd

Evidence supporting conclusions:

- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Ха construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- The conservation objectives would not be undermined for the identified sites in relation to important wintering populations of the designated species during construction, operation and Xb decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Хс similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey Xd populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.



HRA Integrity Matrix 38: Blackwater Estuary SPA

Name of European site:	Blackwater Estuary (Mid-Essex Coast Phase 4) SPA														
EU Code:	UK900924	45													
Distance to Project:	77.69 km	to array													
Likely Effects of Proj	ect									1					
Effect	Habitat los	SS		Disturband birds outsi	ce / displace de SPA	ement of	Pollution (air quality)		Water qua run-off affe	lity: pollutio	n from site at quality	Decreases in water quantity		
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Black-tailed godwit	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Dark-bellied Brent goose	Ха		Xa	Xb	Xb	Xb	Хс		Xc	Xd		Xd	Xe		Xe
Dunlin	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Grey plover	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Hen harrier	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Waterbird assemblage	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Little tern	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Pochard	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe
Ringed plover	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe

Evidence supporting conclusions:

- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Ха construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering Xb populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity
- As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Xc similar thresholds. Air quality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air quality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey Xd populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.
- There will be no impact on water quality or quantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI. Xe End of Matrix 38



HRA Integrity Matrix 39: Blackwater Estuary RAMSAR

Name of European site:	Blackwater Estuary (Mid-Essex Coast Phase 4) RAMSAR																	
EU Code:	UK1100	UK11007																
Distance to Project:	77.81 ki	m to array	/															
Likely Effects of Pro	f Project																	
Effect	Habitat	loss	_	Disturbance / displacement of birds outside SPA			Decreas	Decreases in air quality			uality: pol e run-off a quality	lution ffecting	Decreas quantity	ses in wate	er	Impacts populati inverteb Ramsar	on suppo ons of pla prates out	orting ants and side the
Stage of Development	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D	С	0	D
Black-tailed godwit	Ха		Ха	Xb	Xb	Xb	Xc		Xc	Xd		Xd	Xe		Xe	Xf	Xf	Xf
Dark-bellied brent goose	Ха		Ха	Xb	Xb	Xb	Хс		Хс	Xd		Xd	Xe		Xe	Xf	Xf	Xf
Dunlin	Ха		Ха	Xb	Xb	Xb	Хс		Xc	Xd		Xd	Xe		Xe	Xf	Xf	Xf
Grey plover	Ха		Ха	Xb	Xb	Xb	Хс		Xc	Xd		Xd	Xe		Xe	Xf	Xf	Xf
Waterbird assemblage	Ха		Ха	Xb	Xb	Xb	Хс		Хс	Xd		Xd	Xe		Xe	Xf	Xf	Xf

Evidence supporting conclusions:

- See paragraph 11.6.76 onwards of Volume 5, Report 4: RIAA which highlights the potential impact upon avian features. The majority of habitat loss will be temporary, only during Ха construction. With permanent habitat loss limited to only the footprint of TJBs, junction boxes and the OnSS. There will be no permanent intertidal habitat loss. Any permanent habitat loss will be minimal (refer to Volume 6, Part 3, Chapter 1: Onshore Project description) and could not undermine the conservation objectives. There is, therefore, no AEoI.
- Disturbance and displacement of a predicted small number of individuals will not result in the conservation objectives of the site being undermined in relation to the important wintering Xb populations of the designated species during construction, operation, and decommissioning for the Project alone or in combination. Therefore, there would be no adverse effect on integrity As construction air quality level changes were below threshold, maintenance levels will be considerably below threshold and will not undermine the conservation objectives for species with Xc similar thresholds. Air guality impacts during operation will not have an adverse effect on the relevant designated sites, in relation to air guality during operation for VE alone. Overall, the conservation objectives would not be undermined by any changes in air quality associated with the Project either alone or in combination. Therefore, there would be no adverse effect on integrity and no potential for AEoI.
- Xd The impact of pollution from site run-off on prey availability will be minimal and will not significantly affect the ecological balance of the area. The conservation objectives related to prey populations and their availability for the designated species will remain intact throughout the project's lifecycle, including construction, O&M, and decommissioning phases. Therefore, there will be no adverse effect on the integrity of the ecosystem, ensuring the continued sustainability of the site and its ecological functions.
- There will be no impact on water quality or quantity, in relation to the construction or decommissioning of the Project. There is, therefore, no potential for AEoI. Xe
- Xf The effects of construction and decommissioning activities on the site are expected to be minor as Fisher's estuarine moths are limited to the areas outside of the project site and therefore their habitat is expected to remain intact and undisturbed. There is, therefore, no potential for AEoI.





3 **REFERENCES**

- Bradbury, G., Trinder, M., Furness, B., Banks, A.N., Caldow, R.W. and Hume, D., 2014. Mapping seabird sensitivity to offshore wind farms. PloS one, 9(9).
- Cleasby, I.R., Owen, E., Wilson, L., Wakefield, E.D., O'Connell, P. and Bolton, M., 2020. Identifying important at-sea areas for seabirds using species distribution models and hotspot mapping. Biological Conservation, 241, p.108375.
- Dierschke, V., Furness, R.W. and Garthe, S., 2016. Seabirds and offshore wind farms in European waters: Avoidance and attraction. Biological Conservation, 202, pp.59-68.
- Ellis, J.R., Milligan, S.P. Readdy, L. Taylor, N. and Brown, M.J. (2012), 'Spawning and nursery grounds of selected fish species in UK waters'. Cefas Scientific Series Technical Report 147.
- Fliessbach, K.L., Borkenhagen, K., Guse, N., Markones, N., Schwemmer, P. and Garthe, S., 2019. A ship traffic disturbance vulnerability index for Northwest European seabirds as a tool for marine spatial planning. Frontiers in Marine Science.
- Masden. E.A., Haydon, D.T., Fox A.D., Furmess, R.W. 2010. Barriers to movement: Modelling energetic costs of avoiding marine wind farms amongst breeding seabirds. Marine Pollution Bulletin, 60(7) pp.1085-1091.
- Wildfowl and Wetland Trust (WWT). 2009. Distributions of Cetaceans, Seals, Turtles, Sharks and Ocean Sunfish recorded from Aerial Surveys 2001-2008. WWT Consulting. Report to Department of Energy and Climate Change
- Wright, L. J., Ross-Smith, V. H., Austin, G. E., Massimino, D., Dadam, D., Cook, A. S. C. P., Calbrade, N. A. and Burton, N. H. K. (2012), 'Assessing the risk of offshore wind farm development to migratory birds designated as features of UK Special Protection Areas (and other Annex 1 species)', Strategic Ornithological Support Services, Project SOSS-05, BTO Research Report No. 592.
- Woodward, I., Thaxter, C. B., Owen, E., Cook, A. S. C. P. 2019. Desk-based revision of seabird foraging ranges used for HRA screening. BTO Research Report No. 724.
 ISBN 978-1-912642-12-0Zeale, M., 2009. Barbastelles in the landscape: ecological research and conservation in Dartmoor National Park. SITA Trust.



0333 880 5306 fiveestuaries@rwe.com www.fiveestuaries.co.uk

Five Estuaries Offshore Wind Farm Ltd Windmill Hill Business Park Whitehill Way, Swindon, SN5 6PB Registered in England and Wales company number 12292474