

ENVIRONMENTAL STATEMENT -VOLUME 3 - APPENDIX 6.5 (CLEAN)

Operation Phase Air Quality Assessment Results Tables: Ecological Receptors

Drax Bioenergy with Carbon Capture and Storage

The Infrastructure Planning (Applications: Prescribed Forms and Procedure) Regulations 2009 - 5(2)(a)

Document Reference Number: 6.3.6.5

Applicant: Drax Power Limited

PINS Reference: EN010120



REVISION:03 DATE: February 2023 DOCUMENT OWNER: WSP UK Limited AUTHOR: B. Tuckett-Jones APPROVER: N. Ashworth PUBLIC

TABLE OF CONTENTS

1.	OPEF	RATION PHASE AIR QUALITY ASSESSMENT RESULTS TABLES: ECOLOGICAL	
REC	CEPTO	DRS	1
	1.1.	Impacts on Ecological Receptors	1

TABLES

Table 1.1 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NOx
Table 1.2 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NOx
Table 1.3 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH ₃ 4
Table 1.4 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanSO2
Table 1.5 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual NitrogenDeposition Rate6
Table 1.6 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual AcidDeposition Rate8
Table 1.7 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanNOx (Including Mitigation)
Table 1.8 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NOx(Including Mitigation)
Table 1.9 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH ₃ (Including Mitigation)12
Table 1.10 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanSO2 (Including Mitigation)
Table 1.11 - Modelled Maximum Operational Impacts at Ecological Receptors – AnnualNitrogen Deposition Rate (Including Mitigation)15
Table 1.12 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual AcidDeposition Rate (Including Mitigation)16
Table 1.13 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean NOx 19
Table 1.14 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Daily Mean NOx

Table 1.15 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual MeanNH321
Table 1.16 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean SO2
Table 1.17 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual NitrogenDeposition Rate
Table 1.18 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual AcidDeposition Rate
Table 1.19 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanNOx (Worst Case Emissions Profile)27
Table 1.20 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily MeanNOx (Worst Case Emissions Profile)28
Table 1.21 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanNH3 (Worst Case Emissions Profile)29
Table 1.22 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanSO2 (Worst Case Emissions Profile)
Table 1.23 - Modelled Maximum Operational Impacts at Ecological Receptors – AnnualNitrogen Deposition Rate (Worst Case Emissions Profile)
Table 1.24 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual AcidDeposition Rate (Worst Case Emissions Profile)

1. OPERATION PHASE AIR QUALITY ASSESSMENT RESULTS TABLES: ECOLOGICAL RECEPTORS

1.1. IMPACTS ON ECOLOGICAL RECEPTORS

- 1.1.1. The results of the modelling assessment at each modelled ecological receptor are presented in the below tables for each relevant pollutant and averaging period applicable to the study. The maximum modelled concentration and deposition values are presented, which is based on modelling over all five years of meteorological data (2016-2020). The change in PC and PEC, as a percentage of the relevant critical level / load, is presented for each receptor.
- 1.1.2. The PC impact in the with Proposed Scheme scenario represents the change in concentration / deposition between the Baseline scenario and Proposed Scheme scenario. Where indicated, the tables have all mitigation applied as set out in Chapter 6 (Air Quality) (APP-042) as updated by Air Quality Technical Note 2 (document reference 8.9.5).
- 1.1.3. In all results tables presented in this appendix, the following designated site names are shortened to 'Thorne Moor SPA' and Thorne Moor SSSI', respectively:
 - a. Thorne & Hatfield Moors SPA
 - **b.** Thorne, Crowle and Goole Moors SSSI.

CORE MODEL SCENARIOS

1.1.4. Results pertaining to the core model scenarios are presented in **Tables 1.1 to 1.6**.

Table 1.1 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO_x

		Annual Mean NO _x concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
River Derwent SAC	30	11.91	0.163	0.5%	12.13	40.4%			
Thorne Moor SAC/SPA/SSSI	30	13.21	0.076	0.3%	13.32	44.4%			
Lower Derwent SAC	30	9.92	0.166	0.6%	10.15	33.8%			
Lower Derwent SPA	30	9.92	0.166	0.6%	10.15	33.8%			
Skipwith Common SAC	30	9.76	0.058	0.2%	9.84	32.8%			
Skipwith Common SSSI	30	9.76	0.058	0.2%	9.84	32.8%			
Humber Estuary SAC	30	46.96	0.147	0.5%	47.16	157.2%			
Humber Estuary SPA/SSSI	30	46.96	0.147	0.5%	47.16	157.2%			
Breighton Meadows SSSI	30	9.92	0.166	0.6%	10.15	33.8%			
Eskamhorn Meadows SSSI	30	11.35	0.046	0.2%	11.40	38.0%			
Derwent Ings SSSI	30	9.80	0.134	0.4%	9.99	33.3%			
Went Ings SSSI	30	12.09	0.052	0.2%	12.16	40.5%			
Barn Hill Meadows SSSI	30	12.89	0.152	0.5%	13.09	43.6%			
Burr Closes SSSI	30	10.53	0.062	0.2%	10.61	35.4%			
Common Plantation SINC	30	11.43	0.017	0.1%	11.45	38.2%			
Disused Railway Embankment SINC	30	10.76	0.040	0.1%	10.81	36.0%			
Barmby-on-the-Marsh LWS	30	10.48	0.076	0.3%	10.57	35.2%			
Brockholes SINC	30	11.22	0.019	0.1%	11.24	37.5%			
Meadow East of Orchard Farm SINC	30	10.83	0.009	0.0%	10.84	36.1%			
Barmby Pond LWS	30	9.96	0.127	0.4%	10.12	33.7%			
Cobble Croft Wood SINC	30	11.62	0.027	0.1%	11.65	38.8%			
Hagg Green Lane SINC	30	10.93	0.103	0.3%	11.07	36.9%			

	Annual Mean NO _x concentration (µg/m ³)						
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL	
Sand Pitt Wood & Barffs Close Plantation SINC	30	11.43	0.028	0.1%	11.46	38.2%	
	1%		70%				

Table 1.2 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO_x

	Daily Mean NO _x concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	75	23.82	1.787	2.4%	27.96	37.3%		
Thorne Moor SAC/SPA/SSSI	75	26.42	0.893	1.2%	29.27	39.0%		
Lower Derwent SAC	75	19.84	0.893	1.2%	23.75	31.7%		
Lower Derwent SPA	75	19.84	0.893	1.2%	23.75	31.7%		
Skipwith Common SAC	75	19.52	0.646	0.9%	21.71	28.9%		
Skipwith Common SSSI	75	19.52	0.646	0.9%	21.71	28.9%		
Humber Estuary SAC	75	93.92	1.200	1.6%	96.93	129.2%		
Humber Estuary SPA/SSSI	75	93.92	1.200	1.6%	96.93	129.2%		
Breighton Meadows SSSI	75	19.84	0.893	1.2%	23.75	31.7%		
Eskamhorn Meadows SSSI	75	22.70	1.912	2.5%	25.59	34.1%		
Derwent Ings SSSI	75	19.60	0.831	1.1%	23.37	31.2%		
Went Ings SSSI	75	24.18	0.971	1.3%	26.60	35.5%		
Barn Hill Meadows SSSI	75	25.78	0.962	1.3%	28.30	37.7%		
Burr Closes SSSI	75	21.06	0.761	1.0%	22.99	30.7%		
Common Plantation SINC	75	22.86	1.038	1.4%	24.19	32.3%		
Disused Railway Embankment SINC	75	21.52	1.184	1.6%	23.07	30.8%		
Barmby-on-the-Marsh LWS	75	20.96	1.621	2.2%	23.44	31.3%		

Drax Bioenergy with Carbon Capture and Storage

	Daily Mean NO _x concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Brockholes SINC	75	22.44	1.623	2.2%	24.44	32.6%		
Meadow East of Orchard Farm SINC	75	21.66	0.554	0.7%	22.33	29.8%		
Barmby Pond LWS	75	19.92	1.577	2.1%	23.29	31.1%		
Cobble Croft Wood SINC	75	23.24	1.176	1.6%	25.28	33.7%		
Hagg Green Lane SINC	75	21.86	1.248	1.7%	25.22	33.6%		
Sand Pitt Wood & Barffs Close Plantation SINC	75	22.86	1.630	2.2%	24.97	33.3%		
	10%							

Table 1.3 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH₃

	Annual Mean NH ₃ concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	3	4.57	0.007	0.2%	4.58	152.7%		
Thorne Moor SAC/SPA/SSSI	1	2.59	0.003	0.3%	2.60	259.5%		
Lower Derwent SAC	3	4.57	0.007	0.2%	4.58	152.7%		
Lower Derwent SPA	3	4.57	0.007	0.2%	4.58	152.7%		
Skipwith Common SAC	1	2.58	0.002	0.2%	2.58	258.4%		
Skipwith Common SSSI	1	2.58	0.002	0.2%	2.58	258.4%		
Humber Estuary SAC	3	3.58	0.004	0.1%	3.59	119.5%		
Humber Estuary SPA/SSSI	3	3.58	0.004	0.1%	3.59	119.5%		
Breighton Meadows SSSI	3	3.08	0.007	0.2%	3.09	103.0%		
Eskamhorn Meadows SSSI	3	2.40	0.002	0.1%	2.40	80.1%		
Derwent Ings SSSI	3	4.57	0.005	0.2%	4.58	152.6%		
Went Ings SSSI	3	2.35	0.002	0.1%	2.35	78.4%		

Drax Bioenergy with Carbon Capture and Storage

	Annual Mean NH ₃ concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Barn Hill Meadows SSSI	3	2.32	0.005	0.2%	2.33	77.6%		
Burr Closes SSSI	3	2.50	0.003	0.1%	2.50	83.5%		
Common Plantation SINC	3	2.33	0.001	0.0%	2.33	77.7%		
Disused Railway Embankment SINC	1	2.28	0.002	0.1%	2.28	76.1%		
Barmby-on-the-Marsh LWS	3	2.28	0.003	0.1%	2.28	76.1%		
Brockholes SINC	3	2.28	0.001	0.0%	2.28	76.0%		
Meadow East of Orchard Farm SINC	1	2.33	0.000	0.0%	2.33	77.7%		
Barmby Pond LWS	1	2.28	0.006	0.2%	2.29	76.2%		
Cobble Croft Wood SINC	3	2.33	0.001	0.0%	2.33	77.7%		
Hagg Green Lane SINC	3	3.09	0.004	0.1%	3.10	103.2%		
Sand Pitt Wood & Barffs Close Plantation SINC	3	2.33	0.001	0.0%	2.33	77.7%		
	Criterion (as % of CL)	1%		70%				

Table 1.4 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO2

	Annual Mean SO ₂ concentration (μg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	20	3.93	0.072	0.4%	4.03	20.2%		
Thorne Moor SAC/SPA/SSSI	20	1.34	0.033	0.2%	1.39	7.0%		
Lower Derwent SAC	20	1.70	0.073	0.4%	1.81	9.0%		
Lower Derwent SPA	20	1.70	0.073	0.4%	1.81	9.0%		
Skipwith Common SAC	20	1.42	0.025	0.1%	1.46	7.3%		
Skipwith Common SSSI	20	1.42	0.025	0.1%	1.46	7.3%		
Humber Estuary SAC	20	7.49	0.069	0.3%	7.59	38.0%		

Drax Bioenergy with Carbon Capture and Storage

			Annual Mean SC	02 concentration (μg/m ³)		
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
Humber Estuary SPA/SSSI	20	7.49	0.069	0.3%	7.59	38.0%
Breighton Meadows SSSI	20	1.70	0.073	0.4%	1.81	9.0%
Eskamhorn Meadows SSSI	20	1.29	0.021	0.1%	1.32	6.6%
Derwent Ings SSSI	20	1.69	0.059	0.3%	1.78	8.9%
Went Ings SSSI	20	1.31	0.023	0.1%	1.34	6.7%
Barn Hill Meadows SSSI	20	1.81	0.072	0.4%	1.91	9.5%
Burr Closes SSSI	20	1.23	0.027	0.1%	1.27	6.3%
Common Plantation SINC	20	1.44	0.008	0.0%	1.45	7.2%
Disused Railway Embankment SINC	20	1.32	0.019	0.1%	1.34	6.7%
Barmby-on-the-Marsh LWS	20	1.32	0.036	0.2%	1.36	6.8%
Brockholes SINC	20	1.32	0.009	0.0%	1.33	6.7%
Meadow East of Orchard Farm SINC	20	1.44	0.004	0.0%	1.44	7.2%
Barmby Pond LWS	20	1.32	0.058	0.3%	1.40	7.0%
Cobble Croft Wood SINC	20	1.44	0.013	0.1%	1.46	7.3%
Hagg Green Lane SINC	20	1.43	0.047	0.2%	1.50	7.5%
Sand Pitt Wood & Barffs Close Plantation SINC	20	1.44	0.013	0.1%	1.46	7.3%
		Env. Agency Screening	Criterion (as % of CL)	1%		70%

Table 1.5 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)									
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL				
River Derwent SAC	15	30.22	0.054	0.4%	30.30	202.0%				
Thorne Moor SAC	5	21.31	0.025	0.5%	21.35	426.9%				

Drax Bioenergy with Carbon Capture and Storage

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Thorne Moor SPA	10	21.31	0.025	0.2%	21.35	213.5%			
Thorne Moor SSSI	5	21.31	0.025	0.5%	21.35	426.9%			
Lower Derwent SAC	20	30.22	0.055	0.3%	30.30	151.5%			
Lower Derwent SPA	20	30.22	0.055	0.3%	30.30	151.5%			
Skipwith Common SAC	10	21.12	0.019	0.2%	21.14	211.4%			
Skipwith Common SSSI	10	21.12	0.019	0.2%	21.14	211.4%			
Humber Estuary SAC	20	28.87	0.034	0.2%	28.92	144.6%			
Humber Estuary SPA/SSSI	20	28.87	0.034	0.2%	28.92	144.6%			
Breighton Meadows SSSI	20	23.51	0.055	0.3%	23.58	117.9%			
Eskamhorn Meadows SSSI	10	19.95	0.016	0.2%	19.97	199.7%			
Derwent Ings SSSI	20	30.22	0.043	0.2%	30.29	151.5%			
Went Ings SSSI	15	19.38	0.017	0.1%	19.41	129.4%			
Barn Hill Meadows SSSI	20	20.43	0.039	0.2%	20.48	102.4%			
Burr Closes SSSI	20	20.64	0.020	0.1%	20.67	103.3%			
Common Plantation SINC	10	33.74	0.010	0.1%	33.75	337.5%			
Disused Railway Embankment SINC	10	33.32	0.022	0.2%	33.35	333.5%			
Barmby-on-the-Marsh LWS	10	33.32	0.043	0.4%	33.37	333.7%			
Brockholes SINC	10	19.74	0.007	0.1%	19.75	197.5%			
Meadow East of Orchard Farm SINC	20	19.88	0.003	0.0%	19.88	99.4%			
Barmby Pond LWS	10	19.74	0.044	0.4%	19.80	198.0%			

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Cobble Croft Wood SINC	10	33.74	0.015	0.2%	33.76	337.6%		
Hagg Green Lane SINC	10	40.74	0.058	0.6%	40.82	408.2%		
Sand Pitt Wood & Barffs Close Plantation SINC	10	33.74	0.016	0.2%	33.76	337.6%		
Barlow Common LNR	10	33.74	0.010	0.1%	33.75	337.5%		
Env. Agency Screening C	riterion (as % of CL)	·	1%		70%			

Table 1.6 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate

	Annual Acid Deposition Rate (keq/ha/yr)							
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Thorne Moor SAC	0.462	1.73	0.006	1.3%	1.74	376.9%		
Thorne Moor SSSI	0.462	1.73	0.006	1.3%	1.74	376.9%		
Lower Derwent SAC	0.643	2.40	0.013	2.1%	2.42	376.9%		
Skipwith Common SAC	0.802	1.73	0.005	0.6%	1.73	216.0%		
Skipwith Common SSSI	0.802	1.73	0.005	0.6%	1.73	216.0%		
Breighton Meadows SSSI	0.643	1.92	0.013	2.1%	1.94	302.2%		
Eskamhorn Meadows SSSI	1.998	1.64	0.004	0.2%	1.64	82.2%		
Derwent Ings SSSI	0.643	2.40	0.010	1.6%	2.42	376.5%		
Went Ings SSSI	2.008	1.59	0.004	0.2%	1.60	79.6%		
Barn Hill Meadows SSSI	0.633	1.69	0.010	1.6%	1.70	269.3%		
Burr Closes SSSI	1.248	1.68	0.005	0.4%	1.69	135.2%		
Env. Agency Screening Criterion (as % of CL)				1%		70%		

WITH PROPOSED SCHEME MITIGATION MODEL SCENARIOS

1.1.5. Results pertaining to the core model scenarios, including the With Proposed Scheme Mitigation (as detailed in Section 6.10 of Chapter 6 (Air Quality) (APP-042) and updated in Air Quality Technical Note 2 (document reference 8.9.5) are presented in Tables 1.7 to 1.12.

	Annual Mean NO _x concentration (µg/m ³)							
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation			
River Derwent SAC	30	0.163	0.133	0.5%	0.4%			
Thorne Moor SAC/SPA/SSSI	30	0.076	0.064	0.3%	0.2%			
Lower Derwent SAC	30	0.166	0.138	0.6%	0.5%			
Lower Derwent SPA	30	0.166	0.138	0.6%	0.5%			
Skipwith Common SAC	30	0.058	0.049	0.2%	0.2%			
Skipwith Common SSSI	30	0.058	0.049	0.2%	0.2%			
Humber Estuary SAC	30	0.147	0.125	0.5%	0.4%			
Humber Estuary SPA/SSSI	30	0.147	0.125	0.5%	0.4%			
Breighton Meadows SSSI	30	0.166	0.138	0.6%	0.5%			
Eskamhorn Meadows SSSI	30	0.046	0.034	0.2%	0.1%			
Derwent Ings SSSI	30	0.134	0.114	0.4%	0.4%			
Went Ings SSSI	30	0.052	0.042	0.2%	0.1%			
Barn Hill Meadows SSSI	30	0.152	0.124	0.5%	0.4%			
Burr Closes SSSI	30	0.062	0.050	0.2%	0.2%			
Common Plantation SINC	30	0.017	0.011	0.1%	0.0%			
Disused Railway Embankment SINC	30	0.040	0.029	0.1%	0.1%			
Barmby-on-the-Marsh LWS	30	0.076	0.057	0.3%	0.2%			
Brockholes SINC	30	0.019	0.013	0.1%	0.0%			
Meadow East of Orchard Farm SINC	30	0.009	0.006	0.0%	0.0%			
Barmby Pond LWS	30	0.127	0.098	0.4%	0.3%			
Cobble Croft Wood SINC	30	0.027	0.020	0.1%	0.1%			
Hagg Green Lane SINC	30	0.103	0.083	0.3%	0.3%			

	Annual Mean NO _x concentration (µg/m ³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Sand Pitt Wood & Barffs Close Plantation SINC	30	0.028	0.020	0.1%	0.1%		
Env. Age		1%					

Table 1.8 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO_x (Including Mitigation)

	Daily Mean NO _x concentration (µg/m ³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	75	1.787	1.321	2.4%	1.8%		
Thorne Moor SAC/SPA/SSSI	75	0.893	0.699	1.2%	0.9%		
Lower Derwent SAC	75	0.893	0.673	1.2%	0.9%		
Lower Derwent SPA	75	0.893	0.673	1.2%	0.9%		
Skipwith Common SAC	75	0.646	0.434	0.9%	0.6%		
Skipwith Common SSSI	75	0.646	0.434	0.9%	0.6%		
Humber Estuary SAC	75	1.200	0.872	1.6%	1.2%		
Humber Estuary SPA/SSSI	75	1.200	0.872	1.6%	1.2%		
Breighton Meadows SSSI	75	0.893	0.673	1.2%	0.9%		
Eskamhorn Meadows SSSI	75	1.912	1.438	2.5%	1.9%		
Derwent Ings SSSI	75	0.831	0.670	1.1%	0.9%		
Went Ings SSSI	75	0.971	0.651	1.3%	0.9%		
Barn Hill Meadows SSSI	75	0.962	1.274	1.3%	1.7%		
Burr Closes SSSI	75	0.761	0.561	1.0%	0.7%		
Common Plantation SINC	75	1.038	0.684	1.4%	0.9%		
Disused Railway Embankment SINC	75	1.184	1.069	1.6%	1.4%		
Barmby-on-the-Marsh LWS	75	1.621	1.420	2.2%	1.9%		

Drax Bioenergy with Carbon Capture and Storage

	Daily Mean NO _x concentration (µg/m ³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Brockholes SINC	75	1.623	1.170	2.2%	1.6%		
Meadow East of Orchard Farm SINC	75	0.554	0.385	0.7%	0.5%		
Barmby Pond LWS	75	1.577	1.151	2.1%	1.5%		
Cobble Croft Wood SINC	75	1.176	0.767	1.6%	1.0%		
Hagg Green Lane SINC	75	1.248	0.832	1.7%	1.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	75	1.630	1.083	2.2%	1.4%		
		0%					

Table 1.9 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH₃ (Including Mitigation)

	Annual Mean NH ₃ concentration (µg/m ³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	3	0.007	0.006	0.2%	0.2%		
Thorne Moor SAC/SPA/SSSI	1	0.003	0.003	0.3%	0.3%		
Lower Derwent SAC	3	0.007	0.006	0.2%	0.2%		
Lower Derwent SPA	3	0.007	0.006	0.2%	0.2%		
Skipwith Common SAC	1	0.002	0.002	0.2%	0.2%		
Skipwith Common SSSI	1	0.002	0.002	0.2%	0.2%		
Humber Estuary SAC	3	0.004	0.005	0.1%	0.2%		
Humber Estuary SPA/SSSI	3	0.004	0.005	0.1%	0.2%		
Breighton Meadows SSSI	3	0.007	0.006	0.2%	0.2%		
Eskamhorn Meadows SSSI	3	0.002	0.002	0.1%	0.1%		
Derwent Ings SSSI	3	0.005	0.005	0.2%	0.2%		
Went Ings SSSI	3	0.002	0.002	0.1%	0.1%		

Drax Bioenergy with Carbon Capture and Storage

	Annual Mean NH ₃ concentration (μg/m ³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Barn Hill Meadows SSSI	3	0.005	0.005	0.2%	0.2%		
Burr Closes SSSI	3	0.003	0.002	0.1%	0.1%		
Common Plantation SINC	3	0.001	0.001	0.0%	0.0%		
Disused Railway Embankment SINC	1	0.002	0.001	0.1%	0.0%		
Barmby-on-the-Marsh LWS	3	0.003	0.003	0.1%	0.1%		
Brockholes SINC	3	0.001	0.001	0.0%	0.0%		
Meadow East of Orchard Farm SINC	1	0.000	0.000	0.0%	0.0%		
Barmby Pond LWS	1	0.006	0.004	0.2%	0.1%		
Cobble Croft Wood SINC	3	0.001	0.001	0.0%	0.0%		
Hagg Green Lane SINC	3	0.004	0.004	0.1%	0.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	3	0.001	0.001	0.0%	0.0%		
		1%					

Table 1.10 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO₂ (Including Mitigation)

	Annual Mean SO ₂ concentration (μg/m ³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	20	0.072	0.021	0.4%	0.1%		
Thorne Moor SAC/SPA/SSSI	20	0.033	0.008	0.2%	0.0%		
Lower Derwent SAC	20	0.073	0.020	0.4%	0.1%		
Lower Derwent SPA	20	0.073	0.020	0.4%	0.1%		
Skipwith Common SAC	20	0.025	0.007	0.1%	0.0%		
Skipwith Common SSSI	20	0.025	0.007	0.1%	0.0%		
Humber Estuary SAC	20	0.069	0.019	0.3%	0.1%		

Drax Bioenergy with Carbon Capture and Storage

	Annual Mean SO ₂ concentration (µg/m ³)							
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation			
Humber Estuary SPA/SSSI	20	0.069	0.019	0.3%	0.1%			
Breighton Meadows SSSI	20	0.073	0.020	0.4%	0.1%			
Eskamhorn Meadows SSSI	20	0.021	0.006	0.1%	0.0%			
Derwent Ings SSSI	20	0.059	0.015	0.3%	0.1%			
Went Ings SSSI	20	0.023	0.007	0.1%	0.0%			
Barn Hill Meadows SSSI	20	0.072	0.020	0.4%	0.1%			
Burr Closes SSSI	20	0.027	0.008	0.1%	0.0%			
Common Plantation SINC	20	0.008	0.002	0.0%	0.0%			
Disused Railway Embankment SINC	20	0.019	0.005	0.1%	0.0%			
Barmby-on-the-Marsh LWS	20	0.036	0.011	0.2%	0.1%			
Brockholes SINC	20	0.009	0.003	0.0%	0.0%			
Meadow East of Orchard Farm SINC	20	0.004	0.001	0.0%	0.0%			
Barmby Pond LWS	20	0.058	0.017	0.3%	0.1%			
Cobble Croft Wood SINC	20	0.013	0.004	0.1%	0.0%			
Hagg Green Lane SINC	20	0.047	0.012	0.2%	0.1%			
Sand Pitt Wood & Barffs Close Plantation SINC	20	0.013	0.004	0.1%	0.0%			
		Env. Agency	Screening Criterion (as % of CL)					

Table 1.11 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Including Mitigation)

			Annual Nitrogen Deposition Rate (kg	N/ha/yr)	
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
River Derwent SAC	15	0.054	0.044	0.4%	0.3%
Thorne Moor SAC	5	0.025	0.021	0.5%	0.4%
Thorne Moor SPA	10	0.025	0.021	0.2%	0.2%
Thorne Moor SSSI	5	0.025	0.021	0.5%	0.4%
Lower Derwent SAC	20	0.055	0.045	0.3%	0.2%
Lower Derwent SPA	20	0.055	0.045	0.3%	0.2%
Skipwith Common SAC	10	0.019	0.016	0.2%	0.2%
Skipwith Common SSSI	10	0.019	0.016	0.2%	0.2%
Humber Estuary SAC	20	0.034	0.041	0.2%	0.2%
Humber Estuary SPA/SSSI	20	0.034	0.041	0.2%	0.2%
Breighton Meadows SSSI	20	0.055	0.045	0.3%	0.2%
Eskamhorn Meadows SSSI	10	0.016	0.012	0.2%	0.1%
Derwent Ings SSSI	20	0.043	0.037	0.2%	0.2%
Went Ings SSSI	15	0.017	0.014	0.1%	0.1%
Barn Hill Meadows SSSI	20	0.039	0.041	0.2%	0.2%
Burr Closes SSSI	20	0.020	0.016	0.1%	0.1%
Common Plantation SINC	10	0.010	0.007	0.1%	0.1%
Disused Railway Embankment SINC	10	0.022	0.016	0.2%	0.2%
Barmby-on-the-Marsh LWS	10	0.043	0.032	0.4%	0.3%
Brockholes SINC	10	0.007	0.005	0.1%	0.0%

Drax Bioenergy with Carbon Capture and Storage

			Annual Nitrogen Deposition Rate (kg	N/ha/yr)	
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
Meadow East of Orchard Farm SINC	20	0.003	0.002	0.0%	0.0%
Barmby Pond LWS	10	0.044	0.033	0.4%	0.3%
Cobble Croft Wood SINC	10	0.015	0.011	0.2%	0.1%
Hagg Green Lane SINC	10	0.058	0.046	0.6%	0.5%
Sand Pitt Wood & Barffs Close Plantation SINC	10	0.016	0.011	0.2%	0.1%
Barlow Common LNR	10	0.010	0.007	0.1%	0.1%
			Env. Agency Screening Criterion (as % of CL)		1%

Table 1.12 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Including Mitigation)

			Annual Acid Deposition Rate	(keq/ha/yr)	
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
Thorne Moor SAC	0.462	0.006	0.003	1.3%	0.6%
Thorne Moor SSSI	0.462	0.006	0.003	1.3%	0.6%
Lower Derwent SAC	0.643	0.013	0.006	2.1%	1.0%
Skipwith Common SAC	0.802	0.005	0.002	0.6%	0.3%
Skipwith Common SSSI	0.802	0.005	0.002	0.6%	0.3%
Breighton Meadows SSSI	0.643	0.013	0.006	2.1%	1.0%
Eskamhorn Meadows SSSI	1.998	0.004	0.002	0.2%	0.1%
Derwent Ings SSSI	0.643	0.010	0.005	1.6%	0.8%
Went Ings SSSI	2.008	0.004	0.002	0.2%	0.1%
Barn Hill Meadows SSSI	0.633	0.010	0.006	1.6%	0.9%

Drax Bioenergy with Carbon Capture and Storage

Environmental Statement - Volume 3 - Appendix 6.5

Page 16 of 34

		Annual Acid Deposition Rate (keq/ha/yr)								
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation					
Burr Closes SSSI	1.248	0.005	0.002	0.4%	0.2%					
		Env. Agency S	Screening Criterion (as % of CL)		1%					

CUMULATIVE IMPACTS (WITH PROPOSED SCHEME & OTHER PROJECTS)

1.1.6. Results pertaining to the cumulative impacts, both before mitigation ("No Mitⁿ") and after mitigation ("Mitigⁿ") associated With Proposed Scheme Mitigation is applied (as detailed in Section 6.10 of Chapter 6 (Air Quality) and updated in Air Quality Technical Note 2 (document reference 8.9.5) are presented in Tables 1.13 to 1.18.

				Annual Mean N	NO _x concentra	tion (µg/m³)			
Receptor	Critical Level	Max Cumulat	tive PC Impact	Max PC Impac	ct as % of CL	Max Cumu	lative PEC	Max PEC a	s % of CL
	Chucai Levei	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ
River Derwent SAC	30	0.675	0.647	2.3%	2.2%	12.63	12.60	42.1%	42.0%
Thorne Moor SAC/SPA/SSSI	30	0.348	0.336	1.2%	1.1%	13.58	13.57	45.3%	45.2%
Lower Derwent SAC	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%
Lower Derwent SPA	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%
Skipwith Common SAC	30	0.504	0.495	1.7%	1.7%	10.29	10.28	34.3%	34.3%
Skipwith Common SSSI	30	0.504	0.495	1.7%	1.7%	10.29	10.28	34.3%	34.3%
Humber Estuary SAC	30	0.498	0.474	1.7%	1.6%	47.51	47.48	158.4%	158.3%
Humber Estuary SPA/SSSI	30	0.498	0.474	1.7%	1.6%	47.51	47.48	158.4%	158.3%
Breighton Meadows SSSI	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%
Eskamhorn Meadows SSSI	30	0.488	0.483	1.6%	1.6%	11.84	11.83	39.5%	39.4%
Derwent Ings SSSI	30	0.640	0.619	2.1%	2.1%	10.49	10.47	35.0%	34.9%
Went Ings SSSI	30	0.303	0.296	1.0%	1.0%	12.40	12.40	41.3%	41.3%
Barn Hill Meadows SSSI	30	0.557	0.527	1.9%	1.8%	13.48	13.45	44.9%	44.8%
Burr Closes SSSI	30	0.283	0.276	0.9%	0.9%	10.83	10.82	36.1%	36.1%
Common Plantation SINC	30	0.728	0.723	2.4%	2.4%	12.16	12.15	40.5%	40.5%
Disused Railway Embankment SINC	30	0.558	0.546	1.9%	1.8%	11.32	11.31	37.7%	37.7%
Barmby-on-the-Marsh LWS	30	0.575	0.555	1.9%	1.8%	11.07	11.05	36.9%	36.8%
Brockholes SINC	30	0.480	0.476	1.6%	1.6%	11.70	11.70	39.0%	39.0%
Meadow East of Orchard Farm SINC	30	0.701	0.699	2.3%	2.3%	11.53	11.53	38.4%	38.4%
Barmby Pond LWS	30	0.611	0.585	2.0%	2.0%	10.59	10.57	35.3%	35.2%
Cobble Croft Wood SINC	30	0.697	0.690	2.3%	2.3%	12.32	12.31	41.1%	41.0%
Hagg Green Lane SINC	30	0.704	0.684	2.3%	2.3%	11.67	11.65	38.9%	38.8%

Drax Bioenergy with Carbon Capture and Storage

Receptor				Annual Mean N	IO _x concentrat	tion (µg/m³)			
		Max Cumulative PC Impact Max PC Impact as % of CL		Max Cumulative PEC		Max PEC as % of CL			
		No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ
Sand Pitt Wood & Barffs Close Plantation SINC		0.798	0.790	2.7%	2.6%	12.23	12.22	40.8%	40.7%
Env. Agency Screening	Criterion (as %	of CL)	•	1%	6			709	6

Table 1.14 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Daily Mean NO_x

				Daily Mean N	Ox concentrati	on (µg/m³)			
Receptor		Max Cumula	tive PC Impact	Max PC Impac	ct as % of CL	Max Cumu	lative PEC	Max PEC a	s % of CL
	Critical Level	No Mit ⁿ	Mitig ⁿ						
River Derwent SAC	75	5.459	5.078	7.3%	6.8%	31.44	31.24	41.9%	41.7%
Thorne Moor SAC/SPA/SSSI	75	0.446	0.446	0.6%	0.6%	32.43	32.34	43.2%	43.1%
Lower Derwent SAC	75	0.308	0.308	0.4%	0.4%	27.00	26.90	36.0%	35.9%
Lower Derwent SPA	75	0.308	0.308	0.4%	0.4%	27.00	26.90	36.0%	35.9%
Skipwith Common SAC	75	0.253	0.253	0.3%	0.3%	24.75	24.66	33.0%	32.9%
Skipwith Common SSSI	75	0.253	0.253	0.3%	0.3%	24.75	24.66	33.0%	32.9%
Humber Estuary SAC	75	0.469	0.469	0.6%	0.6%	100.05	99.93	133.4%	133.2%
Humber Estuary SPA/SSSI	75	0.469	0.469	0.6%	0.6%	100.05	99.93	133.4%	133.2%
Breighton Meadows SSSI	75	0.306	0.306	0.4%	0.4%	27.00	26.89	36.0%	35.9%
Eskamhorn Meadows SSSI	75	0.983	0.983	1.3%	1.3%	30.40	29.99	40.5%	40.0%
Derwent Ings SSSI	75	0.308	0.308	0.4%	0.4%	26.63	26.47	35.5%	35.3%
Went Ings SSSI	75	0.245	0.245	0.3%	0.3%	30.27	30.14	40.4%	40.2%
Barn Hill Meadows SSSI	75	0.691	0.691	0.9%	0.9%	32.25	32.14	43.0%	42.9%
Burr Closes SSSI	75	0.346	0.346	0.5%	0.5%	25.99	25.95	34.7%	34.6%
Common Plantation SINC	75	7.240	7.596	9.7%	10.1%	30.30	30.65	40.4%	40.9%
Disused Railway Embankment SINC	75	6.035	5.779	8.0%	7.7%	27.93	27.62	37.2%	36.8%

Drax Bioenergy with Carbon Capture and Storage

				Daily Mean No	Dx concentrati	on (µg/m³)			
Receptor		Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as % of CL	
	Critical Level	No Mit ⁿ	Mitig ⁿ						
Barmby-on-the-Marsh LWS	75	6.310	5.938	8.4%	7.9%	28.32	27.95	37.8%	37.3%
Brockholes SINC	75	5.892	5.440	7.9%	7.3%	28.71	28.26	38.3%	37.7%
Meadow East of Orchard Farm SINC	75	5.436	5.702	7.2%	7.6%	27.18	27.44	36.2%	36.6%
Barmby Pond LWS	75	5.189	4.763	6.9%	6.4%	27.29	27.09	36.4%	36.1%
Cobble Croft Wood SINC	75	7.028	6.706	9.4%	8.9%	31.22	30.90	41.6%	41.2%
Hagg Green Lane SINC	75	5.261	4.843	7.0%	6.5%	29.27	28.95	39.0%	38.6%
Sand Pitt Wood & Barffs Close Plantation SINC	75	7.720	7.948	10.3%	10.6%	31.06	31.02	41.4%	41.4%
	Env. Agency S	creening Criteri	on (as % of CL)	109	%				

Table 1.15 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean NH3

				Annual Mean	NH₃ concentra	ation (µg/m³)			
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as % of CL	
	Childai Levei	No Mit ⁿ	Mitig ⁿ						
River Derwent SAC	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Thorne Moor SAC/SPA/SSSI	1	0.006	0.006	0.6%	0.6%	2.60	2.60	259.8%	259.8%
Lower Derwent SAC	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Lower Derwent SPA	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Skipwith Common SAC	1	0.003	0.003	0.3%	0.3%	2.58	2.58	258.4%	258.4%
Skipwith Common SSSI	1	0.003	0.003	0.3%	0.3%	2.58	2.58	258.4%	258.4%
Humber Estuary SAC	3	0.006	0.007	0.2%	0.2%	3.59	3.59	119.6%	119.7%
Humber Estuary SPA/SSSI	3	0.006	0.007	0.2%	0.2%	3.59	3.59	119.6%	119.7%
Breighton Meadows SSSI	3	0.008	0.007	0.3%	0.2%	3.09	3.09	103.0%	103.0%
Eskamhorn Meadows SSSI	3	0.003	0.003	0.1%	0.1%	2.40	2.40	80.1%	80.1%

Drax Bioenergy with Carbon Capture and Storage

				Annual Mean	NH ₃ concentra	ation (µg/m³)			
Receptor		Max Cumulat	ive PC Impact	Max PC Impac	ct as % of CL	Max Cumul	lative PEC	Max PEC	as % of CL
	Critical Level	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ
Derwent Ings SSSI	3	0.006	0.005	0.2%	0.2%	4.58	4.58	152.7%	152.6%
Went Ings SSSI	3	0.004	0.004	0.1%	0.1%	2.36	2.35	78.5%	78.5%
Barn Hill Meadows SSSI	3	0.006	0.007	0.2%	0.2%	2.33	2.33	77.6%	77.7%
Burr Closes SSSI	3	0.003	0.003	0.1%	0.1%	2.50	2.50	83.5%	83.5%
Common Plantation SINC	3	0.002	0.001	0.1%	0.0%	2.33	2.33	77.7%	77.7%
Disused Railway Embankment SINC	3	0.003	0.002	0.1%	0.1%	2.28	2.28	76.1%	76.1%
Barmby-on-the-Marsh LWS	3	0.004	0.004	0.1%	0.1%	2.29	2.28	76.2%	76.1%
Brockholes SINC	3	0.002	0.002	0.1%	0.1%	2.28	2.28	76.1%	76.1%
Meadow East of Orchard Farm SINC	3	0.001	0.001	0.0%	0.0%	2.33	2.33	77.7%	77.7%
Barmby Pond LWS	3	0.007	0.005	0.2%	0.2%	2.29	2.29	76.3%	76.2%
Cobble Croft Wood SINC	3	0.002	0.002	0.1%	0.1%	2.33	2.33	77.7%	77.7%
Hagg Green Lane SINC	3	0.005	0.004	0.2%	0.1%	3.10	3.10	103.2%	103.2%
Sand Pitt Wood & Barffs Close Plantation SINC	3	0.002	0.002	0.1%	0.1%	2.33	2.33	77.7%	77.7%
	Env. Agency Se	inv. Agency Screening Criterion (as % of CL)			6	70%			0%

Table 1.16 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean SO₂

		•	A	nnual Mean SO	2 concentratio	n (µg/m³)										
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impact as % of CL		Max Cumulative PEC		Max PEC as % of CL								
		No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ							
River Derwent SAC	20	0.074	0.022	0.4%	0.1%	4.04	3.98	20.2%	19.9%							
Thorne Moor SAC/SPA/SSSI	20	0.041	0.017	0.2%	0.1%	1.40	1.38	7.0%	6.9%							
Lower Derwent SAC	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%							
Lower Derwent SPA	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%							

Drax Bioenergy with Carbon Capture and Storage

			Δ	nnual Mean SC	2 concentratio	n (µg/m³)			
Receptor		Max Cumulat	ive PC Impact	Max PC Impa	ct as % of CL	Max Cumul	ative PEC	Max PEC as % of CL	
	Critical Level	No Mit ⁿ	Mitig ⁿ						
Skipwith Common SAC	20	0.026	0.008	0.1%	0.0%	1.46	1.44	7.3%	7.2%
Skipwith Common SSSI	20	0.026	0.008	0.1%	0.0%	1.46	1.44	7.3%	7.2%
Humber Estuary SAC	20	0.073	0.023	0.4%	0.1%	7.59	7.54	38.0%	37.7%
Humber Estuary SPA/SSSI	20	0.073	0.023	0.4%	0.1%	7.59	7.54	38.0%	37.7%
Breighton Meadows SSSI	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%
Eskamhorn Meadows SSSI	20	0.024	0.009	0.1%	0.0%	1.32	1.30	6.6%	6.5%
Derwent Ings SSSI	20	0.061	0.017	0.3%	0.1%	1.78	1.74	8.9%	8.7%
Went Ings SSSI	20	0.028	0.011	0.1%	0.1%	1.35	1.33	6.7%	6.7%
Barn Hill Meadows SSSI	20	0.075	0.024	0.4%	0.1%	1.91	1.86	9.6%	9.3%
Burr Closes SSSI	20	0.028	0.009	0.1%	0.0%	1.27	1.25	6.3%	6.3%
Common Plantation SINC	20	0.009	0.004	0.0%	0.0%	1.45	1.44	7.3%	7.2%
Disused Railway Embankment SINC	20	0.021	0.008	0.1%	0.0%	1.34	1.33	6.7%	6.7%
Barmby-on-the-Marsh LWS	20	0.038	0.013	0.2%	0.1%	1.37	1.34	6.8%	6.7%
Brockholes SINC	20	0.011	0.005	0.1%	0.0%	1.33	1.33	6.7%	6.6%
Meadow East of Orchard Farm SINC	20	0.006	0.003	0.0%	0.0%	1.45	1.44	7.2%	7.2%
Barmby Pond LWS	20	0.060	0.019	0.3%	0.1%	1.40	1.36	7.0%	6.8%
Cobble Croft Wood SINC	20	0.014	0.005	0.1%	0.0%	1.46	1.45	7.3%	7.2%
Hagg Green Lane SINC	20	0.049	0.014	0.2%	0.1%	1.50	1.47	7.5%	7.3%
Sand Pitt Wood & Barffs Close Plantation SINC	20	0.014	0.005	0.1%	0.0%	1.46	1.45	7.3%	7.2%
	Env. Agency S	creening Criteri	on (as % of CL)	19	/₀			70%	

Table 1.17 - Modelled Maximum Cumulative Impa	acts at Ecological Receptors – Annual Nitrogen Deposition Rate

Annual Nitrogen Deposition Rate (kgN/ha/yr)									
Receptor	Critical Load	Max Cumul	ative PC Impact	Max PC Imp	act as % of CL	Max Cumulative PEC		Max PEC as % of CL	
	Critical Load	No Mit ⁿ	Mitig ⁿ						
River Derwent SAC	15	0.106	0.098	0.7%	0.7%	30.35	30.34	202.3%	202.3%
Thorne Moor SAC	5	0.061	0.063	1.2%	1.3%	21.38	21.38	427.6%	427.6%
Thorne Moor SPA	10	0.061	0.063	0.6%	0.6%	21.38	21.38	213.8%	213.8%
Thorne Moor SSSI	5	0.061	0.063	1.2%	1.3%	21.38	21.38	427.6%	427.6%
Lower Derwent SAC	20	0.105	0.098	0.5%	0.5%	30.35	30.34	151.8%	151.7%
Lower Derwent SPA	20	0.105	0.098	0.5%	0.5%	30.35	30.34	151.8%	151.7%
Skipwith Common SAC	10	0.067	0.064	0.7%	0.6%	21.19	21.19	211.9%	211.9%
Skipwith Common SSSI	10	0.067	0.064	0.7%	0.6%	21.19	21.19	211.9%	211.9%
Humber Estuary SAC	20	0.073	0.086	0.4%	0.4%	28.96	28.98	144.8%	144.9%
Humber Estuary SPA/SSSI	20	0.073	0.086	0.4%	0.4%	28.96	28.98	144.8%	144.9%
Breighton Meadows SSSI	20	0.105	0.098	0.5%	0.5%	23.63	23.62	118.2%	118.1%
Eskamhorn Meadows SSSI	10	0.058	0.061	0.6%	0.6%	20.01	20.02	200.1%	200.2%
Derwent Ings SSSI	20	0.096	0.091	0.5%	0.5%	30.34	30.34	151.7%	151.7%
Went Ings SSSI	15	0.049	0.046	0.3%	0.3%	19.44	19.43	129.6%	129.6%
Barn Hill Meadows SSSI	20	0.082	0.088	0.4%	0.4%	20.52	20.54	102.6%	102.7%
Burr Closes SSSI	20	0.045	0.041	0.2%	0.2%	20.69	20.69	103.5%	103.4%
Common Plantation SINC	10	0.158	0.155	1.6%	1.5%	33.90	33.90	339.0%	339.0%
Disused Railway Embankment SINC	10	0.126	0.129	1.3%	1.3%	33.45	33.45	334.5%	334.5%
Barmby-on-the-Marsh LWS	10	0.141	0.141	1.4%	1.4%	33.47	33.47	334.7%	334.7%
Brockholes SINC	10	0.056	0.057	0.6%	0.6%	19.80	19.80	198.0%	198.0%
Meadow East of Orchard Farm SINC	20	0.075	0.076	0.4%	0.4%	19.96	19.96	99.8%	99.8%
Barmby Pond LWS	10	0.095	0.085	1.0%	0.9%	19.85	19.84	198.5%	198.4%

		Annual Nitrogen Deposition Rate (kgN/ha/yr)									
Receptor	Critical Load	Max Cumulative PC Impact		Max PC Impac	ct as % of CL	Max Cumu	lative PEC	Max PEC a	Max PEC as % of CL		
		No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ		
Cobble Croft Wood SINC	10	0.155	0.151	1.6%	1.5%	33.90	33.89	339.0%	338.9%		
Hagg Green Lane SINC	10	0.176	0.173	1.8%	1.7%	40.94	40.93	409.4%	409.3%		
Sand Pitt Wood & Barffs Close Plantation SINC	10	0.176	0.171	1.8%	1.7%	33.92	33.91	339.2%	339.1%		
Barlow Common LNR	10	0.173	0.170	1.7%	1.7%	33.91	33.91	339.1%	339.1%		
	Env. Agency Screening Criterion (as % of CL)			19	6			70)%		

Table 1.18 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Acid Deposition Rate

		Annual Acid Deposition Rate (keq/ha/yr)										
Receptor	Critical Level	Max Cumu	Max Cumulative PC Impact		Max PC Impact as % of CL		Max Cumulative PEC		Max PEC as % of CL			
	Critical Level	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	Max F No Mit ⁿ 375.8% 377.8% 377.5% 216.4% 216.4% 302.8% 82.3% 377.1% 79.7% 269.9% 135.3%	Mitig ⁿ			
Thorne Moor SAC	0.4623	0.010	0.007	2.1%	1.5%	1.74	1.74	375.8%	377.1%			
Thorne Moor SSSI	0.4623	0.010	0.007	2.1%	1.5%	1.75	1.74	377.8%	377.1%			
Lower Derwent SAC	0.643	0.017	0.010	2.6%	1.6%	2.43	2.42	377.5%	376.4%			
Skipwith Common SAC	0.802	0.008	0.006	1.0%	0.7%	1.74	1.73	216.4%	216.1%			
Skipwith Common SSSI	0.802	0.008	0.006	1.0%	0.7%	1.74	1.73	216.4%	216.1%			
Breighton Meadows SSSI	0.643	0.017	0.010	2.6%	1.6%	1.95	1.94	302.8%	301.7%			
Eskamhorn Meadows SSSI	1.998	0.007	0.005	0.3%	0.3%	1.64	1.64	82.3%	82.2%			
Derwent Ings SSSI	0.643	0.014	0.009	2.2%	1.4%	2.42	2.42	377.1%	376.2%			
Went Ings SSSI	2.008	0.007	0.005	0.4%	0.2%	1.60	1.60	79.7%	79.6%			
Barn Hill Meadows SSSI	0.633	0.014	0.010	2.2%	1.5%	1.71	1.70	269.9%	269.3%			
Burr Closes SSSI	1.248	0.007	0.004	0.5%	0.3%	1.69	1.69	135.3%	135.1%			
	Env. Agency S	creening Crite	erion (as % of CL)		1%		1		70%			

SENSITIVITY TEST: WORST CASE EMISSIONS PROFILE

- 1.1.7. Results pertaining to the worst-case emissions profile sensitivity test are presented in **Tables 1.19 to 1.24**, based on emissions from the Proposed Scheme alone.
- 1.1.8. For all pollutant concentrations and deposition rates, it is evident that the modelled maximum PC impacts attributed to the with Proposed Scheme scenario are lower at all receptors relative to the core model scenarios. This is a function of all four biomass units in the Baseline scenario switching from 'mid-merit' operation (full load for 4,000 hours per year) to continuous operation (full load for 8,760 hours per year), resulting in more pollutants being emitted and thus more pronounced changes (increases) in concentrations / deposition rates relative to the With Proposed Scheme scenario. In the With Proposed Scheme scenario, operation changes from 'mid-merit' to continuous full load at the two non-BECCS biomass units only (BECCS units already assumed to operate at continuous full load in core modelling scenario), meaning the changes (increases and decreases) in concentrations / deposition rates are relatively small compared to the Baseline.
- 1.1.9. As a consequence, the maximum modelled impacts of the Proposed Scheme decrease at all receptors under the worst-case emissions profile scenario relative to the core modelling. Whilst some modelled maximum PEC concentrations do increase under worst case emissions in both the Baseline and With Proposed Scheme scenarios, there are no material changes relative to the core modelling equivalents, meaning that the respective assessment significance criteria are not exceeded.
- 1.1.10. The results confirm that the assessment of likely significant effects reported in **Chapter 6 (Air Quality)** is not affected when considering the worst-case emissions profiles in both the Baseline and With Proposed Scheme scenarios. Given that the modelled maximum impacts are lower under a worst-case emissions profile, there was no need to repeat the test in relation to cumulative impacts, as the core modelling results for the cumulative scenarios represent the most conservative results in terms of potential impacts.

Table 1.19 - Modelled Maximum Operational Impacts	at Ecological Receptors – Annual Mean NO _x (Worst Case Emissions Profile)

	Annual Mean NO _x concentration (µg/m ³)									
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL				
River Derwent SAC	30	11.91	0.055	0.2%	12.10	40.3%				
Thorne Moor SAC/SPA/SSSI	30	13.21	0.026	0.1%	13.33	44.4%				
Lower Derwent SAC	30	9.92	0.058	0.2%	10.11	33.7%				
Lower Derwent SPA	30	9.92	0.058	0.2%	10.11	33.7%				
Skipwith Common SAC	30	9.76	0.023	0.1%	9.84	32.8%				
Skipwith Common SSSI	30	9.76	0.023	0.1%	9.84	32.8%				
Humber Estuary SAC	30	46.96	0.056	0.2%	47.14	157.1%				
Humber Estuary SPA/SSSI	30	46.96	0.056	0.2%	47.14	157.1%				
Breighton Meadows SSSI	30	9.92	0.058	0.2%	10.11	33.7%				
Eskamhorn Meadows SSSI	30	11.35	0.012	0.0%	11.38	37.9%				
Derwent Ings SSSI	30	9.80	0.048	0.2%	9.98	33.3%				
Went Ings SSSI	30	12.09	0.018	0.1%	12.15	40.5%				
Barn Hill Meadows SSSI	30	12.89	0.055	0.2%	13.04	43.5%				
Burr Closes SSSI	30	10.53	0.020	0.1%	10.59	35.3%				
Common Plantation SINC	30	11.43	0.004	0.0%	11.44	38.1%				
Disused Railway Embankment SINC	30	10.76	0.009	0.0%	10.78	35.9%				
Barmby-on-the-Marsh LWS	30	10.48	0.021	0.1%	10.53	35.1%				
Brockholes SINC	30	11.22	0.004	0.0%	11.23	37.4%				
Meadow East of Orchard Farm SINC	30	10.83	0.002	0.0%	10.83	36.1%				
Barmby Pond LWS	30	9.96	0.036	0.1%	10.06	33.5%				
Cobble Croft Wood SINC	30	11.62	0.007	0.0%	11.64	38.8%				
Hagg Green Lane SINC	30	10.93	0.034	0.1%	11.05	36.8%				

	Annual Mean NO _x concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Sand Pitt Wood & Barffs Close Plantation SINC	30	11.43	0.007	0.0%	11.45	38.2%		
		Env. Agency Screening C	criterion (as % of CL)	1%		70%		

Table 1.20 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO_x (Worst Case Emissions Profile)

	Daily Mean NO _x concentration (µg/m ³)									
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL				
River Derwent SAC	75	23.82	1.787	2.4%	27.96	37.3%				
Thorne Moor SAC/SPA/SSSI	75	26.42	0.893	1.2%	29.27	39.0%				
Lower Derwent SAC	75	19.84	0.893	1.2%	23.75	31.7%				
Lower Derwent SPA	75	19.84	0.893	1.2%	23.75	31.7%				
Skipwith Common SAC	75	19.52	0.646	0.9%	21.71	28.9%				
Skipwith Common SSSI	75	19.52	0.646	0.9%	21.71	28.9%				
Humber Estuary SAC	75	93.92	1.200	1.6%	96.93	129.2%				
Humber Estuary SPA/SSSI	75	93.92	1.200	1.6%	96.93	129.2%				
Breighton Meadows SSSI	75	19.84	0.893	1.2%	23.75	31.7%				
Eskamhorn Meadows SSSI	75	22.70	1.912	2.5%	25.59	34.1%				
Derwent Ings SSSI	75	19.60	0.831	1.1%	23.37	31.2%				
Went Ings SSSI	75	24.18	0.971	1.3%	26.60	35.5%				
Barn Hill Meadows SSSI	75	25.78	0.962	1.3%	28.30	37.7%				
Burr Closes SSSI	75	21.06	0.761	1.0%	22.99	30.7%				
Common Plantation SINC	75	22.86	1.038	1.4%	24.19	32.3%				
Disused Railway Embankment SINC	75	21.52	1.184	1.6%	23.07	30.8%				
Barmby-on-the-Marsh LWS	75	20.96	1.621	2.2%	23.44	31.3%				

Drax Bioenergy with Carbon Capture and Storage

	Daily Mean NO _x concentration (µg/m ³)								
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Brockholes SINC	75	22.44	1.623	2.2%	24.44	32.6%			
Meadow East of Orchard Farm SINC	75	21.66	0.554	0.7%	22.33	29.8%			
Barmby Pond LWS	75	19.92	1.577	2.1%	23.29	31.1%			
Cobble Croft Wood SINC	75	23.24	1.176	1.6%	25.28	33.7%			
Hagg Green Lane SINC	75	21.86	1.248	1.7%	25.22	33.6%			
Sand Pitt Wood & Barffs Close Plantation SINC	75	22.86	1.630	2.2%	24.97	33.3%			
	10%								

Table 1.21 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH₃ (Worst Case Emissions Profile)

		•	`	³ concentration (μg/m ³))	
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
River Derwent SAC	3	4.57	0.002	0.1%	4.58	152.6%
Thorne Moor SAC/SPA/SSSI	1	2.59	0.001	0.1%	2.60	259.6%
Lower Derwent SAC	3	4.57	0.002	0.1%	4.58	152.6%
Lower Derwent SPA	3	4.57	0.002	0.1%	4.58	152.6%
Skipwith Common SAC	1	2.58	0.001	0.1%	2.58	258.4%
Skipwith Common SSSI	1	2.58	0.001	0.1%	2.58	258.4%
Humber Estuary SAC	3	3.58	0.002	0.1%	3.59	119.6%
Humber Estuary SPA/SSSI	3	3.58	0.002	0.1%	3.59	119.6%
Breighton Meadows SSSI	3	3.08	0.002	0.1%	3.09	103.0%
Eskamhorn Meadows SSSI	3	2.40	0.001	0.0%	2.40	80.1%
Derwent Ings SSSI	3	4.57	0.002	0.1%	4.58	152.6%
Went Ings SSSI	3	2.35	0.001	0.0%	2.35	78.4%

Drax Bioenergy with Carbon Capture and Storage

			Annual Mean NH	₃ concentration (µg/m³))	
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
Barn Hill Meadows SSSI	3	2.32	0.002	0.1%	2.33	77.6%
Burr Closes SSSI	3	2.50	0.001	0.0%	2.50	83.4%
Common Plantation SINC	3	2.33	0.000	0.0%	2.33	77.7%
Disused Railway Embankment SINC	1	2.28	0.000	0.0%	2.28	76.0%
Barmby-on-the-Marsh LWS	3	2.28	0.001	0.0%	2.28	76.1%
Brockholes SINC	3	2.28	0.000	0.0%	2.28	76.0%
Meadow East of Orchard Farm SINC	1	2.33	0.000	0.0%	2.33	77.7%
Barmby Pond LWS	1	2.28	0.001	0.0%	2.29	76.2%
Cobble Croft Wood SINC	3	2.33	0.000	0.0%	2.33	77.7%
Hagg Green Lane SINC	3	3.09	0.001	0.0%	3.10	103.2%
Sand Pitt Wood & Barffs Close Plantation SINC	3	2.33	0.000	0.0%	2.33	77.7%
	Env. Agency Screening Criterion (as % of CL)					

Table 1.22 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO₂ (Worst Case Emissions Profile)

	Annual Mean SO ₂ concentration (µg/m ³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	20	3.93	0.021	0.1%	4.03	20.1%		
Thorne Moor SAC/SPA/SSSI	20	1.34	0.009	0.0%	1.40	7.0%		
Lower Derwent SAC	20	1.70	0.021	0.1%	1.80	9.0%		
Lower Derwent SPA	20	1.70	0.021	0.1%	1.80	9.0%		
Skipwith Common SAC	20	1.42	0.008	0.0%	1.46	7.3%		
Skipwith Common SSSI	20	1.42	0.008	0.0%	1.46	7.3%		
Humber Estuary SAC	20	7.49	0.021	0.1%	7.58	37.9%		

Drax Bioenergy with Carbon Capture and Storage

Receptor	Annual Mean SO ₂ concentration (µg/m ³)								
	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Humber Estuary SPA/SSSI	20	7.49	0.021	0.1%	7.58	37.9%			
Breighton Meadows SSSI	20	1.70	0.021	0.1%	1.80	9.0%			
Eskamhorn Meadows SSSI	20	1.29	0.005	0.0%	1.31	6.5%			
Derwent Ings SSSI	20	1.69	0.016	0.1%	1.78	8.9%			
Went Ings SSSI	20	1.31	0.007	0.0%	1.34	6.7%			
Barn Hill Meadows SSSI	20	1.81	0.022	0.1%	1.89	9.4%			
Burr Closes SSSI	20	1.23	0.008	0.0%	1.26	6.3%			
Common Plantation SINC	20	1.44	0.002	0.0%	1.44	7.2%			
Disused Railway Embankment SINC	20	1.32	0.004	0.0%	1.33	6.7%			
Barmby-on-the-Marsh LWS	20	1.32	0.009	0.0%	1.35	6.7%			
Brockholes SINC	20	1.32	0.002	0.0%	1.32	6.6%			
Meadow East of Orchard Farm SINC	20	1.44	0.001	0.0%	1.44	7.2%			
Barmby Pond LWS	20	1.32	0.015	0.1%	1.37	6.9%			
Cobble Croft Wood SINC	20	1.44	0.003	0.0%	1.45	7.2%			
Hagg Green Lane SINC	20	1.43	0.013	0.1%	1.49	7.5%			
Sand Pitt Wood & Barffs Close Plantation SINC	20	1.44	0.003	0.0%	1.45	7.2%			
	1%		70%						

Table 1.23 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Worst Case Emissions Profile)

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	15	30.22	0.017	0.1%	30.29	201.9%		
Thorne Moor SAC	5	21.31	0.008	0.2%	21.35	427.0%		

Beconter	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Thorne Moor SPA	10	21.31	0.008	0.1%	21.35	213.5%		
Thorne Moor SSSI	5	21.31	0.008	0.2%	21.35	427.0%		
Lower Derwent SAC	20	30.22	0.017	0.1%	30.29	151.5%		
Lower Derwent SPA	20	30.22	0.017	0.1%	30.29	151.5%		
Skipwith Common SAC	10	21.12	0.007	0.1%	21.14	211.4%		
Skipwith Common SSSI	10	21.12	0.007	0.1%	21.14	211.4%		
Humber Estuary SAC	20	28.87	0.017	0.1%	28.93	144.7%		
Humber Estuary SPA/SSSI	20	28.87	0.017	0.1%	28.93	144.7%		
Breighton Meadows SSSI	20	23.51	0.017	0.1%	23.57	117.9%		
Eskamhorn Meadows SSSI	10	19.95	0.004	0.0%	19.96	199.6%		
Derwent Ings SSSI	20	30.22	0.014	0.1%	30.29	151.4%		
Went Ings SSSI	15	19.38	0.005	0.0%	19.40	129.4%		
Barn Hill Meadows SSSI	20	20.43	0.017	0.1%	20.49	102.4%		
Burr Closes SSSI	20	20.64	0.006	0.0%	20.66	103.3%		
Common Plantation SINC	10	33.74	0.002	0.0%	33.75	337.5%		
Disused Railway Embankment SINC	10	33.32	0.005	0.1%	33.33	333.3%		
Barmby-on-the-Marsh LWS	10	33.32	0.011	0.1%	33.35	333.5%		
Brockholes SINC	10	19.74	0.001	0.0%	19.74	197.4%		
Meadow East of Orchard Farm SINC	20	19.88	0.001	0.0%	19.88	99.4%		
Barmby Pond LWS	10	19.74	0.012	0.1%	19.78	197.8%		
Cobble Croft Wood SINC	10	33.74	0.004	0.0%	33.75	337.5%		
Hagg Green Lane SINC	10	40.74	0.017	0.2%	40.81	408.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	10	33.74	0.004	0.0%	33.75	337.5%		

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Barlow Common LNR	10	33.74	0.002	0.0%	33.75	337.5%			
Env. Agency Screening Criterion (as % of CL)				1%		70%			

Table 1.24 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Worst Case Emissions Profile)

Receptor	Annual Acid Deposition Rate (keq/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Thorne Moor SAC	0.462	1.73	0.002	0.4%	1.74	377.2%			
Thorne Moor SSSI	0.462	1.73	0.002	0.4%	1.74	377.2%			
Lower Derwent SAC	0.643	2.40	0.004	0.6%	2.42	376.6%			
Skipwith Common SAC	0.802	1.73	0.002	0.2%	1.73	216.0%			
Skipwith Common SSSI	0.802	1.73	0.002	0.2%	1.73	216.0%			
Breighton Meadows SSSI	0.643	1.92	0.004	0.6%	1.94	301.9%			
Eskamhorn Meadows SSSI	1.998	1.64	0.001	0.0%	1.64	82.1%			
Derwent Ings SSSI	0.643	2.40	0.003	0.5%	2.42	376.4%			
Went Ings SSSI	2.008	1.59	0.001	0.1%	1.60	79.5%			
Barn Hill Meadows SSSI	0.633	1.69	0.004	0.6%	1.70	269.2%			
Burr Closes SSSI	1.248	1.68	0.001	0.1%	1.69	135.1%			
	1%		70%						