

# ENVIRONMENTAL STATEMENT -VOLUME 3 - APPENDIX 6.5

## **Operation Phase Air Quality Assessment Results Tables: Ecological Receptors**

## **Drax Bioenergy with Carbon Capture and Storage**

The Infrastructure Planning (Applications: Prescribed Forms and Procedure) Regulations 2009 - 5(2)(a)

**Document Reference Number: 6.3.6.5** 

Applicant: Drax Power Limited

PINS Reference: EN010120



REVISION:05 DATE: July 2023 DOCUMENT OWNER: WSP UK Limited AUTHOR: B Tuckett-Jones APPROVER: N. Ashworth PUBLIC

## **TABLE OF CONTENTS**

1.	OPEF	RATION PHASE AIR QUALITY ASSESSMENT RESULTS TABLES: ECOLOGICAL	
RE	CEPTO	DRS	1
	1.1.	Impacts on Ecological Receptors	1

## TABLES

Table 1.1 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean         NOx         3
Table 1.2 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NOx
Table 1.3 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH <sub>3</sub>
Table 1.4 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanSO2
Table 1.5 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual NitrogenDeposition Rate8
Table 1.6 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual AcidDeposition Rate10
Table 1.7 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanNOx (Including Mitigation)
Table 1.8 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NOx(Including Mitigation)13
Table 1.9 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH <sub>3</sub> (Including Mitigation)14
Table 1.10 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanSO2 (Including Mitigation)15
Table 1.11 - Modelled Maximum Operational Impacts at Ecological Receptors – AnnualNitrogen Deposition Rate (Including Mitigation)16
Table 1.12 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual AcidDeposition Rate (Including Mitigation)18
Table 1.13 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean         NOx         20
Table 1.14 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Daily Mean NOx

Table 1.15 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual MeanNH322
Table 1.16 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual MeanSO223
Table 1.17 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual NitrogenDeposition Rate25
Table 1.18 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual AcidDeposition Rate26
Table 1.19 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanNOx (Worst Case Emissions Profile)28
Table 1.20 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily MeanNOx (Worst Case Emissions Profile)29
Table 1.21 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH <sub>3</sub> (Worst Case Emissions Profile)
Table 1.22 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual MeanSO2 (Worst Case Emissions Profile)31
Table 1.23 - Modelled Maximum Operational Impacts at Ecological Receptors – AnnualNitrogen Deposition Rate (Worst Case Emissions Profile)
Table 1.24 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual AcidDeposition Rate (Worst Case Emissions Profile)

## 1. OPERATION PHASE AIR QUALITY ASSESSMENT RESULTS TABLES: ECOLOGICAL RECEPTORS

## 1.1. IMPACTS ON ECOLOGICAL RECEPTORS

- 1.1.1. The results of the modelling assessment at each modelled ecological receptor are presented in the below tables for each relevant pollutant and averaging period applicable to the study. The maximum modelled concentration and deposition values are presented, which is based on modelling over all five years of meteorological data (2016-2020). The change in PC and PEC, as a percentage of the relevant critical level / load, is presented for each receptor.
- 1.1.2. The PC impact in the with Proposed Scheme scenario represents the change in concentration / deposition between the Baseline scenario and Proposed Scheme scenario.
- 1.1.3. In all results tables presented in this appendix, the following designated site names are shortened to 'Thorne Moor SPA' and Thorne Moor SSSI', respectively:
  - a. Thorne & Hatfield Moors SPA
  - **b.** Thorne, Crowle and Goole Moors SSSI.
- 1.1.4. In the latest version of this Appendix, the results reflect advice provided by Natural England via email that the May 2023 updates on the APIS website need not be incorporated into the assessment at this stage of the DCO process. Following this advice, the following change is made:
  - Background NOx concentrations over the Humber Estuary SAC/SPA/SSSI reflect APIS data from 2017-2019 within the 15km study area for the Proposed Development. In comparison to Revs 01 to 03 of this Appendix, this represents a reduction in NOx concentrations since the original data presented reflected the maximum NOx concentrations *anywhere* within the Humber Estuary designated sites, and a slight increase in comparison to Rev 04 since this reflected the May 2023 updated (and reduced) 2019 – 2021 concentration data. Notwithstanding this update, as in Rev 04 of this Appendix, the NOx PEC over the Humber Estuary designated sites remains within the critical level both with and without the Proposed Development.
- 1.1.5. Further, the increased critical load for acid deposition over Lower Derwent Valley SAC and Barn Hill Meadows SSSI is retained from Rev 04 of this Appendix, and also applied to Breighton Meadows SSSI, Derwent Ings SSSI in this revision, This critical load was determined by the project ecologists and agreed with Natural England outwith the APIS update process. The rationale for this is presented in the Lower Derwent Valley Habitats and Soil Analysis Technical Note (REP3-009), which forms Appendix 8 to the HRA Report (REP6-021, Rev04 submitted at Deadline 9).
- 1.1.6. Finally the cumulative daily mean NOx process contributions (PCs) have been corrected in Table 1.14 for some designated sites where the annual mean NOx PCs

had erroneously been inserted into the Table. This has no impact on the assessment since the amended PCs remain below the EA screening criterion (10% of the critical level) and the Predicted Environmental Concentrations (PECs), which did not require updating, are below the critical levels.

#### CORE MODEL SCENARIOS

1.1.7. Results pertaining to the core model scenarios are presented in **Tables 1.1 to 1.6**.

#### Table 1.1 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO<sub>x</sub>

		Annual Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
River Derwent SAC	30	11.91	0.163	0.5%	12.13	40.4%			
Thorne Moor SAC/SPA/SSSI	30	13.21	0.076	0.3%	13.32	44.4%			
Lower Derwent SAC	30	9.92	0.166	0.6%	10.15	33.8%			
Lower Derwent SPA	30	9.92	0.166	0.6%	10.15	33.8%			
Skipwith Common SAC	30	9.76	0.058	0.2%	9.84	32.8%			
Skipwith Common SSSI	30	9.76	0.058	0.2%	9.84	32.8%			
Humber Estuary SAC	30	19.22ª	0.147	0.5%	19.42	64.7%			
Humber Estuary SPA/SSSI	30	19.22ª	0.147	0.5%	19.42	64.7%			
Breighton Meadows SSSI	30	9.92	0.166	0.6%	10.15	33.8%			
Eskamhorn Meadows SSSI	30	11.35	0.046	0.2%	11.40	38.0%			
Derwent Ings SSSI	30	9.80	0.134	0.4%	9.99	33.3%			
Went Ings SSSI	30	12.09	0.052	0.2%	12.16	40.5%			
Barn Hill Meadows SSSI	30	12.89	0.152	0.5%	13.09	43.6%			
Burr Closes SSSI	30	10.53	0.062	0.2%	10.61	35.4%			
Common Plantation SINC	30	11.43	0.017	0.1%	11.45	38.2%			
Disused Railway Embankment SINC	30	10.76	0.040	0.1%	10.81	36.0%			
Barmby-on-the-Marsh LWS	30	10.48	0.076	0.3%	10.57	35.2%			
Brockholes SINC	30	11.22	0.019	0.1%	11.24	37.5%			
Meadow East of Orchard Farm SINC	30	10.83	0.009	0.0%	10.84	36.1%			
Barmby Pond LWS	30	9.96	0.127	0.4%	10.12	33.7%			
Cobble Croft Wood SINC	30	11.62	0.027	0.1%	11.65	38.8%			
Hagg Green Lane SINC	30	10.93	0.103	0.3%	11.07	36.9%			

	Annual Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Sand Pitt Wood & Barffs Close Plantation SINC	30	11.43	0.028	0.1%	11.46	38.2%		
Env. Agency Screening Criterion (as % of CL) 1% 70%								
a. Background NOx concentration for Humber Estuary SAC/SPA/SSSI taken from maximum within study area (within 15km of Proposed Scheme, 2017-2019), from								

## Table 1.2 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NOx

	Daily Mean NO <sub>x</sub> concentration (μg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	75	23.82	1.787	2.4%	27.96	37.3%		
Thorne Moor SAC/SPA/SSSI	75	26.42	0.893	1.2%	29.27	39.0%		
Lower Derwent SAC	75	19.84	0.893	1.2%	23.75	31.7%		
Lower Derwent SPA	75	19.84	0.893	1.2%	23.75	31.7%		
Skipwith Common SAC	75	19.52	0.646	0.9%	21.71	28.9%		
Skipwith Common SSSI	75	19.52	0.646	0.9%	21.71	28.9%		
Humber Estuary SAC	75	38.44ª	1.200	1.6%	41.45	55.3%		
Humber Estuary SPA/SSSI	75	38.44ª	1.200	1.6%	41.45	55.3%		
Breighton Meadows SSSI	75	19.84	0.893	1.2%	23.75	31.7%		
Eskamhorn Meadows SSSI	75	22.70	1.912	2.5%	25.59	34.1%		
Derwent Ings SSSI	75	19.60	0.831	1.1%	23.37	31.2%		
Went Ings SSSI	75	24.18	0.971	1.3%	26.60	35.5%		
Barn Hill Meadows SSSI	75	25.78	0.962	1.3%	28.30	37.7%		
Burr Closes SSSI	75	21.06	0.761	1.0%	22.99	30.7%		
Common Plantation SINC	75	22.86	1.038	1.4%	24.19	32.3%		
Disused Railway Embankment SINC	75	21.52	1.184	1.6%	23.07	30.8%		

#### Drax Bioenergy with Carbon Capture and Storage

	Daily Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Barmby-on-the-Marsh LWS	75	20.96	1.621	2.2%	23.44	31.3%		
Brockholes SINC	75	22.44	1.623	2.2%	24.44	32.6%		
Meadow East of Orchard Farm SINC	75	21.66	0.554	0.7%	22.33	29.8%		
Barmby Pond LWS	75	19.92	1.577	2.1%	23.29	31.1%		
Cobble Croft Wood SINC	75	23.24	1.176	1.6%	25.28	33.7%		
Hagg Green Lane SINC	75	21.86	1.248	1.7%	25.22	33.6%		
Sand Pitt Wood & Barffs Close Plantation SINC	75	22.86	1.630	2.2%	24.97	33.3%		
	10%		·					
a. Background NOx concentration for Humber Estuary SAC/SPA/SSSI taken from maximum within study area (within 15km of Proposed Scheme, 2017-2019), from								

## Table 1.3 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH<sub>3</sub>

	Annual Mean NH <sub>3</sub> concentration (µg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	3	4.57	0.007	0.2%	4.58	152.7%		
Thorne Moor SAC/SPA/SSSI	1	2.59	0.003	0.3%	2.60	259.5%		
Lower Derwent SAC	3	4.57	0.007	0.2%	4.58	152.7%		
Lower Derwent SPA	3	4.57	0.007	0.2%	4.58	152.7%		
Skipwith Common SAC	1	2.58	0.002	0.2%	2.58	258.4%		
Skipwith Common SSSI	1	2.58	0.002	0.2%	2.58	258.4%		
Humber Estuary SAC	3	3.58	0.004	0.1%	3.59	119.5%		
Humber Estuary SPA/SSSI	3	3.58	0.004	0.1%	3.59	119.5%		
Breighton Meadows SSSI	3	3.08	0.007	0.2%	3.09	103.0%		
Eskamhorn Meadows SSSI	3	2.40	0.002	0.1%	2.40	80.1%		

#### Drax Bioenergy with Carbon Capture and Storage

	Annual Mean NH <sub>3</sub> concentration (μg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Derwent Ings SSSI	3	4.57	0.005	0.2%	4.58	152.6%		
Went Ings SSSI	3	2.35	0.002	0.1%	2.35	78.4%		
Barn Hill Meadows SSSI	3	2.32	0.005	0.2%	2.33	77.6%		
Burr Closes SSSI	3	2.50	0.003	0.1%	2.50	83.5%		
Common Plantation SINC	3	2.33	0.001	0.0%	2.33	77.7%		
Disused Railway Embankment SINC	1	2.28	0.002	0.1%	2.28	76.1%		
Barmby-on-the-Marsh LWS	3	2.28	0.003	0.1%	2.28	76.1%		
Brockholes SINC	3	2.28	0.001	0.0%	2.28	76.0%		
Meadow East of Orchard Farm SINC	1	2.33	0.000	0.0%	2.33	77.7%		
Barmby Pond LWS	1	2.28	0.006	0.2%	2.29	76.2%		
Cobble Croft Wood SINC	3	2.33	0.001	0.0%	2.33	77.7%		
Hagg Green Lane SINC	3	3.09	0.004	0.1%	3.10	103.2%		
Sand Pitt Wood & Barffs Close Plantation SINC	3	2.33	0.001	0.0%	2.33	77.7%		
		Criterion (as % of CL)	1%		70%			

## Table 1.4 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO<sub>2</sub>

	Annual Mean SO <sub>2</sub> concentration (µg/m <sup>3</sup> )						
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL	
River Derwent SAC	20	3.93	0.072	0.4%	4.03	20.2%	
Thorne Moor SAC/SPA/SSSI	20	1.34	0.033	0.2%	1.39	7.0%	
Lower Derwent SAC	20	1.70	0.073	0.4%	1.81	9.0%	
Lower Derwent SPA	20	1.70	0.073	0.4%	1.81	9.0%	
Skipwith Common SAC	20	1.42	0.025	0.1%	1.46	7.3%	

#### Drax Bioenergy with Carbon Capture and Storage

Environmental Statement - Volume 3 - Appendix 6.5

Page 6 of 34

		Annual Mean SO <sub>2</sub> concentration (μg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Skipwith Common SSSI	20	1.42	0.025	0.1%	1.46	7.3%			
Humber Estuary SAC	20	7.49	0.069	0.3%	7.59	38.0%			
Humber Estuary SPA/SSSI	20	7.49	0.069	0.3%	7.59	38.0%			
Breighton Meadows SSSI	20	1.70	0.073	0.4%	1.81	9.0%			
Eskamhorn Meadows SSSI	20	1.29	0.021	0.1%	1.32	6.6%			
Derwent Ings SSSI	20	1.69	0.059	0.3%	1.78	8.9%			
Went Ings SSSI	20	1.31	0.023	0.1%	1.34	6.7%			
Barn Hill Meadows SSSI	20	1.81	0.072	0.4%	1.91	9.5%			
Burr Closes SSSI	20	1.23	0.027	0.1%	1.27	6.3%			
Common Plantation SINC	20	1.44	0.008	0.0%	1.45	7.2%			
Disused Railway Embankment SINC	20	1.32	0.019	0.1%	1.34	6.7%			
Barmby-on-the-Marsh LWS	20	1.32	0.036	0.2%	1.36	6.8%			
Brockholes SINC	20	1.32	0.009	0.0%	1.33	6.7%			
Meadow East of Orchard Farm SINC	20	1.44	0.004	0.0%	1.44	7.2%			
Barmby Pond LWS	20	1.32	0.058	0.3%	1.40	7.0%			
Cobble Croft Wood SINC	20	1.44	0.013	0.1%	1.46	7.3%			
Hagg Green Lane SINC	20	1.43	0.047	0.2%	1.50	7.5%			
Sand Pitt Wood & Barffs Close Plantation SINC	20	1.44	0.013	0.1%	1.46	7.3%			
		Env. Agency Screening (	Criterion (as % of CL)	1%		70%			

#### Table 1.5 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)										
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL					
River Derwent SAC	15	30.22	0.054	0.4%	30.30	202.0%					
Thorne Moor SAC	5	21.31	0.025	0.5%	21.35	426.9%					
Thorne Moor SPA	10	21.31	0.025	0.2%	21.35	213.5%					
Thorne Moor SSSI	5	21.31	0.025	0.5%	21.35	426.9%					
Lower Derwent SAC	20	30.22	0.055	0.3%	30.30	151.5%					
Lower Derwent SPA	20	30.22	0.055	0.3%	30.30	151.5%					
Skipwith Common SAC	10	21.12	0.019	0.2%	21.14	211.4%					
Skipwith Common SSSI	10	21.12	0.019	0.2%	21.14	211.4%					
Humber Estuary SAC	20	28.87	0.034	0.2%	28.92	144.6%					
Humber Estuary SPA/SSSI	20	28.87	0.034	0.2%	28.92	144.6%					
Breighton Meadows SSSI	20	23.51	0.055	0.3%	23.58	117.9%					
Eskamhorn Meadows SSSI	10	19.95	0.016	0.2%	19.97	199.7%					
Derwent Ings SSSI	20	30.22	0.043	0.2%	30.29	151.5%					
Went Ings SSSI	15	19.38	0.017	0.1%	19.41	129.4%					
Barn Hill Meadows SSSI	20	20.43	0.039	0.2%	20.48	102.4%					
Burr Closes SSSI	20	20.64	0.020	0.1%	20.67	103.3%					
Common Plantation SINC	10	33.74	0.010	0.1%	33.75	337.5%					
Disused Railway Embankment SINC	10	33.32	0.022	0.2%	33.35	333.5%					
Barmby-on-the-Marsh LWS	10	33.32	0.043	0.4%	33.37	333.7%					
Brockholes SINC	10	19.74	0.007	0.1%	19.75	197.5%					

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Meadow East of Orchard Farm SINC	20	19.88	0.003	0.0%	19.88	99.4%		
Barmby Pond LWS	10	19.74	0.044	0.4%	19.80	198.0%		
Cobble Croft Wood SINC	10	33.74	0.015	0.2%	33.76	337.6%		
Hagg Green Lane SINC	10	40.74	0.058	0.6%	40.82	408.2%		
Sand Pitt Wood & Barffs Close Plantation SINC	10	33.74	0.016	0.2%	33.76	337.6%		
Barlow Common LNR	10	33.74	0.010	0.1%	33.75	337.5%		
Env. Agency Screening C	riterion (as % of CL)	·	1%		70%			

Table 1.6 - Modelled Maximum O	perational Impacts at Eco	ological Receptors – Annual	Acid Deposition Rate

	Annual Acid Deposition Rate (keq/ha/yr)							
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Thorne Moor SAC	0.462	1.73	0.006	1.3%	1.74	376.9%		
Thorne Moor SSSI	0.462	1.73	0.006	1.3%	1.74	376.9%		
Lower Derwent SAC	4.856	2.40	0.013	0.3%	2.42	49.9%		
Skipwith Common SAC	0.802	1.73	0.005	0.6%	1.73	216.0%		
Skipwith Common SSSI	0.802	1.73	0.005	0.6%	1.73	216.0%		
Breighton Meadows SSSI	4.856	1.92	0.013	0.3%	1.94	40.0%		
Eskamhorn Meadows SSSI	1.998	1.64	0.004	0.2%	1.64	82.2%		
Derwent Ings SSSI	4.856	2.40	0.010	0.2%	2.42	49.8%		
Went Ings SSSI	2.008	1.59	0.004	0.2%	1.60	79.6%		
Barn Hill Meadows SSSI	4.856	1.69	0.010	0.2%	1.70	35.1%		
Burr Closes SSSI	1.248	1.68	0.005	0.4%	1.69	135.2%		
	Env. Agency Screening Criterion (as % of CL)					70%		

#### WITH PROPOSED SCHEME MITIGATION MODEL SCENARIOS

1.1.8. Results pertaining to the core model scenarios, including the With Proposed Scheme Mitigation (as detailed in Section 6.10 of Chapter 6 (Air Quality) (APP-042)) are presented in Tables 1.7 to 1.12.

		Annual Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation			
River Derwent SAC	30	0.163	0.133	0.5%	0.4%			
Thorne Moor SAC/SPA/SSSI	30	0.076	0.064	0.3%	0.2%			
Lower Derwent SAC	30	0.166	0.138	0.6%	0.5%			
Lower Derwent SPA	30	0.166	0.138	0.6%	0.5%			
Skipwith Common SAC	30	0.058	0.049	0.2%	0.2%			
Skipwith Common SSSI	30	0.058	0.049	0.2%	0.2%			
Humber Estuary SAC	30	0.147	0.125	0.5%	0.4%			
Humber Estuary SPA/SSSI	30	0.147	0.125	0.5%	0.4%			
Breighton Meadows SSSI	30	0.166	0.138	0.6%	0.5%			
Eskamhorn Meadows SSSI	30	0.046	0.034	0.2%	0.1%			
Derwent Ings SSSI	30	0.134	0.114	0.4%	0.4%			
Went Ings SSSI	30	0.052	0.042	0.2%	0.1%			
Barn Hill Meadows SSSI	30	0.152	0.124	0.5%	0.4%			
Burr Closes SSSI	30	0.062	0.050	0.2%	0.2%			
Common Plantation SINC	30	0.017	0.011	0.1%	0.0%			
Disused Railway Embankment SINC	30	0.040	0.029	0.1%	0.1%			
Barmby-on-the-Marsh LWS	30	0.076	0.057	0.3%	0.2%			
Brockholes SINC	30	0.019	0.013	0.1%	0.0%			
Meadow East of Orchard Farm SINC	30	0.009	0.006	0.0%	0.0%			
Barmby Pond LWS	30	0.127	0.098	0.4%	0.3%			
Cobble Croft Wood SINC	30	0.027	0.020	0.1%	0.1%			
Hagg Green Lane SINC	30	0.103	0.083	0.3%	0.3%			

	Annual Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )					
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation	
Sand Pitt Wood & Barffs Close Plantation SINC	30	0.028	0.020	0.1%	0.1%	
Env. Age		1%				

## Table 1.8 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO<sub>x</sub> (Including Mitigation)

	Daily Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	75	1.787	1.321	2.4%	1.8%		
Thorne Moor SAC/SPA/SSSI	75	0.893	0.699	1.2%	0.9%		
Lower Derwent SAC	75	0.893	0.673	1.2%	0.9%		
Lower Derwent SPA	75	0.893	0.673	1.2%	0.9%		
Skipwith Common SAC	75	0.646	0.434	0.9%	0.6%		
Skipwith Common SSSI	75	0.646	0.434	0.9%	0.6%		
Humber Estuary SAC	75	1.200	0.872	1.6%	1.2%		
Humber Estuary SPA/SSSI	75	1.200	0.872	1.6%	1.2%		
Breighton Meadows SSSI	75	0.893	0.673	1.2%	0.9%		
Eskamhorn Meadows SSSI	75	1.912	1.438	2.5%	1.9%		
Derwent Ings SSSI	75	0.831	0.670	1.1%	0.9%		
Went Ings SSSI	75	0.971	0.651	1.3%	0.9%		
Barn Hill Meadows SSSI	75	0.962	1.274	1.3%	1.7%		
Burr Closes SSSI	75	0.761	0.561	1.0%	0.7%		
Common Plantation SINC	75	1.038	0.684	1.4%	0.9%		
Disused Railway Embankment SINC	75	1.184	1.069	1.6%	1.4%		
Barmby-on-the-Marsh LWS	75	1.621	1.420	2.2%	1.9%		

#### Drax Bioenergy with Carbon Capture and Storage

	Daily Mean NO <sub>x</sub> concentration (μg/m <sup>3</sup> )						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Brockholes SINC	75	1.623	1.170	2.2%	1.6%		
Meadow East of Orchard Farm SINC	75	0.554	0.385	0.7%	0.5%		
Barmby Pond LWS	75	1.577	1.151	2.1%	1.5%		
Cobble Croft Wood SINC	75	1.176	0.767	1.6%	1.0%		
Hagg Green Lane SINC	75	1.248	0.832	1.7%	1.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	75	1.630	1.083	2.2%	1.4%		
		10%					

## Table 1.9 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH<sub>3</sub> (Including Mitigation)

	Annual Mean NH <sub>3</sub> concentration (µg/m <sup>3</sup> )						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	3	0.007	0.006	0.2%	0.2%		
Thorne Moor SAC/SPA/SSSI	1	0.003	0.003	0.3%	0.3%		
Lower Derwent SAC	3	0.007	0.006	0.2%	0.2%		
Lower Derwent SPA	3	0.007	0.006	0.2%	0.2%		
Skipwith Common SAC	1	0.002	0.002	0.2%	0.2%		
Skipwith Common SSSI	1	0.002	0.002	0.2%	0.2%		
Humber Estuary SAC	3	0.004	0.005	0.1%	0.2%		
Humber Estuary SPA/SSSI	3	0.004	0.005	0.1%	0.2%		
Breighton Meadows SSSI	3	0.007	0.006	0.2%	0.2%		
Eskamhorn Meadows SSSI	3	0.002	0.002	0.1%	0.1%		
Derwent Ings SSSI	3	0.005	0.005	0.2%	0.2%		
Went Ings SSSI	3	0.002	0.002	0.1%	0.1%		

#### Drax Bioenergy with Carbon Capture and Storage

	Annual Mean NH <sub>3</sub> concentration (μg/m <sup>3</sup> )						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Barn Hill Meadows SSSI	3	0.005	0.005	0.2%	0.2%		
Burr Closes SSSI	3	0.003	0.002	0.1%	0.1%		
Common Plantation SINC	3	0.001	0.001	0.0%	0.0%		
Disused Railway Embankment SINC	1	0.002	0.001	0.1%	0.0%		
Barmby-on-the-Marsh LWS	3	0.003	0.003	0.1%	0.1%		
Brockholes SINC	3	0.001	0.001	0.0%	0.0%		
Meadow East of Orchard Farm SINC	1	0.000	0.000	0.0%	0.0%		
Barmby Pond LWS	1	0.006	0.004	0.2%	0.1%		
Cobble Croft Wood SINC	3	0.001	0.001	0.0%	0.0%		
Hagg Green Lane SINC	3	0.004	0.004	0.1%	0.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	3	0.001	0.001	0.0%	0.0%		
		1%					

## Table 1.10 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO<sub>2</sub> (Including Mitigation)

	Annual Mean SO <sub>2</sub> concentration (μg/m <sup>3</sup> )						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	20	0.072	0.021	0.4%	0.1%		
Thorne Moor SAC/SPA/SSSI	20	0.033	0.008	0.2%	0.0%		
Lower Derwent SAC	20	0.073	0.020	0.4%	0.1%		
Lower Derwent SPA	20	0.073	0.020	0.4%	0.1%		
Skipwith Common SAC	20	0.025	0.007	0.1%	0.0%		
Skipwith Common SSSI	20	0.025	0.007	0.1%	0.0%		
Humber Estuary SAC	20	0.069	0.019	0.3%	0.1%		

#### Drax Bioenergy with Carbon Capture and Storage

	Annual Mean SO₂ concentration (µg/m³)						
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Humber Estuary SPA/SSSI	20	0.069	0.019	0.3%	0.1%		
Breighton Meadows SSSI	20	0.073	0.020	0.4%	0.1%		
Eskamhorn Meadows SSSI	20	0.021	0.006	0.1%	0.0%		
Derwent Ings SSSI	20	0.059	0.015	0.3%	0.1%		
Went Ings SSSI	20	0.023	0.007	0.1%	0.0%		
Barn Hill Meadows SSSI	20	0.072	0.020	0.4%	0.1%		
Burr Closes SSSI	20	0.027	0.008	0.1%	0.0%		
Common Plantation SINC	20	0.008	0.002	0.0%	0.0%		
Disused Railway Embankment SINC	20	0.019	0.005	0.1%	0.0%		
Barmby-on-the-Marsh LWS	20	0.036	0.011	0.2%	0.1%		
Brockholes SINC	20	0.009	0.003	0.0%	0.0%		
Meadow East of Orchard Farm SINC	20	0.004	0.001	0.0%	0.0%		
Barmby Pond LWS	20	0.058	0.017	0.3%	0.1%		
Cobble Croft Wood SINC	20	0.013	0.004	0.1%	0.0%		
Hagg Green Lane SINC	20	0.047	0.012	0.2%	0.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	20	0.013	0.004	0.1%	0.0%		
		Env. Agency	Screening Criterion (as % of CL)				

## Table 1.11 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Including Mitigation)

	Annual Nitrogen Deposition Rate (kgN/ha/yr)									
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation					
River Derwent SAC	15	0.054	0.044	0.4%	0.3%					
Thorne Moor SAC	5	0.025	0.021	0.5%	0.4%					

#### Drax Bioenergy with Carbon Capture and Storage

		Annual	Nitrogen Deposition Rate (k	gN/ha/yr)	
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
Thorne Moor SPA	10	0.025	0.021	0.2%	0.2%
Thorne Moor SSSI	5	0.025	0.021	0.5%	0.4%
Lower Derwent SAC	20	0.055	0.045	0.3%	0.2%
Lower Derwent SPA	20	0.055	0.045	0.3%	0.2%
Skipwith Common SAC	10	0.019	0.016	0.2%	0.2%
Skipwith Common SSSI	10	0.019	0.016	0.2%	0.2%
Humber Estuary SAC	20	0.034	0.041	0.2%	0.2%
Humber Estuary SPA/SSSI	20	0.034	0.041	0.2%	0.2%
Breighton Meadows SSSI	20	0.055	0.045	0.3%	0.2%
Eskamhorn Meadows SSSI	10	0.016	0.012	0.2%	0.1%
Derwent Ings SSSI	20	0.043	0.037	0.2%	0.2%
Went Ings SSSI	15	0.017	0.014	0.1%	0.1%
Barn Hill Meadows SSSI	20	0.039	0.041	0.2%	0.2%
Burr Closes SSSI	20	0.020	0.016	0.1%	0.1%
Common Plantation SINC	10	0.010	0.007	0.1%	0.1%
Disused Railway Embankment SINC	10	0.022	0.016	0.2%	0.2%
Barmby-on-the-Marsh LWS	10	0.043	0.032	0.4%	0.3%
Brockholes SINC	10	0.007	0.005	0.1%	0.0%
Meadow East of Orchard Farm SINC	20	0.003	0.002	0.0%	0.0%
Barmby Pond LWS	10	0.044	0.033	0.4%	0.3%
Cobble Croft Wood SINC	10	0.015	0.011	0.2%	0.1%
Hagg Green Lane SINC	10	0.058	0.046	0.6%	0.5%

		Annual	Nitrogen Deposition Rate (k	gN/ha/yr)	
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
Sand Pitt Wood & Barffs Close Plantation SINC	10	0.016	0.011	0.2%	0.1%
Barlow Common LNR	10	0.010	0.007	0.1%	0.1%
		Env. Agency Scree	ening Criterion (as % of CL)	19	%

## Table 1.12 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Including Mitigation)

	Critical LoadMitigation0.4620.0060.4620.0064.8560.0130.8020.0050.8020.0054.8560.0131.9980.0044.8560.0102.0080.0044.8560.0101.2480.005		Annual Acid Deposition Rate	(keq/ha/yr)	
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
Thorne Moor SAC	0.462	0.006	0.003	1.3%	0.6%
Thorne Moor SSSI	0.462	0.006	0.003	1.3%	0.6%
Lower Derwent SAC	4.856	0.013	0.006	0.3%	0.1%
Skipwith Common SAC	0.802	0.005	0.002	0.6%	0.3%
Skipwith Common SSSI	0.802	0.005	0.002	0.6%	0.3%
Breighton Meadows SSSI	4.856	0.013	0.006	0.3%	0.1%
Eskamhorn Meadows SSSI	1.998	0.004	0.002	0.2%	0.1%
Derwent Ings SSSI	4.856	0.010	0.005	0.2%	0.1%
Went Ings SSSI	2.008	0.004	0.002	0.2%	0.1%
Barn Hill Meadows SSSI	4.856	0.010	0.006	0.2%	0.1%
Burr Closes SSSI	1.248	0.005	0.002	0.4%	0.2%
		Env. Agency S	Screening Criterion (as % of CL)		1%

#### **CUMULATIVE IMPACTS (WITH PROPOSED SCHEME & OTHER PROJECTS)**

1.1.9. Results pertaining to the cumulative impacts, both before mitigation ("No Mit<sup>n</sup>") and after mitigation ("Mitig<sup>n</sup>") associated With Proposed Scheme Mitigation is applied (as detailed in **Section 6.10 of Chapter 6 (Air Quality)**) are presented in **Tables 1.13 to 1.18**.

	Annual Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )										
Receptor	Critical Level	Max Cumulat	tive PC Impact	Max PC Impa	ct as % of CL	Max Cumu	lative PEC	Max PEC a	s % of CL		
	Chical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>								
River Derwent SAC	30	0.675	0.647	2.3%	2.2%	12.63	12.60	42.1%	42.0%		
Thorne Moor SAC/SPA/SSSI	30	0.348	0.336	1.2%	1.1%	13.58	13.57	45.3%	45.2%		
Lower Derwent SAC	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%		
Lower Derwent SPA	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%		
Skipwith Common SAC	30	0.504	0.495	1.7%	1.7%	10.29	10.28	34.3%	34.3%		
Skipwith Common SSSI	30	0.504	0.495	1.7%	1.7%	10.29	10.28	34.3%	34.3%		
Humber Estuary SAC	30	0.498	0.474	1.7%	1.6%	19.77	19.74	65.9%	65.8%		
Humber Estuary SPA/SSSI	30	0.498	0.474	1.7%	1.6%	19.77	19.74	65.9%	65.8%		
Breighton Meadows SSSI	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%		
Eskamhorn Meadows SSSI	30	0.488	0.483	1.6%	1.6%	11.84	11.83	39.5%	39.4%		
Derwent Ings SSSI	30	0.640	0.619	2.1%	2.1%	10.49	10.47	35.0%	34.9%		
Went Ings SSSI	30	0.303	0.296	1.0%	1.0%	12.40	12.40	41.3%	41.3%		
Barn Hill Meadows SSSI	30	0.557	0.527	1.9%	1.8%	13.48	13.45	44.9%	44.8%		
Burr Closes SSSI	30	0.283	0.276	0.9%	0.9%	10.83	10.82	36.1%	36.1%		
Common Plantation SINC	30	0.728	0.723	2.4%	2.4%	12.16	12.15	40.5%	40.5%		
Disused Railway Embankment SINC	30	0.558	0.546	1.9%	1.8%	11.32	11.31	37.7%	37.7%		
Barmby-on-the-Marsh LWS	30	0.575	0.555	1.9%	1.8%	11.07	11.05	36.9%	36.8%		
Brockholes SINC	30	0.480	0.476	1.6%	1.6%	11.70	11.70	39.0%	39.0%		
Meadow East of Orchard Farm SINC	30	0.701	0.699	2.3%	2.3%	11.53	11.53	38.4%	38.4%		
Barmby Pond LWS	30	0.611	0.585	2.0%	2.0%	10.59	10.57	35.3%	35.2%		
Cobble Croft Wood SINC	30	0.697	0.690	2.3%	2.3%	12.32	12.31	41.1%	41.0%		
Hagg Green Lane SINC	30	0.704	0.684	2.3%	2.3%	11.67	11.65	38.9%	38.8%		

Drax Bioenergy with Carbon Capture and Storage

				Annual Mean N	IO <sub>x</sub> concentrat	tion (µg/m³)								
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as % of CL						
		No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>					
Sand Pitt Wood & Barffs Close Plantation SINC	30	0.798	0.790	2.7%	2.6%	12.23	12.22	40.8%	40.7%					
Env. Agency Screening	Criterion (as %	of CL)	•	1%	6			709	6					

## Table 1.14 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Daily Mean NO<sub>x</sub>

		•		Daily Mean N	O <sub>x</sub> concentrat	tion (µg/m³)			
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	ct as % of CL	Max Cumul	ative PEC	Max PEC a	as % of CL
	Critical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	41.9%         43.2%         36.0%         36.0%         36.0%         33.0%         33.0%         59.4%         36.0%         36.0%         33.0%         35.5%	Mitig <sup>n</sup>
River Derwent SAC	75	5.459	5.078	7.3%	6.8%	31.44	31.24	41.9%	41.7%
Thorne Moor SAC/SPA/SSSI	75	4.336	4.141	5.8%	5.5%	32.43	32.34	43.2%	43.1%
Lower Derwent SAC	75	4.382	4.199	5.8%	5.6%	27.00	26.90	36.0%	35.9%
Lower Derwent SPA	75	4.382	4.199	5.8%	5.6%	27.00	26.90	36.0%	35.9%
Skipwith Common SAC	75	3.932	3.615	5.2%	4.8%	24.75	24.66	33.0%	32.9%
Skipwith Common SSSI	75	3.932	3.615	5.2%	4.8%	24.75	24.66	33.0%	32.9%
Humber Estuary SAC	75	4.609	4.365	6.1%	5.8%	44.57	44.45	59.4%	59.3%
Humber Estuary SPA/SSSI	75	4.609	4.365	6.1%	5.8%	44.57	44.45	59.4%	59.3%
Breighton Meadows SSSI	75	4.382	4.198	5.8%	5.6%	27.00	26.89	36.0%	35.9%
Eskamhorn Meadows SSSI	75	6.777	6.347	9.0%	8.5%	30.40	29.99	40.5%	40.0%
Derwent Ings SSSI	75	4.234	4.158	5.6%	5.5%	26.63	26.47	35.5%	35.3%
Went Ings SSSI	75	5.111	5.040	6.8%	6.7%	30.27	30.14	40.4%	40.2%
Barn Hill Meadows SSSI	75	4.591	4.478	6.1%	6.0%	32.25	32.14	43.0%	42.9%
Burr Closes SSSI	75	3.736	3.537	5.0%	4.7%	25.99	25.95	34.7%	34.6%
Common Plantation SINC	75	7.240	7.596	9.7%	10.1%	30.30	30.65	40.4%	40.9%
Disused Railway Embankment SINC	75	6.035	5.779	8.0%	7.7%	27.93	27.62	37.2%	36.8%

#### Drax Bioenergy with Carbon Capture and Storage

				Daily Mean N	Ox concentrat	ion (µg/m³)			
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as % of CL	
	Critical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>						
Barmby-on-the-Marsh LWS	75	6.310	5.938	8.4%	7.9%	28.32	27.95	37.8%	37.3%
Brockholes SINC	75	5.892	5.440	7.9%	7.3%	28.71	28.26	38.3%	37.7%
Meadow East of Orchard Farm SINC	75	5.436	5.702	7.2%	7.6%	27.18	27.44	36.2%	36.6%
Barmby Pond LWS	75	5.189	4.763	6.9%	6.4%	27.29	27.09	36.4%	36.1%
Cobble Croft Wood SINC	75	7.028	6.706	9.4%	8.9%	31.22	30.90	41.6%	41.2%
Hagg Green Lane SINC	75	5.261	4.843	7.0%	6.5%	29.27	28.95	39.0%	38.6%
Sand Pitt Wood & Barffs Close Plantation SINC	75	7.720	7.948	10.3%	10.6%	31.06	31.02	41.4%	41.4%
	Env. Agency So	creening Criterio	on (as % of CL)	109	%				

## Table 1.15 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean NH3

				Annual Mean	NH₃ concentra	ation (µg/m³)			
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC	as % of CL
	Critical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>						
River Derwent SAC	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Thorne Moor SAC/SPA/SSSI	1	0.006	0.006	0.6%	0.6%	2.60	2.60	259.8%	259.8%
Lower Derwent SAC	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Lower Derwent SPA	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Skipwith Common SAC	1	0.003	0.003	0.3%	0.3%	2.58	2.58	258.4%	258.4%
Skipwith Common SSSI	1	0.003	0.003	0.3%	0.3%	2.58	2.58	258.4%	258.4%
Humber Estuary SAC	3	0.006	0.007	0.2%	0.2%	3.59	3.59	119.6%	119.7%
Humber Estuary SPA/SSSI	3	0.006	0.007	0.2%	0.2%	3.59	3.59	119.6%	119.7%
Breighton Meadows SSSI	3	0.008	0.007	0.3%	0.2%	3.09	3.09	103.0%	103.0%
Eskamhorn Meadows SSSI	3	0.003	0.003	0.1%	0.1%	2.40	2.40	80.1%	80.1%

#### Drax Bioenergy with Carbon Capture and Storage

				Annual Mean	NH <sub>3</sub> concentra	ation (µg/m³)			
Receptor		Max Cumulat	ive PC Impact	Max PC Impac	ct as % of CL	Max Cumul	lative PEC	Max PEC	as % of CL
	Critical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mitn           152.7%           78.5%           77.6%           83.5%           77.7%           76.1%           76.1%           76.3%           77.7%           103.2%           77.7%	Mitig <sup>n</sup>
Derwent Ings SSSI	3	0.006	0.005	0.2%	0.2%	4.58	4.58	152.7%	152.6%
Went Ings SSSI	3	0.004	0.004	0.1%	0.1%	2.36	2.35	78.5%	78.5%
Barn Hill Meadows SSSI	3	0.006	0.007	0.2%	0.2%	2.33	2.33	77.6%	77.7%
Burr Closes SSSI	3	0.003	0.003	0.1%	0.1%	2.50	2.50	83.5%	83.5%
Common Plantation SINC	3	0.002	0.001	0.1%	0.0%	2.33	2.33	77.7%	77.7%
Disused Railway Embankment SINC	3	0.003	0.002	0.1%	0.1%	2.28	2.28	76.1%	76.1%
Barmby-on-the-Marsh LWS	3	0.004	0.004	0.1%	0.1%	2.29	2.28	76.2%	76.1%
Brockholes SINC	3	0.002	0.002	0.1%	0.1%	2.28	2.28	76.1%	76.1%
Meadow East of Orchard Farm SINC	3	0.001	0.001	0.0%	0.0%	2.33	2.33	77.7%	77.7%
Barmby Pond LWS	3	0.007	0.005	0.2%	0.2%	2.29	2.29	76.3%	76.2%
Cobble Croft Wood SINC	3	0.002	0.002	0.1%	0.1%	2.33	2.33	77.7%	77.7%
Hagg Green Lane SINC	3	0.005	0.004	0.2%	0.1%	3.10	3.10	103.2%	103.2%
Sand Pitt Wood & Barffs Close Plantation SINC	3	0.002	0.002	0.1%	0.1%	2.33	2.33	77.7%	77.7%
	Env. Agency Se	creening Criterie	on (as % of CL)	1%	6			70%	

## Table 1.16 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean SO<sub>2</sub>

	Annual Mean SO <sub>2</sub> concentration (µg/m <sup>3</sup> )										
Receptor		Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as % of			
	Critical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>								
River Derwent SAC	20	0.074	0.022	0.4%	0.1%	4.04	3.98	20.2%	19.9%		
Thorne Moor SAC/SPA/SSSI	20	0.041	0.017	0.2%	0.1%	1.40	1.38	7.0%	6.9%		
Lower Derwent SAC	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%		
Lower Derwent SPA	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%		

#### Drax Bioenergy with Carbon Capture and Storage

			٩	nnual Mean SC	2 concentratio	n (µg/m³)			
Receptor	Critical Laval	Max Cumulat	ive PC Impact	Max PC Impa	ct as % of CL	Max Cumul	ative PEC	Max PEC as	% of CL
	Critical Level	No Mit <sup>n</sup>	Mitig <sup>n</sup>						
Skipwith Common SAC	20	0.026	0.008	0.1%	0.0%	1.46	1.44	7.3%	7.2%
Skipwith Common SSSI	20	0.026	0.008	0.1%	0.0%	1.46	1.44	7.3%	7.2%
Humber Estuary SAC	20	0.073	0.023	0.4%	0.1%	7.59	7.54	38.0%	37.7%
Humber Estuary SPA/SSSI	20	0.073	0.023	0.4%	0.1%	7.59	7.54	38.0%	37.7%
Breighton Meadows SSSI	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%
Eskamhorn Meadows SSSI	20	0.024	0.009	0.1%	0.0%	1.32	1.30	6.6%	6.5%
Derwent Ings SSSI	20	0.061	0.017	0.3%	0.1%	1.78	1.74	8.9%	8.7%
Went Ings SSSI	20	0.028	0.011	0.1%	0.1%	1.35	1.33	6.7%	6.7%
Barn Hill Meadows SSSI	20	0.075	0.024	0.4%	0.1%	1.91	1.86	9.6%	9.3%
Burr Closes SSSI	20	0.028	0.009	0.1%	0.0%	1.27	1.25	6.3%	6.3%
Common Plantation SINC	20	0.009	0.004	0.0%	0.0%	1.45	1.44	7.3%	7.2%
Disused Railway Embankment SINC	20	0.021	0.008	0.1%	0.0%	1.34	1.33	6.7%	6.7%
Barmby-on-the-Marsh LWS	20	0.038	0.013	0.2%	0.1%	1.37	1.34	6.8%	6.7%
Brockholes SINC	20	0.011	0.005	0.1%	0.0%	1.33	1.33	6.7%	6.6%
Meadow East of Orchard Farm SINC	20	0.006	0.003	0.0%	0.0%	1.45	1.44	7.2%	7.2%
Barmby Pond LWS	20	0.060	0.019	0.3%	0.1%	1.40	1.36	7.0%	6.8%
Cobble Croft Wood SINC	20	0.014	0.005	0.1%	0.0%	1.46	1.45	7.3%	7.2%
Hagg Green Lane SINC	20	0.049	0.014	0.2%	0.1%	1.50	1.47	7.5%	7.3%
Sand Pitt Wood & Barffs Close Plantation SINC	20	0.014	0.005	0.1%	0.0%	1.46	1.45	7.3%	7.2%
	Env. Agency S	creening Criteri	on (as % of CL)	19	/₀			70%	

Table 1.17 - Modelled Maximum Cumulative Impa	cts at Ecological Receptors – Annual Nitrogen Deposition Rate

		Annual Nitrogen Deposition Rate (kgN/ha/yr)										
Receptor	Critical Load	Max Cumu	lative PC Impact	Max PC Imp	pact as % of CL	Max Cur	nulative PEC	Max PE	C as % of CL			
	Critical Load	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>			
River Derwent SAC	15	0.106	0.098	0.7%	0.7%	30.35	30.34	202.3%	202.3%			
Thorne Moor SAC	5	0.061	0.063	1.2%	1.3%	21.38	21.38	427.6%	427.6%			
Thorne Moor SPA	10	0.061	0.063	0.6%	0.6%	21.38	21.38	213.8%	213.8%			
Thorne Moor SSSI	5	0.061	0.063	1.2%	1.3%	21.38	21.38	427.6%	427.6%			
Lower Derwent SAC	20	0.105	0.098	0.5%	0.5%	30.35	30.34	151.8%	151.7%			
Lower Derwent SPA	20	0.105	0.098	0.5%	0.5%	30.35	30.34	151.8%	151.7%			
Skipwith Common SAC	10	0.067	0.064	0.7%	0.6%	21.19	21.19	211.9%	211.9%			
Skipwith Common SSSI	10	0.067	0.064	0.7%	0.6%	21.19	21.19	211.9%	211.9%			
Humber Estuary SAC	20	0.073	0.086	0.4%	0.4%	28.96	28.98	144.8%	144.9%			
Humber Estuary SPA/SSSI	20	0.073	0.086	0.4%	0.4%	28.96	28.98	144.8%	144.9%			
Breighton Meadows SSSI	20	0.105	0.098	0.5%	0.5%	23.63	23.62	118.2%	118.1%			
Eskamhorn Meadows SSSI	10	0.058	0.061	0.6%	0.6%	20.01	20.02	200.1%	200.2%			
Derwent Ings SSSI	20	0.096	0.091	0.5%	0.5%	30.34	30.34	151.7%	151.7%			
Went Ings SSSI	15	0.049	0.046	0.3%	0.3%	19.44	19.43	129.6%	129.6%			
Barn Hill Meadows SSSI	20	0.082	0.088	0.4%	0.4%	20.52	20.54	102.6%	102.7%			
Burr Closes SSSI	20	0.045	0.041	0.2%	0.2%	20.69	20.69	103.5%	103.4%			
Common Plantation SINC	10	0.158	0.155	1.6%	1.5%	33.90	33.90	339.0%	339.0%			
Disused Railway Embankment SINC	10	0.126	0.129	1.3%	1.3%	33.45	33.45	334.5%	334.5%			
Barmby-on-the-Marsh LWS	10	0.141	0.141	1.4%	1.4%	33.47	33.47	334.7%	334.7%			
Brockholes SINC	10	0.056	0.057	0.6%	0.6%	19.80	19.80	198.0%	198.0%			
Meadow East of Orchard Farm SINC	20	0.075	0.076	0.4%	0.4%	19.96	19.96	99.8%	99.8%			
Barmby Pond LWS	10	0.095	0.085	1.0%	0.9%	19.85	19.84	198.5%	198.4%			

		Annual Nitrogen Deposition Rate (kgN/ha/yr)									
Receptor	Critical Load	Max Cumulative PC Impact		Max PC Impact as % of CL		Max Cumulative PEC		Max PEC as % of CL			
		No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	Mitig <sup>n</sup>		
Cobble Croft Wood SINC	10	0.155	0.151	1.6%	1.5%	33.90	33.89	339.0%	338.9%		
Hagg Green Lane SINC	10	0.176	0.173	1.8%	1.7%	40.94	40.93	409.4%	409.3%		
Sand Pitt Wood & Barffs Close Plantation SINC	10	0.176	0.171	1.8%	1.7%	33.92	33.91	339.2%	339.1%		
Barlow Common LNR	10	0.173	0.170	1.7%	1.7%	33.91	33.91	339.1%	339.1%		
	Env. Agency Screening Criterion (as % of CL)			1%				70%			

## Table 1.18 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Acid Deposition Rate

				Annual Aci	d Deposition Rat	te (keq/ha/yr)				
Receptor	Critical Load	Max Cumu	Max Cumulative PC Impact		Max PC Impact as % of CL		Max Cumulative PEC		Max PEC as % of CL	
	Critical Load	No Mit <sup>n</sup>	Mitig <sup>n</sup>	No Mit <sup>n</sup>	o Mitign         Mitign         No Mitn         Mitign         I           .1%         1.5%         1.74         1.74         3           .1%         1.5%         1.75         1.74         3           .1%         1.5%         1.75         1.74         3           .4%         0.2%         2.43         2.42         5           .0%         0.7%         1.74         1.73         2           .0%         0.7%         1.74         1.73         2           .0%         0.7%         1.74         1.73         2           .0%         0.2%         1.95         1.94         4           .3%         0.2%         2.42         2.42         4           .3%         0.2%         1.60         1.64         5           .3%         0.2%         1.60         1.60         7           .3%         0.2%         1.71         1.70         3	No Mit <sup>n</sup>	Mitig <sup>n</sup>			
Thorne Moor SAC	0.4623	0.010	0.007	2.1%	1.5%	1.74	1.74	375.8%	377.1%	
Thorne Moor SSSI	0.4623	0.010	0.007	2.1%	1.5%	1.75	1.74	377.8%	377.1%	
Lower Derwent SAC	4.856	0.017	0.010	0.4%	0.2%	2.43	2.42	50.0%	49.8%	
Skipwith Common SAC	0.802	0.008	0.006	1.0%	0.7%	1.74	1.73	216.4%	216.1%	
Skipwith Common SSSI	0.802	0.008	0.006	1.0%	0.7%	1.74	1.73	216.4%	216.1%	
Breighton Meadows SSSI	0.643	0.017	0.010	0.4%	0.2%	1.95	1.94	40.1%	40.0%	
Eskamhorn Meadows SSSI	1.998	0.007	0.005	0.3%	0.3%	1.64	1.64	82.3%	82.2%	
Derwent Ings SSSI	0.643	0.014	0.009	0.3%	0.2%	2.42	2.42	49.9%	49.8%	
Went Ings SSSI	2.008	0.007	0.005	0.4%	0.2%	1.60	1.60	79.7%	79.6%	
Barn Hill Meadows SSSI	4.856	0.014	0.010	0.3%	0.2%	1.71	1.70	35.2%	35.1%	
Burr Closes SSSI	1.248	0.007	0.004	0.5%	0.3%	1.69	1.69	135.3%	135.1%	
	Env. Agency So	creening Crite	erion (as % of CL)		1%				70%	

#### SENSITIVITY TEST: WORST CASE EMISSIONS PROFILE

- 1.1.10. Results pertaining to the worst-case emissions profile sensitivity test are presented in **Tables 1.19 to 1.24**, based on emissions from the Proposed Scheme alone.
- 1.1.11. For all pollutant concentrations and deposition rates, it is evident that the modelled maximum PC impacts attributed to the with Proposed Scheme scenario are lower at all receptors relative to the core model scenarios. This is a function of all four biomass units in the Baseline scenario switching from 'mid-merit' operation (full load for 4,000 hours per year) to continuous operation (full load for 8,760 hours per year), resulting in more pollutants being emitted and thus more pronounced changes (increases) in concentrations / deposition rates relative to the With Proposed Scheme scenario. In the With Proposed Scheme scenario, operation changes from 'mid-merit' to continuous full load at the two non-BECCS biomass units only (BECCS units already assumed to operate at continuous full load in core modelling scenario), meaning the changes (increases and decreases) in concentrations / deposition rates are relatively small compared to the Baseline.
- 1.1.12. As a consequence, the maximum modelled impacts of the Proposed Scheme decrease at all receptors under the worst-case emissions profile scenario relative to the core modelling. Whilst some modelled maximum PEC concentrations do increase under worst case emissions in both the Baseline and With Proposed Scheme scenarios, there are no material changes relative to the core modelling equivalents, meaning that the respective assessment significance criteria are not exceeded.
- 1.1.13. The results confirm that the assessment of likely significant effects reported in Chapter 6 (Air Quality) is not affected when considering the worst-case emissions profiles in both the Baseline and With Proposed Scheme scenarios. Given that the modelled maximum impacts are lower under a worst-case emissions profile, there was no need to repeat the test in relation to cumulative impacts, as the core modelling results for the cumulative scenarios represent the most conservative results in terms of potential impacts.

Table 1.19 - Modelled Maximum Operational Impacts	at Ecological Receptors – Annual Mean NO <sub>x</sub> (Worst Case Emissions Profile)

	Annual Mean NO <sub>x</sub> concentration (μg/m <sup>3</sup> )									
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL				
River Derwent SAC	30	11.91	0.055	0.2%	12.10	40.3%				
Thorne Moor SAC/SPA/SSSI	30	13.21	0.026	0.1%	13.33	44.4%				
Lower Derwent SAC	30	9.92	0.058	0.2%	10.11	33.7%				
Lower Derwent SPA	30	9.92	0.058	0.2%	10.11	33.7%				
Skipwith Common SAC	30	9.76	0.023	0.1%	9.84	32.8%				
Skipwith Common SSSI	30	9.76	0.023	0.1%	9.84	32.8%				
Humber Estuary SAC	30	19.22	0.056	0.2%	19.40	64.7%				
Humber Estuary SPA/SSSI	30	19.22	0.056	0.2%	19.40	64.7%				
Breighton Meadows SSSI	30	9.92	0.058	0.2%	10.11	33.7%				
Eskamhorn Meadows SSSI	30	11.35	0.012	0.0%	11.38	37.9%				
Derwent Ings SSSI	30	9.80	0.048	0.2%	9.98	33.3%				
Went Ings SSSI	30	12.09	0.018	0.1%	12.15	40.5%				
Barn Hill Meadows SSSI	30	12.89	0.055	0.2%	13.04	43.5%				
Burr Closes SSSI	30	10.53	0.020	0.1%	10.59	35.3%				
Common Plantation SINC	30	11.43	0.004	0.0%	11.44	38.1%				
Disused Railway Embankment SINC	30	10.76	0.009	0.0%	10.78	35.9%				
Barmby-on-the-Marsh LWS	30	10.48	0.021	0.1%	10.53	35.1%				
Brockholes SINC	30	11.22	0.004	0.0%	11.23	37.4%				
Meadow East of Orchard Farm SINC	30	10.83	0.002	0.0%	10.83	36.1%				
Barmby Pond LWS	30	9.96	0.036	0.1%	10.06	33.5%				
Cobble Croft Wood SINC	30	11.62	0.007	0.0%	11.64	38.8%				
Hagg Green Lane SINC	30	10.93	0.034	0.1%	11.05	36.8%				

Receptor	Annual Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )							
	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Sand Pitt Wood & Barffs Close Plantation SINC	30	11.43	0.007	0.0%	11.45	38.2%		
	Env. Agency Screening Criterion (as % of CL)			1%		70%		

## Table 1.20 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO<sub>x</sub> (Worst Case Emissions Profile)

	Daily Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )									
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL				
River Derwent SAC	75	23.82	1.787	2.4%	27.96	37.3%				
Thorne Moor SAC/SPA/SSSI	75	26.42	0.893	1.2%	29.27	39.0%				
Lower Derwent SAC	75	19.84	0.893	1.2%	23.75	31.7%				
Lower Derwent SPA	75	19.84	0.893	1.2%	23.75	31.7%				
Skipwith Common SAC	75	19.52	0.646	0.9%	21.71	28.9%				
Skipwith Common SSSI	75	19.52	0.646	0.9%	21.71	28.9%				
Humber Estuary SAC	75	38.44	1.200	1.6%	41.45	55.3%				
Humber Estuary SPA/SSSI	75	38.44	1.200	1.6%	41.45	55.3%				
Breighton Meadows SSSI	75	19.84	0.893	1.2%	23.75	31.7%				
Eskamhorn Meadows SSSI	75	22.70	1.912	2.5%	25.59	34.1%				
Derwent Ings SSSI	75	19.60	0.831	1.1%	23.37	31.2%				
Went Ings SSSI	75	24.18	0.971	1.3%	26.60	35.5%				
Barn Hill Meadows SSSI	75	25.78	0.962	1.3%	28.30	37.7%				
Burr Closes SSSI	75	21.06	0.761	1.0%	22.99	30.7%				
Common Plantation SINC	75	22.86	1.038	1.4%	24.19	32.3%				
Disused Railway Embankment SINC	75	21.52	1.184	1.6%	23.07	30.8%				
Barmby-on-the-Marsh LWS	75	20.96	1.621	2.2%	23.44	31.3%				

#### Drax Bioenergy with Carbon Capture and Storage

Environmental Statement - Volume 3 - Appendix 6.5

Page 29 of 34

	Daily Mean NO <sub>x</sub> concentration (µg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Brockholes SINC	75	22.44	1.623	2.2%	24.44	32.6%		
Meadow East of Orchard Farm SINC	75	21.66	0.554	0.7%	22.33	29.8%		
Barmby Pond LWS	75	19.92	1.577	2.1%	23.29	31.1%		
Cobble Croft Wood SINC	75	23.24	1.176	1.6%	25.28	33.7%		
Hagg Green Lane SINC	75	21.86	1.248	1.7%	25.22	33.6%		
Sand Pitt Wood & Barffs Close Plantation SINC	75	22.86	1.630	2.2%	24.97	33.3%		
	10%							

## Table 1.21 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH<sub>3</sub> (Worst Case Emissions Profile)

		•	Annual Mean NH	<sup>3</sup> concentration (μg/m <sup>3</sup> )		
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
River Derwent SAC	3	4.57	0.002	0.1%	4.58	152.6%
Thorne Moor SAC/SPA/SSSI	1	2.59	0.001	0.1%	2.60	259.6%
Lower Derwent SAC	3	4.57	0.002	0.1%	4.58	152.6%
Lower Derwent SPA	3	4.57	0.002	0.1%	4.58	152.6%
Skipwith Common SAC	1	2.58	0.001	0.1%	2.58	258.4%
Skipwith Common SSSI	1	2.58	0.001	0.1%	2.58	258.4%
Humber Estuary SAC	3	3.58	0.002	0.1%	3.59	119.6%
Humber Estuary SPA/SSSI	3	3.58	0.002	0.1%	3.59	119.6%
Breighton Meadows SSSI	3	3.08	0.002	0.1%	3.09	103.0%
Eskamhorn Meadows SSSI	3	2.40	0.001	0.0%	2.40	80.1%
Derwent Ings SSSI	3	4.57	0.002	0.1%	4.58	152.6%
Went Ings SSSI	3	2.35	0.001	0.0%	2.35	78.4%

#### Drax Bioenergy with Carbon Capture and Storage

		Annual Mean NH <sub>3</sub> concentration (μg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Barn Hill Meadows SSSI	3	2.32	0.002	0.1%	2.33	77.6%			
Burr Closes SSSI	3	2.50	0.001	0.0%	2.50	83.4%			
Common Plantation SINC	3	2.33	0.000	0.0%	2.33	77.7%			
Disused Railway Embankment SINC	1	2.28	0.000	0.0%	2.28	76.0%			
Barmby-on-the-Marsh LWS	3	2.28	0.001	0.0%	2.28	76.1%			
Brockholes SINC	3	2.28	0.000	0.0%	2.28	76.0%			
Meadow East of Orchard Farm SINC	1	2.33	0.000	0.0%	2.33	77.7%			
Barmby Pond LWS	1	2.28	0.001	0.0%	2.29	76.2%			
Cobble Croft Wood SINC	3	2.33	0.000	0.0%	2.33	77.7%			
Hagg Green Lane SINC	3	3.09	0.001	0.0%	3.10	103.2%			
Sand Pitt Wood & Barffs Close Plantation SINC	3	2.33	0.000	0.0%	2.33	77.7%			
	1%		70%						

#### Table 1.22 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO<sub>2</sub> (Worst Case Emissions Profile)

	Annual Mean SO <sub>2</sub> concentration (µg/m <sup>3</sup> )							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	20	3.93	0.021	0.1%	4.03	20.1%		
Thorne Moor SAC/SPA/SSSI	20	1.34	0.009	0.0%	1.40	7.0%		
Lower Derwent SAC	20	1.70	0.021	0.1%	1.80	9.0%		
Lower Derwent SPA	20	1.70	0.021	0.1%	1.80	9.0%		
Skipwith Common SAC	20	1.42	0.008	0.0%	1.46	7.3%		
Skipwith Common SSSI	20	1.42	0.008	0.0%	1.46	7.3%		
Humber Estuary SAC	20	7.49	0.021	0.1%	7.58	37.9%		

#### Drax Bioenergy with Carbon Capture and Storage

Receptor	Annual Mean SO <sub>2</sub> concentration (μg/m <sup>3</sup> )								
	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Humber Estuary SPA/SSSI	20	7.49	0.021	0.1%	7.58	37.9%			
Breighton Meadows SSSI	20	1.70	0.021	0.1%	1.80	9.0%			
Eskamhorn Meadows SSSI	20	1.29	0.005	0.0%	1.31	6.5%			
Derwent Ings SSSI	20	1.69	0.016	0.1%	1.78	8.9%			
Went Ings SSSI	20	1.31	0.007	0.0%	1.34	6.7%			
Barn Hill Meadows SSSI	20	1.81	0.022	0.1%	1.89	9.4%			
Burr Closes SSSI	20	1.23	0.008	0.0%	1.26	6.3%			
Common Plantation SINC	20	1.44	0.002	0.0%	1.44	7.2%			
Disused Railway Embankment SINC	20	1.32	0.004	0.0%	1.33	6.7%			
Barmby-on-the-Marsh LWS	20	1.32	0.009	0.0%	1.35	6.7%			
Brockholes SINC	20	1.32	0.002	0.0%	1.32	6.6%			
Meadow East of Orchard Farm SINC	20	1.44	0.001	0.0%	1.44	7.2%			
Barmby Pond LWS	20	1.32	0.015	0.1%	1.37	6.9%			
Cobble Croft Wood SINC	20	1.44	0.003	0.0%	1.45	7.2%			
Hagg Green Lane SINC	20	1.43	0.013	0.1%	1.49	7.5%			
Sand Pitt Wood & Barffs Close Plantation SINC	20	1.44	0.003	0.0%	1.45	7.2%			
	1%		70%						

### Table 1.23 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Worst Case Emissions Profile)

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	15	30.22	0.017	0.1%	30.29	201.9%		
Thorne Moor SAC	5	21.31	0.008	0.2%	21.35	427.0%		

Becenter	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Thorne Moor SPA	10	21.31	0.008	0.1%	21.35	213.5%			
Thorne Moor SSSI	5	21.31	0.008	0.2%	21.35	427.0%			
Lower Derwent SAC	20	30.22	0.017	0.1%	30.29	151.5%			
Lower Derwent SPA	20	30.22	0.017	0.1%	30.29	151.5%			
Skipwith Common SAC	10	21.12	0.007	0.1%	21.14	211.4%			
Skipwith Common SSSI	10	21.12	0.007	0.1%	21.14	211.4%			
Humber Estuary SAC	20	28.87	0.017	0.1%	28.93	144.7%			
Humber Estuary SPA/SSSI	20	28.87	0.017	0.1%	28.93	144.7%			
Breighton Meadows SSSI	20	23.51	0.017	0.1%	23.57	117.9%			
Eskamhorn Meadows SSSI	10	19.95	0.004	0.0%	19.96	199.6%			
Derwent Ings SSSI	20	30.22	0.014	0.1%	30.29	151.4%			
Went Ings SSSI	15	19.38	0.005	0.0%	19.40	129.4%			
Barn Hill Meadows SSSI	20	20.43	0.017	0.1%	20.49	102.4%			
Burr Closes SSSI	20	20.64	0.006	0.0%	20.66	103.3%			
Common Plantation SINC	10	33.74	0.002	0.0%	33.75	337.5%			
Disused Railway Embankment SINC	10	33.32	0.005	0.1%	33.33	333.3%			
Barmby-on-the-Marsh LWS	10	33.32	0.011	0.1%	33.35	333.5%			
Brockholes SINC	10	19.74	0.001	0.0%	19.74	197.4%			
Meadow East of Orchard Farm SINC	20	19.88	0.001	0.0%	19.88	99.4%			
Barmby Pond LWS	10	19.74	0.012	0.1%	19.78	197.8%			
Cobble Croft Wood SINC	10	33.74	0.004	0.0%	33.75	337.5%			
Hagg Green Lane SINC	10	40.74	0.017	0.2%	40.81	408.1%			
Sand Pitt Wood & Barffs Close Plantation SINC	10	33.74	0.004	0.0%	33.75	337.5%			

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Barlow Common LNR	10	33.74	0.002	0.0%	33.75	337.5%			
Env. Agency Screening Criterion (as % of CL)			1%		70%				

## Table 1.24 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Worst Case Emissions Profile)

Receptor	Annual Acid Deposition Rate (keq/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Thorne Moor SAC	0.462	1.73	0.002	0.4%	1.74	377.2%			
Thorne Moor SSSI	0.462	1.73	0.002	0.4%	1.74	377.2%			
Lower Derwent SAC	4.856	2.40	0.004	0.1%	2.42	49.9%			
Skipwith Common SAC	0.802	1.73	0.002	0.2%	1.73	216.0%			
Skipwith Common SSSI	0.802	1.73	0.002	0.2%	1.73	216.0%			
Breighton Meadows SSSI	4.856	1.92	0.004	0.1%	1.94	40.0%			
Eskamhorn Meadows SSSI	1.998	1.64	0.001	0.0%	1.64	82.1%			
Derwent Ings SSSI	4.856	2.40	0.003	0.1%	2.42	49.8%			
Went Ings SSSI	2.008	1.59	0.001	0.1%	1.60	79.5%			
Barn Hill Meadows SSSI	4.856	1.69	0.004	0.1%	1.70	35.1%			
Burr Closes SSSI	1.248	1.68	0.001	0.1%	1.69	135.1%			
	1%		70%						