

West Burton Solar Project

Environmental Statement Appendix 9.2: Preliminary Ecological Appraisal

Prepared by: Clarkson & Woods
August 2021

PINS reference: EN010132
Document reference: APP/WB6.3.9.2
APFP Regulation 5(2)(a)



PRELIMINARY ECOLOGICAL APPRAISAL
WEST BURTON SOLAR PROJECT

carried out by



commissioned by

WEST BURTON SOLAR PROJECT LTD.


AUGUST 2021



CONTENTS

KEY ECOLOGICAL CONSTRAINTS AND OPPORTUNITIES.....	4
1 INTRODUCTION	6
1.2 Report Aims.....	6
1.3 Appraisal Scope and Limitations	6
1.4 Consultation	7
1.5 Site Description Summary	7
1.6 Surveys Carried out to Date	8
2 DESK STUDY	8
2.1 Local Planning Policy	8
2.2 Protected and Designated Sites	9
2.3 Ancient Woodland	11
2.4 Biodiversity Opportunities Mapping.....	11
3 HABITAT SURVEY	13
3.2 Common Habitat Constraints and Opportunities.....	13
3.3 West Burton 1 Habitat Assessment	20
3.4 West Burton 2 Habitat Assessment	21
3.5 West Burton 3 Habitat Assessment	24
4 SPECIES INFORMATION COLLATED TO DATE.....	27
4.2 Badgers.....	27
4.3 Bats.....	28
4.4 Otter.....	32
4.5 Water Vole.....	33
4.6 Dormouse	34
4.7 Great Crested Newts and Other Amphibians	34
4.8 Reptiles.....	37
4.9 Birds	38
4.10 Invertebrates	42
4.11 Other Protected Species and Species of Conservation Concern	42
5 FURTHER WORK AND NEXT STEPS	44
5.1 Recommended and Optional Further Surveys	44
5.2 Anticipated Reporting/Design Milestones.....	45
5.3 Construction and Landscape Environmental Management Plans (CEMP and LEMP)	45
5.4 Cumulative Impacts	46
5.5 Future Baseline and Decommissioning Effects.....	46
APPENDIX A: WILDLIFE LEGISLATION SUMMARY.....	47
APPENDIX B – SUMMARY OF METHODOLOGIES	52
APPENDIX C – DESIGNATED SITES MAPS.....	54
APPENDIX D – SPECIES RECORDS WITHIN 2KM OF WEST BURTON 1 (BROXHOLME)	57
APPENDIX E – SPECIES RECORDS WITHIN 2KM OF WEST BURTON 2 (INGLEBY).....	60
APPENDIX F – SPECIES RECORDS WITHIN 2KM OF WEST BURTON 3 (BELLWOOD AND BRAMPTON)	67
APPENDIX G: LOCAL PLANNING POLICY	75
APPENDIX H – PHASE 1 HABITATS MAPS.....	84



Project title:	West Burton Solar Project		
Document title:	Preliminary Ecological Appraisal	Project number:	7479
Client:	West Burton Project Ltd.	Author:	Harry Fox
Version 1	Final Report	Issued on:	August 2021
Quality Assurance	Checked by:	Approved by:	
	Tom Clarkson	Tom Clarkson	
			

The information, data and advice which has been prepared and provided is true, and has been prepared and provided in accordance with the Chartered Institute of Ecology and Environmental Management's (CIEEM) Code of Professional Conduct. We confirm that the opinions expressed are our true and professional bona fide opinions. This report and its contents remain the property of Clarkson and Woods Ltd. until payment has been made in full.



KEY ECOLOGICAL CONSTRAINTS AND OPPORTUNITIES

Item	Key Constraints	Key Opportunities
Local Policy: Central Lincolnshire Local Plan	<ul style="list-style-type: none"> • CLLP Policy LP20: Green Infrastructure (GI) Network Protection, integration, enhancement and creation of GI wherever possible. • CLLP Policy LP21: Protecting Biodiversity and Geodiversity • Protection, management and delivery of net gain for biodiversity, focusing on Habitats and Species of Principal Importance. 	<p>Policy compliance achievable through:</p> <ul style="list-style-type: none"> • Arable reversion to grassland. • Hedgerow planting, in-filling and positive aftercare. • Positive grassland management within adequate buffer zones. • Protection of hedgerows and watercourses during construction. • Strategic use of BOM in guiding habitat enhancements (West Burton 1-3). • Locally-appropriate tree planting in screening. Pre-emptive replacement of ash affected by dieback.
Local Biodiversity Action Plan: Key habitats and species	<ul style="list-style-type: none"> • Arable field margins • Hedgerows and hedgerow trees • Lowland meadows • Ponds, lakes and reservoirs • Rivers, canals and drains • Bats • Farmland birds • Newts • Water vole 	<p>Positive effects likely to arise on all features through:</p> <ul style="list-style-type: none"> • Adequate buffering of hedgerows, ditches and watercourses. • Blend of habitat enhancement options within buffer zones. • Selective grassland enhancement options within array. • Habitat feature provision for bats, reptiles, amphibians for birds as discussed individually.
Biodiversity Net Gain	<ul style="list-style-type: none"> • Recent amendments to the Environment Bill will extend obligation to deliver 10% net gain to NSIPs. • West Burton 1: Two fields in NW are included within BOM 'opportunity for creation'. • West Burton 2: Eastern half of Site included within BOM 'opportunity for creation' with one field 'ecological network'. • West Burton 3: Four fields in south are included within BOM 'opportunity for creation'. 	<p>High confidence in deliverability of BNG due to:</p> <ul style="list-style-type: none"> • Large scale reversion of arable to grassland. • Cost-effective positive management of field margin buffers. • Hedgerow enhancements and tree planting. • Discrete grassland habitat creation options. <p>Will require habitat mapping (pre-construction state mapping complete) and completion of Habitat Unit change using Defra Metric 3.0 using iterations of landscape proposals and habitat management plans. Siting should be influenced by inclusion within BOM</p>
Designated Sites	<ul style="list-style-type: none"> • Mr. Rose's Hay Meadow LWS and Torksey Grassland LWS – Located just off the southern boundary of West Burton 3. 	
Arable fields	<ul style="list-style-type: none"> • Only constraints relate to ground nesting birds. 	<p>BNG and Policy contribution can be maximised through adoption of sensitive grassland management (see Section 3.2):</p> <ul style="list-style-type: none"> • 'Shade cutting' rather than wholesale mowing • Conservation grazing rates and timings • Selective meadow restoration • 'Aftermath' grazing • Cut-and-collect rather than leaving arisings
Field Margins	<ul style="list-style-type: none"> • Habitat of Principal Importance (HPI) and on Lincs and Notts BAPs. • Some are species rich and broad on all sites. • Many on all Sites hold potential for reptiles. 	<ul style="list-style-type: none"> • Significant BOM overlap at West Burton 1-3. • River Till corridor a significant enhancement opportunity. • Semi-improved grassland in West Burton 2 stand to gain from conservation management.



Item	Key Constraints	Key Opportunities
Hedgerows	<ul style="list-style-type: none"> HPI and on Lincs and Notts BAPs. Most contain at least occasional mature trees. Abundant nesting birds – constraint to removal. Key habitat for bats. <p>Minimum buffer zones from hedgerow edge to security fence proposed:</p> <ul style="list-style-type: none"> Species-poor hedgerows or hedgerows without trees: 8m Species-rich hedgerows or hedgerows with trees: 10m 	<ul style="list-style-type: none"> In-filling or replanting defunct hedgerows New hedgerow planting along bare field boundaries Possible new hedgerows in strategic locations for maximum green infrastructure/connectivity benefit. Pre-emptive replacement of large number of ash-dieback affected trees.
Ditches and Watercourses	<ul style="list-style-type: none"> HPI (rivers) and on Lincs and Notts BAPs. Minimum buffer zones from banktop to security fence proposed of 8m up to 30m depending on significance. 	<ul style="list-style-type: none"> River Till and Toft Dyke corridor grassland mosaic enhancement – West Burton 1/4
Badgers	<p>This information has been removed to a confidential appendix to the Environmental Statement</p>	
Bats	<ul style="list-style-type: none"> Hedgerows and trees of moderate value while arable fields of low value. Potential for roosts within hedgerow trees and buildings. Potentially at risk of fragmentation. <p>Minimum buffer zones from feature edge to security fence proposed:</p> <ul style="list-style-type: none"> Ditches, species-poor hedgerows and hedgerows without trees: 8m Minor watercourses (streams, beckes), species-rich hedgerows and hedgerows with trees of low or negligible roost potential: 10m Woodland, in-field trees, hedgerows with trees of moderate or high roost potential: 20m Rivers, possible ancient woodland (West Burton 2) confirmed roosts in buildings or trees: 30m 	<ul style="list-style-type: none"> Grassland management (under array and at buffer zones) will significantly enhance foraging potential. Standalone and tree-mounted roosting features.
Otters and water voles	<ul style="list-style-type: none"> All Sites contained habitat of potential value to otters and water voles as well as local records. Buffering of ditches and watercourses to avoid disturbance and habitat damage. 	<ul style="list-style-type: none"> Periodic ditch and grassland margin maintenance. Deepening and wetting of ditches to improve connectivity.
Amphibians (incl. GCN) and Reptiles	<ul style="list-style-type: none"> Two ponds positive for GCN eDNA at West Burton 3. Potential for licensing constraints and adoption of precautionary methods within 250m of positive ponds. All Sites contained habitat suitable for reptiles and amphibians in hedgerows, watercourses and field margins. Precautions/supervision during any habitat clearance required. 	<ul style="list-style-type: none"> Selective deepening of on Site ponds to enhance their value. Construction of new ponds in locations suitable for linking known populations.
Birds	<ul style="list-style-type: none"> Significant numbers of skylark and other ground nesting birds at all Sites, particularly West Burton 3. Displacement of territories by solar array anticipated. Avoidance of disturbance and damage to nests during breeding season. 	<ul style="list-style-type: none"> Targeted management of field margin buffers and grassland under panels for birds such as quail, partridge and turtle dove as well as foraging skylark and yellow wagtail. Nesting and roosting boxes and standalone habitat features.
Invertebrates	<ul style="list-style-type: none"> Low to moderate habitat suitability for invertebrates limited to field margins, hedgerows and ditches/watercourses at all Sites. 	<ul style="list-style-type: none"> Targeted management of field margins to include scrub and ruderal vegetation mosaic.



1 INTRODUCTION

- 1.1.1 Clarkson and Woods Ltd. was commissioned by West Burton Solar Project Ltd. to carry out a Preliminary Ecological Appraisal across three parcels of land known as West Burton 1, 2 and 3 which are situated in the West Lindsey District of Central Lincolnshire. These parcels are referred to hereafter as 'the Sites', or individually as given above. Proposals are understood to be in an early design stage and comprise the development of an NSIP-scale solar park, containing solar energy production and storage components.
- 1.1.2 This Preliminary Ecological Appraisal discusses the results collected during an Extended Phase 1 Habitats Survey carried out in April and May 2021 by Clarkson and Woods Ltd, supplemented by partial datasets from breeding bird surveys, bat surveys and great crested newt eDNA surveys carried out in spring and summer 2021.

1.2 Report Aims

- 1.2.1 The aims of this report are:
- To describe the habitats present within the Sites and their potential to support protected or otherwise notable species and habitats capable of being material considerations within the planning process.
 - To set out the results of a desk study based on third party ecological records from the Site and its surroundings supplied by the Lincolnshire Environmental Records Centre (LERC) and Nottinghamshire Biological and Geological Record Centre (NBGRC) in the context of Local Planning Policy.
 - To outline any key potential ecological constraints to development of the Site.
 - To broadly discuss avoidance, mitigation or compensation measures likely to be required to minimise potential ecological impacts.
 - To identify where further surveys to establish baseline conditions or develop mitigation or compensatory measures may be required.
 - To identify where further consultation with statutory bodies, planning authorities or other key consultees would be advantageous to determine a robust and acceptable assessment scope.
 - To outline options for ecological enhancement and Biodiversity Net Gain and how they may be secured, managed and monitored.

1.3 Appraisal Scope and Limitations

- 1.3.1 The appraisal recorded habitat information from within the red line boundaries (the option land boundaries) only. However, a desk-based general assessment of the surrounding landscape was made, supported by extensive visual appraisal from public rights of way in the land immediately surrounding the Sites. This information has been factored into the appraisal of habitat suitability for certain species and advice on opportunities for Biodiversity Net Gain.
- 1.3.2 No appraisal of proposed cable routes is contained within this report.
- 1.3.3 To date, no consultation with statutory or non-statutory third parties has been carried out. Considering the potential for impacts upon a number of protected and notable species combined with the desired timescales applied to the project, it is recommended that the indicated scope and approach to further survey is consulted on with local authorities, their nature conservation consultees and Natural England at an early stage.
- 1.3.4 Under CIEEM guidelines, PEA reports are not considered suitable on their own for inclusion with an eventual DCO application. However, information has been provided below with a view to support and enhance the masterplanning process.
- 1.3.5 It is anticipated that the results of further detailed survey work will be reported separately in due course and will serve to underpin an eventual Preliminary Environmental Information Report and Environmental Impact Assessment.
- 1.3.6 Records obtained from LERC and NBGRC are not exhaustive or complete and an absence of records for a species does not preclude their possible presence.
- 1.3.7 The appraisal has been prepared by Harry Fox, an experienced ecologist, who is a full member of the Chartered Institute of Ecology and Environmental Management (CIEEM). The report has been subject to quality assurance review by appropriately experienced senior consultants who are full members of CIEEM.

1.3.8 Unless the client indicates to the contrary, information on the presence of species collected during the surveys will be passed on to LERC and NBGRC following submission of a planning application in order to augment their records for the area. This is in line with the CIEEM code of professional conduct¹.




Figure 1. Locations of the Proposed Development Sites West Burton 1-3

1.4 Consultation

1.4.1 The following statutory bodies will be consulted in due course:

- **Natural England** – Advisor assigned at onset of consultation. Paid-for Discretionary Advice Service available outside of statutory consultation process should this prove advantageous.
- **West Lindsey District Council** – No district ecology officer. Ecology issues dealt with by planning officer team with reference to Natural England Standing Advice. Therefore, pre-application consultation response likely to be very limited.
- **Lincolnshire County Council** – No County ecologist – ecology matters likely referred to Environmental Services Team and Wildlife Trusts/NE Standing Advice referenced.
- **Bassetlaw District Council** – No district ecology officer. BDC refer to Nottinghamshire Wildlife Trust to provide consultation opinion on their behalf.
- **Nottinghamshire County Council** – County ecology officer present: [REDACTED] [ottsc.gov.uk](mailto:ottsc@nottinghamshire.gov.uk)

1.4.2 It is recommended due to the specific impacts and constraints at the sites that the following organisations are consulted with at the appropriate stage:

- **RSPB** and **British Trust for Ornithology** (BTO) – Both organisations consult on planning applications (particularly RSPB) and run targeted conservation programmes in the local agricultural landscape to which a contribution may be a viable mitigation option for impacts on breeding birds.
- Lincolnshire Wildlife Trust

1.5 Site Description Summary

1.5.1 West Burton 1, 2 and 3 are located within the West Lindsey District, Lincolnshire and are situated within 8km of each other close to the settlements of Broxholme (West Burton 1), Ingleby (West Burton 2) and Brampton (West Burton 3). The Sites have been mapped in Figure 1. West Burton 1, 2 and 3 predominantly comprise large, open and generally flat arable fields characterised by winter-sown cereal crops with some fields of permanent

¹ Code of Professional Conduct. CIEEM, January 2019.



pasture (West Burton 2), bounded by a network of managed hedgerows and ditches with narrow field margins, where present.

- 1.5.2 These Sites' habitats are very much typical of the surrounding landscapes which are dominated by arable farmland and occasional pasture grassland that is interspersed with small settlements and farmsteads linked by minor and single track roads. The landscape surrounding West Burton 1 – 3 is mostly flat but to the east of the Sites at the 'Lincoln Cliff', a significant north-south escarpment, located 3km east of West Burton 1. The River Trent is located west of West Burton 1 – 3 and is located 1.4km from West Burton 3 at its closest point as it flows north towards the Humber Estuary. While no woodland is present within the Sites, several small stands of managed and unmanaged woodland are present adjacent and in the surrounding landscape, often the result of historical game management. Standing water is generally absent from the Sites and the surroundings following the in-filling of traditional livestock drinking ponds, save for a very small number of agricultural pools/pits, decoy ponds or managed recreational fisheries. Flowing water occurs occasionally in proximity to the Sites, with the River Till running adjacent to the eastern boundary of West Burton 2 and 0.4km west of West Burton 1 the River Trent running 1.4km west of West Burton 3. Various feeder streams for the above watercourses are managed as agricultural drainage ditches within or adjacent to the Sites which regularly dry out.

1.6 Surveys Carried out to Date

1.6.1 To date, the following surveys have been carried out across all the above sites in 2021:

- An Extended Phase 1 Habitats Survey to characterise habitats present and potential for protected species (April)
- Four breeding bird survey visits (late May to late June)
- One nocturnal bird survey visit, to focus on quail and owls (late June to early July)
- Water testing for presence of great crested newt (GCN) DNA from all ponds within the Site and adjacent land within 250m under same ownership (June)
- Two deployments of static bat detectors at a low to moderate density (June and July)

1.6.2 Surveys currently planned to be carried out at the Sites are:

- Four wintering bird surveys to record any significant numbers of migratory or resident wildfowl, waders and other birds reliant on arable farmland during the winter (November 2021 to February 2022)
- Three further deployments of static bat detectors (July to September 2021)
- Further water testing for GCN eDNA beyond option and current landowner boundaries (mid-April 2021 to end June 2021).

2 DESK STUDY

2.1 Local Planning Policy

2.1.1 The following nature conservation-related policies taken from the Central Lincolnshire Local Plan are considered pertinent to West Burton 1 - 3 and the proposals. The text of each policy is given in turn in Appendix C at the end of this report.

Central Lincolnshire Local Plan (Adopted April 2017)

- Policy LP19: Renewable Energy Proposals
- Policy LP20: Green Infrastructure Network
- Policy LP21: Biodiversity and Geodiversity

Central Lincolnshire Local Plan (Under Consultation - Anticipated adoption of revised plan in April 2022)

- Policy S13: Renewable Energy
- Policy S58: Green Infrastructure Network
- Policy S59: Protecting Biodiversity and Geodiversity
- Policy S60: Biodiversity Opportunity and Delivering Measurable Net Gains
- Policy S65: Trees, Woodland and Hedgerows



2.1.2 In the local area, Neighborhood Areas have been designated for the purposes of creating Neighborhood Plans. The Saxilby with Ingleby Neighborhood Plan (2016 – 2036) was pertinent to **West Burton 2**, with no other Neighborhood Plans relevant to the Sites. Relevant policies are as follows and are also detailed in Appendix C.

- Policy 11: Minimising The Impact Of Development On The Natural Environment
- Policy 12: Green Infrastructure
- Policy 13: Development Along The Fossdyke Canal
- Policy 15: Designated Local Green Spaces

2.1.3 The following habitats and species have been identified within Lincolnshire Biodiversity Action Plan (BAP) 2011-2020 (3rd Edition) and are considered relevant to the Site. As mentioned above, it is anticipated that alongside the re-drafting and eventual adoption of the new Central Lincolnshire Local Plan, the Lincolnshire BAP will be replaced by a Local Nature Recovery Strategy.

Habitats

- Arable field margins
- Grazing marsh
- Hedgerows and hedgerow trees
- Lowland calcareous grassland
- Lowland meadows
- Lowland dry acid grassland
- Ponds, lakes and reservoirs
- Rivers, canals and drains
- Lowland mixed deciduous woodland
- Wet woodland

Species

- Bats
- Farmland birds
- Freshwater fish
- Greater water-parsnip
- Newts
- Water vole
- White-clawed crayfish
- Invasive non-native species

Habitats

2.2 Protected and Designated Sites

2.2.1 Statutory and non-statutory sites designated for nature conservation were identified within the desk study and are summarised for each Site in Tables 1 and 2 below. Appendix C provides maps showing the relationship between the designated sites and the development Sites.

2.2.2 Many of these sites present potential ecological opportunities for the enhancement of local biodiversity and ecological connectivity.

2.2.3 'International' designated sites are statutory sites designated in response to international law or conventions, including Special Protection Areas (SPAs), Special Areas of Conservation (SACs) and Ramsar wetland sites. The search radius from each Site for these sites used was 10Km

2.2.4 National sites are statutorily protected sites which include Sites of Special Scientific Interest (SSSIs) and National Nature Reserves (NNRs). The search radius from each Site for these sites used was 5Km. Local Nature Reserves (LNRs) are statutorily protected local sites and thus are searched for within 5Km.

2.2.5 Local sites are predominantly non-statutory sites designated by Local Planning Authorities (in this case West Lindsey in collaboration with the Greater Lincolnshire Nature Partnership), including Sites of Nature Conservation Interest (SNCIs) and Local Wildlife Sites (LWSs). The search radius from each Site for these sites used was 2Km.

West Burton 1

2.2.6 No designated sites were identified in proximity to West Burton 1 within the desk study.

West Burton 2

2.2.7 One SSSI, Doddington Clay Woods SSSI, was identified during the desk study which was located 4.7km south of West Burton 2 and supports several types of woodland that are scarce in Britain. Three non-statutorily designated Sites were identified within 2km of West Burton 2 which comprise a disused railway embankment



that supports a diverse range of flora within a mosaic of habitats and two sites of species-rich agricultural grassland (see Table 1 below).

Table 1: Designated Sites in Proximity to West Burton 2

Site Name	Size (Ha)	Distance and Direction from Site	Reason for Designation
National Sites			
1. Doddington Clay Woods SSSI	23.9	4.7km south	Old Hag and Little Sale Woods are two ancient semi-natural woodlands containing representative examples of several stand types now scarce in lowland Britain, such as wet ash-wych elm woodland, acid birch-ash-lime woodland and lowland maple-ash-lime woodland. They also have a rich and varied ground flora. Old Hag Wood is the site of a large heronry and also supports a varied community of other breeding birds including woodcock and five species of warbler.
Local Sites			
2. Torksey Common to Sykes Junction Disused Railway LWS	6.4	0.5km west	Disused railway embankments with a range of habitats including grassland, scrub, trees and arable supporting a diverse assemblage of flora.
3. Mr. Rose's Hay Meadow LWS	5	0.7km north-west	The main hay meadow is quite species-rich, but dominated by coarse vegetation resulting from annual application of artificial fertiliser and infrequent aftermath grazing. Calcareous species are present in places.
4. Torksey Grassland LWS	3.6	1.6km north-west	This is a first class, quite large, pasture field, the major part being herb-rich neutral to acidic grassland on the dry ridges between furrows, but there is also some excellent damp and wet grassland. Boundary features are a significant element of the site, including old trees, ditches, hedges and dry acidic banks with anthills.

West Burton 3

2.2.8 No statutorily protected sites were found during the desk study within the search radiuses set out above for International and National designated sites. Seven non-statutorily designated Sites were identified within 2km of West Burton 3 and are described in Table 2 below.

Table 2: Designated Sites in Proximity to West Burton 3

Site Name	Size (Ha)	Distance and Direction from Site	Reason for Designation
Local Sites			
1. Mr. Rose's Hay Meadow LWS	5	30m south-east	The main hay meadow is quite species-rich, but dominated by coarse vegetation resulting from annual application of artificial fertiliser and infrequent aftermath grazing. Calcareous species are present in places.
2. Torksey Grassland LWS	3.6	100m south	This is a first class, quite large, pasture field, the major part being herb-rich neutral to acidic grassland on the dry ridges between furrows, but there is also some excellent damp and wet grassland. Boundary features are a significant element of the site, including old trees, ditches, hedges and dry acidic banks with anthills.
3. Torksey Marsh LWS	4.8	0.7km south	This is a very interesting site supporting a lot of bare ground, colonising vegetation and three ponds, as well as short stretch of disused railway line. All the ponds are of wildlife interest, supporting a wide range of plants, insects, birds and other fauna. The combination of diverse ponds, large expanses of wet ground, smaller areas of dry ground, an infertile sandy soil; and the recent management episode have all contributed to the current significant wildlife interest of the site.
4. Torksey Common to Sykes Junction Disused Railway LWS	6.4	0.7km south	Disused railway embankments with a range of habitats including grassland, scrub, trees and arable supporting a diverse assemblage of flora
5. Torksey Road Verge LWS	0.1	0.7km south	This 100 m stretch of un-managed verge on the north side of the road is mostly damp. However, a small area of dry habitat occurs at the western end, which is suitable for the nationally scarce blue fescue, the presence of which was the reason for designating this site as a Protected Roadside Verge.




6. Torksey Disused Railway LWS	0.7	0.8km south-west	The western end of this site retains a significant amount of sparse acidic grassland with characteristic species. The remainder of the site has little of the sparse sward mentioned above, and instead has plentiful common knapweed, bird's-foot-trefoil and field scabious. Scrub is prominent here.
7. Trent Port Wetland LWS	4.1	0.9km west	This un-managed triangular area of floodplain to the east of the River Trent comprises coarse neutral grassland and scattered scrub surrounding a large expanse of shallow water and wetland vegetation.

2.3 Ancient Woodland

- 2.3.1 Defra's Magic Map Application was used to identify stands of ancient woodland within 2km of the Sites.
- 2.3.2 No stands of ancient woodland were identified within 2Km of West Burton 1 and 2.
- 2.3.3 Burton Wood, listed as an Ancient & Semi-Natural Wood and Replanted Ancient Wood, was located 1.1Km north of West Burton 3.

2.4 Biodiversity Opportunities Mapping

- 2.4.1 Central Lincolnshire Local Plan Policy S60 relates to the delivery of measurable net gains for biodiversity within the county. Biodiversity Opportunity Mapping (BOM) has been created to show which areas and habitats are of greatest potential strategic value for enhancement in order to achieve this goal. This study built on a previous Central Lincolnshire Green Infrastructure Study and factors in potential beneficial outcomes for the local economy and society as well as nature. Key drivers for the inclusion of land within the mapping included agri-environment scheme targeting, restoring, buffering and connecting Local Wildlife Sites, and targets under Lincolnshire's Biodiversity Action Plan.
- 2.4.2 Figure 3 overleaf show the layout of BOM in relation to the Sites (within approximately 2Km).
- 2.4.3 **West Burton 1 and 2** fall within and close to the Biodiversity Opportunity Area known as River Till and Fosdyke Navigation Biodiversity Opportunity Area.
- 2.4.4 **West Burton 3** falls approximately 250m east of the Trent Vale Biodiversity Opportunity Area.
- 2.4.5 According to "Central Lincolnshire Policy S60: Biodiversity Opportunity and Net Gain Evidence Report", dated June 2021, work has begun on the preparation of a Local Nature Recovery Strategy (LNRS) for Lincolnshire which will replace the BAP. The LNRS will be a new system of spatial strategies for nature to support the delivery of biodiversity net gain and provide a tool for the public authorities to guide their approach. The LNRS will map the most valuable habitats for nature and provide specific proposals for effecting net gain opportunities. This will build upon the existing Biodiversity Opportunity Mapping and Areas work.



- Key
- Red line boundary
 - 2km site buffer
 - Ecological network - high quality
 - Ecological network - opportunity for management
 - Opportunity for creation
 - Opportunity for creation - more joined up

CLARKSON & WOODS
ECOLOGICAL CONSULTANTS

Project
West Burton 1 - 3

Title
Biodiversity Opportunity Mapping

Project Number
7479

Date
24/08/2021

Scale 0 1 2 km

Figure 3. Biodiversity Opportunity Mapping for West Burton 1-3.



3 HABITAT SURVEY

3.1.1 The findings of the habitat survey are discussed in this section, beginning with an overview of habitats common to each Site and a discussion of general opportunities for Biodiversity Net Gain. Thereafter, habitat features and findings particular to each Site are discussed in turn, with suggestions for Site-specific enhancements. Phase 1 habitat maps of each Site are given in Appendix H (supplied as a separate volume) and referred to in the text, along with target notes relating to specific habitat features. Each boundary is given a reference code (D# for ditch and H# for hedgerow), however hedgerows with ditches are referred to with an H# code only.

3.2 Common Habitat Constraints and Opportunities

Arable Fields

3.2.1 The arable fields occupied the vast majority of the Site's areas and were intensively farmed monocultures which are likely to receive periodic fertiliser and pesticide treatments. Vegetated field boundaries were sparse and historical field boundaries can be expected to have been progressively removed over preceding years since the industrialisation of farming. The arable fields across all Sites are therefore generally botanically poor and contained little particular ecological interest, save for their value to a relatively small number of ground-nesting bird species and arable specialists including hunting raptors (several of which are notable species of conservation concern) and brown hare, as described later in this document.

3.2.2 The removal of arable fields is unlikely to result in any intrinsic loss of ecological importance, particularly given the local abundance of this habitat. The arrays and the creation of grassland should help to promote local ecological diversity.

Opportunities for Enhancement and Biodiversity Net Gain

3.2.3 Considerable opportunities for the enhancement of these fields' ecological value compatible with a solar array are available. The reversion from intensive agriculture to low (or no) inputs (fertiliser and soil improvers) grassland alone would be expected to provide a modest net gain in plant and invertebrate species diversity. When multiplied over the large combined area of the Sites, this effect is likely to be significant at a County or District scale.

3.2.4 The benefit described above is able to be further enhanced through favourable and ecologically-led approaches to the ongoing management of the grassland. It is recommended that if grazing is desired, it forms a component of an overall management plan where grassland cutting and meadow management is also present, whereby some areas are not grazed. The establishment of a network of species-rich meadow within the ongoing site management would help realise especially significant net gain. Lowland meadows are a Habitat of Principal Importance under the NERC Act (2006) and are a Lincolnshire BAP priority. Areas identified within the Lincolnshire Biodiversity Opportunities Mapping (especially within **West Burton 2 and 3**) would be well suited to creation of this habitat. Furthermore, the proximity to nesting habitat for skylark and ground nesting birds (either on or off-site, if secured) could be another consideration for the most beneficial siting of high-value grassland management.

3.2.5 Further options for grassland habitat management and creation which could be incorporated under panels are given in 3.2.25 below, in relation to field margins and buffers.

3.2.6 While grazing is not necessarily incompatible with net gain for biodiversity or the creation of ecologically valuable grassland, grazing too often or too densely carries the risk of depleting botanical diversity through the raising of nutrient levels, favouring of fewer vigorous species, and inhibition of flowering and seed-setting. Ideal grazing regimes would include the limiting of number of animals per hectare/acre to 'conservation grazing' or Higher Level Stewardship (agri-environment scheme) rates, the seasonal restriction of animals from the land to allow flowering and recovery, or the use of sheep in 'aftermath' grazing in short periods following hay cuts.

3.2.7 Cutting or mowing can be carried out relatively quickly and cost-effectively, although cutting under panels can present a problem where weeds and scrubby vegetation takes hold. This should be treated through spraying or specialist cutting – advances are being made in these areas within solar arrays.

- 3.2.8 Where possible cutting should be carried out using a cut-and-collect system so as to minimise nutrient build up in the soil which stifles species diversity. Cutting regimes are often dictated by the perceived need to keep the sward height low to minimise shading risk. This can be simply avoided through the use of a 'shade cut', as shown in Figure 4 below, which aims to cut the first 50-100cm of grass out from the toe of each string during spring and mid-summer, while maintaining the invertebrate, bird and mammal value of the remaining grassland.



Figure 4. Photographs to show a 'Shade Cut' along the first 1m of grassland from the toe edge of the array, leaving flowering and seeding meadow grassland elsewhere.

- 3.2.9 In order to calculate a reasonably accurate forecast of Biodiversity Net Gain as a result of development, it will be necessary to formulate an operational land management plan which integrates the above broad management options. As different management techniques will have different ecological outcomes or targets, the management plan will be able to provide representative information on which a calculation can be based. The management approaches and management plan will therefore need to be formulated in due course and ideally in advance of completing a Net Gain assessment.
- 3.2.10 Regardless of chosen management regimes, the preparation of the fields before reversion to grassland will be key and must aim to minimise the impact of competition between desirable, sown species and unsown agricultural weeds and cereals.² This should be done through application of herbicide and, ideally, full cultivation followed by an additional herbicide treatment. Sowing of well-selected (locally-derived and appropriate) seed mixes (and to a lesser extent plug planting, in specific areas) would be carried out in the autumn. This should be followed by regular spring mowing with removal of arisings to control annual weed and nutrient levels in the following year, before establishing the final management regime, whether cut or grazed, from year three onwards. These are basic principles, which should be further investigated and tailored to site-specific conditions.
- 3.2.11 All habitat restoration and management approaches should be subject to periodic ecological monitoring to establish their success or otherwise to guide future management. This would be set out within a management plan (e.g. Landscape and Ecological Management Plan (LEMP)).
- 3.2.12 Solar development will drive a diversification of local habitats toward that of historical land use patterns where agriculture in the region was characterised by a mix of arable and pasture farming, which supported a greater abundance of wildlife. It is possible that, other concerns notwithstanding, the reversion of large areas of

² Blakesley, D. and Buckley, G.P. (2016) *Grassland Restoration and Management*. Exeter: Pelagic Publishing, UK



intensive arable to grassland, especially if managed with an emphasis on ecological benefit, would be perceived favourably in the local area.

Hedgerows

- 3.2.13 Hedgerows and Hedgerow Trees are a Habitat of Principal Importance and listed on the Lincolnshire BAP.
- 3.2.14 The hedgerow network is extensive across the majority of the Sites and is generally well-managed and species-poor, although several sections of species-rich hedgerow are present. It is also generally intact, with few gaps.
- 3.2.15 Roughly half of the hedgerows were accompanied by drainage ditches or streams, most of which were dry or partially wetted and were relatively narrow features.
- 3.2.16 Roughly half of the hedgerows contained at least sporadic mature and semi-mature trees. Trees were predominantly restricted to outer boundary hedgerows, while minor internal hedgerows were normally devoid of trees. Typical tree species recorded included ash (showing extensive signs of dieback), field maple, oak, rowan, holly, elder and grey willow. Woody shrub species most frequently recorded in hedgerows were hawthorn, blackthorn, and field rose.
- 3.2.17 Should any loss of hedgerow or boundary feature be required, it should be replaced on a 2:1 basis through supplementary planting in appropriate locations nearby.
- 3.2.18 The hedgerow network is probably the single most valuable habitat feature within the Sites and should be protected adequately during construction and operation with sufficient buffers. As a general rule, and in line with recommendations for watercourses and field margins below, recommended minimum buffer widths from hedgerow edge to the security fence are:
- Species-poor hedgerows or hedgerows without trees: 8m
 - Species-rich hedgerows or hedgerows with trees: 10m
 - Woodland: 20m
 - Ancient woodland: 30m
- 3.2.19 Perhaps the most pertinent driver of buffer width is the hedgerow's value to bats, therefore recommended buffer widths are likely to vary and increase according to the value of the hedgerows and trees present to bats, as discussed further in the species section.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.2.20 Much of the hedgerow network will require periodic cutting to maintain a reasonable height and structure. While specific hedgerows may require different management, cutting should generally be carried out on a 3-yearly rotation, with only either side or the top being cut each year. Significant net gains can be had by allowing the currently highly-managed hedgerows to fill out and broaden, encouraging a height of 3m or more, where currently they are often below 1.5m.
- 3.2.21 Additional hedgerow, tree or shrub planting would also provide significant net gains for biodiversity while contributing to visual screening. This can take the form both of in-filling of gaps in defunct or patchy hedgerows or new hedgerows laid at bare fenced boundaries. Additionally, it may be possible to reinstate a small number of old historical hedgerows which have been grubbed out in the past where the scheme allows (e.g. where advantageous for screening or at easements for PROW and services etc.). Maps such as those on [REDACTED] can be consulted for this. The planting of a small number of new hedgerows parallel to current ones to create a double hedgerow would contribute significantly to Green Infrastructure policies and aid the connectivity across sites if strategically located.
- 3.2.22 Species suitable for additional planting due to their abundance locally include blackthorn, hawthorn, elder, field maple, field and dog rose, grey willow, oak and dogwood. Site specific planting recommendations are given in the appropriate sections below.
- 3.2.23 It may be appropriate and well-received if an emphasis is placed on planting long-lived standard native trees, especially oak, sycamore and disease-resistant elm (but also potentially field maple, birch, lime, rowan, and alder) in order to replace the many ash trees which can be expected to be lost in the next five years due to ash dieback.



Field Margins and Semi-Improved Grassland

- 3.2.24 Arable field margins are a Habitat of Principal Importance and listed on the Lincolnshire BAP.
- 3.2.25 The uncultivated arable field margins across the Sites are predominantly absent or very narrow (<2m wide), apart from some areas in **West Burton 2 and West Burton 3** which have been purposefully left wide, in places approximately 5-6m or more. Generally they are species poor and poor in terms of structure, being mown most years in order to halt any scrub encroachment from hedgerows. Parcels of richer grassland habitat have been individually noted within the corresponding Site maps, although these are infrequent.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.2.26 Considerable cost-effective opportunities for the enhancement of field margins to become wider and more diverse are present. Diversification of grassland management maximises the available niches for invertebrates to lay eggs, overwinter and feed and in turn drive opportunities for diversification up the food chain. Furthermore, widening of existing margins as ecological buffer zones has the beneficial effect of enhancing the neighbouring hedgerows and ditches they frequently run parallel with. This in turn increases the interconnectedness of habitats within the site and within the neighbouring landscape, a key tenet of the NPPF and local planning policy.
- 3.2.27 The field margins lend themselves to being incorporated into wider buffer zones between hedgerows/field boundaries and the security fence line. Within these, a variety of straightforward management options can be pursued and ideally a mosaic of several techniques would be incorporated into the management of each Site according to Site-specific species conservation opportunities (dealt with separately in sections below). Management would ideally seek to avoid a uniform, regularly-mown grassland habitat as this reduces habitat structure and species diversity and instead follow a low-maintenance regime. Management options include:
- Tussocky grassland, mown no more than once per year (arising can be left in situ). This can be extended to once per two or three years on a rotational basis where monitoring indicates. A very low-maintenance technique providing habitat for small mammals, invertebrates and winter bird seed sources. See Figure 5.
 - Sown and annually mown (arising removed using cut-and-collect systems) species-rich meadow, potentially with aftermath grazing. Promotes low-growing flowering plants key for spring and summer invertebrate lifestages. See Figure 6.
 - Sown wild bird-seed crop (millet, quinoa, kale, linseed, teasel etc). Requires annual or bi-annual cultivation. Provides excellent autumn and winter food for birds.
 - Encouragement of a scattered scrub/ruderal vegetation habitat mosaic on a three-year rotational cut basis. Provides invertebrate overwintering habitats as well as year-round foraging habitat for many bird species. See Figure 7.
 - Pollen and nectar strips. Fine grassland dominated by low-growing nectar rich species such as clover, bird's-foot trefoil and sainfoin. Requires cultivation and/or sowing approximately every 3 years. See Figure 8.
- 3.2.28 It is recommended that these field margin buffer zones measure a minimum of approximately 7-10m from boundary (e.g. nearest hedgerow edge) to security fence in order to realise most ecological benefits³. Specific ecological constraints can be expected to increase this recommendation as discussed accordingly in the Site-specific species sections.
- 3.2.29 Locations within **West Burton 1-3** which appear on the Biodiversity Opportunities Mapping would be well suited to the more diverse habitat management options and mosaics. It is considered that sympathetically managed grassland buffer zones would constitute Arable Field Margin habitat in line with the Lincolnshire BAP.

³ BRE (2014) Biodiversity Guidance for Solar Developments. Eds G E Parker and L Greene.



Figure 5. Low-maintenance tussocky grassland can provide excellent habitat for small mammals.



Figure 6. Species-rich meadow can be created through well-timed cutting, aftermath grazing and collection of arisings.



Figure 7. Ruderal-encroached grassland can form ecologically valuable habitat in field margins.



Figure 8. Low-growing nectar-rich mixes (clover picotred) are cost-effective under panels and are of value to invertebrates.

Ditches and Watercourses

- 3.2.30 Rivers are a Habitat of Principal Importance while Rivers, Canals and Drains are listed on the Lincolnshire BAP.
- 3.2.31 The River Till (**West Burton 2** and to a lesser extent, **West Burton 1**) were small but relatively significant watercourses associated with the Sites and were fed by various drainage ditches present at field boundaries. Most of the wetted ditches and becks/streams held emergent vegetation and grassy banks, some of which were relatively diverse. The River Till and the larger watercourses featured wide grassy margins which formed large field headlands and were seen to be relatively diverse and provide key habitat for birds, small mammals and invertebrates.
- 3.2.32 Water quality appeared to vary, and in many cases was relatively poor owing to the presence of agricultural run-off. Water quality can be expected to significantly improve post-development due to the anticipated reversion to permanent grassland under the array (reduced sediment run-off) and cessation of application of fertilisers and pesticides.
- 3.2.33 Wetted ditches and watercourses are likely also to be key habitats for otter and water vole, both being legally protected species recorded near to or within all Sites. This will need to be considered when carrying out any engineering works close to or within ditches or river corridors.

3.2.34 Buffer zones along wet ditches and watercourses should be wider than many other simpler boundaries (such as defunct hedgerows or fences) owing to their elevated greater value to wildlife and the pollutant/sediment-attenuating properties of dense grassland vegetation and rich soils. Appropriate buffer widths from feature to security fence should range from 8 to 30m depending on the significance of the watercourse and associated protected species habitat value (e.g. bats, otters, water voles). 8m as a minimum offset from watercourses (including drainage ditches) is a standard Environment Agency and Internal Drainage Board requirement in order to preserve maintenance access and limit risk of pollution events. Significant watercourses clearly attract a wider buffer. These measurements are also discussed in the relevant Site-specific sections below.

Opportunities for Enhancement and Biodiversity Net Gain

3.2.35 The Green Infrastructure value of these features would be maximised through the creation of a wide buffering grassland habitat swathe, contributing to local policy aims and strengthening the value of the watercourse corridor. Habitat management options as listed for arable field margins could be implemented, as well as scattered tree planting.

Ponds and Standing Water

3.2.36 Ponds are a Habitat of Principal Importance and listed on the Lincolnshire BAP.

3.2.37 Few ponds were present at the Sites, most having been filled following the decline of pasture and mixed farming in favour of arable intensification. Those which remain on the Sites tend to be formed by wider, pooled sections of drainage ditches, are agricultural sumps/slurry pits, or are associated with woodland or woodland edge as shooting decoys. **West Burton 2** features the most actual in-field ponds, located within semi-improved grassland fields, while **West Burton 3** also had a small number of substantial waterbodies.

3.2.38 Ponds should receive a buffer of at least 10m unless other ecological constraints are present.

Opportunities for Enhancement and Biodiversity Net Gain

3.2.39 Ponds are of significant ecological value, and as a strong, high-quality pond network is absent within the local landscape, any creation of such features would be beneficial and likely to be favourably received by the LPA. Ponds could be created within field margin buffer zones and have a role to play in flood risk alleviation and water attenuation. These could take the form of linear ponds such as deepened swales as shown in Figure 9 below.



Figure 9. Swales can form intermittently drying linear pond features of value to wildlife if sufficiently deep.



3.3 West Burton 1 Habitat Assessment

Habitat Map and Target Notes

3.3.1 Please see Appendix H (separate document) for a Phase 1 habitat survey map for West Burton 1. Table 4, below, gives a description of the features referred to on the map by numbered Target Notes.

Table 4: Target Notes For West Burton 1 (Constraints and Opportunities)

No.	Description
TN1	3m tall ruderal margin on a south facing bank adjacent a wet ditch
TN2(x4)	Wide grassy margin of value for invertebrates and reptiles
TN3	Two white willows, growing very close to one another
TN4	Single white willow
TN5	Appears to have been in-filled with whip planting in recent years
TN6(x3)	Tree rooted halfway down a steep bank to the relatively large ditch (drain)

Habitat Overview

3.3.2 West Burton 1 is the smallest Site and measures approximately 90ha, being characterised by five large wheat fields (presumed winter sown) surrounded by a network of moderately managed, species-poor hedgerows with ditches. Most of the hedgerows contained at least intermittent mature trees. Field margins were generally narrow and comprised unmanaged grass, although in certain locations these were wider, up to 6m, with encroaching ruderal vegetation and tussocky in nature. No agricultural buildings were present on or adjacent to the Site. Two small copses or woodland stands were present adjacent to the Site (west and east boundary respectively). No ponds were present on Site.

Arable Fields and Field Margins

3.3.3 The arable fields are all of low botanical interest and general ecological value, being monocultures, save for their value to ground nesting birds and hares, predominantly.

3.3.4 Field margins were mostly narrow, being up to 2m wide, although in certain locations, such as H8, H11 and H12, as indicated by TN2, these measured up to 6m wide, although mostly on the northern sides to hedgerows. All field margins were left unmanaged and had become coarse and tussocky. These comprised species including Timothy-grass, cocks-foot, cough grass, creeping bent, red fescue, thistles, hogweed, willowherb species, common nettle, redshank and hedge bindweed. Ragwort, meadowsweet, chamomile, bittersweet and kidney vetch were occasionally recorded.

3.3.5 Consequently, several of the existing grassy field margins mentioned above hold some moderate ecological value and should be retained and incorporated into buffer zones extending from their corresponding hedgerows/ditches wherever possible.

Opportunities for Enhancement and Biodiversity Net Gain

3.3.6 As West Burton 1 was dominated by cultivated land and did not include any discrete semi-improved grassland or pasture fields, there are few locations where traditional meadow creation would be considered a natural succession of existing habitats, although widening and low-key management to tussocky grassland of all currently narrow margins in order to match those noted as bearing wildlife value would be beneficial.

3.3.7 Field 1 is located within a BOM zone listed as 'opportunity for creation', presumably on account of its proximity to the River Till and priority habitat in the form of plantation woodland to the north east of the Site. Therefore, Field 1 would be the best single field candidate for high-value meadow grassland creation, although arguably this could extend to Field 3 and the western half of Field 5 for the same reason.

3.3.8 All cereal fields would benefit from their reversion to permanent grassland receiving ecologically-sympathetic management as set out in Section 3.2.

Hedgerows

3.3.9 While all hedgerows were considered species-poor, the majority featured at least intermittent mature and semi-mature trees with accompanying, mostly wet, drainage ditches and had been allowed to grow to



approximately 2.5m in width and height making them valuable nonetheless. All hedgerows were likely to be over 30 years in age and receive trimming or flailing at least every two years.

- 3.3.10 Hedgerows were invariably dominated by hawthorn and blackthorn, with other woody species integral to the hedgerow including elder, hazel, goat willow, dogwood, field and dog rose and bramble. Occasional trees were typically made up of semi-mature elm (dead), white willow, oak and field maple, with immature ash and sycamore.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.3.11 No gappy or defunct hedgerows are present, although managed hedgerow lengths could be supplemented by planting new whips which would encourage structural diversity. Planting an occasional hedgerow parallel to an existing one, to create a double hedgerow, would also strengthen the existing green infrastructure network and improve habitat connectivity across the Site, for example along H12 or H7.
- 3.3.12 Pre-emptive replacement of ash trees as described in Section 3.2 would be a good opportunity at West Burton 1.
- 3.3.13 Generally, the management of hedgerows in order to encourage a tall and bushy form, with incremental and rotational trimming, is advised as per Section 3.2.

Ditches

- 3.3.14 The ditches on site were predominantly wet or damp, with grassy banks and associated with hedgerows. The majority of ditches were small and measured approximately 2m wide at the banktop and 1-1.5m deep. D1 and H12 featured ditches which were approximately 4m wide at banktop and 2-3m deep. In parallel to the north of D1 was a larger watercourse with engineered embankments which was a tributary of the River Till further to the west.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.3.15 Few specific enhancements for the Site's ditches are recommended over and above that of periodic inspection and maintenance wherever necessary in order to ensure proper drainage function. However, it is recommended that ditches are not overly dredged or cleared unless they are causing a drainage issue or at a frequency in line with EA/IDB recommendations. Grassy buffers would help to maintain water quality and mitigate pollution risks.

3.4 West Burton 2 Habitat Assessment

Habitat Map and Target Notes

- 3.4.1 Please refer to Appendix H (separate document) for a Phase 1 habitat survey maps for West Burton 2. Table 5, below, gives a description of the features referred to on the map by numbered Target Notes.

Table 5: Target Notes For West Burton 2 (Constraints and Opportunities)

No.	Description
TN1	Small pile of tipped rubble overgrown – suitable hibernaculum
TN3	Small pocket of broadleaved woodland with lots of dead and decaying wood
TN5	Several 'mammal' burrows in ditch bank, presumed to be rat (rat droppings present)
TN6	Fenced hardstanding with newly planted hawthorn hedgerow
TN7	Large rabbit warren underneath hedgerow
TN8	Scrubby widened hedgerow with several piles of tipped rubble present. Good reptile hibernacula
TN9	Orchard belonging to farmhouse adjacent to Red Line Boundary
TN10	Cuckoo seen in tree
TN12	Species-poor semi-improved field margin with ruderals
TN13	Scrub encroachment
TN14	Small ox-bow lake surrounded by scrub
TN15	Double hedgerow with dry ditch



No.	Description
TN16	17m wide tussocky field margin with high reptile potential
TN18	25m stubble set-a-side – after the 5m field margin
TN19	Small block of broadleaved woodland
TN20	Tussocky field margin
TN21	20m wide tussocky field margin with some scrub encroachment
TN22	20m of stubble margin
TN23	Coarse grassland margin

Habitat Overview

- 3.4.2 West Burton 2 measures approximately 330ha and is characterised by moderately large or large winter and spring sown cereal fields with a small proportion of permanent pasture fields supporting semi-improved grassland. The fields are separated by drainage ditches of widely varying sizes and habitat value and a network of hedgerows, often species-rich and with intermittent mature trees. Field margins generally supported moderately rich, tussocky uncultivated or managed grassland with some scrub encroachment and were generally narrow, although in many cases have been allowed to occupy up to 5-8m widths and up to 20m in some instances. Several ponds were present adjacent to and within the Site, particularly within the north-east field. The River Till borders the Site immediately to the east and it is understood that a margin of approximately 200m along this eastern boundary is designated as flood alleviation land, used by the Environment Agency to protect Lincoln and surrounding settlements from a 1-in-100-year flood event.
- 3.4.3 The eastern half of the Site (south and east from F59) is designated within Lincolnshire's Biodiversity Opportunities Mapping as being 'opportunity for creation', while F62 is designated as 'opportunity for management' on account of its elevated current biodiversity value.

Arable Fields

- 3.4.4 The cereal fields are all of low botanical interest and general ecological value save for their value to ground nesting birds and hares, discussed in the species sections below.

Semi Improved Grassland and Field Margins

- 3.4.5 Two fields (F53, F54) of species-poor semi improved grassland were present in the centre of the Site and had been heavily grazed by sheep. These were characterised by species such as perennial ryegrass and sweet vernal grass with occasional ruderals such as common nettle and creeping thistle. One field (F62) of tall (ungrazed or cultivated), species-poor semi improved rank grassland (part of the flood alleviation land) was present in the north-east corner of the Site, adjacent to the River Till, which contained numerous ponds and it is possible that it is periodically inundated with floodwater, although further information from the EA and the landowner would enable a fuller understanding of this. The field was dominated by perennial ryegrass, meadow foxtail and sweet vernal grass with curled dock.
- 3.4.6 Field margins were generally narrow across the Site, although in many cases were 5-8m and up to 20m within F84, F50, F77 and F82 and supported uncultivated semi-improved grassland that had occasionally been allowed to become tussocky with some encroachment of scrub, particularly within the eastern fields adjacent to the River Till. Dominant species were perennial ryegrass, cock's foot, meadow foxtail, timothy, rough-stalked meadow grass, false oat-grass with hogweed, teasel, cow parsley and willowherbs.
- 3.4.7 Grassy field margins should be retained and incorporated into buffer zones extending from their corresponding hedgerows/ditches wherever possible.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.4.8 Field within the eastern half of the Site could be prioritised for receiving species-rich or valuable grassland creation measures as a response to their inclusion within Lincolnshire's BOM. F62 in particular is considered to hold the potential to be significantly enhanced to a species-rich traditional lowland meadow through cessation of regular grazing and introduction of a single hay cut (cut-and-collect) potentially with aftermath grazing. This should have the effect of stifling ryegrass dominance and allowing finer grasses and flowering plants to compete. The sward can be further diversified through over sowing within an appropriate meadow seed mix. Depending on the regime of inundation it receives (awaiting further EA information), this treatment



should be adjusted to reflect its wetter conditions through skewing a seeding mix towards wet/floodplain meadow or marshy grassland.

- 3.4.9 All cereal fields would benefit from their reversion to permanent grassland receiving ecologically-sympathetic management as set out in Section 3.2.
- 3.4.10 Field margins are wide in places, particularly in F84 alongside the River Till embankments, and should be retained and enhanced. Enhancements in these margins would lend themselves to simple tussocky grassland management, with desirable encroachment by ruderal and scattered scrub habitats, in line with the conditions of the similar habitat fragments found on Site mentioned above.
- 3.4.11 The margins along the western edge of F63 (TN16), southern edge of F50, the northern, southern and western edges of F77 (alongside the woodland strip and D2) and western edge of F82 (alongside D5) are also particularly wide and should be preserved and enhanced. These locations would lend themselves to wild bird cover crop (millet, kale, quinoa etc) or ruderal-encroached tussocky grassland in line with their current condition. The above treatments at F50 and F77 would help buffer the strip of woodland located between the two fields and enhance its connectivity and Green Infrastructure at the Site.

Hedgerows and Woodland

- 3.4.12 Hedgerows were absent from many field boundaries, with fields being demarcated with drainage ditches and grassy field margins in these instances. Hedgerows, where present, were a roughly even mixture of species-poor and species-rich with dominant species being hawthorn and blackthorn, with rose, field maple, grey willow, ash, crab apple, elder all regularly present. Hedgerows generally received minimal management, causing many to have become quite tall and bushy, improving their ecological value. Several internal hedgerows were gappy and classed as defunct. Hedgerows should be adequately buffered as set out in Section 3.2.
- 3.4.13 Woodland was present on Site in the form of a relatively diverse strip of broadleaved woodland separating F50 and F77 (known locally as Codder Lane Belt) which was seen to contain a good variety of species, including mature ash, oak with an understorey of field maple and dog rose and ground flora featuring ramsons, cleavers, bluebell, garlic mustard, yellow archangel, dog's mercury and ground ivy. While not officially designated as ancient woodland, several of the above species are indicators of ancient woodland and therefore the habitat should be buffered and maintained as such.
- 3.4.14 Further woodland fragments were present at the corner junction of F55, F60 and F61 (TN19), TN3 south of F77 and around the northern boundaries of F40 and F41, with another copse alongside the north western boundary of F48.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.4.15 The woodland blocks given above should receive a buffer of at least 20m, potentially 30m for the long woodland strip at F50/77 due to the presence of ancient woodland species. These buffers should be managed to enhance their interconnectivity with hedgerows and neighbouring off-site woodland. Scrub and tussocky grassland would be desirable within these zones.
- 3.4.16 The gappy hedgerows (H2, H3, H6, H27, H28, H41, H56, H66,) would lend themselves to being made intact through new planting, including standard trees managed to become emergent above the surrounding hedgerow as per existing trees.
- 3.4.17 Bare ditches could have hedgerows or individual trees planted, for instance along D2, D3, D4, D5, D6, D7, D9, D11, D16, D21, D24 and D25. However, this should be carefully considered as it may be more appropriate to encourage wide tussocky grassland margins. The copse of woodland at TN3 could be offered greater habitat connectivity through the planting of a hedgerow along D4, combined with a wide buffer strip along H18.
- 3.4.18 Many ash trees were suffering from dieback and would benefit from pre-emptive replacement as described in Section 3.2.

Ditches

- 3.4.19 The ditches on Site were mostly wet, with a small number of dry ditches. The most significant drainage watercourses were D1 and D2 in the west of the Site, D9 (with H29 and H48), D16 (with D17-19) and D25 (with H61). These measured up to 10m wide and 2m deep in places, with tussocky grassland banks colonised by tall



grasses, ruderals and marginal wetland plant species. Generally, the ditches at West Burton 2 were of good quality and species diversity so should be protected as far as possible.

Opportunities for Enhancement and Biodiversity Net Gain

3.4.20 Few specific enhancements for the Site's ditches are recommended over and above that of periodic inspection and maintenance wherever necessary in order to ensure proper drainage function. However, it is recommended that ditches are not routinely dredged or cleared unless they are causing a drainage issue. Grassy buffers would help to maintain water quality and mitigate pollution risks. The ditches should be buffered as previously mentioned in Section 3.2, potentially with more significant drainage watercourses (listed above) receiving a slightly wider buffer.

Ponds and Standing Water

3.4.21 Eleven ponds were present within the Site boundary, ten of which were present within F62. These ponds were generally shallow and likely to dry annually with good water quality and a high proportion of emergent vegetation comprising tall grasses, rush, sedge and bulrush.

3.4.22 The remaining ponds were situated along the boundaries within F52 and F53 (P4-6) and comprised permanent ponds with poor water quality.

3.4.23 It is recommended that F62, which contains a high density of in-field ponds, is not developed and is enhanced to create a species-rich marshy lowland meadow with ephemeral ponds.

3.4.24 A buffer of 10m from the pond edges to security fences is considered appropriate for all on-Site and adjacent pond features.

Opportunities for Enhancement and Biodiversity Net Gain

3.4.25 Pond creation is not considered to be a priority at West Burton 2. However, the ponds present would all benefit from positive management, including selective deepening and the planting of marginal and emergent aquatic plants. Ongoing monitoring and reactive management would help to significantly enhance the ecological contribution made by them. It can be expected that water quality would improve following the reversion of arable to grassland and the completion of construction.

3.5 West Burton 3 Habitat Assessment

Habitat Map and Target Notes

3.5.1 Please refer to Appendix H (separate document) for a Phase 1 habitat survey maps for West Burton 3. Table 6, below, gives a description of the features referred to on the map by numbered Target Notes.

Table 6: Target Notes For West Burton 3 (Constraints and Opportunities)

No.	Description
TN1	Dense scrub and felled logs in corner of field
TN2	Rough grass margin good for invertebrates and reptiles
TN3	1.5m wide rough grassland margin good for reptiles
TN4(3x)	Pile of brash/debris – reptile potential
TN5	Common lizard (1x) seen here
TN6(8x)	Multiple skylark recorded here
TN7	Grassy bank of ditch good for reptiles – grass snake seen here
TN8	Cuckoo seen here
TN9	Patch of uncultivated ground – good for reptiles
TN10(3x)	Good water vole habitat
TN11	Tall ruderals and tussocky grassland – good for reptiles and water voles
TN13	Additional pond feature noted here
TN14	Good GCN and reptile habitat within tall ruderal vegetation plus log and brash piles
TN15	Diverse habitat buffer at woodland edge
TN16	Ash tree with large raptor nest



TN17	Strip of broadleaved woodland just outside of red line boundary
TN19	Tussocky grassland and scattered scrub margin along woodland edge
TN21	Rough grassland suitable for reptiles
TN25	Significant main sett and lots of mammal paths in wood
TN27	Planted wild bird foraging cover strip to be retained and enhanced
TN28	Pair of skylarks seen
TN29	Woodland strip with species rich grassland bank

Habitat Overview

- 3.5.2 West Burton 3 measures approximately 375ha and is characterised by medium and large sized arable fields with a small number of species-poor semi-improved grassland fields. The Site was bisected roughly north to south by a railway line, the embankments of which were heavily vegetated with woodland, scrub and ruderal vegetation. Most of the arable fields were winter-sown wheat, with some linseed and bean crops located to the east and south (Brampton). Within the north east of the Site was an area of woodland and ponds associated with a scheduled monument and moat features – this designation is understood to extend to field boundaries in the south of the Site (western edge of F8 and F30 and eastern edge of F31). Woodland also bordered much of the Site's western boundary and occupied a corner of the southernmost field. The Site featured a strong hedgerow network, with most Site perimeter boundaries featuring tall hedgerows with mature trees. Drainage ditches were occasional, mostly being located within the centre and northeast of the Site.
- 3.5.3 Fields F22, F8 and F30 are all included within the Lincolnshire Biodiversity Opportunities Mapping as 'opportunity for creation'. The woodland parcel just off the southern corner off the Site, containing ponds P4 and P5 is designated as an 'opportunity for management'.

Arable and Improved Grassland Fields

- 3.5.4 The arable and improved grassland fields are all of low botanical interest and general ecological value apart from their value to ground nesting birds and hares. The semi-improved grassland of F26 and F27 was towards the managed and species-poor end of the category, likely being used as a fodder crop, as in F3 and F6.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.5.5 Considerable opportunities for reversion to grassland or meadows exist at West Burton 3 in line with general grassland creation advice previously discussed in Section 3.2. No fields are considered priorities for meadow restoration in terms of their current species composition, but the overlap between F22, F8 and F30 with the BOM designation would provide a good place to apply higher-value grassland management techniques for greatest contribution to local policy objectives.

Field Margins and Semi-Improved Grassland

- 3.5.6 Uncultivated grassy field margins were generally very poor in terms of extent (0-2m from field boundaries), especially in the east of the Site (Brampton). Species diversity and structure were also poor, with approximately half evidently receiving a mow or cut annually, with others left to become rough. Species typically present included cock's-foot, Yorkshire fog, perennial ryegrass, meadow foxtail, meadowsweet, docs, ground ivy, cow parsley, hogweed and nettles.
- 3.5.7 A small number of field margins within fields F5-8 in the north west of the Site (Bellwood) were slightly wider and more diverse, measuring up to 4m wide in places and with greater value and structure for reptiles and amphibians.
- 3.5.8 Other wide grassland margins or patches of uncultivated rough vegetation were present at TN25 (west of F8 and F30 associated with broadleaved woodland), TN16 in F22, TN27 in F31, alongside woodland in the west of F13, TN21 in F28 and along the railway embankments (e.g. at TN19).

Opportunities for Enhancement and Biodiversity Net Gain

- 3.5.9 The field margins on Site would benefit significantly from reduced management and extension in width to create either tussocky grassland, species-rich meadow habitat, cultivated wild bird cover crop or scrub-ruderal grassland mosaic. Those listed above would be good candidates for enhancement treatments.



- 3.5.10 Small patches of semi-improved grassland and tall ruderal scrub were present in corners of some fields which were difficult to cultivate or maintain and as such had become tall and tussocky. Although they hold little botanical interest, they offer invertebrate habitat and habitat for small mammals which are hunted by birds of prey. These are located at TN9, TN16 (and surrounding P9), margins surrounding P12, the southwest corner of F9 and along the western boundary of F13 where the land slopes away towards woodland and is dominated by scrub and tall ruderal vegetation.

Hedgerows

- 3.5.11 Relatively few species rich hedgerows were present, although most featured mature trees. Only very few, short lengths of hedgerows were considered defunct or gappy.
- 3.5.12 The strongest tree lines and largest, most mature hedgerows were present along the wooded western perimeter of F11, F13 and F15, along H13 and H15, H27, H36, H38, H49, adjacent woodland containing P3-5 and the woodland strip at TN29. These should be considered to attract a wider buffer than the standard minimum owing to their elevated conservation value. Woodland should be buffered by at least 20m.
- 3.5.13 Common mature and semi-mature trees present included ash, oak, beech, horse chestnut, hazel, rowan, alder, sycamore and English elm. Woody hedgerow species were generally hawthorn and blackthorn with occasional elder, *Viburnum opulus*, field rose and spindle.
- 3.5.14 Internal hedgerows, i.e. those which did not describe the Site perimeter were generally species poor and without emergent mature trees, being dominated by hawthorn and blackthorn. These were generally well managed and maintained to between 1.5m to 2m in height and width.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.5.15 The west side of **West Burton 3** presented relatively few options for hedgerow planting, although several were considered species-poor and lacking trees, therefore these should be targeted for enhancement.
- 3.5.16 The land east of the railway featured several ditches without any associated hedgerow. The planting of such a feature on one banktop, with approximately a 2m setback could be a suitable enhancement. Examples of suitable locations would be D4, D5 and D8, with D15 in the south of the Site.
- 3.5.17 Bare access tracks, for example those between F9 and F10, F11 and F12 and between F8 and F31 would be appropriate locations for new hedgerow planting, potentially double hedgerows to create an avenue feature.
- 3.5.18 Pre-emptive compensation for ash trees suffering from dieback as described in Section 3.2 would be a good opportunity at **West Burton 3**.

Ditches and Ponds

- 3.5.19 Significant ditches are only present within the north west and north east of the Site and a limited number in the Site centre. The strongest watercourse feature was D2, which was a stream flowing north to south, connecting with D3 in the centre of the Site.
- 3.5.20 Several ponds are present on Site, including a significant pond at P9, a decoy pond at P12, and several immediately adjacent to the Site within woodland in the north eastern corner, the southern corner and the western boundary. One ditch contained a pond-like feature which could be easily enhanced (TN13).
- 3.5.21 Positive records of great crested newt were returned within Ponds 4 (just off Site) and 9.

Opportunities for Enhancement and Biodiversity Net Gain

- 3.5.22 Few specific enhancements for the Site's ditches are recommended over and above that of periodic inspection and maintenance wherever necessary in order to ensure proper drainage function. However, it is recommended that ditches are not routinely dredged or cleared unless they are causing a drainage issue. Grassy buffers would help to maintain water quality and mitigate pollution risks.
- 3.5.23 Ditches noted as having good habitat for water voles (TN10x3) could be priorities for enhancement through removal of debris, widening of banktop grass buffers and planting of marginal and emergent vegetation.
- 3.5.24 While pond creation is not expressly necessary, it would be a welcomed enhancement to create one or more waterbodies in appropriate locations between the two ponds containing GCN records. The woodland strip



and associated grassland bank along the western edges of F8 and F30 would be a suitable location, as would within the southern half of F22 or the grassland at TN16.

- 3.5.25 Positive management of all ponds present on Site (or immediately adjacent if possible, see Pond 10 and Ponds 21-24), including selective deepening, clearance of tipped material and planting of emergent and marginal vegetation would significantly enhance their value to aquatic wildlife.

4 SPECIES INFORMATION COLLATED TO DATE

- 4.1.1 This section sets out the results of preliminary species survey work and an appraisal of the Sites' value to various protected and notable species. It also gives recommendations and suggestions for mitigation of potential impacts and opportunities for biodiversity net gain. In the interests of brevity and to avoid repetition, the site-specific results and recommendations are given together under each species' sub-heading in turn.

4.2 Badgers

Desk Study Information

- 4.2.1 This information has been removed to a confidential appendix to the Environmental Statement.

Field Survey Results

- 4.2.2 This information has been removed to a confidential appendix to the Environmental Statement.

Potential Constraints, Mitigation and Further Work

- 4.2.3 An operational solar array would most likely present at worst a neutral impact on badgers provided that appropriate protective measures outlined below are undertaken during construction and maintenance. Potentially, the diversification of habitats by introduction of permanent grassland may help to provide better foraging opportunities for badger in the long term.
- 4.2.4 The grassland habitats beneath the array are highly likely remain conducive to foraging by badgers (whether grazed or cut) and access to other woodland and farmland likely to remain unimpeded.
- 4.2.5 The perimeter fencing of the array is not considered to pose a limitation to badger dispersal unless it is deeply buried and of a tight mesh size which is not typical of solar arrays. For this reason, buried fencing is not advised as it would risk leading to its excavation by the badgers in the long term and potential fragmentation of badger social groups.
- 4.2.6 The use of badger gates in perimeter fencing is also not recommended although is something that is commonly encountered. This is considered unnecessary unless fencing is significantly buried and in our experience of monitoring arrays across the UK we have not encountered a single badger gate in a section of linear fencing which showed any evidence of use. By contrast we have recorded multiple locations where badgers squeeze beneath fencing (often adjacent to a badger gate). Badger gates represent an unnecessary expense and likely just compromise the integrity of the fencing should the intention be to graze areas with livestock.

Protection and Avoidance of Setts

- 4.2.7 Badgers and their setts are legally protected from disturbance and damage when active (likely to be occupied). Badgers are unlikely to pose a significant constraint to the development at the Site given the general lack of activity at the Site and potential for impact onto significant setts. Constraints are likely only to apply to the construction phase of the development.
- 4.2.8 As badgers are liable to dig new setts at any time, a pre-construction survey (approximately 3-6months prior) of woodland edges and hedgerows within approximately 30m of any development activities is recommended to ensure any new setts can be mitigated for in advance of commencement. Any setts capable of being impacted should be examined to determine whether they are active or disused. Disused setts generally do not pose a constraint. Such investigation work may require monitoring using cameras over a (minimum) three-week period.



- 4.2.9 To ensure that construction and operational maintenance works do not cause unlawful impacts on badgers and setts, a 20-30m buffer zone should be established from the perimeter of any active sett. The size of the buffer zone should reflect the status and activity levels within the sett and the nature of the local topography and the direction of tunnels associated with the sett entrances. Within this buffer zone, there should be no movement of plant, excavations or installation of array structures or buried cabling for the life of the scheme. Protective fencing and signage should be installed at the beginning of the construction phase.
- 4.2.10 If it is not possible to retain an active sett within the proposals, or maintain adequate buffer zones, it is likely to be possible to close (either temporarily or permanently) them under a licence from Natural England. For any main setts, it is probable that an alternative badger sett will need to be constructed in a suitable nearby location in order to ensure sufficient alternative shelter. The artificial sett will also need to be created well in advance of closure operations and uptake by the badgers will need to have been demonstrated by means of video surveillance or similar. It is therefore advisable to undertake artificial sett creation at least six months in advance of sett closure. Sett closure under licence can only take place between the months of July and November inclusive so as to avoid impacts on dependent young underground.
- 4.2.11 Badgers will forage within grassland creating shallow pits and scrapes down to approximately 15-20mm when excavating earthworms and grubs. To date we have not come across any examples of badger activity causing issues with buried cabling on active solar arrays. We believe that the standard armouring surrounding buried cabling is sufficiently robust enough to not be damaged by badger foraging or digging activity.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.2.12 The substitution of grassland within areas previously supporting arable land will provide a greater diversity of habitats. Badgers are reliant upon a diversity of foraging opportunities, exploiting different habitat types and areas through the year in response to availability.
- 4.2.13 The grasslands within arrays generally present good opportunities for forage, the stability and undisturbed nature of soils promotes earth worm abundance, and invertebrate and small mammal populations are generally improved within arrays, all of which provide foraging opportunities for badger.
- 4.2.14 Consideration might be given to the incorporation of fruiting trees (crab apple, apple and pear for example) within marginal areas as windfall fruits provide an important foraging resource in the autumn when badgers are looking to build weight for the winter period.

4.3 Bats

Desk Study Information

- 4.3.1 For **West Burton 1**, approximately 60 records for four species were recorded within the desk study data, none of which were recorded within the red line boundary and the vast majority beyond 250m of the Site. The most commonly recorded species was common pipistrelle, with the remaining three species (soprano pipistrelle, brown-long eared bat, and noctule bat) having only one record each. This represents a low diversity of species, all of which can be expected to roost within buildings and/or trees in the local area. The species present in the data were generally common and widespread. Most records were made post-2000.
- 4.3.2 For **West Burton 2** approximately 160 records for six species were recorded within the desk study data, none of which were recorded within the red line boundary and the vast majority beyond 250m of the Site. The most commonly recorded species was common pipistrelle, followed by Daubenton's bat, brown long-eared, noctule bat, soprano pipistrelle and natterer's bat.
- 4.3.3 For **West Burton 3**, approximately 230 records for six species were recorded within the desk study data. Two records of an unidentified bat are located within the red line boundary with the vast majority of the remaining records located beyond 250m from the Site. The most commonly recorded species was common pipistrelle, followed by soprano pipistrelle, brown long-eared bat, noctule bat, Daubenton's bat and natterer's bat.
- 4.3.4 Bats are Species of Principal Importance under the NERC Act (2006) and are listed on the Lincolnshire BAP.



Field Survey Results

Bat Detector Survey

- 4.3.5 16 bat detector locations were utilised, with two at **West Burton 1**, six at **West Burton 2**, eight at **West Burton 3**.
- 4.3.6 A preliminary inspection of data gathered from the first two detector deployments (June and July) indicated that a relatively moderate diversity of species was present across the Sites.
- 4.3.7 The majority of activity was made up of common and soprano pipistrelle, noctule bat and several *Myotis* species, which was expected. Brown long-eared bat is another relatively common species which featured regularly within the assemblage.
- 4.3.8 Two rarer species featured sporadically and in very low numbers, which were barbastelle and Nathusius' pipistrelle. The Sites are located at the northern edge of the range for these two species. Barbastelle are rare and Nathusius' pipistrelle uncommon in Lincolnshire according to the Lincolnshire BAP. Both species are considered to be most closely linked with woodland edge habitats and tree roosts although they will occasionally roost in buildings. A significant colony of barbastelle bats is known in Norfolk. Nathusius' pipistrelle bats are known to migrate long distances and have strongholds in the east and south east of England. Leisler's bat may also be present within the dataset. This is a rarer species but is difficult to fully separate from noctule bats by call so further analysis will be necessary.
- 4.3.9 It is considered likely that roosts for all the species recorded within the data occur either in trees within the Sites, or in trees and buildings in proximity to the Sites.

Habitat Appraisal

- 4.3.10 Initial fieldwork determined that the suitability of habitats for bats across the option land was generally low, being dominated by monoculture arable and a simple network of managed hedgerows. The arable and relatively small proportion of pasture are intensively farmed environments, receiving pesticide treatments, and would be expected to support a lower abundance and diversity of prey items upon which bats feed.
- 4.3.11 The linear natural features along which bats tend to navigate and disperse, as well as forage in preference to monoculture arable, were generally highly managed and restricted in size and structure. Woodland stands were sparse within the landscape and generally poorly interlinked, with historic hedgerow removal resulting in large open expanses of arable.
- 4.3.12 Mature trees are only sporadically present within the hedgerow networks and field edges, along with at the edges of any woodland adjacent to the option land. In-field trees are absent from the option land. Many of these trees hold potential for roosting by bats.
- 4.3.13 A relatively small number of agricultural buildings and farm dwellings (of varying levels of use and disuse) were present adjacent to the red line boundary
- 4.3.14 At **West Burton 1**, few hedgerows contained trees, mainly at the Site perimeter, especially mature ash with signs of dieback. No in-field trees were present. No buildings were present on or immediately adjacent to the Site. A small number of ditches were present at the Site perimeter only, and a tributary of the River Till ran along the northernmost outline.
- 4.3.15 At **West Burton 2**, the hedgerow network was moderately diverse, with many hedgerows containing trees and a small number of species-rich hedgerows. However, there were many fields bounded by bare ditches only. A strip of well-established broadleaved woodland ran along the middle of the western half of the Site. Woodland also bordered several fields in the north. Buildings associated with the settlement of Ingleby and Ingleby Grange (mostly agricultural) may hold potential to support roosting bats and some potential for fragmentation of movement is possible considering their proximity to the Site. There were also a small number of in-field trees.
- 4.3.16 At **West Burton 3**, the hedgerow network was extensive and without gaps and approximately half of the hedgerows contained trees. Some stands of well established broadleaved woodland was located just off Site to the west, in the north east of the site and just off the southern corner. Several farm buildings and a small number of farmhouses were located immediately off Site including several buildings capable of being surrounded by the proposed development. This Site also contained a small number of in-field trees.



Potential Constraints, Mitigation and Further Work

4.3.17 It is unclear to what extent roosting, foraging and dispersing bats are affected by large scale solar development as research evidence is sparse. Arrays have been demonstrated to increase invertebrate abundance in comparison with surrounding arable landscapes⁴ which is likely to be of benefit to foraging bats, particularly around the perimeters of the arrays. Whether bats use or avoid the centres or arrays and forage within or commute along array strings is currently ambiguous. Montag *et al* found non-significant reduction in abundance of bats from within the centres of arrays compared with surrounding arable fields. Evidence has also been proposed by Natural England which suggested bats may confuse panels with sources of water and attempt to drink from them. This builds on studies undertaken at the Max Planck institute in Zurich. This however is now widely dismissed as the experiments involved drought stressed juvenile bats and did not represent real world conditions. There is currently no evidence to indicate bats attempt to drink from or collide with panels, nor is there any indication of significant change in the sizes or abundance of populations of bats in proximity to established array sites, although research on the subject is sparse. As such the most reasonable assumption at this stage is that arrays are broadly neutral upon foraging and commuting bats with the potential to offer enhancement where they are able to promote night flying invertebrate abundance.

Roosts in Buildings and Trees

4.3.18 Clarkson and Woods should be consulted to review any proposals to prune or fell any mature or semi-mature trees, or remove built structures, within or adjacent to the option land.

4.3.19 Buildings immediately adjacent to the red line boundaries which are considered at risk of fragmentation of bat roosts by their proximity to the array should be inspected for bat roosts should be carried out to determine the potential for impacts from an array of this scale. Daytime inspections can take place at any time of year to determine levels of potential. Structures with roost potential can be followed up with emergence surveys or static detector surveys completed between May and September.

4.3.20 It may be prudent to carry out close inspections (via a climbing survey) of any semi-mature and mature trees situated in locations at risk of being encircled or at least partially enclosed by solar array. Severely decaying trees, especially ash, would be avoided. This would establish the potential for impacts upon any roosts therein. Close inspections should be preceded by ground-based inspections to ascertain levels of potential for roosting from negligible to high. Alternatively, a pre-emptive buffer of c.30+m may be appropriate. Such inspection work can be carried out at any time of year, with the potential for follow-up emergence surveys within the months of May and September inclusive.

4.3.21 Likely mitigation for roosts present in trees and buildings will revolve around adequate buffering from development in order to avoid fragmentation of populations.

Habitat Buffers

4.3.22 Pending the detailed results revealed by the static detector surveys and above further surveys, it is likely that few constraints are posed by bats, as long as steps are taken within the design of the scheme to sufficiently buffer the linear vegetated features (hedgerows of differing habitat value, ditches, watercourses and woodland edges) and any adjacent buildings containing bat roosts from the nearest array structures.

4.3.23 For development of this scale, cumulative impacts (both in combination with the other Sites and West Burton Solar Project and other potential forthcoming solar schemes) upon the already limited local dispersal route network and access to foraging habitat are possible and will need to be carefully assessed.

4.3.24 It would be prudent to apply an absolute minimum buffer zone of 8m between all such above key habitat features and the nearest panels. It can be expected that this would increase to around habitat of elevated value to bats, such as hedgerows with trees, buildings with roost potential (or confirmed roosts), woodland edges and watercourses such as the River Till and other rivers and streams. This reflects their importance to navigating and foraging bats in sustaining population movement and long-term genetic flow.

⁴ Montag, H., Parker, G.T., Clarkson, T. (2016) *The effects of solar farms on local biodiversity: a comparative study*. Clarkson and Woods and Wychwood Biodiversity, UK.



4.3.25 The following is therefore recommended as buffers from habitat edges to nearest array structures (in line with Section 3.2), subject to consultation.

- Ditches, species-poor hedgerows and hedgerows without trees: **8m**
- Minor watercourses (streams, becks), species-rich hedgerows and hedgerows with trees of low or negligible roost potential: **10m**
- Woodland, in-field trees, hedgerows with trees of moderate or high roost potential: **20m**
- Rivers, confirmed roosts in buildings or trees: **30m**

Lighting

4.3.26 Lighting can act as a significant barrier to the movement of bats, potentially also causing unlawful obstruction of roost accesses within trees or adjacent buildings. Any construction phase lighting should be carefully considered and positioned. Details of, and the need for, construction phase lighting should be reviewed by Clarkson and Woods as early as possible. Solar development does not typically require permanent lighting installation, however the need for any such lighting at substations or the proposed battery facility should be reviewed by Clarkson and Woods.

Opportunities for Enhancement and Biodiversity Net Gain

4.3.27 Suggested strategic focal locations for habitat creation and enhancement will follow as part of the bat survey report once bat survey data has been analysed. Bats are Species of Principal Importance and listed on the Lincolnshire BAP, therefore enhancements for them would be favourably received.

4.3.28 Habitat creation opportunities will revolve around the planting of new linear features such as hedgerows and tree lines within the local landscape. Replacement of former, grubbed out hedgerows (through examination of historical maps) could be a valuable technique where the scheme allows. This would benefit dispersal and navigation (providing connectivity and green infrastructure) as well as foraging resources (and in turn, increased reproductive success and population viability).

4.3.29 The most significant habitat enhancement opportunities revolve around the management of the following locations sympathetically for bats in order to maximise their productivity for invertebrates.

- Buffers between boundary habitats and the array
- Grassland habitat beneath the array
- Any off site mitigation land

4.3.30 Sympathetic management for bats generally involves leaving plants to flower before any cutting or mowing, encouragement of a tussocky sward at margins through rotational (less than annual) cutting, and grazing at a low "conversation" density of animals. It is likely that a blended approach to these management techniques would be appropriate across the option sites, to be tailored according to local nature conservation priorities and the results of the surveys.

4.3.31 Roosting opportunities should be incorporated into the scheme through the installation of tree and building-mounted bat roost boxes. A rate of approximately 1-2 boxes per 10ha of development land would be appropriate.

4.3.32 Specialist, bespoke roost buildings could be created in key flyways, for example close to the River Till or stands of woodland at intersections in the hedgerow network or at eventual habitat enhancement zones. Such features, also known as 'wildlife towers' (see Figure 10 below) would comprise small, free-standing timber, brick or block buildings with crevice and void-roosting opportunities on the vertical faces and roof pitches. Alternatively, buildings associated with the array infrastructure could be modified to include roosting features such as roost boxes, but also wooden waney-edge cladding.



Figure 10. Example of a wildlife tower and waney-edge cladding modifications for bats.

4.4 Otter

Desk Study Information

- 4.4.1 For **West Burton 1**, fourteen records of otter were present within 2km of the Site, all of which were located along the River Till and its tributaries more than 250m from the Site boundary.
- 4.4.2 For **West Burton 2**, twenty two records of otters were present within 2km of the Site, four of which were located within the red line boundary and were associated with the River Till and its tributaries. The remaining records were located beyond 250m from the Site boundary.
- 4.4.3 For **West Burton 3**, seventeen records of otters were present within 2km of the Site, one of which was located 240m west of the Site boundary with the remaining sixteen records located more than 250m from the Site boundary.
- 4.4.4 Otter are a Species of Principal Importance under the NERC Act (2006).

Field Survey Results

- 4.4.5 Habitat for otters was restricted to river corridors, wet ditches and streams present on or adjacent to the sites. No direct observations of holts or field signs for otters were encountered during the initial walkover survey.
- 4.4.6 **West Burton 1** lay approximately 500m east of the River Till and south of a substantial tributary. Several small, wetted ditches were located within the Site.
- 4.4.7 **West Burton 2** contained no significant watercourses (rivers and streams), although just beyond the eastern boundary ran the embanked River Till and field boundaries were generally characterised by moderately wide, wet drainage ditches, typically without hedgerows.
- 4.4.8 **West Burton 3** contained a limited number of wetted ditches although a significant stream was present (D2 and D3) flowing north to south through the centre of the Site. Further substantial ditches were present in the north east of the Site.

Potential Constraints, Mitigation and Further Work

- 4.4.9 Otters, as well as their resting places, are legally protected. Should any habitat clearance, excavation or engineering works be required within 5m of any ditch and 10m of any watercourse, a prior survey of the affected area for signs of otters and its suitability should be undertaken. In the event that evidence of any otter shelter is discovered (either in advance through a specific otter survey or during supervised works), works may require a licence from Natural England in order to proceed. In the absence of evidence of a holt or other shelter, the potential for disturbance or damage to habitat should be mitigated for by carrying works out under an Ecological Watching Brief attended by an experienced ecologist.
- 4.4.10 Otters are able to range over considerable distances and use small streams and ditches occasionally for dispersal and reaching inland waterbodies for hunting. Consequently, the potential for otters within field



boundary features should not be entirely ruled out at any of the Sites. The most effective design based mitigation would be to adopt sufficient buffers (>10m) between watercourses and the nearest zone of development activity.

- 4.4.11 It may be prudent to supplement the baseline by investigating the presence of these species within the watercourses on site, although further consultation with the LPA and Natural England would help determine whether this would be advantageous. Such a survey would consist of walking along on bank of each watercourse or ditch and examining, either in detail or intermittently where access permits, for signs of otter such as spraints, tracks and feeding remains. This can be carried out at any time of year but is best done in the period autumn or spring when activity is greatest and water levels generally lower.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.4.12 All sites were considered relatively well connected to significant river and stream networks, with the River Till being located in close proximity to both **West Burton 1** and **West Burton 2**, while the River Trent is associated with **West Burton 3**, as well as extensive waterbodies associated with the Lincoln Golf Course just off the south west corner of the Site.
- 4.4.13 Habitat enhancements for otter are mostly limited to the favourable management of river and stream banks to encourage a dense growth of vegetation cover in the form of tussocky grassland, as well as thick shrubs and mature trees. Consequently, new tree planting schemes could include a small degree of planting of alder, willow and birch whips at stream and river banks. Grassland field margins should be left to grow long and tussocky within approximately 5-10m from streams and rivers where possible.
- 4.4.14 Depending on the results of the field surveys, further opportunities to provide habitat links and improve connectivity between watercourses potentially by deepening or wetting ditches and planting scrub and trees may be possible. Any new waterbodies (for example as GCN enhancement) and swales may also contribute positively to otter conservation.
- 4.4.15 The potential for pollution events and discharge of sediments and excess agricultural and soil runoff during construction should be avoided through best practice construction measures.

4.5 Water Vole

Desk Study Information

- 4.5.1 For **West Burton 1**, 30 records of water vole were present within 2km of the Site, all of which were located more than 250m from the Site boundary.
- 4.5.2 For **West Burton 2**, 92 records of water vole were present within 2km of the Site, ten of which were located within the red line boundary between 1990 and 2012. 58 records were located beyond 250m of the Site with the exact location of a further 24 records not provided.
- 4.5.3 For **West Burton 3**, 62 records of water vole were present within 2km of the Site, three of which were located within 250m of the Site boundary and the remaining records location more than 250m from the Site boundary.
- 4.5.4 Water voles are a Species of Principal Importance under the NERC Act (2006) and listed on the Lincolnshire BAP.

Field Survey Results

- 4.5.5 As with otters, suitable habitat for water vole was restricted to river corridors, wet ditches and streams present on or adjacent to the Sites. Habitat requirements for water vole are simpler than for otter, just requiring shelter (diggable earth banks), aquatic vegetation and reliable access to water. Considering the abundance of suitable habitat across much of each Site in the form of strong vegetated ditch networks, target noted habitat for this species and the large number of nearby records, **all Sites**, are likely to support water vole to some extent. It is likely that **West Burton 2** would contain the highest population of water voles.

Potential Constraints, Mitigation and Further Work

- 4.5.6 Water voles are legally protected from harm as well as disturbance while within burrows. As with otters, should any habitat clearance, excavation or engineering works be required within 5m of any ditch and 10m of any watercourse, a prior survey of the affected area for signs of water voles and its suitability should be undertaken.



In the event that evidence of any burrows is discovered (either in advance through a specific water vole survey or during supervised works), works may require a licence from Natural England in order to proceed. In the absence of water voles signs, the potential for minor disturbance or damage to habitat should be mitigated for by carrying works out under an Ecological Watching Brief attended by an experienced ecologist. The most effective design-based mitigation for water voles would be to adopt sufficient buffers (>10m) between watercourses and the nearest zone of development activity.

- 4.5.7 It is recommended to supplement the baseline by investigating the presence of water vole within the watercourses on Site. Such a survey could be combined with an investigation for otters and would consist of walking along on bank of each watercourse or ditch and examining, either in detail or intermittently where access permits, for signs such tracks, droppings, feeding remains, runs and burrows. This can be carried out at any time of year but is best done in the period March to October inclusive when activity is greatest and water levels generally lower. Two visits are required, one in the spring and one in the autumn.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.5.8 Enhancements for water voles are similar to those given for otters and revolve around the preservation of stream and river banks, protection from disturbance and damage by buffering and avoidance of pollution events.

4.6 Dormouse

- 4.6.1 Dormice are not known to be present in the Lincoln to Gainsborough area and are only very locally distributed in Lincolnshire at all. No records for dormice were revealed by the desk study. Habitats on the Sites were considered poor for dormice, being restricted to managed simple hedgerow networks alone. It is highly unlikely that the Sites could be functionally linked to any populations of dormice, therefore this species is not considered a potential constraint to development.

4.7 Great Crested Newts and Other Amphibians

Desk Study Information

- 4.7.1 For **West Burton 1**, 7 records of toad were present in the dataset, the closest being located 900m west of the Site. 65 great crested newt records are present beyond 250m of the Site, the closest being 1.4km north-west of the Site. A small number of other amphibian records (smooth newt and common frog) were revealed between 250m and 2km from the Site.
- 4.7.2 For **West Burton 2**, 11 records of toad were present in the dataset, the closest being located 460m north of the Site. 23 great crested newt records are present beyond 250m of the Site, the closest being 1.9km south-west of the Site. 28 records of common frog and 22 records of smooth newt were revealed between 250m and 2km from the Site.
- 4.7.3 For **West Burton 3**, 18 records of toad were present in the dataset, one of which was located within 250m of the Site. 6 great crested newt records are present with exact locations not provided. 18 records of common frog and 9 records of smooth newt were revealed between 250m and 2km from the Site.
- 4.7.4 Great crested newt and common toad are Species of Principal Importance under the NERC Act (2006) and newts are listed on the Lincolnshire BAP.

Field Survey Results

- 4.7.5 At **West Burton 1**, **no ponds** were visited to test for GCN environmental DNA. The only pond within 250m of the Site boundary was located on private property.
- 4.7.6 At **West Burton 2**, 13 ponds were visited to test for GCN and **none were positive**. Seven of these ponds were dry at the time of survey.
- 4.7.7 At **West Burton 3**, 11 ponds were visited to test for GCN and **two were positive (Pond 4 and Pond 9)**. Three ponds were found to be dry, one was deemed unsafe (agricultural slurry pit) and one gave an 'indeterminate' result due to high sediment or pollutant content. See Figure 11 below.




Figure 11. GCN Positive Ponds – West Burton 3 Ponds 4 and 9.

- 4.7.8 An indeterminate result occurs where factors such as the presence of contaminants or silt make DNA extraction difficult, as in the case of slurry pits, or waterbodies subject to accumulated leachate or agricultural runoff.
- 4.7.9 Several waterbodies were found to be dry. This is considered partly as a result of the period of warm weather at the time of surveys, and the fact that many of the mapped waterbodies were in actual fact ephemeral field ponds or are subject to regular drying. Drying out in three or more years in every ten is considered to significantly reduce the suitability of a pond for GCN.
- 4.7.10 GCN records are very sparsely distributed within the West Lindsey district, reflecting the fact that the intensive agricultural land-use which characterises the landscape provides generally poor habitat for this species. Nevertheless, the West Burton project sites are considered to be consistently sub-optimal for GCN in terms of intrinsic habitat value and local population densities.

Potential Constraints and Mitigation

- 4.7.11 Legal protection afforded to GCN extends to their habitat (breeding and resting places), which includes both aquatic and terrestrial types. Arable and actively cut grassland or grazed pasture, which make up the vast majority of the option land, are considered sub-optimal habitats. Scrub, tussocky or uncultivated grassland, woodland and hedgerows are all optimal (as well as wetland and other aquatic habitat). These habitats typically occur within field margins and boundaries and at field headlands, or in a relatively low number of uncultivated fields. Hard standing and bare ground are considered unsuitable.
- 4.7.12 Despite the majority of the option land - or certainly the likely development footprints - occupying sub-optimal habitat, a zoned approach to the risk of unlawful habitat clearance or direct disturbance to GCN should be adopted, in accordance best practice guidance. This recognises the fact that the likelihood of encountering newts within potentially suitable habitat decreases with distance from ponds known to support them. Table 8



provides general constraints during the construction phase and working methods recommended for all Sites containing or adjacent to positive GCN ponds.

Table 8. Summary of Constraints and Working Methods in Proximity to GCN Breeding Ponds

Zone (Distance from perimeter of nearest known breeding pond)	Temporary or Permanent Loss of, or Disturbance to:	
	Optimal Habitat	Sub-Optimal Habitat
0-100m	<ul style="list-style-type: none"> • Licence from Natural England likely to be required – <i>see further information below</i>. • Newt exclusion exercise likely required, involving installation of partially buried fencing and pitfall traps, to be checked daily for 30+ days to declare habitat clear of GCN in advance of works commencing. • Constrained to active season (March to October inclusive, weather depending) in order to avoid impacts on hibernating individuals. • Ecological Clerk of Works (ECoW) required to supervise. • Destructive Search methodology to precede works – consists of a staged cutting (mowing or strimming) of vegetation before being methodically removed using an excavator. 	<ul style="list-style-type: none"> • Licence from Natural England potentially required, but unlikely. To be informed through pre-application consultation with LPA and NE. • Due to negligible hibernation potential within these habitats, works likely to be constrained to winter period (November to February inclusive, weather depending). • Ecological Clerk of Works (ECoW) required to give tool-box talk to contractors.
101-250m	<ul style="list-style-type: none"> • Licence only required where approx. 5000m² (0.5ha) impacted. • Additional constraints as above. 	<ul style="list-style-type: none"> • Licensing constraints unlikely - to be informed through pre-application consultation with LPA and NE. • Potential for restriction to winter working methodology.
251m +	<ul style="list-style-type: none"> • Licence only required where approx. 50,000m² (5ha) impacted. • Additional constraints as above. 	<ul style="list-style-type: none"> • Licensing constraints highly unlikely.

- 4.7.13 The above construction phase constraints will be the subject of discussion with LPA consultees and Natural England. An acceptable approach to construction during the DCO process will need to be established, therefore the information given in Table 6 above is indicative at this stage subject to amendment. The final, agreed approach to construction and licensing will be detailed within an eventual EIA (and its great crested newt survey report technical appendix) and Construction Ecological Management Plan, or similar document.
- 4.7.14 Currently, licensing for great crested newts in this region generally involves recourse to a traditional mitigation licence. This typically requires the need for an exclusion, trapping and translocation exercise where suitable habitats in close proximity to breeding ponds are to be lost or temporarily affected. This is seasonally constrained and may require 30 or more days to undertake prior to construction commencement. Licence determination post-construction also takes a statutory 30-day period.
- 4.7.15 An alternative option exists, known as the Low Impact Class Licence, which is applicable for developments where impacts in proximity to breeding ponds are considered to be small, and do not affect the ponds themselves. These licences are streamlined and far less onerous to apply for and have determined. Should the scheme be designed to minimise impacts to suitable habitats within 100m as far as possible, this licence type may be available. Further consultation will be necessary to determine this.
- 4.7.16 Finally, it is probable that by the time the scheme is consented, Lincolnshire will be added to the regions eligible to use the District Licence scheme for GCN mitigation. This scheme permits all but the most damaging impacts to breeding ponds and habitat in return for a tailored and proportionate financial contribution to local great crested newt conservation schemes.



Further Work

- 4.7.17 To underpin the DCO application and finalisation of ES, CEMP and any future licence, water testing of ponds within 250m of the site should be carried out. Best efforts to gain access to third party land should be made. Samples can only be taken between the months of **mid-April to end June each year**.
- 4.7.18 It is recommended that a proportion of the indeterminate or dry ponds encountered during the 2021 surveys are re-visited in 2022 for completeness and to demonstrate best efforts.
- 4.7.19 Survey requirements for the cable routes should be determined and planned for the 2022 survey season.
- 4.7.20 Recommendations and constraints given above would apply to any newly confirmed breeding ponds.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.7.21 Construction of new waterbodies within 250m of known breeding ponds would improve the long-term viability of currently sparse and poorly connected local populations. This would contribute substantially to local and national green infrastructure policy and the restoration of local biodiversity.
- 4.7.22 Planting of new hedgerows, woodland strips and scrub/shrub vegetation in locations strategic to improving corridors for dispersal between existing (and any new) ponds would serve to improve green infrastructure for amphibians and long-term population sustainability.
- 4.7.23 Management of field edges, hedgerow/woodland/ditch/watercourse buffer zones, wayleaves and easements within 250m of known breeding ponds to create coarse, tussocky grassland or meadow habitat would also contribute to the above aims.
- 4.7.24 Sympathetic management of fields beneath arrays within 250m of known breeding ponds to form a taller, more diverse grassland sward (managed through low-density/intensity conservation grazing or collection of a late-season hay cut).
- 4.7.25 As set out in Section 3.4, basic water and habitat quality enhancements at the ponds on Site at or immediately adjacent to **West Burton 2 and West Burton 3** would be of benefit for any amphibian populations present. This includes selective deepening and planting.

4.8 Reptiles

Desk Study Information

- 4.8.1 At **West Burton 1**, 3 historical (pre-2000) records for common lizard were located within 2km of the Site, as well as 18 records for grass snake (6 post 2000) located beyond 250m from the Site.
- 4.8.2 At **West Burton 2**, 1 historical (pre-2000) record for common lizard was located within 2km of the Site, as well as 29 records for grass snake (9 post 2000) two of which were located within the red line boundary for the Site and three of which were located within 250m of the Site.
- 4.8.3 At **West Burton 3**, 1 record for common lizard was located within 250m of the Site, as well as 19 records for grass snake (6 post 2000) all of which were located beyond 250m of the Site. 5 records of slow worm were recorded within 2km of the Site with the exact locations not provided.
- 4.8.4 Reptiles are Species of Principal Importance under the NERC Act (2006).

Field Survey Results

- 4.8.5 Habitats for reptiles are generally limited in quality and extent across all the sites, being restricted to hedgerow bases, tussocky field margins and woodland edges only. Almost universally, the development will be sited on land of poor habitat quality for reptiles. Furthermore, the desk study data shows a lack of records for reptile species within 2km of the sites, with an absence generally within 250m.
- 4.8.6 **West Burton 1** contained occasional habitat of potential suitability for reptiles in field margins and the banks of more substantial watercourses/ditches only.
- 4.8.7 At **West Burton 2**, Most of the grass margins were uncultivated or managed and had become tussocky. Several of these, including the feature at TN8 were noted to be of particularly good value to reptiles, although none were seen.



- 4.8.8 At **West Burton 3**, A grass snake was seen in the vegetated banks of a stream at D2, while a common lizard was seen within the hedgerow margin in F8 (TN5). Margins were mostly narrow but were uncultivated and unmanaged and had become tussocky. Several of these were noted to be good for reptiles.

Potential Constraints, Mitigation and Further Work

- 4.8.9 Reptiles are legally protected from reckless and intentional harm, therefore it is recommended that all field margins and hedgerows, as well as target noted locations of discrete reptile habitat are retained and protected wherever possible.
- 4.8.10 Given the limited records, habitat quality and extent within the development footprint, it is unlikely that a targeted reptile survey would be necessary. If any Site were to benefit from one, it would probably be **West Burton 3**, as a population has been observed and removal of habitats features conducive to reptiles (hedgerows and field margins) may be required for access and logistical purposes. Elsewhere, should proposals seek to significantly remove or alter boundary features, the requirement for a reptile survey may need to be re-assessed. Further consultation with LPAs would determine acceptability of this approach.
- 4.8.11 It should be possible to avoid any impacts on reptiles through the installation of sufficient protective fencing, adherence to a construction methodology which avoids damage to such habitats and the avoidance of any widening of field accesses. A suitable buffer of at least 5m from these habitats would ensure accidental damage during construction and ongoing maintenance is avoided.
- 4.8.12 A best practice approach to habitat clearance and management is considered appropriate. Where habitat suitable for reptiles (all field margins, hedgerows, tussocky grassland and river corridors) is proposed for clearance, a Reasonable Avoidance Method Statement should be followed. Depending on the amount of land affected, this is likely to involve the phased removal of vegetation in order to dissuade reptiles from that area, followed by a destructive search supervised by an ecologist. Should particularly large areas of habitat be earmarked for removal, a survey and translocation exercise may be a last resort, although such an approach is considered unlikely to be required.
- 4.8.13 Should any of the arable fields become dominated by a long or tussocky sward, either through the cessation of cultivation or cutting prior to development, site clearance/preparation may need to be carried out in a sensitive manner. This is to avoid impacts to any reptiles which may have dispersed onto the development footprint as the habitat has increased in suitability. A suitable habitat cutting/clearance methodology (Risk Avoidance Method Statement) would be set out in an eventual Construction Environmental Management Plan.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.8.14 Optimal reptile habitat includes tussocky grassland, scattered scrub and ruderal vegetation interspersed with physical features conducive to basking on and hibernating in. Considerable net gains for
- 4.8.15 The local area is unlikely to support significant populations of reptile species and therefore enhancements specifically for these species are of a low priority, however the following basic measures are suggested.
- 4.8.16 The creation of a number of appropriately located reptile hibernaculum would improve the Sites' habitat suitability by providing features within which to hibernate during the winter and to bask during the summer. The construction of these habitat piles using partially buried dead wood, earth and stone would also provide invertebrate prey items. Further advice on numbers and locations can be given as the proposals evolve.
- 4.8.17 The reversion of intensive agriculture to diverse grassland is encouraged as this would improve the plant species diversity and habitat structure within the Sites. In turn, this would provide improved foraging and hibernation habitat for reptiles. Advice on the favourable management of the grasslands on Site for the benefit of reptiles and other wildlife would be agreed with you and provided within a Landscape Environmental Management Plan.

4.9 Birds

Desk Study Information

- 4.9.1 At **West Burton 1**, numerous records of 40 species of notable birds, or birds of conservation concern, were revealed by the Desk Study, none of which were located within the Red Line Boundary of the Site, as detailed



in Appendix B. One record of both kingfisher and skylark were recorded within 250m of the Site with the remaining records located between 250m and 2km from the Site. The recorded notable species comprise farmland birds such as corn bunting, quail, barn owl and turtle dove as well as waders and raptors.

- 4.9.2 For **West Burton 2**, numerous records of 44 species of birds were recorded, as detailed in Appendix C. Reed bunting has been recorded within the Red Line Boundary of the Site with kingfisher and lapwing recorded within 250m of the Site since 2000. All other bird species were recorded beyond 250m from the Site, including curlew, tree sparrow and yellowhammer.
- 4.9.3 For **West Burton 3**, numerous records of 57 bird species were recorded within 2km of the Site as detailed in Appendix D. One record of house sparrow was located within the Red Line Boundary and starling and song thrush have been recorded within 250m of the Site. All other records were located beyond 250m of the Site, including species such as yellowhammer, yellow wagtail, lapwing and barn owl.
- 4.9.4 Farmland birds are listed on the Lincolnshire BAP and many species are Species of Principal Importance under the NERC Act (2006).

Field Survey Results

- 4.9.5 Four daytime breeding bird surveys and one dusk, nocturnal bird survey (with a focus on quail) has been carried out. Winter bird surveys are scheduled for November 2021 to February 2022.
- 4.9.6 In general, considering the broad similarities in habitat arrangement, topography, field size and agricultural management, the breeding bird species assemblage is consistent across the option sites. Results can be broadly divided into those for ground-nesting birds, birds of hedgerows and boundaries and other bird species.

Ground-nesting Birds

Skylark

- 4.9.7 This is a red-listed species on account of its declining population trend as a result of agricultural intensification and land-use change. It is also a Species of Principal Importance (SPI) under the NERC Act 2006. Skylark are a resident species whose numbers swell each winter from an influx of visitors from northern Europe. Skylark require long, unbroken sightlines in grassland (including arable or set-aside up to 40cm high) of at least approximately 200m for predator avoidance.
- 4.9.8 Skylark were recorded on **all Sites** in varying densities. Particularly dense populations were located at **West Burton 2** and **West Burton 3** as these featured some of the largest arable fields within a similarly open landscape. In addition, some of the barley was planted in the spring, allowing for greater nesting success on second broods (due to the lower sward height) and better wintering habitat in the form of stubbles. Together, West Burton supports significant populations of skylark, although this would be expected to be in line with population densities in the local landscape.
- 4.9.9 Winter-sown wheat - as is ubiquitous across most of the Sites - is considered to be a suitable but sub-optimal habitat for skylark on account of its growth above 40cm at a time when skylark are looking to have second or third broods in the mid-late summer. It can reasonably be assumed that a large proportion of the nests present, if not all, would be displaced from solar arrays. There is no robust, long-term evidence indicating that skylark nest within solar arrays, although the reversion from arable to grassland in solar development has been shown to improve foraging opportunities for skylark which are able to include array land within their adjacent territories. This effect is likely to increase nesting and breeding success in adjacent suitable (non-array) habitats. Some nesting may persist within buffers and wayleaves, although it is considered that this reflects a tendency for site-fidelity which may persist for approximately one to three seasons post-construction.

Yellow wagtail

- 4.9.10 For the same reasons as skylark, yellow wagtail are also red listed, and a SPI. Yellow wagtail migrate to the UK from Africa each spring. Yellow wagtail are a far less numerous bird than skylark and were recorded across all Sites at significantly lower rates than skylark. The site supporting greater numbers was **West Burton 2**. As for skylark, it is likely that yellow wagtail nests would be displaced through solar development, although solar development could be expected to improve foraging opportunities for birds with nearby territories.



Grey Partridge

- 4.9.11 This is a red listed species and an SPI, typical of lowland arable farmland although having suffered marked recent declines. Grey partridge were recorded in low to moderate numbers across **all Sites**. These are not thought to be the result of introduction for shooting. The effects of solar development on grey partridge is unknown. Preferring field edges and proximity to sources of cover, grey partridge may continue to use solar arrays, although potentially at the edges and in lower overall densities. It may also transpire that solar array may provide a desirable shelter from nearby game shooting and therefore provide a valuable refuge for the population.

Birds of Field Boundary Habitats

- 4.9.12 Significant populations of birds typical of hedgerows, woodland edges, scrub and river corridors in a lowland agricultural setting were recorded throughout the Sites, principally yellowhammer, linnet, common whitethroat, lesser whitethroat, tree sparrow, reed bunting and great spotted woodpecker. Many of these birds will forage within arable field edges or nest in ditches, hedgerow bases or grassy margins as well as the hedgerows themselves. It is expected that the assemblage and abundance would not be significantly affected provided that sufficient buffering is designed into the schemes. These species have been seen to persist on established small and medium-scale solar arrays, although impacts are largely untested at this scale. Given the scale of proposals and likely unbroken expanse of array, it would be prudent to instigate an increased degree of buffering compared to small and medium sized array schemes.

Other Birds

- 4.9.13 Curlew and lapwing are red listed species and also SPIs. These waders were recorded at **West Burton 2**, close to the banks of the River Till. Solar development can be expected to displace nesting locations for these species for the same reasons of predator surveillance as listed for skylark.
- 4.9.14 Turtle dove are a red listed species and an SPI and were not observed at any site. This species is increasingly rare and in danger of extinction in the UK. Turtle dove rely on uncultivated land and arable weeds for seeds, as well as tall hedgerows, open woodland and scrub. Again, no research exists on the effect of solar development on turtle dove, however opportunities exist for the enhancement of foraging habitat and planting of nesting habitat for this key species of local conservation concern.
- 4.9.15 Barn owl, little owl, short-eared owl and tawny owl were all recorded during the evening surveys, with barn owl being recorded at almost every site in good numbers. Tawny owl and little owl were only recorded in stands of woodland adjacent to the option land. Barn owl and short-eared owl were the most likely owl species to be recorded within the arable fields themselves. River banks, especially at the River Till were regularly-used foraging corridors for these species. The impacts of solar development on owls are unclear as barn owls in particular as associated with open hunting habitat. However, it is likely that tussocky margins and buffers, as well as sympathetically managed grassland beneath arrays (longer grassland suitable for voles and other small rodents) would support a far greater abundance of prey items than intensive arable.
- 4.9.16 Buzzard, peregrine, hobby, kestrel, marsh harrier and red kite were all observed during the bird surveys. Nesting buzzard were regularly recorded within woodland edge at the majority of the sites.

Potential Constraints and Options for Mitigation

- 4.9.17 On account of their status as birds listed on Schedule 1 of the Wildlife and Countryside Act 1981 (as amended) nests of hobby, peregrine, barn owl, quail and red kite will need to be protected from disturbance during any development activity. Consequently, pre-commencement precautionary survey work is likely to be required to establish risks immediately prior to the construction phase.
- 4.9.18 Similarly, all nests for other species are protected from harm, therefore any potential nesting habitat clearance will need to be carried out either during the period September to February inclusive, under the supervision of an ecologist, or following further survey to confirm absence.
- 4.9.19 In order to ensure that boundary habitats remain suitable for use by the species recorded, as well as being able to be re-visited and discovered, it is recommended that sufficient buffers to the nearest arrays are implemented. The size of these should be coordinated with other constraints, for example bats, in due course following the completion of survey work and analysis. The following is therefore likely to be recommended,



subject to consultation. Hedgerows: 10m. Ditches and minor watercourses: 15m. Woodland, in-field trees and major watercourses: 20m. Ancient woodland: 30m

Skylark and Other Ground Nesting Birds

- 4.9.20 Impacts on ground nesting birds can be mitigated for either by the creation of newly-available (i.e. not already suitable) compensatory nesting habitat, or the enhancement of existing habitat by the improvement of foraging opportunities causing an increase in carrying capacity and likely knock-on nesting success. Given the scale of likely impacts on these species, mitigation should be achieved through a blend of different mitigation techniques and land management approaches on Site and, potentially locally off-Site. As has been described, solar arrays are not considered suitable nesting habitat for ground nesting species which require long sightlines for predator monitoring, therefore mitigation for these will need to occupy contiguous blocks of land free of solar array and other structures.
- 4.9.21 On Site, land unviable for development could be managed specifically for ground nesting birds, ideally reverting from intensive arable to non-rotational set-aside or meadow for the greatest capacity to absorb displaced territories. Careful site selection will be necessary as suitable mitigation land for skylark in particular usually requires a radius of >200m from all vegetation and structures above 50cm in height.
- 4.9.22 Off site, winter sown cereals can be reverted to spring (March) sown crop to enable existing birds to successfully rear a second or third brood. This technique should be supplemented through the inclusion of 'bird foraging plots' whereby 5x5m squares of unsown land are introduced at a rate of at least 2 per hectare into fields by temporarily halting the seed drill during sowing. This has the effect of increasing invertebrate food item abundance, improving the breeding success, number of young reared and densities of territories able to be supported. Additionally, agricultural land can be reverted from unsuitable or sub-optimal habitat to meadow, long cut-rotation silage (>7weeks), and have reduced application of inorganic fertiliser and insecticide. Again, only large, open fields with vegetation below 40-50cm during the majority of the breeding season would be considered suitable.
- 4.9.23 The precise quantum of land required to achieve an acceptable mitigation for the species can be calculated once bird survey data has been analysed. This would then be refined according to the combination of mitigation techniques listed above that are employed.
- 4.9.24 Furthermore, consultation with Natural England and Local Authorities would be key in establishing an acceptable approach. Indeed, Local Authorities (as well as consultees such as the RSPB and BTO) may be in a position to assist with recommending local conservation initiatives to which the schemes can contribute. The above mitigation techniques can be expected to be of benefit to a wide variety of birds, not limited to the listed ground nesting species.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.9.25 Beyond the mitigation options for ground-nesting birds outlined above, substantial nesting and foraging habitat can be created through the planting of new hedgerows, lines of trees and scrub, as well as the management of buffers, wayleaves and other easements for invertebrate and seed eating species. These measures can be tailored to each site and particular bird species of note.
- 4.9.26 Buffer areas and easements can be managed preferentially for different species. Where raptors such as owls and kestrels are targeted, tussocky grassland valuable for small rodents can be encouraged. This can be diversified with ruderal and flowering meadow plants to be of greater benefit to invertebrate-eating species such as whitethroat, skylark and yellow wagtail. Hobby can be targeted through the inclusion of waterbodies to encourage dragonflies. Further options would be discussed within the dedicated bird survey reports.
- 4.9.27 An additional consideration for siting grassland enhancement measures would be the proximity to any on or off-site land secured for skylark mitigation. The success of skylark nesting enhancement can be further improved by better access to productive foraging grounds. As young skylarks are almost exclusively fed on invertebrates, it would be of benefit to have these treatments adjacent to known or targeted skylark nesting habitats. While arrays are not known to support optimally nesting skylarks, they have been found to support foraging skylarks.
- 4.9.28 Nesting opportunities should be incorporated into the scheme through the installation of tree and building-mounted bird boxes. A rate of approximately 1-2 boxes per 10ha of development land would be appropriate.



- 4.9.29 Specialist boxes for raptors and owls can be installed in appropriate key locations within the schemes.

Further Survey Considerations

- 4.9.30 Wintering bird surveys will need to be carried out to determine the potential for impacts upon wetland birds, winter migrants and bird associated with the Humber Estuary SPA.
- 4.9.31 It may be prudent to carry out further daytime inspection of buildings and mature trees adjacent to the sites capable of being encircled or at least partially surrounded by arrays, to determine any impacts on movements or access to habitat by birds such as barn owls nesting or roosting within them.

4.10 Invertebrates

- 4.10.1 Habitat quality for invertebrates within the development sites is generally low, owing to the intensive agricultural land use and regularity of pesticide use. Boundary habitats are also generally poor for invertebrates, while the River Till corridor, waterbodies and watercourses represent some elevated habitat value. The desk study data on invertebrates will be fully analysed in due course, alongside further consultation, to determine whether any further targeted invertebrate survey may be useful. At this stage, this is considered unlikely.

Desk Study Information

- 4.10.2 At **West Burton 1**, the Desk Study revealed historic records of three notable invertebrate species including small heath and wall butterflies and large-mouthed valve snail.
- 4.10.3 At **West Burton 2**, the Desk Study revealed records of numerous notable invertebrate species including small heath and wall butterflies, large-mouthed valve snail and 49 species of moth, which are listed in full in Appendix C.
- 4.10.4 At **West Burton 3**, the Desk Study revealed records of numerous notable invertebrate species including small heath and wall butterflies, mud snail and 42 species of moth, which are listed in full in Appendix D.

Field Survey Results

- 4.10.5 Habitat quality for invertebrates within the development sites is generally low, owing to the intensive agricultural land use and regularity of pesticide use. Boundary habitats are also generally of lower to moderate value for invertebrates, while the species rich hedgerows, trees, River Till corridor, waterbodies and watercourses represent relatively elevated habitat value.

Potential Constraints, Mitigation and Further Work

- 4.10.6 The desk study data on invertebrates did not raise any concerns regarding the need for further survey.

Opportunities for Enhancement and Biodiversity Net Gain

- 4.10.7 The creation of more diverse grassland over time (both under panels and within field margin buffer zones) should provide an increase in habitat value for invertebrates. Alternatively, a new meadow or diverse grassland can be created by cultivation and over seeding, followed by monitoring and timed cutting as described in the Habitats section. The final approach can be discussed for inclusion within an Ecological Management Plan.

4.11 Other Protected Species and Species of Conservation Concern

Desk Study Information

West Burton 1

- 4.11.1 29 records of brown hare are present between 250m and 2km from the Site.
- 4.11.2 2 records of feral ferret are located 1.3km from the Site.
- 4.11.3 58 records of hedgehog are located within 2km of the Site, one of which was located within the Red Line Boundary and 3 records within 250m of the Site.



- 4.11.4 1 historic record of harvest mouse within 2km of the Site.
- 4.11.5 21 records of European eel (2 post-2000) and 24 records of spiny loach (11 post-2000) within 2km of the Site.

West Burton 2

- 4.11.6 53 records of brown hare are present within 2km of the Site, one of which was located within the Red Line Boundary.
- 4.11.7 2 records of feral ferret are located 1.6km from the Site.
- 4.11.8 131 records of hedgehog are located within 2km of the Site, four of which were located within the Red Line Boundary and 8 records within 250m of the Site.
- 4.11.9 3 historic records of harvest mouse within 2km of the Site.
- 4.11.10 65 records of European eel (15 post-2000) and 25 records of spiny loach (11 post-2000) are present within 2km of the Site.
- 4.11.11 The only flowering plant records present are for Tubular Water-dropwort, of which there were 3 records within 2km of the Site.

West Burton 3

- 4.11.12 127 records of brown hare are present within 2km of the Site, four of which were located within the Red Line Boundary.
- 4.11.13 3 records of feral ferret are present within 2km of the Site, two of which were located within the Red Line Boundary.
- 4.11.14 167 records of hedgehog are located within 2km of the Site, 14 of which were located within the Red Line Boundary and 20 records within 250m of the Site.
- 4.11.15 5 records of harvest mouse are present within 2km of the Site, 2 of which are post-2000.
- 4.11.16 21 records of European eel (11 post-2000) are present within 2km of the Site.
- 4.11.17 The only flowering plant records present are for annual knawel, of which there were 1 record within 2km of the Site.

Field Survey Results

- 4.11.18 At **all sites**, large numbers of brown hare were noted within the fields. **All sites** were conducive to the presence of species such as hedgehog, polecat and other small mammals within hedgerows and field margins. Harvest mice are assumed to be present to some degree. The larger watercourses are likely to support several species of fish and other aquatic life.

Potential Constraints, Mitigation and Further Work

- 4.11.19 It is unlikely that significant effects on any of these species would arise from the development provided that steps are taken to protect existing boundary features and maximise their habitat value through simple and sympathetic management practices for the life of the scheme. Mitigation measures given for other species above would serve species mentioned here well. It has been observed that brown hare, in particular, appear to benefit from solar array installations and favour the shelter and longer grass associated with them in preference to pasture grassland. Security fencing is not considered likely to impede movement by these species as long as the mesh size is large enough (e.g. standard deer fencing).



5 FURTHER WORK AND NEXT STEPS

5.1 Recommended and Optional Further Surveys

5.1.1 As derived from the above species and habitats discussions, the following further surveys are either recommended or suggested pending the outcome of consultation on the current proposed survey and assessment scope.

Further Survey to Inform DCO Application			
Species/Item	Survey Type	Timing	Comments
Great Crested Newts	Water sampling	Mid-April and end-June 2022	Survey of all accessible ponds on third party land within 250m of red line boundaries, plus on-site dry ponds. Survey of ponds in proximity to cable route likely required.
Birds	Wintering birds	Monthly visits between November 2021 and February 2022	Scheduled
Birds	Tree and building inspection	Any time of year, best March to September	Can be carried out now or subject to consultation with LPA/NE.
Bats	Static bat detector survey	Two remaining deployments (August and September 2021)	Ongoing
Bats	Tree and building inspection	Daytime work: any time of year. Emergence Survey (if needed) May to September inclusive.	Can be carried out now or subject to consultation with LPA/NE.
Otters and Water Voles	Watercourse inspection	One or two visits, ideally autumn and spring	Can be carried out now or subject to consultation with LPA/NE. Cable routes to be considered also.
Reptiles	Presence/absence survey of boundary habitats	Spring and early autumn best	Unlikely to be required but can be carried out now or subject to consultation with LPA/NE.
Cable Routes	Phase 1 Walkover	Any time of year for walkover	Further survey for GCN, otters and water voles, designated habitats as a minimum likely to be required.



5.2 Anticipated Reporting/Design Milestones

- Input into Pre-App/Early Consultation Docs. Basis formed by PEA but with relevant additions – **September 2021**
- Opinions received on proposed survey scope and early mitigation approach – **August-October 2021**
- Breeding Bird Survey Report – **October 2021** –
Expanded thereafter following completion of any tree/building surveys.
Will enable finalisation of on and off-Site mitigation requirements for skylark and associated species.
- Bat Survey Report – **October/November 2021**
Expanded thereafter following completion of any tree/building surveys.
Will enable finalisation of buffer widths from hedgerows and trees to security fence.
- Preliminary Biodiversity Net Gain Analysis – **October/November 2021** –
Will facilitate habitat management plan and landscape enhancement design.
- Wintering Bird Survey Report – **March/April 2022** –
May have implications for on and off-Site bird mitigation if not already catered for.
- Otter and Water Vole Survey Report (if required by consultees – considered likely) - **March/April 2022**
Interim report can be provided on basis of 2021 data for purposes of PEIR, scoping and consultation.
Will help refine recommendations for watercourse buffering and habitat management.
- Great Crested Newt Survey Report following 2022 survey of off-Site ponds – **May 2022**
Interim report can be provided on basis of 2021 data.
Will refine constraints in proximity to some ponds.
- PEIR – Spring 2022
- ES Chapter – Summer/Autumn 2022
- Construction Ecological Management Plan (or similar) – **TBC in support of PEIR/ES**
- Landscape and Ecological Management Plan (or similar) – **TBC in support of PEIR/ES**
- Final Biodiversity Net Gain Analysis Report - **TBC in support of PEIR/ES**

5.3 Construction and Landscape Environmental Management Plans (CEMP and LEMP)

5.3.1 The PEIR and ES will likely need to be supported by a document setting out how construction-phase impacts upon sensitive ecological receptors will be avoided and minimised. Typically, a Construction Environmental Management Plan is prepared in collaboration with other environmental and landscape disciplines and an ecology chapter produced. Alternatively, a specific Construction Ecological Protection Plan can be produced as a standalone document.

5.3.2 This document would set out the following:

- Details of protective and permanent fencing including distances from habitat features etc.
- Working methods adopted to avoid accidental damage (including root compaction, contamination and pollution) to retained features such as trees, hedgerows and watercourses.
- Examples of and a plan to show where signage will be installed.
- The roles of different site personnel in protecting and maintaining retained habitat during construction.
- The role of an Ecological Clerk of Works to ensure inspections are carried out and that activities carrying a risk of harm to protected and notable species and habitats can be appropriately planned and carried out.
- Steps taken to prevent the spread of invasive non-native species potentially present.
- Considerations for the minimisation of damage to the ground during the winter months.

5.3.3 The achievement and success of Biodiversity Net Gain is likely to be contingent on the efforts made in the long term management of the Site's habitats. A Landscape and Ecological Management Plan (LEMP) would set out the agreed habitat creation and planting to be undertaken during and immediately after the construction phase as well as an ecologically-sensitive management schedule for a period of at least 20years. Details on the installation of features of value to wildlife including reptile hibernacula, invertebrate habitats and bird and bat habitat boxes will also be given alongside a monitoring and maintenance schedule. The LEMP is likely to be a requirement of an eventual PEIR/ES in order to demonstrate how proposed mitigation and enhancement will be secured and the various roles and responsibilities for carrying this out.



5.4 Cumulative Impacts

5.4.1 An assessment of cumulative impacts arising from between the sub-sites, between Cottam and West Burton applications and with other large-scale solar in the District will be an essential part of the PEIR/ES. Given the similarities of habitat and value to protected and notable species between the Sites and other applications, the potential for significant cumulative impacts on certain receptors, especially ground nesting birds. This factor will be a key consideration when formulating acceptable mitigation (i.e. its location, quantity and habitat management), not least for ground nesting birds. Preliminary recommendations given in this document, such as buffer widths etc., attempt to take this effect (and the effect of the project scale) into account and apply a precautionary approach.

5.5 Future Baseline and Decommissioning Effects

5.5.1 An assessment of a potential future baseline will be necessary as part of the PEIR/ES in understanding possible effects of decommissioning. Fundamentally, it is impossible to accurately predict the nature of future legal and planning constraints related to ecology in 30-40 years' time. However, on the basis of the current legal and policy situation, it is likely that the biodiversity value of the Sites within the red lines boundaries will overall increase moderately over time and in response to Biodiversity Net Gain-led management principles.

5.5.2 The majority of land where new habitats of value will be created, and colonisation by species of conservation concern most likely to take place, will be at the Sites' boundaries and relatively separated from array infrastructure. This means that future constraints would likely remain similarly distributed to how they are at present. It is considered that the likely DCO requirement (and that of Policy S13 of Central Lincolnshire Local Plan) of an eventual reversion to pre-construction state following decommissioning is compatible with the management of the Sites up to that point as grassland of varying management types.

5.5.3 It is worth noting our experience to date that PINS have been broadly accepting of the view that whilst a robust strategy to protecting valuable ecological features will be required they have also agreed that it is difficult, if not impossible, to prepare or write an ecological strategy to decommissioning now as the conditions and legislative framework at this future point will direct how it would proceed.



APPENDIX A: WILDLIFE LEGISLATION SUMMARY

BADGERS

Badgers and their setts are protected under the Protection of Badgers Act 1992 (as amended) against damage or destruction of a sett, or disturbance, death or injury to the badgers. The Act defines a sett as "any structure or place which displays signs indicating current use by a badger". The definition of current use is subject to considerable debate. Natural England have produced guidance on the definition of current use. (*Badgers and Development – A guide to best practice and development. Natural England 2011*). Given the ambiguity surrounding the definition in all circumstances we would recommend an assessment of current use is always undertaken by a qualified ecologist. Natural Resources Wales (NRW) have a slightly different definition of current use. Please see the NRW website for further information. Penalties for offences against badgers or their setts include fines of up to £5,000 and/or up to six months in prison.

Disturbance of badgers could be caused by any digging activity or scrub clearance within 30 metres of an occupied sett and therefore every case needs to be assessed individually. Felling of trees close to a badger sett may also cause disturbance in some situations. Some activities such as pile driving may cause disturbance at even greater distances, and should be discussed with Natural England or NRW.

Licences are issued by Natural England (or NRW in Wales) to allow the disturbance of badgers, and the destruction of their setts in certain circumstances, in relation to development. Full planning permission must be obtained before a licence application will be considered. Although licences can be applied for at any time of year, disturbance of badgers or exclusion of badgers from a sett can only take place between 1 July and 30 November, to avoid the breeding season when dependant young may be underground. This restriction may be relaxed in some cases where a sett is seasonal and badgers can be shown to be absent from a sett at that time of year.

This report contains information of a confidential nature relating to the location of badger setts. Public access to this data should be restricted to those who have a legitimate need to assess the information and to know the exact situation of the setts rather than simply that badgers are present.

BATS

All 17 species of bat known to breed in England and Wales, and their roost sites, are protected under the Conservation of Habitats and Species Regulations 2017, known as the 'Habitats Regulations'. This makes it an offence to deliberately kill or injure a bat, or to deliberately disturb a bat such that its ability to hibernate, breed or rear young, or such that the species' distribution, were significantly affected. It is also an offence to damage or destroy any breeding site or resting place. Intentional or reckless disturbance of bats in their resting places, and damage to or obstruction of resting places are also offences under the Wildlife and Countryside Act 1981 (as amended). Under UK law a bat roost is "any structure or place which any wild [bat]...uses for shelter or protection". As bats tend to reuse the same roosts, legal opinion is that the roost is protected whether or not the bats are present at the time. Penalties for offences against bats or their roosts include fines of up to £5,000 and/or up to six months in prison.

As a result, development works which are likely to involve the loss of or alteration to roost sites, or which could result in killing of or injury to bats, need to take place under licence. Works which could disturb bats may also be licensable, though this needs to be assessed on a case by case basis, as bats' sensitivity to disturbance varies depending on normal background levels, and the definition of disturbance offences under the Habitats Regulations is complex. In practice this means that works involving modification or loss of roosts (typically in buildings, trees or underground sites) or significant disturbance to bats in roosts are likely to be licensable.

Licences can be obtained from Natural England or the Welsh Government to permit works that would otherwise be illegal, provided it can be demonstrated that the proposed works are needed to protect public health or safety, or for other reasons of overriding public interest including social and economic reasons. It is also necessary to demonstrate that there is no satisfactory alternative to the proposed works, and that the conservation status of bats in the area will be maintained. Appropriate mitigation and post-construction monitoring are therefore a requirement of all licences.

AMPHIBIANS

Great Britain supports seven native amphibian species. The four most widespread species; smooth and palmate newts, common frog, and common toad, receive partial protection under the Wildlife and Countryside Act 1981 (as amended) which prohibits sale, barter, exchange, transporting for sale and advertising to sell or to buy. The great crested newt, pool frog and natterjack toad are also fully protected in England and Wales under the Conservation of Habitats and Species Regulations 2017. Penalties for offences against amphibian species include fines of up to £5,000 and/or up to six months in prison.

Four amphibian species (great crested newt, pool frog, common toad, natterjack toad) are listed as priority species under the UK Biodiversity Action Plan, and are therefore considered to be Species of Principal Importance in England and Wales (excluding the pool frog, which does not occur in Wales) under the Natural Environment and Rural Communities (NERC) Act 2006. All public bodies including local and regional authorities have a duty under this legislation to have regard for the conservation of biodiversity.



GREAT CRESTED NEWTS

Great crested newts are protected in England and Wales under the Conservation of Habitats and Species Regulations 2017, known as the 'Habitats Regulations'. This makes it an offence to deliberately kill or injure a great crested newt, or to deliberately disturb a great crested newt such that its ability to hibernate, breed or rear young, or such that the species' distribution, were significantly affected. It is also an offence to damage or destroy any breeding site or resting place for great crested newts. Intentional or reckless disturbance of great crested newts in places of shelter (ponds or terrestrial refuges), and damage to or obstruction of places of shelter are also offences under the Wildlife and Countryside Act 1981 (as amended). Penalties for offences against great crested newts include fines of up to £5,000 and/or up to six months in prison.

As a result, development works which are likely to involve the loss of ponds or terrestrial habitat, or which could result in killing of or injury to great crested newts, need to take place under licence. Works which could disturb great crested newts may also be licensable, though this is rarely the case unless loss of great crested newt habitat is also proposed, and should be assessed on a case by case basis. In practice this means that works involving any removal of or significant modification to ponds or terrestrial habitats (typically rough grassland, scrub, hedgerow bases and woodland) supporting great crested newts are likely to be licensable.

Licences can be obtained from Natural England or the Welsh Government to permit works that would otherwise be illegal, provided it can be demonstrated that the proposed works are needed to protect public health or safety, or for other reasons of overriding public interest including social and economic reasons. It is also necessary to demonstrate that there is no satisfactory alternative to the proposed works, and that the conservation status of great crested newts in the area will be maintained. Appropriate mitigation and post-construction monitoring are therefore a requirement of all licences.

REPTILES

All six native reptile species receive protection under the Wildlife and Countryside Act 1981 (as amended). The four more common species (common lizard *Zootoca vivipara*, slow-worm *Anguis fragilis*, adder *Vipera berus* and grass snake *Natrix natrix*) receive partial protection which makes it an offence to intentionally kill or injure a reptile. The two other reptile species (smooth snake *Coronella austriaca* and sand lizard *Lacerta agilis*), both of which are rare with very restricted UK ranges receive full protection under the Conservation of Habitats and Species Regulations 2017. Penalties for offences against reptile species include fines of up to £5,000 and/or up to six months in prison.

Works such as site clearance or topsoil stripping which could result in killing or injury of reptiles could be considered result in an offence unless measures are taken to minimise the risk of this occurring. Any inadvertent impacts on common reptile species despite these mitigation measures being in place would be considered an 'incidental result of an otherwise lawful operation' which 'could not reasonably have been avoided' and therefore not an offence. Works which could affect smooth snakes or sand lizards, or their habitats, would need to take place under licence from Natural England or Natural Resources Wales. However sites supporting smooth snakes or sand lizards are very rarely affected by development proposals.

In practice, mitigation for impacts of development on common reptiles generally comprise one or more of the following techniques: displacement, in which reptiles are encouraged to move to suitable retained habitat by changing the management of areas affected by development; exclusion, where reptile-resistant fencing is provided between a development site and suitable retained habitat allowing reptiles to be trapped from the development footprint and released elsewhere on the site; and translocation, where animals are trapped from a development site and released on another suitable site nearby. Reptile mitigation proposals, particularly those involving translocation of animals, should be agreed in advance with the local planning authority.

BIRDS

All British birds, their nests and eggs (with certain exceptions) are protected under the Wildlife & Countryside Act 1981 (as amended) which makes it an offence to: intentionally kill, injure or take a wild bird; intentionally take, damage or destroy nests which are in use or being built; intentionally take or destroy birds' eggs; or possess live or dead wild birds or eggs. A number of species receive additional protection through inclusion on Schedule 1 of the Wildlife and Countryside Act; for these it is also an offence to intentionally or recklessly disturb birds while nest building, or at a nest containing eggs or young, or to disturb the dependant young of such a bird. Penalties for offences against bird species include fines of up to £5,000 and/or up to six months in prison.

General licences for control of some bird species are issued by Natural England and Natural Resources Wales in order to prevent damage or disease, or to preserve public health or public safety, but it is not possible to obtain a licence for control of birds or removal of eggs/nests for development purposes. Consequently if nesting birds are present on a development site when works are programmed to start it is usually necessary to delay works, at least in the areas supporting nests, until any chicks have fledged and left the nest. It is usually possible, once chicks have hatched, for an experienced ecologist to predict approximately when they are likely to fledge, in order to inform programming of works on site.

OTTERS

Otters and their holts are protected in England and Wales under the Conservation of Habitats and Species Regulations 2017, known as the 'Habitats Regulations'. This makes it an offence to deliberately kill or injure an otter, or to deliberately disturb an otter such that its ability to breed or rear young, or such that the species' distribution, were significantly affected. It is also an offence to damage or destroy any breeding site or resting place. Intentional or reckless disturbance of otters in their holts, and damage to or obstruction of



holts are also offences under the Wildlife and Countryside Act 1981 (as amended). Penalties for offences against otters or their holts include fines of up to £5,000 and/or up to six months in prison.

Any development works which are likely to involve the loss of holts, or which could result in killing of or injury to otters (which are only likely to occur extremely rarely), need to take place under licence. Works which could disturb otters may also be licensable, though this is also rarely the case as the majority of developments on watercourses and coastal areas where otters are present can be carried out in a way which avoids significant disturbance.

Where it is necessary, licences can be obtained from Natural England or the Welsh Government to permit works that would otherwise be illegal, provided it can be demonstrated that the proposed works are needed to protect public health or safety, or for other reasons of overriding public interest including social and economic reasons. It is also necessary to demonstrate that there is no satisfactory alternative to the proposed works, and that the conservation status of otters in the area will be maintained. Appropriate mitigation and post-construction monitoring are therefore a requirement of all licences.

WATER VOLES

Water voles *Arvicola amphibius* receive protection under the Wildlife and Countryside Act 1981 (as amended), which makes it an offence to: intentionally kill, injure, or take a water vole; intentionally or recklessly disturb a water vole whilst in its place of shelter; intentionally or recklessly damage, obstruct or destroy a water vole's place of shelter; or intentionally or recklessly obstruct access to a place of shelter. Penalties for offences against water voles include fines of up to £5,000 and/or up to six months in prison.

Works such as watercourse re-profiling, installing culverts, or topsoil stripping close to watercourses and ponds which could result in destruction or obstruction of burrows could be considered reckless, and/or could be considered intentional if water voles are killed or injured, unless measures are taken to minimise the risk of this occurring. Any inadvertent impacts on water voles despite these mitigation measures being in place would be considered an 'incidental result of an otherwise lawful operation' which 'could not reasonably have been avoided' and therefore not an offence.

In practice, mitigation for impacts of development on water voles generally comprise one or more of the following techniques: displacement, in which water voles are encouraged to move to suitable retained habitat by changing the management of areas affected by development; exclusion, where water vole-resistant fencing is provided between a development site and suitable retained habitat allowing animals to be trapped from the development footprint and released elsewhere on the site; and translocation, where animals are trapped from a development site and released on another suitable site nearby. Water vole mitigation proposals, particularly those involving translocation of animals, should be agreed in advance with Natural England or Natural Resources Wales.

PLANNING POLICY IN RELATION TO BIODIVERSITY

The National Planning Policy Framework (NPPF), was published in March 2012 and revised in July 2021. Additional guidance can be found online at <http://planningguidance.planningportal.gov.uk/blog/guidance/>. The NPPF simplifies and collates a number of previous planning documents and outlines the government's objective towards biodiversity.

The NPPF identifies ways in which the planning system should contribute to and enhance the natural and local environment (Paragraph 174), including:

- (a) protecting and enhancing valued landscapes, sites of biodiversity or geological value and soils (in a manner commensurate with their statutory status or identified quality in the development plan);
- (b) recognising the intrinsic character and beauty of the countryside, and the wider benefits from natural capital and ecosystem services – including the economic and other benefits of the best and most versatile agricultural land, and of trees and woodland;
- (d) minimising impacts on and providing net gains for biodiversity, including by establishing coherent ecological networks that are more resilient to current and future pressures;
- (e) preventing new and existing development from contributing to, being put at unacceptable risk from, or being adversely affected by, unacceptable levels of soil, air, water or noise pollution or land instability. Development should, wherever possible, help to improve local environmental conditions such as air and water quality, taking into account relevant information such as river basin management plans; and
- (f) remediating and mitigating despoiled, degraded, derelict, contaminated and unstable land, where appropriate. protecting and enhancing valued landscapes, geological conservation interests and soils;

It also emphasises the importance of conserving biodiversity and areas covered by landscape designations (Paragraph 176):

Great weight should be given to conserving landscape and scenic beauty in National Parks, the Broads and Areas of Outstanding Natural Beauty, which have the highest status of protection in relation to landscape and scenic beauty. The conservation of wildlife and cultural heritage are important considerations in all these areas, and should be given great weight in National Parks and the Broads. The scale and extent of development within all these designated areas should be limited, while development within their setting should be sensitively located and designed to avoid or minimise adverse impacts on the designated areas.

When determining planning applications, the NPPF states that local planning authorities should aim to conserve and enhance biodiversity (Paragraph 175) by applying principles including:

- (a) if significant harm to biodiversity resulting from a development cannot be avoided (through locating on an alternative site with less harmful impacts), adequately mitigated, or, as a last resort, compensated for, then planning permission should be refused;



- (b) development on land within or outside a Site of Special Scientific Interest, and which is likely to have an adverse effect on it (either individually or in combination with other developments), should not normally be permitted. The only exception is where the benefits of the development in the location proposed clearly outweigh both its likely impact on the features of the site that make it of special scientific interest, and any broader impacts on the national network of Sites of Special Scientific Interest;
- (c) development resulting in the loss or deterioration of irreplaceable habitats (such as ancient woodland and ancient or veteran trees) should be refused, unless there are wholly exceptional reasons⁶ and a suitable compensation strategy exists; and
- (d) development whose primary objective is to conserve or enhance biodiversity should be supported; while opportunities to improve biodiversity in and around developments should be integrated as part of their design, especially where this can secure measurable net gains for biodiversity or enhance public access to nature where this is appropriate..

The following should be given the same protection as habitats sites:

- (a) potential Special Protection Areas and possible Special Areas of Conservation;
- (b) listed or proposed Ramsar sites⁷; and
- (c) sites identified, or required, as compensatory measures for adverse effects on habitats sites, potential Special Protection Areas, possible Special Areas of Conservation, and listed or proposed Ramsar sites.

There is a general presumption in favour of sustainable development within the NPPF. It is noted in Paragraph 182 that this presumption does not apply where the plan or project is likely to have a significant effect on a habitat site (either alone or in combination with other plans or projects), unless an appropriate assessment has concluded that the plan or project will not adversely affect the integrity of the habitats site.

The Natural Environment and Rural Communities Act (2006) states that a public authority must, "in exercising its functions, have regard, so far as is consistent with the proper exercise of those functions, to the purpose of conserving biodiversity; Conserving biodiversity includes, in relation to a living organism or type of habitat, restoring or enhancing a population or habitat". DEFRA issued further guidance on implementation of this act in the document; Guidance for Local Authorities on Implementing the Biodiversity Duty (May 2007), which notes that "Conserving biodiversity includes restoring and enhancing species populations and habitats, as well as protecting them".

ECOLOGICAL ENHANCEMENTS

The Natural Environment and Rural Communities Act (2006) states that a public authority must, "in exercising its functions, have regard, so far as is consistent with the proper exercise of those functions, to the purpose of conserving biodiversity; Conserving biodiversity includes, in relation to a living organism or type of habitat, restoring or enhancing a population or habitat". DEFRA issued further guidance on implementation of this act in the document; Guidance for Local Authorities on Implementing the Biodiversity Duty (May 2007), which notes that "Conserving biodiversity can include restoring or enhancing a population or habitat".

In England, the National Planning Policy Framework (NPPF), issued in July 2021, states that the planning system should contribute to "minimising impacts on and providing net gains for biodiversity, including by establishing coherent ecological networks that are more resilient to current and future pressures;. It also states that "opportunities to incorporate biodiversity in and around developments should be encouraged, especially where this can secure measurable net gains for biodiversity".

UK BIODIVERSITY ACTION PLANS

The UK Biodiversity Action Plan (UK BAP) 2011 is a policy first published in 1994 to protect biodiversity and stems from the 1992 Rio Biodiversity Earth Summit. The policy is continuously revised to combine new and existing conservation initiatives to conserve and enhance species and habitats, promote public awareness and contribute to international conservation efforts. Each plan details the status, threats and unique conservation strategies for the species or habitat concerned, to encourage spread and promote population numbers.

Species or habitats identified as priorities under the UK Biodiversity Action Plan receive some status in the planning process through their identification as Species/Habitats of Principal Importance in England and Wales, under the Natural Environment and Rural Communities (NERC) Act 2006 (as amended).

Current planning guidance in England, the National Planning Policy Framework, does not specifically refer to Species or Habitats of Principal Importance, though it includes guidance for conservation of biodiversity in general. Supplementary guidance is available online at <http://planningguidance.planningportal.gov.uk/blog/guidance/> and this guidance indicates that it is 'useful to consider' the potential effects of a development on the habitats or species on the Natural Environment and Rural Communities Act 2006 section 41 list.

PROTECTED PLANTS

All wild plants receive some protection under the Wildlife and Countryside Act 1981 (as amended), which makes it an offence for any unauthorised person to intentionally uproot any wild plant. Additionally, certain rare species of plants listed on Schedule 8 of the Act are given greater protection. For these species, it is an offence to intentionally pick, uproot or destroy them, or to possess or sell them (live or dead), or anything derived from them. Penalties for offences under this legislation include fines of up to £5,000 and/or up to six months in prison.



Schedule 8 of the Act is reviewed every 5 years, but currently it includes 185 species or sub-species of vascular plants, bryophytes (mosses, liverworts and hornworts), lichens and stoneworts (see www.jncc.gov.uk for current list), all protected due to their rarity and/or restricted distributions.

Works which could result in uprooting or destruction of plants listed on Schedule 8 of the Act could result in an offence being committed, unless measures are taken to minimise the risk of this occurring. Any inadvertent impacts on Schedule 8 plants despite these mitigation measures being in place, and impacts on other plant species during development works, would be considered an 'incidental result of an otherwise lawful operation' which 'could not reasonably have been avoided' and therefore not an offence.

In practice, the mitigation measures required on the very rare occasions when Schedule 8 plants are affected by development proposals will be determined by the ecological requirements of the species concerned, and any mitigation strategy should be agreed in advance with Natural England or Natural Resources Wales.

THE HEDGEROWS REGULATIONS

In England and Wales the Hedgerows Regulations (1997) as amended confer a level of protection on hedgerows (though hedgerows within or bordering domestic gardens are excluded), particularly those hedgerows classified as 'Important' under the legislation. The Regulations require those wishing to remove hedgerows to submit a Hedgerow Removal Notice to the Local Planning Authority (LPA), which will then determine whether the hedgerow affected is classified as 'Important' under the Regulations. If it is, the LPA will either approve the proposed hedgerow removal, or issue a retention notice. It is an offence to remove or destroy a hedgerow which is subject to a retention notice, or to remove one without a removal notice.

Routine management of hedgerows, removal of hedgerows for development which has been granted planning consent, and certain other situations are allowed under the Regulations, which also specifically exclude hedgerows within or bordering domestic gardens. Determination of whether a hedgerow should be classified as 'Important' is based on a number of criteria including assessment of its likely historic value (e.g. old parish boundary or part of an ancient monument), ecological value (e.g. presence of protected species, and/or diversity of tree/shrub species in the hedgerow), and landscape value (e.g. associated with a public footpath, or being associated with hedgebanks, ditches, hedgerow trees etc).

Ancient and species-rich hedgerows are listed as a priority habitat in the UK Biodiversity Action Plan (2011)

JAPANESE KNOTWEED

Japanese knotweed *Fallopia japonica* is a non-native invasive species listed on Schedule 9 of the Wildlife and Countryside Act 1981 (as amended). This Act states that it is an offence to plant or otherwise cause this species to grow in the wild. Penalties for offences under this legislation include fines of up to £25,000 and/or up to six months in prison.

In addition to this legislation, all parts of the plant and soil contaminated with plant fragments, is classified as contaminated waste under the Environmental Protection Act 1990, and will require a special waste licence and/or waste transfer note under the Environmental Protection (Duty of Care) Regulations 1991 (as amended).

The Environment Agency has produced a 'Code of Practice for the Management, Destruction and Disposal of Japanese Knotweed' (2001), which provides guidance for developers.

HIMALAYAN BALSAM

Himalayan balsam *Impatiens glandulifera* is a non-native invasive species listed on Schedule 9 of the Wildlife and Countryside Act 1981 (as amended). This Act states that it is an offence to plant or otherwise cause this species to grow in the wild. Penalties for offences under this legislation include fines of up to £25,000 and/or up to six months in prison.

Advice on management and control of Himalayan balsam is provided in the Environment Agency's leaflet 'Managing Invasive Non-native Plants' (2010).



APPENDIX B – SUMMARY OF METHODOLOGIES

Desk Study Methodology

Statutory designated sites for nature conservation were identified using the Natural England/DEFRA web-based MAGIC map database (www.MAGIC.gov.uk). International-level sites such as Special Areas of Conservation (SACs) and Special Protection Areas (SPAs) within 10km from the Site were searched for. National-level sites such as National Nature Reserves (NNRs) and Sites of Special Scientific Interest (SSSIs) within 5km of the Site were searched for.

The Lincolnshire Environmental/ Biological Records Centre (LERC) was consulted for records of protected species and species of conservation concern within 2km of the Site as well as details of locally-designated and non-statutory sites for nature conservation within 2km of the Site.

Ordnance Survey maps (1:25,000) and aerial images of the Site were examined online [REDACTED] and [REDACTED] to allow a better understanding of the context of the Site and its connections to potentially important habitats, known species records and protected sites.

The data presented within this report constitutes a summary of the data obtained from the local records centre. Should additional detail be required on any of the records described within this report Clarkson and Woods Ltd. should be contacted.

Species of Conservation Concern are defined as those appearing in any of the following; Priority Habitats and Species under Section 41 of the Natural Environment and Rural Communities Act (2006); red or amber-listed birds within the British Trust for Ornithology's Birds of Conservation Concern (2015); and any specific local conservation priority species such as those listed in Red Data Books.

Habitat Survey Methodology

A habitat survey was carried out based on standard field methodology set out in the *Handbook for Phase 1 Habitat Survey* (2010 edition)⁵. The survey was co-ordinated and led by Harry Fox BSc MCIEEM, Principal Ecologist. Harry has 13 years' experience undertaking ecological surveys and has a BSc in ecology. Harry was assisted by the following personnel in completing the Phase 1 surveys:

- Peter Timms BSc MSc MCIEEM – Senior Ecologist
- Henry Sturgess BSc MCIEEM – Senior Ecologist
- Belinda Howell BSc MCIEEM – Senior Ecologist
- Joel Wright BSc MSc MCIEEM – Senior Ecologist
- Mike Hockey BSc ACIEEM – Senior Ecologist
- Charlie Durigan BSc MSc PgCert ACIEEM - Ecologist

Botanical names follow Stace (1997)⁶ for higher plants and Edwards (1999)⁷ for bryophytes.

Badgers

A search was made for badger *Meles meles* setts, and any sett entrances found were checked for signs of use by badgers or other mammals. Setts were classified into the following categories; Main, Subsidiary, Annexe or Outlying⁸. Sett entrances found were counted and mapped to record tunnel direction and their relative level of usage.

Field signs such as 'snuffle holes' (holes dug by badgers when searching for invertebrates), pathways through vegetation, 'latrines' (small pits in which badgers deposit their faeces) and 'day nests' (nests of bedding material made by badgers for sleeping above ground) were also mapped, if found.

Areas with dense ground cover (hedges, scrub, woodland etc. were examined closely. If impenetrable vegetation prevented entry then the perimeter was examined in order to detect badger paths suggesting a hidden sett within the area. It cannot be guaranteed that all the entrances have been located, especially if a small sett is currently inactive or used seasonally and concealed in an area of thick scrub. Badgers may dig new holes and create new setts in a very short space of time.

Bats

The assessment of the suitability of the site for foraging and roosting bats was based on current guidance set out by the Bat Conservation Trust⁹.

The habitats within the sites were appraised for their suitability for use by foraging and commuting bats. In particular, the connectivity of the habitats on site to those lying beyond was taken into account. Vegetated linear features are typically important for many species to navigate around the landscape, while the presence of woodland, scrub, gardens, grassland and wetland features increases a site's foraging resource value to bats. The potential for noise or lighting disturbance which may affect commuting links was also recorded.

⁵ Nature Conservancy Council. (1990 - 2010 edition). *Handbook for Phase 1 Habitat Survey – A Technique for Environmental Audit*, Joint Nature Conservation Committee

⁶ Stace, C. (1997). *New Flora of the British Isles Second Edition*. Cambridge University Press

⁷ Edwards, S.R. (1999). *English Names for British Bryophytes*. BBS, Cardiff

⁸ Lewns, P., Clarkson, T. & Lewns, D. (2019). *Badger Survey and Mitigation Guidelines (The Mammal Society Mitigation Guidance Series)*. Eds. Fiona Mathews and Paul Chanin. The Mammal Society, London. (as yet unpublished)

⁹ Collins, J. (ed) (2016) *Bat Surveys for Professional Ecologists: Good Practice Guidelines* (3rd edn). The Bat Conservation Trust, London. ISBN-13 978-1-872745-96-1.



It was considered impractical to conduct walked evening transect of all option land given their extent. In accordance with best practice guidance, it was elected that baseline data would be most effectively collected through the use of static bat detectors. An elevated number of detectors and deployments compared to that recommended within The Bat Conservation Trusts' Good Practice Guidelines was used in lieu of walked transect surveys. The guidelines also recommend that, "if the habitat has been classified as having low suitability for bats, an ecologist should make a professional judgment on how to proceed based on all of the evidence available. It may or may not be appropriate for bat activity survey to be carried out in low suitability habitats." It was therefore considered that 42 static bat detector locations spread across all option land, installed at field boundaries and surveyed once per month between June and September inclusive, would enable the proportionate collection of an adequate baseline. It was considered impractical to install detectors within the centres of fields on account of ongoing agricultural activities such as crop spraying and harvesting. In any case, these arable habitats are of comparatively the lowest value to bats within the option sites and the field edges were considered the most conducive to bat activity.

Otter

A brief search was made along the banks of water courses and water bodies and their adjacent habitats for otter *Lutra lutra* signs including spraints, tracks, castling, and rolling. The banks of any water courses were searched for the presence or potential for holts or other sheltering areas.

Water Vole

The banks of the water course were searched for water vole *Arvicola amphibius* signs including latrines, burrow entrances, feeding stations, 'runways' and footprints. Surveys and field recording followed the protocol set out within the Water Vole Mitigation Handbook¹⁰

GCN and Toads

All waterbodies within 250m / 500m of the Sites were identified using Ordnance Survey maps and aerial imagery. Waterbodies within the site ownership were assessed during the field survey for their suitability to support amphibian species where access was possible.

Where suitable water bodies were identified on accessible land a Habitat Suitability Index (HSI) score was calculated for each one following the methodology described by Oldham et al¹¹. HSI scores give a relative indication of the likelihood that a water body would support breeding great crested newts. Factors which increase these scores include the presence of other ponds nearby, water quality, pond size, absence of fish/waterfowl, vegetation cover and shading.

Terrestrial habitats were also assessed for their suitability for foraging and sheltering great crested newts. This species requires habitats such as grassland, scrub, woodland and hedgerows for dispersal and hibernation. Further hibernation features include buried rubble and logs, or mammal burrows.

Where eDNA surveys were taken, a standard methodology was followed according to Natural England best practice guidance and ADAS' laboratory requirements, carried out between the period of 15th April and 30th June.

Reptiles

Features on the Sites were assessed for their potential to provide suitable habitats for use by reptile species. These include rough, tussocky grassland, scrub, disturbed land or refugia such as wood piles, rubble or compost heaps. Where present, suitable existing refugia were inspected for sheltering reptiles, and the ground was scanned whilst walking to look for basking species.

Birds

Any buildings and vegetation were surveyed for signs of use by nesting birds and any birds seen or heard during the survey were noted. The site's potential to support bird species of particular conservation concern (i.e. Schedule 1, NERC S41 and Red List species) was assessed, taking into consideration the bird species assemblage observed during the survey, the habitats present on and around the site, the context of the site in the wider landscape and the results of the desk study.


¹⁰ Dean, M., Strachan, R., Gow, D. and Andrews, R. (2016). *The Water Vole Mitigation Handbook (The Mammal Society Mitigation Guidance Series)*. Eds. Fiona Mathews and Paul Chanin. The Mammal Society, London.

¹¹ Oldham, R.S., Keeble L., Swan M.J.S. & Jeffcote M. (2000). Evaluating the suitability of habitat for the Great Crested Newt (*Triturus cristatus*). *Herpetological Journal* 10 (4), 143-155.



APPENDIX C – DESIGNATED SITES MAPS

West Burton 1



Key:

- Sites of Special Scientific Interest (SSSIs)
- Local Nature Reserves (LNRs)
- Local Wildlife Sites (LWSs)
- 5km Buffer
- 2km Buffer
- Red Line Boundary
- 1 Site Reference

Un-numbered sites are those which occur outside of the scope of the relevant search radius for their level of designation, i.e. 2Km for non-statutory sites and 5Km for statutory sites.



Project
West Burton 1

Title
Designated Sites


Project Number
7479

Scale
See Scalebar








Date
12/08/2021



West Burton 2



Key:

-  Sites of Special Scientific Interest (SSSIs)
-  Local Nature Reserves (LNRs)
-  Local Wildlife Sites (LWSs)
-  5km Buffer
-  2km Buffer
-  Red Line Boundary
-  Site Reference

Un-numbered sites are those which occur outside of the scope of the relevant search radius for their level of designation, i.e. 2km for local sites and 5km for national sites



Project
West Burton 2

Title
Designated Sites


Project Number
7479

Scale
See Scalebar

Date
12/08/2021



West Burton 3



Key:

- Sites of Special Scientific Interest (SSSIs)
- Local Nature Reserves (LNRs)
- Local Wildlife Sites (LWSs)
- 5km Buffer
- 2km Buffer
- Red Line Boundary
- 1 Site Reference

Un-numbered sites are those which occur outside of the scope of the relevant search radius for their level of designation, i.e. 2Km for non-statutory sites and 5Km for statutory sites.

CLARKSON & WOODS
ECOLOGICAL CONSULTANTS

Project West Burton 3	
Title Designated Sites	
Project Number 7479	
Scale See Scalebar	Date 12/08/2021



APPENDIX D – SPECIES RECORDS WITHIN 2KM OF WEST BURTON 1 (BROXHOLME)

Records of Protected and Notable Species Derived from the Desk Study Data Search (LERC)

Group	Scientific Name	Common Name	Records	Location	Date
Amphibians	<i>Bufo bufo</i>	Common Toad	7 records within 2km	The only known record location is 900m west of the site. Exact location unknown for all other records – within 2km of the site.	6 records pre 2000 (1977) 1 record post 2000 (2016)
Amphibians	<i>Rana temporaria</i>	Common Frog	17 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~1.4km north-west of the site, with 3 individuals recorded in 2008 (Grid Reference SK902802 – Sturton by Stow)	11 records pre 2000 6 records post 2000
Amphibians	<i>Triturus cristatus</i>	Great Crested Newt	65 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~1.5km north-west of the site (Grid Reference SK902803) with up to 59 individuals recorded between 2008 and 2017.	6 records pre 2000 59 records post 2000
Amphibians	<i>Lissotriton vulgaris</i>	Smooth Newt	17 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~1.5km north-west of the site (Grid Reference SK902802) with 8 individuals recorded in 2008.	7 records pre 2000 10 records post 2000
Reptiles	<i>Zootoca vivipara</i>	Common Lizard	3 records within 2km	Exact location unknown – within 2km of the site.	3 records pre 2000 (1977) 0 records post 2000
Reptiles	<i>Natrix Helvetica</i>	Grass Snake	18 records within 2km	All records are located beyond 250m of the site. Closest of which is located 680m north-east of the site (Grid Reference SK924794) in 2008.	12 records pre 2000 6 records post 2000
Terrestrial Mammal	<i>Arvicola amphibius</i>	European Water Vole	30 records within 2km	All records are located beyond 250m of the site. Closest of which is located 460m north-east of the site (Grid reference SK923792) in 2011.	9 records pre 2000 21 records post 2000
Terrestrial Mammal	<i>Lepus europaeus</i>	Brown Hare	29 records within 2km	All records are located beyond 250m of the site. Closest of which is located 335m north of the site (Grid Reference SK912795) in 2010.	13 records pre 2000 16 Records post 2000
Terrestrial Mammal	<i>Meles meles</i>	Eurasian Badger	16 records within 2km	All records are located beyond 250m of the site. Closest of which is located 260m north of the site (Grid Reference SK915794) in 2018.	0 records pre 2000 16 records post 2000
Terrestrial Mammal	<i>Lutra lutra</i>	European Otter	14 records within 2km	All records are located beyond 250m of the site. 13 records are located along the River Till corridor. The closest are located ~760m north-west of the site (Grid Reference SK907797), recorded in 1999 and 2009.	8 records pre 2000 6 records post 2000
Terrestrial Mammal	<i>Mustela putorius subsp. furo</i>	Feral Ferret	2 records within 2km	All records are located beyond 250m of the site. Both records are located 1.3km north-west of the site (Grid Reference SK901799) in 2013.	0 records pre 2000 2 records post 2000 (2013)
Terrestrial Mammal	<i>Erinaceus europaeus</i>	West European Hedgehog	58 records within 2km	One record dated 2014 is located within the red line boundary (Grid Reference SK917785). Three records are located within 250m of the site. 54 records are located beyond 250m of the site.	13 records pre 2000 45 records post 2000
Terrestrial Mammal	<i>Micromys minutus</i>	Harvest Mouse	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1977) 0 records post 2000
Bats	<i>Plecotus auritus</i>	Brown Long-eared Bat	1 record within 2km	Record is located ~1.9km west of the site (Grid Reference SK8979) and is dated 2010.	0 records pre 2000 1 record post 2000 (2010)
Bats	<i>Pipistrellus pipistrellus sensu stricto</i>	Common Pipistrelle	22 records within 2km	All records are located beyond 250m of the site. Closest of which are located ~1.3km north-west of the site (Grid Reference SK898794) and are dated 2010 and 2012.	0 records pre 2000 22 records post 2000
Bats	<i>Nyctalus noctula</i>	Noctule Bat	1 record within 2km	Record is located ~1.9km east of the site (Grid Reference SK9479) and is dated 1987.	1 record pre 2000 (1987) 0 records post 2000
Bats	<i>Pipistrellus</i>	Pipistrelle Bat species	11 records within 2km	Four records are located within 250m of the site to the south-west, recorded in 2008. Seven records are located beyond 250m of the site.	0 records pre 2000 11 records post 2000
Bats	<i>Pipistrellus pygmaeus</i>	Soprano Pipistrelle	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2013)
Bats	Unidentified Bat	Unidentified Bat	19 records within 2km	Two records are located within 250m of the site to the south-west, recorded in 2008 and 2015. 17 records are located beyond 250m of the site.	7 records pre 2000 12 records post 2000



Birds	Tyto alba	Barn Owl	99 records within 2km	All records are located beyond 250m of the site. Closest of which are located ~860m south-west of the site (Grid Reference SK904777), dated 2008 and 2009.	2 records pre 2000 (1998) 97 records post 2000
Birds	Phoenicurus ochruros	Black Redstart	2 records within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1998) 1 record post 2000 (2016)
Birds	Fringilla montifringilla	Brambling	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2013)
Birds	Pyrrhula pyrrhula	Bullfinch	6 records within 2km	All records are located beyond 250m of the site. Closest known location record is ~1.6km south of the site (Grid Reference SK911763) dated 1970.	3 records pre 2000 3 records post 2000
Birds	Emberiza calandra	Corn Bunting	10 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 10 records post 2000
Birds	Cuculus canorus	Cuckoo	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1976) 0 records post 2000
Birds	Numenius arquata	Curlew	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	Anser albifrons subsp. albifrons	European Greater White-fronted Goose	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2015)
Birds	Turdus pilaris	Fieldfare	10 records within 2km	The only known record location is 385m south-west of the site. Exact location unknown for all other records – within 2km of the site.	0 records pre 2000 10 records post 2000
Birds	Tringa ochropus	Green Sandpiper	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	Tringa nebularia	Greenshank	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000 (2002)
Birds	Perdix perdix	Grey Partridge	25 records within 2km	Two records are located 1.65km south of the site (Grid Reference SK911763) in 1970. Exact location unknown for all other records – within 2km of the site.	4 records pre 2000 21 records post 2000
Birds	Anser anser	Greylag Goose	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2017)
Birds	Circus cyaneus	Hen Harrier	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2015)
Birds	Falco subbuteo	Hobby	7 records within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1998) 6 records post 2000
Birds	Pernis apivorus	Honey-buzzard	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2009)
Birds	Passer domesticus	House Sparrow	11 records within 2km	All records are located beyond 250m of the site. The two closest known location records are 385m to the south-west, dated 2008 and 2009.	2 records pre 2000 9 records post 2000
Birds	Alcedo atthis	Kingfisher	9 records within 2km	One record is located within 250m of the site to the north (Grid Reference SK911790), associated with Scampton CP ditch in 2009. Eight records are located beyond 250m of the site.	1 record pre 2000 (1976) 8 records post 2000
Birds	Vanellus vanellus	Lapwing	18 records within 2km	The closest known record location is 895m south-west of the site (Grid Reference SK904776), dated 2016. Exact location unknown for all other records – within 2km of the site.	4 records pre 2000 14 records post 2000
Birds	Linaria cannabina	Linnet	7 records within 2km	The only known record location is ~1.6km south of the site, dated 1970. Exact location unknown for all other records – within 2km of the site.	2 records pre 2000 5 records post 2000
Birds	Circus aeruginosus	Marsh Harrier	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2007)
Birds	Falco columbarius	Merlin	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2015)
Birds	Pandion haliaetus	Osprey	3 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 3 records post 2000
Birds	Falco peregrinus	Peregrine	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	Coturnix coturnix	Quail	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1999) 0 records post 2000
Birds	Milvus milvus	Red Kite	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000



					2 records post 2000 (2017)
Birds	<i>Turdus iliacus</i>	Redwing	5 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 5 records post 2000
Birds	<i>Emberiza schoeniclus</i>	Reed Bunting	16 records within 2km	All records are located beyond 250m of the site. Closest known location record is 830m west of the site, dated 2016.	3 records pre 2000 13 records post 2000
Birds	<i>Philomachus pugnax</i>	Ruff	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2002)
Birds	<i>Alauda arvensis</i>	Skylark	9 records within 2km	One record is located within 250m of the site, dated 2016. The only other known record location is ~1.6km south of the site, dated 1970. Exact locations unknown for all other records – within 2km of the site.	3 records pre 2000 6 records post 2000
Birds	<i>Gallinago gallinago</i>	Snipe	4 records within 2km	Exact location unknown – within 2km of the site.	3 records pre 2000 1 record post 2000 (2004)
Birds	<i>Turdus philomelos</i>	Song Thrush	15 records within 2km	The only known record locations are ~380m south-west of the site in 2009 and 1.7km south-west in 1970. Exact locations unknown for all other records – within 2km of the site.	3 records pre 2000 12 records post 2000
Birds	<i>Muscicapa striata</i>	Spotted Flycatcher	2 records within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1970) 1 record post 2000 (2009)
Birds	<i>Sturnus vulgaris</i>	Starling	18 records within 2km	All records are located beyond 250m of the site. The two closest known location records are 385m south-west of the site, dated 2008 and 2009.	3 records pre 2000 15 records post 2000
Birds	<i>Apus apus</i>	Swift	6 records within 2km	The only known location records are 660m north, dated 2011 and 1.6km south of the site in 1970. Exact locations unknown for all other records – within 2km of the site.	2 records pre 2000 4 records post 2000
Birds	<i>Passer montanus</i>	Tree Sparrow	22 records within 2km	The only known location record is ~1.7km south of the site, dated 1970. Exact locations unknown for all other records – within 2km of the site.	2 records pre 2000 20 records post 2000
Birds	<i>Streptopelia turtur</i>	Turtle Dove	8 records within 2km	The only known location record is ~1.7km south of the site, dated 1970. Exact locations unknown for all other records – within 2km of the site.	3 records pre 2000 5 records post 2000
Birds	<i>Cygnus cygnus</i>	Whooper Swan	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000 (2010)
Birds	<i>Motacilla flava</i>	Yellow Wagtail	11 records within 2km	The only known location record is ~1.7km south of the site, dated 1970. Exact locations unknown for all other records – within 2km of the site.	2 records pre 2000 9 records post 2000
Birds	<i>Emberiza citrinella</i>	Yellowhammer	12 records within 2km	The only known location records are 1.4km north-west in 2018 and ~1.7km south of the site, dated 1970. Exact locations unknown for all other records – within 2km of the site.	3 records pre 2000 9 records post 2000
bony fish (Actinopterygii)	<i>Anguilla anguilla</i>	European Eel	21 records within 2km	All records are located beyond 250m of the site. Closest of which is located 475m north of the site, with 4 individuals recorded in 1995 (Grid Reference SK917795). The remaining 17 records are associated with the River Till.	19 records pre 2000 2 records post 2000
bony fish (Actinopterygii)	<i>Cobitis taenia</i>	Spined Loach	24 records within 2km	All records are located beyond 250m of the site. Closest of which is located 695m north-west of the site, with 4 individuals recorded in 1994 (Grid Reference SK906795). All records are associated with the River Till.	13 records pre 2000 11 records post 2000
insect - butterfly	<i>Coenonympha pamphilus</i>	Small Heath	2 records within 2km	Exact location unknown – within 2km of the site.	2 records pre 2000 0 records post 2000
insect - butterfly	<i>Lasiommata megera</i>	Wall	3 records within 2km	Exact location unknown – within 2km of the site.	3 records pre 2000 0 records post 2000
Mollusc	<i>Valvata marcostoma</i>	Large-mouthed Valve Snail	1 record within 2km	The record is located 1.4km south-west of the site (Grid Reference SK904768), dated 1980.	1 record pre 2000 (1980) 0 records post 2000



APPENDIX E – SPECIES RECORDS WITHIN 2KM OF WEST BURTON 2 (INGLEBY)

Records of Protected and Notable Species Derived from the Desk Study Data Search (LERC)

Group	Scientific Name	Common Name	Records	Location	Date
Amphibians	<i>Bufo bufo</i>	Common Toad	11 record within 2km	The only known record locations are 460m north of the site in 2016, 645m south in 2009 and 1.2km south-east of the site in 2011. Exact locations unknown for all other records – within 2km of the site.	8 records pre 2000 3 records post 2000
Amphibians	<i>Rana temporaria</i>	Common Frog	28 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1km south of the site (Grid Reference SK914753), dated 2011.	20 records pre 2000 8 records post 2000
Amphibians	<i>Triturus cristatus</i>	Great Crested Newt	23 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1.9km south-west of the site (Grid Reference SK873747), dated 2014. 13 records are located ~1.9km north of the site (Grid Reference SK901802) dated between 2008 and 2009.	9 records pre 2000 14 records post 2000
Amphibians	<i>Lissotriton vulgaris</i>	Smooth Newt	22 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~640m south of the site (Grid Reference SK893760), dated 2009.	11 record pre 2000 11 record post 2000
Reptiles	<i>Zootoca vivipara</i>	Common Lizard	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 (1977)
Reptiles	<i>Natrix helvetica</i>	Grass Snake	29 records within 2km	Two records dated 2011 are located within the red line boundary (Grid Reference SK905780). Three records are located within 250m of the site and are associated with the River Till corridor. 24 records are located beyond 250m of the site.	20 records pre 2000 9 records post 2000
Terrestrial Mammal	<i>Arvicola amphibius</i>	European Water Vole	92 records within 2km	Ten records are located within 250m of the red line boundary, dated between 1990 and 2012. 58 records are located beyond 250m of the site. Exact location unknown for the remaining 24 records – within 2km of the site.	18 records pre 2000 74 records post 2000
Terrestrial Mammal	<i>Lepus europaeus</i>	Brown Hare	53 records within 2km	One record is located within 250m of the red line boundary (Grid Reference SK892785) in 2000 (road kill). All other records are located beyond 250m of the site.	23 records pre 2000 30 records post 2000
Terrestrial Mammal	<i>Meles meles</i>	Eurasian Badger	57 records within 2km	5 records are located within 250m of the site, dated 2004 and 2011. 34 records are records of badger setts, dated between 1996 and 2016. Exact Locations for these are unknown – within 2km of the site. 18 records are located beyond 250m of the site.	2 records pre 2000 55 records post 2000
Terrestrial Mammal	<i>Lutra lutra</i>	European Otter	22 records within 2km	Four records are located within 250m of the red line boundary, dated between 1994 and 2009. All are associated with the River Till corridor. 18 records are located beyond 250m of the site.	11 record pre 2000 11 record post 2000
Terrestrial Mammal	<i>Mustela putorius subsp. furo</i>	Feral Ferret	2 records within 2km	Both records are located 1.6km north of the site (Grid Reference SK901799), dated 2013 (road kill).	0 records pre 2000 2 records post 2000
Terrestrial Mammal	<i>Erinaceus europaeus</i>	West European Hedgehog	131 record within 2km	Four records, dated 2015, are located within the red line boundary. (Grid References SK891781 and SK891773; all records are road kill). 8 records are located within 250m of the site, dated between 2015 and 2018. 119 records are located beyond 250m of the site.	23 records pre 2000 108 records post 2000
Terrestrial Mammal	<i>Micromys minutus</i>	Harvest Mouse	3 records within 2km	Exact location unknown – within 2km of the site.	3 records pre 2000 (1977) 0 records post 2000
Bats	<i>Plecotus auritus</i>	Brown Long-eared Bat	4 records within 2km	All records are located beyond 250m of the site. Closest of which is ~515m north-west of the site (Grid Reference SK873785), dated 2012.	0 records pre 2000 4 records post 2000



Bats	<i>Pipistrellus pipistrellus sensu stricto</i>	Common Pipistrelle	36 records within 2km	All records are located beyond 250m of the site. Closest known location records are 1.1km north of the site (Grid Reference SK882796), dated 2004 and 2006.	0 records pre 2000 35 records post 2000
Bats	<i>Myotis daubentonii</i>	Daubenton's Bat	7 record within 2km	All records are located beyond 250m of the site. Closest of which are 4 records located 1.1km north of the site (Grid Reference SK882796), dated between 2004 and 2006.	0 records pre 2000 7 record post 2000
Bats	<i>Myotis nattereri</i>	Natterer's Bat	1 record within 2km	Record is located 1.4km south of the site (Grid Reference SK895748), dated 2005.	0 records pre 2000 1 record post 2000 (2005)
Bats	<i>Nyctalus noctula</i>	Noctule Bat	2 records within 2km	Both records are located beyond 250m of the site. Closest of which is located 1km south of the site (Grid Reference SK895752), dated 2010.	0 records pre 2000 2 record post 2000
Bats	<i>Pipistrellus</i>	Pipistrelle Bat species	24 records within 2km	Two records dated 2007 are located within 250m of the site (Grid Reference SK892779). 22 records are located beyond 250m of the site.	1 record pre 2000 23 records post 2000
Bats	<i>Pipistrellus pygmaeus</i>	Soprano Pipistrelle	2 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 record post 2000
Bats	Unidentified Bat	Unidentified Bat	83 records within 2km	Five records are located within 250m of the site, dated between 1999 and 2015. 78 records are located beyond 250m of the site.	20 records pre 2000 63 records post 2000
Birds	<i>Tyto alba</i>	Barn Owl	103 records within 2km	Two records are located within 250m of the site (Grid Reference SK904777), dated 2008 and 2009. 101 records are located beyond 250m of the site.	2 records pre 2000 101 record post 2000
Birds	<i>Phoenicurus ochruros</i>	Black Redstart	1 record within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 1 record post 2000
Birds	<i>Motacilla flava flava</i>	Blue-headed Wagtail	2 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 record post 2000
Birds	<i>Fringilla montifringilla</i>	Brambling	4 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 record post 2000
Birds	<i>Pyrrhula pyrrhula</i>	Bullfinch	12 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. The only other known location record is ~1km south of the site, dated 2009. Exact location unknown for all other records – within 2km of the site.	5 records pre 2000 7 records post 2000
Birds	<i>Emberiza calandra</i>	Corn Bunting	3 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 3 records post 2000
Birds	<i>Crex crex</i>	Corncrake	2 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 record post 2000
Birds	<i>Cuculus canorus</i>	Cuckoo	7 records within 2km	The only known location record is 700m south of the site (Grid Reference SK890758), dated 2006). Exact location unknown for all other records – within 2km of the site.	2 record pre 2000 5 records post 2000
Birds	<i>Numenius arquata</i>	Curlew	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	<i>Turdus pilaris</i>	Fieldfare	19 record within 2km	The only known location record is 620m east of the site (Grid Reference SK911776), dated 2009. Exact location unknown for all other records – within 2km of the site.	0 records pre 2000 19 record post 2000
Birds	<i>Tringa ochropus</i>	Green Sandpiper	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	<i>Tringa nebularia</i>	Greenshank	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	<i>Perdix perdix</i>	Grey Partridge	34 record within 2km	Two records dated 1970 are located within 250m of the site (Grid Reference SK911763). Exact location unknown for all other records – within 2km of the site.	7 records pre 2000 27 record post 2000
Birds	<i>Anser anser</i>	Greylag Goose	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000
Birds	<i>Circus cyaneus</i>	Hen Harrier	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000
Birds	<i>Falco subbuteo</i>	Hobby	3 record within 2km	Exact location unknown – within 2km of the site.	2 record pre 2000 1 record post 2000



Birds	<i>Pernis apivorus</i>	Honey-buzzard	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	<i>Passer domesticus</i>	House sparrow	94 records within 2km	One record dated 1970 is located within 250m of the site (Grid Reference SK911763). All other records are located beyond 250m of the site.	4 records pre 2000 90 records post 2000
Birds	<i>Alcedo atthis</i>	Kingfisher	22 records within 2km	Two records are located within 250m of the site (Grid Reference SK904777), dated 2008 and 2009. 20 records are located beyond 250m of the site.	2 records pre 2000 20 records post 2000
Birds	<i>Vanellus vanellus</i>	Lapwing	26 records within 2km	Three records are located within 250m of the site, dated between 1970 and 2016. All other records are located beyond 250m of the site.	7 records pre 2000 19 records post 2000
Birds	<i>Acanthis cabaret</i>	Lesser Redpoll	4 records within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 3 records post 2000
Birds	<i>Linaria cannabina</i>	Linnet	13 records within 2km	The only known location record is within 250m of the site (Grid Reference SK911763), dated 1970. Exact location unknown for all other records – within 2km of the site.	3 records pre 2000 10 records post 2000
Birds	<i>Circus aeruginosus</i>	Marsh Harrier	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000
Birds	<i>Falco columbarius</i>	Merlin	3 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 3 records post 2000
Birds	<i>Pandion haliaetus</i>	Osprey	6 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 6 records post 2000
Birds	<i>Falco peregrinus</i>	Peregrine	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	<i>Coturnix coturnix</i>	Quail	2 records within 2km	Exact location unknown – within 2km of the site.	2 records pre 2000 0 records post 2000
Birds	<i>Milvus milvus</i>	Red Kite	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	<i>Turdus iliacus</i>	Redwing	6 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 6 records post 2000
Birds	<i>Emberiza schoeniclus</i>	Reed Bunting	23 records within 2km	Two records dated 2016 are located within the red line boundary (Grid Reference SK906766). Two records are located within 250m of the site, dated 1970 and 2016. Exact location unknown for all other records – within 2km of the site.	4 records pre 2000 19 records post 2000
Birds	<i>Turdus torquatus</i>	Ring Ouzel	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	<i>Philomachus pugnax</i>	Ruff	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	<i>Alauda arvensis</i>	Skylark	14 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. Exact location unknown for all other records – within 2km of the site.	6 records pre 2000 8 records post 2000
Birds	<i>Gallinago gallinago</i>	Snipe	7 records within 2km	Exact location unknown – within 2km of the site.	5 records pre 2000 2 records post 2000
Birds	<i>Turdus philomelos</i>	Song Thrush	30 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. All other records are located beyond 250m of the site.	6 records pre 2000 24 records post 2000



Birds	Muscicapa striata	Spotted Flycatcher	4 records within 2km	The only known location record is 1.8km north-west of the site (Grid Reference SK857786), dated 1977. Exact location unknown for all other records – within 2km of the site.	3 records pre 2000 1 record post 2000
Birds	Sturnus vulgaris	Starling	97 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. All other records are located beyond 250m of the site.	6 records pre 2000 91 record post 2000
Birds	Apus apus	Swift	11 record within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. All other records are located beyond 250m of the site.	3 records pre 2000 8 records post 2000
Birds	Passer montanus	Tree Sparrow	41 record within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. All other records are located beyond 250m of the site.	3 records pre 2000 38 records post 2000
Birds	Streptopelia turtur	Turtle Dove	14 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. All other records are located beyond 250m of the site.	4 records pre 2000 10 records post 2000
Birds	Numenius phaeopus	Whimbrel	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 0 records post 2000
Birds	Cygnus cygnus	Whooper Swan	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 record post 2000
Birds	Motacilla flava	Yellow Wagtail	19 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. Exact location unknown for all other records – within 2km of the site.	3 records pre 2000 16 records post 2000
Birds	Emberiza citrinella	Yellowhammer	22 records within 2km	One record is located within 250m of the site (Grid Reference SK911763), dated 1970. All other records are located beyond 250m of the site.	4 records pre 2000 18 records post 2000
Bony fish	Anguilla anguilla	European Eel	65 records within 2km	8 records are located within 250m of the site, dated between 1991 and 2017, all associated with the River Till. All other records are located beyond 250m of the site.	50 records pre 2000 15 records post 2000
Bony fish	Cobitis taenia	Spined Loach	25 records within 2km	Two records are located within the red line boundary (Grid Reference SK903768), dated 1978. 18 records are located within 250m of the site boundary, all associated with the River Till. 5 records are located beyond 250m of the site.	14 records pre 2000 11 record post 2000
Butterflies	Coenonympha pamphilus	Small Heath	3 records within 2km	All records are located beyond 250m of the site. Closest of which is located 965m north-west (Grid Reference SK868787), dated 2006.	1 record pre 2000 2 records post 2000
Butterflies	Lasiommata megera	Wall	7 records within 2km	All records are located beyond 250m of the site. Closest of which is located 630m south (Grid Reference SK884757), dated 1986.	5 records pre 2000 2 records post 2000
Moths	Ennomos quercinaria	August Thorn	2 records within 2km	The two records are located 2km south of the site (Grid Reference SK897742), dated 2003 and 2008.	0 records pre 2000 2 records post 2000
Moths	Agrochola lychnidis	Beaded Chestnut	15 records within 2km	All 15 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 15 records post 2000
Moths	Timandra comae	Blood-vein	19 records within 2km	All 19 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 19 records post 2000
Moths	Lycia hirtaria	Brindled Beauty	27 records within 2km	All 27 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 27 records post 2000



Moths	<i>Ceramica pisi</i>	Broom Moth	3 records within 2km	The 3 records are located 2km south of the site (Grid Reference SK897742), dated between 2005 and 2010.	0 records pre 2000 3 records post 2000
Moths	<i>Agrochola macilenta</i>	Brown-spot Pinion	13 records within 2km	All 13 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 13 records post 2000
Moths	<i>Spilarctia luteum</i>	Buff Ermine	20 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~805m north of the site (Grid Reference SK877792), dated 2006.	0 records pre 2000 20 records post 2000
Moths	<i>Atethmia centrago</i>	Centre-barred Sallow	12 records within 2km	All 12 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 12 records post 2000
Moths	<i>Tyria jacobaeae</i>	Cinnabar	20 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~1.5km south of the site (Grid Reference SK894748), dated 2011.	0 records pre 2000 20 records post 2000
Moths	<i>Helotropha leucostigma</i>	Crescent	7 records within 2km	All 7 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2010.	0 records pre 2000 7 records post 2000
Moths	<i>Pelurga comitata</i>	Dark Spinach	8 records within 2km	All 8 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2010.	0 records pre 2000 8 records post 2000
Moths	<i>Xanthorhoe ferrugata</i>	Dark-barred Twin-spot Carpet	24 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1.5km north of the site (Grid Reference SK874799), dated 1999.	1 record pre 2000 23 records post 2000
Moths	<i>Aporophyla lutulenta</i>	Deep-brown Dart	3 records within 2km	All 3 records are located 2km south of the site (Grid Reference SK897742), dated between 2006 and 2011.	0 records pre 2000 3 records post 2000
Moths	<i>Melanchra persicariae</i>	Dot Moth	12 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1.5km north of the site (Grid Reference SK873798), dated 2001.	0 records pre 2000 12 records post 2000
Moths	<i>Apamea remissa</i>	Dusky Brocade	11 record within 2km	All 11 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 11 record post 2000
Moths	<i>Euxoa nigricans</i>	Dusky Dart	3 records within 2km	All 3 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 3 records post 2000
Moths	<i>Ennomos fuscantaria</i>	Dusky Thorn	8 records within 2km	All 8 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 8 records post 2000
Moths	<i>Cirrhia gilvago</i>	Dusky-lemon Sallow	5 records within 2km	All 5 records are located 2km south of the site (Grid Reference SK897742), dated between 2006 and 2011.	0 records pre 2000 5 records post 2000
Moths	<i>Amphipoea oculea</i>	Ear Moth	1 record within 2km	Record is located 2km south of the site (Grid Reference SK897742), dated 2004.	0 records pre 2000 1 record post 2000 (2004)
Moths	<i>Tholera decimalis</i>	Feathered Gothic	6 records within 2km	All 6 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 6 records post 2000
Moths	<i>Agrochola helvola</i>	Flounced Chestnut	2 records within 2km	Both records are located 2km south of the site (Grid Reference SK897742), dated 2011.	0 records pre 2000 2 records post 2000
Moths	<i>Arctia caja</i>	Garden Tiger	6 records within 2km	All 6 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2009.	0 records pre 2000 6 records post 2000
Moths	<i>Hepialus humuli</i>	Ghost Moth	13 records within 2km	All 13 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 13 records post 2000
Moths	<i>Allophyes oxyacanthae</i>	Green-brindled Crescent	16 records within 2km	All 16 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 16 records post 2000
Moths	<i>Acronicta psi</i>	Grey Dagger	12 records within 2km	All 12 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2010.	0 records pre 2000 12 records post 2000
Moths	<i>Xestia agathina</i>	Heath Rustic	3 records within 2km	All 3 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2009.	0 records pre 2000 3 records post 2000
Moths	<i>Acronicta rumicis</i>	Knot Grass	20 records within 2km	All 20 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 20 records post 2000



Moths	<i>Malacosoma neustria</i>	Lackey	2 records within 2km	Both records are located 2km south of the site (Grid Reference SK897742), dated 2003 and 2004.	0 records pre 2000 2 records post 2000
Moths	<i>Apamea anceps</i>	Large Nutmeg	11 record within 2km	All 11 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 11 record post 2000
Moths	<i>Rhizodra lutosa</i>	Large Wainscot	16 records within 2km	All 16 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 16 records post 2000
Moths	<i>Chiasmia clathrata</i>	Latticed Heath	12 records within 2km	All 12 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 12 records post 2000
Moths	<i>Brachylomia viminalis</i>	Minor Shoulder-knot	6 records within 2km	All 6 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2010.	0 records pre 2000 6 records post 2000
Moths	<i>Caradrina morpheus</i>	Mottled Rustic	14 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1.5km north of the site (Grid Reference SK874799), dated 1999.	1 record pre 2000 13 records post 2000
Moths	<i>Amphipyra tragopoginis</i>	Mouse Moth	14 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1.5km north of the site (Grid Reference SK874799), dated 1999.	1 record pre 2000 13 records post 2000
Moths	<i>Watsonalla binaria</i>	Oak Hook-tip	12 records within 2km	All 12 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 12 records post 2000
Moths	<i>Trichiura crataegi</i>	Pale Eggar	11 record within 2km	All 11 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 11 record post 2000
Moths	<i>Orthosia gracilis</i>	Powdered Quaker	52 records within 2km	All 52 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 52 records post 2000
Moths	<i>Litoligia literosa</i>	Rosy Minor	6 records within 2km	All 6 records are located 2km south of the site (Grid Reference SK897742), dated between 2005 and 2011.	0 records pre 2000 6 records post 2000
Moths	<i>Hydraecia micacea</i>	Rosy Rustic	14 records within 2km	All 14 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2010.	0 records pre 2000 14 records post 2000
Moths	<i>Mesapamea secalis</i>	Rustic	10 records within 2km	All 10 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 10 records post 2000
Moths	<i>Cirrhia icteritia</i>	Sallow	16 records within 2km	All 16 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 16 records post 2000
Moths	<i>Ennomos erosaria</i>	September Thorn	6 records within 2km	All 6 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2010.	0 records pre 2000 6 records post 2000
Moths	<i>Scotopteryx chenopodiata</i>	Shaded Broad-bar	11 record within 2km	All records are located beyond 250m of the site. Closest of which are two records located 1.5km south of the site (Grid Reference SK894748), dated 2008 and 2019.	0 records pre 2000 11 record post 2000
Moths	<i>Leucania comma</i>	Shoulder-striped Wainscot	12 records within 2km	All 12 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 12 records post 2000
Moths	<i>Ecliptopera silaceata</i>	Small Phoenix	7 records within 2km	All 7 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 7 records post 2000
Moths	<i>Diarsia rubi</i>	Small Square-spot	13 records within 2km	All records are located beyond 250m of the site. Closest of which is located 1.5km north of the site (Grid Reference SK874799), dated 1999.	1 record pre 2000 12 records post 2000
Moths	<i>Asteroscopus sphinx</i>	Sprawler	6 records within 2km	All 6 records are located 2km south of the site (Grid Reference SK897742), dated between 2004 and 2011.	0 records pre 2000 6 records post 2000
Moths	<i>Chesias legatella</i>	Streak	3 records within 2km	All 3 records are located 2km south of the site (Grid Reference SK897742), dated between 2005 and 2011.	0 records pre 2000 3 records post 2000



Moths	<i>Spilosoma lubricipeda</i>	White Ermine	40 records within 2km	All 40 records are located 2km south of the site (Grid Reference SK897742), dated between 2003 and 2011.	0 records pre 2000 40 records post 2000
Mollusc	<i>Valvata macrostoma</i>	Large-mouthed Valve Snail	1 record within 2km	Record is located within 250m of the site (Grid Reference SK904768), dated 1980.	1 record pre 2000 0 records post 2000
Flowering Plant	<i>Oenanthe fistulosa</i>	Tubular Water-dropwort	3 records within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 2 records post 2000



APPENDIX F – SPECIES RECORDS WITHIN 2KM OF WEST BURTON 3 (BELLWOOD AND BRAMPTON)

Records of Protected and Notable Species Derived from the Desk Study Data Search (LERC)

Group	Scientific Name	Common Name	Records	Location	Date
Amphibians	Bufo bufo	Common Toad	18 record within 2km	One record is located within 250m of the site. Two records are located 365m north-west and 1.1km west of the site, dated 2019 and 2018, respectively. Exact location unknown for all other records – within 2km of the site.	15 records pre 2000 3 records post 2000
Amphibians	Rana temporaria	Common Frog	50 records within 2km	All records are located beyond 250m of the site. Closest records include 3 records located 345m north-west of the site, dated 2018 (Grid Reference SK843821) and 21 records located 260m north-west of the site, dated between 2009 and 2019 (Grid Reference SK842820).	20 records pre 2000 30 records post 2000
Amphibians	Triturus cristatus	Great Crested Newt	6 records within 2km	Exact location unknown – within 2km of the site.	5 records pre 2000 1 records post 2000
Amphibians	Lissotriton vulgaris	Smooth Newt	9 records within 2km	Exact location unknown – within 2km of the site.	6 record pre 2000 3 record post 2000
Reptiles	Zootoca vivipara	Common Lizard	1 record within 2km	The record is located within 250m of the site (Grid Reference SK856813), dated 2012.	0 records pre 2000 1 record post 2000
Reptiles	Natrix helvetica	Grass Snake	19 records within 2km	All records are located beyond 250m of the site. Closest of which is located ~325m west of the site (Grid Reference SK841809) in 2018.	13 records pre 2000 6 records post 2000
Reptiles	Anguis fragilis	Slow-worm	5 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 5 records post 2000
Terrestrial Mammal	Arvicola amphibius	European Water Vole	62 records within 2km	Three records are located within 250m of the Site (Grid References SK844803 and SK843804), dated 2015 and 2016. All other records are located beyond 250m of the site.	11 records pre 2000 51 records post 2000
Terrestrial Mammal	Lepus europaeus	Brown Hare	127 records within 2km	Four records, dated between 2010 and 2018 are located within the red line boundary (Grid References SK847816 and SK862811). 16 records are located within 250m of the site. 107 records are located beyond 250m of the site.	21 records pre 2000 116 records post 2000
Terrestrial Mammal	Meles meles	Eurasian Badger	70 records within 2km	6 records are located within the red line boundary, dated between 2010 and 2015 (all road kill). 3 records are located within 250m of the site, dated between 2007 and 2013 (all road kill). 30 badger sett records are within 2km of the site. Exact locations are unknown.	8 records pre 2000 62 records post 2000
Terrestrial Mammal	Lutra lutra	European Otter	17 records within 2km	One record is located within 250m of the site (Grid Reference SK842810), dated 2005.	4 record pre 2000 13 record post 2000



				16 records are located beyond 250m of the site.	
Terrestrial Mammal	<i>Mustela putorius subsp. furo</i>	Feral Ferret	3 records within 2km	Two records, dated 2012 are located within the red line boundary (Grid Reference SK854815; both road kill). One record is located beyond 250m of the site.	0 records pre 2000 3 records post 2000
Terrestrial Mammal	<i>Erinaceus europaeus</i>	West European Hedgehog	167 record within 2km	14 records are located within the red line boundary, dated between 2010 and 2015. All are road kill associated with Stow Park Road at the north of the site. 20 records are located within 250m of the site, dated between 2005 and 2018. 133 records are located beyond 250m of the site.	21 records pre 2000 146 records post 2000
Terrestrial Mammal	<i>Micromys minutus</i>	Harvest Mouse	5 records within 2km	The only known location records are 650m north-west of the site (Grid Reference SK839820), dated 2007 and ~1.5km north-west, dated 2009 (Grid Reference SK835829). Exact location unknown for the other 3 records – within 2km of the site.	3 records pre 2000 2 records post 2000
Bats	<i>Plecotus auritus</i>	Brown Long-eared Bat	14 records within 2km	The closest known location record is ~715m south-east of the site (Grid Reference SK873785), dated 2012. Exact location unknown for 9 records – within 2km of the site.	2 records pre 2000 12 records post 2000
Bats	<i>Pipistrellus pipistrellus sensu stricto</i>	Common Pipistrelle	87 records within 2km	Two records are located within 250m of the site, dated 2007 (Grid Reference SK845818). All other records are located beyond 250m of the site.	0 records pre 2000 87 records post 2000
Bats	<i>Myotis daubentonii</i>	Daubenton's Bat	7 record within 2km	All records are located beyond 250m of the site. Closest of which is ~1km west of the site, dated 2004 (Grid Reference SK834813).	0 records pre 2000 7 record post 2000
Bats	<i>Myotis nattereri</i>	Natterer's Bat	4 record within 2km	Exact location unknown – within 2km of the site.	2 records pre 2000 2 record post 2000
Bats	<i>Nyctalus noctula</i>	Noctule Bat	10 records within 2km	All records are located beyond 250m of the site. Closest of which is 4 records located 345m north-west of the site, dated between 2006 and 2015 (Grid Reference SK843821).	0 records pre 2000 10 record post 2000
Bats	<i>Pipistrellus</i>	Pipistrelle Bat species	20 records within 2km	Two records are located within 250m of the site, dated 2009 (Grid Reference SK844817). 18 records are located beyond 250m of the site.	1 record pre 2000 19 records post 2000
Bats	<i>Pipistrellus pygmaeus</i>	Soprano Pipistrelle	42 record within 2km	The only known location records are 1.2km north-west of the site, dated 2008 (Grid Reference SK839829) and 1.3km south-west of the site, dated 2014 (SK835791). Exact location unknown for all other records – within 2km of the site.	0 records pre 2000 42 record post 2000
Bats	Unidentified Bat	Unidentified Bat	47 records within 2km	Two records, dated 2007 are located within the red line boundary (Grid Reference SK855815). 45 records are located beyond 250m of the site.	40 records pre 2000 7 records post 2000



Birds	Tyto alba	Barn Owl	182 records within 2km	Only 5 record locations are known – one record located ~370m north-west of the site, dated 2015 (Grid Reference SK842820) and 4 records located 1.8km south-west of the site, dated between 2015 and 2019 (Grid Reference SK836780). Exact location unknown for all other records – within 2km of the site.	2 records pre 2000 180 record post 2000
Birds	Cygnus columbianus	Bewick's Swan	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 0 record post 2000
Birds	Phoenicurus ochruros	Black Redstart	4 record within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 4 record post 2000
Birds	Limosa limosa	Black-tailed Godwit	2 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 record post 2000
Birds	Motacilla flava flava	Blue-headed Wagtail	2 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 record post 2000
Birds	Fringilla montifringilla	Brambling	12 record within 2km	Exact location unknown – within 2km of the site.	1 records pre 2000 11 record post 2000
Birds	Pyrrhula pyrrhula	Bullfinch	129 records within 2km	The only known location records are 3 records 375m north-west of the site, dated between 2009 and 2015 (Grid Reference SK842820) and 1 record 740m west of the site, dated 2009 (Grid Reference SK838817). Exact location unknown for all other records – within 2km of the site.	4 records pre 2000 125 records post 2000
Birds	Cettia cetti	Cetti's Warbler	12 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 12 records post 2000
Birds	Loxia curvirostra	Common Crossbill	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 records post 2000
Birds	Melanitta nigra	Common Scoter	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 records post 2000
Birds	Emberiza calandra	Corn Bunting	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 records post 2000
Birds	Crex crex	Corncrake	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	Cuculus canorus	Cuckoo	31 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 31 records post 2000
Birds	Turdus pilaris	Fieldfare	147 records within 2km	Exact location unknown – within 2km of the site.	4 records pre 2000 143 records post 2000
Birds	Bucephala clangula	Goldeneye	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	Locustella naevia	Grasshopper Warbler	30 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 30 records post 2000
Birds	Tringa ochropus	Green Sandpiper	60 records within 2km	The only known record location is 1.1km west of the site, dated 2010 (Grid Reference SK833811). Exact location unknown for all other records – within 2km of the site.	0 records pre 2000 60 records post 2000
Birds	Tringa nebularia	Greenshank	18 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 18 records post 2000
Birds	Perdix perdix	Grey Partridge	61 record within 2km	The only known record location is 270m south of the site, dated 1977 (Grid Reference SK857786). Exact location unknown for all other records – within 2km of the site.	2 records pre 2000 59 record post 2000
Birds	Anser anser	Greylag Goose	51 record within 2km	The only known record location is 825m south of the site, dated 2006 (Grid Reference SK845787).	0 records pre 2000 51 record post 2000



				Exact location unknown for all other records – within 2km of the site.	
Birds	Coccothraustes coccothraustes	Hawfinch	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	Circus cyaneus	Hen Harrier	2 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 record post 2000
Birds	Falco subbuteo	Hobby	60 record within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 60 record post 2000
Birds	Pernis apivorus	Honey-buzzard	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	Passer domesticus	House sparrow	94 records within 2km	7 records are located within the red line boundary and were recorded in 2008 and 2009. All other records are beyond 250m of the site.	3 records pre 2000 85 records post 2000
Birds	Alcedo atthis	Kingfisher	109 records within 2km	All records are located beyond 250m of the site. Closest known location record is 870m south of the site, dated 2007.	3 records pre 2000 106 records post 2000
Birds	Calcarius lapponicus	Lapland Bunting	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	Vanellus vanellus	Lapwing	180 records within 2km	The only known location records are ~825m south of the site, dated 2006 and ~955m south of the site, dated 1977. Exact location unknown for all other records – within 2km of the site.	11 records pre 2000 169 records post 2000
Birds	Acanthis cabaret	Lesser Redpoll	33 records within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 33 records post 2000
Birds	Linaria cannabina	Linnet	50 records within 2km	Exact location unknown – within 2km of the site.	1 records pre 2000 50 records post 2000
Birds	Circus aeruginosus	Marsh Harrier	12 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 12 record post 2000
Birds	Falco columbarius	Merlin	17 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 17 records post 2000
Birds	Pandion haliaetus	Osprey	9 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 9 records post 2000
Birds	Falco peregrinus	Peregrine	44 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 44 records post 2000
Birds	Coturnix coturnix	Quail	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Birds	Milvus milvus	Red Kite	7 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 7 records post 2000
Birds	Tringa totanus	Redshank	68 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 68 records post 2000
Birds	Turdus iliacus	Redwing	86 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 86 records post 2000
Birds	Emberiza schoeniclus	Reed Bunting	62 records within 2km	The only known record location is 370m north-west of the site, with 2 records dated 2014 and 2015. Exact location unknown for all other records – within 2km of the site.	3 records pre 2000 59 records post 2000
Birds	Turdus torquatus	Ring Ouzel	4 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 4 records post 2000
Birds	Philomachus pugnax	Ruff	16 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 16 records post 2000



Birds	Aythya marila	Scaup	1 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 1 records post 2000
Birds	Alauda arvensis	Skylark	64 records within 2km	All records are located beyond 250m of the site. Closest of which located 270m south of the site, dated 1977 (Grid Reference SK857786).	7 records pre 2000 57 records post 2000
Birds	Gallinago gallinago	Snipe	160 records within 2km	Exact location unknown – within 2km of the site.	3 records pre 2000 157 records post 2000
Birds	Turdus philomelos	Song Thrush	46 records within 2km	One record is located within 250m of the site, dated 2009. All other records are beyond 250m of the site.	2 records pre 2000 44 records post 2000
Birds	Muscicapa striata	Spotted Flycatcher	57 records within 2km	The only known location record is 270m south of the site, dated 1977 (Grid Reference SK857786). Exact location unknown for all other records – within 2km of the site.	1 records pre 2000 56 record post 2000
Birds	Sturnus vulgaris	Starling	104 records within 2km	7 records are located within 250m of the site, dated 2008 and 2009. All other records are beyond 250m of the site.	3 records pre 2000 101 record post 2000
Birds	Apus apus	Swift	67 records within 2km	All records are beyond 250m of the site. Closest of which is 375m north-west of the site, dated 2009 (Grid Reference SK842820).	5 records pre 2000 62 records post 2000
Birds	Anthus trivialis	Tree Pipit	2 records within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 2 records post 2000
Birds	Passer montanus	Tree Sparrow	162 records within 2km	The only known location record is 375m north-west of the site, dated 2015 (Grid Reference SK842820). Exact location unknown for all other records – within 2km of the site.	1 records pre 2000 161 records post 2000
Birds	Streptopelia turtur	Turtle Dove	41 records within 2km	The only known location record is 970m south of the site, dated 2006 (Grid Reference SK860778). Exact location unknown for all other records – within 2km of the site.	1 records pre 2000 40 records post 2000
Birds	Numenius phaeopus	Whimbrel	15 record within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 15 records post 2000
Birds	Haliaeetus albicilla	White-tailed Eagle	2 records within 2km	Exact location unknown – within 2km of the site.	0 record pre 2000 2 records post 2000
Birds	Cygnus cygnus	Whooper Swan	10 record within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 10 record post 2000
Birds	Lullula arborea	Woodlark	5 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 5 record post 2000
Birds	Motacilla flava	Yellow Wagtail	88 records within 2km	Exact location unknown – within 2km of the site.	1 records pre 2000 87 records post 2000
Birds	Emberiza citrinella	Yellowhammer	94 records within 2km	All records are located beyond 250m of the site. Closest of which located 270m south of the site, dated 1977 (Grid Reference SK857786).	16 records pre 2000 78 records post 2000
Bony fish	Anguilla anguilla	European Eel	21 records within 2km	All records are located beyond 250m of the site. Closest of which located 275m west of the site, dated 2018 (Grid Reference SK842806).	10 records pre 2000 11 records post 2000



Butterflies	<i>Coenonympha pamphilus</i>	Small Heath	9 records within 2km	Two records are located within 250m of the site, dated 2006 (Grid Reference SK868787). All other records are beyond 250m of the site.	3 record pre 2000 6 records post 2000
Butterflies	<i>Lasiommata megera</i>	Wall	7 records within 2km	All records are beyond 250m of the site. Closest of which are 10 records located 345m north-west of the site, dated between 2001 and 2006 (Grid Reference SK843821).	5 records pre 2000 27 records post 2000
Moths	<i>Agrochola lychnidis</i>	Beaded Chestnut	29 records within 2km	All records are beyond 250m of the site. Closest of which are 27 records located 345m north-west of the site, between 2003 and 2020 (Grid Reference SK843821).	0 records pre 2000 29 records post 2000
Moths	<i>Timandra comae</i>	Blood-vein	11 records within 2km	All 11 records are located 345m north-west of the site, dated between 2003 and 2011 (Grid Reference SK843821).	0 records pre 2000 11 records post 2000
Moths	<i>Lycia hirtaria</i>	Brindled Beauty	6 records within 2km	All 6 records are located 345m north-west of the site, dated between 2004 and 2011 (Grid Reference SK843821).	0 records pre 2000 6 records post 2000
Moths	<i>Ceramica pisi</i>	Broom Moth	2 records within 2km	Both records are located 345m north-west of the site, dated 2006 (Grid Reference SK843821).	0 records pre 2000 2 records post 2000
Moths	<i>Spilarctia luteum</i>	Buff Ermine	32 records within 2km	All records are beyond 250m of the site. Closest of which are 30 records located 345m north-west of the site, dated between 2003 and 2011 (Grid Reference SK843821).	0 records pre 2000 32 records post 2000
Moths	<i>Atethmia centrigo</i>	Centre-barred Sallow	15 records within 2km	All 15 records are located 345m north-west of the site, dated between 2003 and 2011 (Grid Reference SK843821).	0 records pre 2000 15 records post 2000
Moths	<i>Tyria jacobaeae</i>	Cinnabar	24 records within 2km	All records are beyond 250m of the site. Closest of which are 10 records located 345m north-west of the site, dated between 2006 and 2011 (Grid Reference SK843821).	0 records pre 2000 24 records post 2000
Moths	<i>Helotropha leucostigma</i>	Crescent	2 records within 2km	Both records are located 345m north-west of the site, dated 2003 and 2009 (Grid Reference SK843821).	0 records pre 2000 2 records post 2000
Moths	<i>Pelurga comitata</i>	Dark Spinach	4 records within 2km	All 4 records are located 345m north-west of the site, dated between 2005 and 2008 (Grid Reference SK843821).	0 records pre 2000 4 records post 2000
Moths	<i>Xanthorhoe ferrugata</i>	Dark-barred Twin-spot Carpet	10 records within 2km	One record is located within 250m of the site, dated 1999. All other records are beyond 250m of the site.	2 record pre 2000 8 records post 2000
Moths	<i>Melanchra persicariae</i>	Dot Moth	16 records within 2km	All records are beyond 250m of the site. Closest of which are 15 records located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 16 records post 2000
Moths	<i>Apamea remissa</i>	Dusky Brocade	13 record within 2km	All 13 records are located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 13 record post 2000
Moths	<i>Euxoa nigricans</i>	Dusky Dart	5 records within 2km	All 5 records are located 345m north-west of the site, dated between 2005 and 2011.	0 records pre 2000 5 records post 2000
Moths	<i>Ennomos fuscantaria</i>	Dusky Thorn	21 records within 2km	All records are beyond 250m of the site. Closest of which is 16 records located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 21 records post 2000
Moths	<i>Amphipoea oculea</i>	Ear Moth	2 records within 2km	Both records are located beyond 250m of the site. Closest of which is located 345m north-west of the site, dated 2010.	0 records pre 2000 2 record post 2000



Moths	<i>Tholera decimalis</i>	Feathered Gothic	7 records within 2km	All 7 records are located 345m north-west of the site, dated between 2005 and 2011.	0 records pre 2000 7 records post 2000
Moths	<i>Agrochola helvola</i>	Forester	1 record within 2km	Record is located 345m north-west of the site, dated 2004.	0 records pre 2000 1 record post 2000
Moths	<i>Arctia caja</i>	Garden Tiger	1 record within 2km	Record is located 345m north-west of the site, dated 2007.	0 records pre 2000 1 record post 2000
Moths	<i>Hepialus humuli</i>	Ghost Moth	3 records within 2km	All 3 records are located 345m north-west of the site, dated between 2004 and 2006.	0 records pre 2000 3 records post 2000
Moths	<i>Allophyes oxyacanthae</i>	Green-brindled Crescent	15 records within 2km	All records are beyond 250m of the site. Closest of which is 8 records located 345m north-west of the site, dated between 2005 and 2011.	0 records pre 2000 15 records post 2000
Moths	<i>Xestia agathina</i>	Heath Rustic	1 record within 2km	Record is located 345m north-west of the site, dated 2003.	0 records pre 2000 1 record post 2000
Moths	<i>Acronicta rumicis</i>	Knot Grass	22 records within 2km	All records are beyond 250m of the site. Closest of which is 16 records located 345m north-west of the site, dated between 2003 and 2010.	0 records pre 2000 22 records post 2000
Moths	<i>Malacosoma neustria</i>	Lackey	5 records within 2km	All 5 records are located 345m north-west of the site, dated between 2003 and 2010.	0 records pre 2000 5 records post 2000
Moths	<i>Apamea anceps</i>	Large Nutmeg	4 record within 2km	All records are beyond 250m of the site. Closest of which is 3 records located 345m north-west of the site, dated between 2006 and 2008.	0 records pre 2000 4 record post 2000
Moths	<i>Rhizedra lutosa</i>	Large Wainscot	11 records within 2km	All 11 records are located 345m north-west of the site, dated between 2006 and 2011.	0 records pre 2000 11 records post 2000
Moths	<i>Chiasmia clathrata</i>	Latticed Heath	3 records within 2km	All 3 records are located 345m north-west of the site, dated between 2006 and 2011.	0 records pre 2000 3 records post 2000
Moths	<i>Caradrina morpheus</i>	Mottled Rustic	41 records within 2km	All records are beyond 250m of the site. Closest of which is 34 records located 345m north-west of the site, dated between 2003 and 2011.	1 record pre 2000 41 records post 2000
Moths	<i>Amphipyra tragopoginis</i>	Mouse Moth	36 records within 2km	All records are beyond 250m of the site. Closest of which is 28 records located 345m north-west of the site, dated between 2003 and 2011.	1 record pre 2000 35 records post 2000
Moths	<i>Watsonalla binaria</i>	Oak Hook-tip	5 records within 2km	All 5 records are located 345m north-west of the site, dated between 2003 and 2010.	0 records pre 2000 5 records post 2000
Moths	<i>Trichiura crataegi</i>	Pale Eggar	1 record within 2km	Record is located 345m north-west of the site, dated 2005.	0 records pre 2000 1 record post 2000
Moths	<i>Orthosia gracilis</i>	Powdered Quaker	24 records within 2km	All records are beyond 250m of the site. Closest of which is 28 records located 345m north-west of the site.	0 records pre 2000 24 records post 2000
Moths	<i>Litoligia literosa</i>	Rosy Minor	5 records within 2km	All 5 records are located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 5 records post 2000
Moths	<i>Hydraecia micacea</i>	Rosy Rustic	22 records within 2km	All records are beyond 250m of the site. Closest of which is 19 records located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 22 records post 2000



Moths	<i>Mesapamea secalis</i>	Rustic	43 records within 2km	All records are beyond 250m of the site. Closest of which is 31 records located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 43 records post 2000
Moths	<i>Cirrhia icteritia</i>	Sallow	11 records within 2km	All records are beyond 250m of the site. Closest of which is 10 records located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 11 records post 2000
Moths	<i>Scotopteryx chenopodiata</i>	Shaded Broad-bar	4 records within 2km	All records are beyond 250m of the site. Closest of which is 2 records located 345m north-west of the site, dated 2008.	0 records pre 2000 4 records post 2000
Moths	<i>Leucania comma</i>	Shoulder-striped Wainscot	9 records within 2km	All 9 records are located 345m north-west of the site, dated between 2004 and 2011.	0 records pre 2000 9 records post 2000
Moths	<i>Ecliptopera silaceata</i>	Small Phoenix	9 records within 2km	All records are beyond 250m of the site. Closest of which is 7 records located 345m north-west of the site, dated between 2003 and 2010.	0 records pre 2000 9 records post 2000
Moths	<i>Diarsia rubi</i>	Small Square-spot	21 records within 2km	All records are beyond 250m of the site. Closest of which is 16 records located 345m north-west of the site, dated between 2003 and 2010.	1 record pre 2000 20 records post 2000
Moths	<i>Asteroscopus sphinx</i>	Sprawler	2 records within 2km	Both records are located 345m north-west of the site, dated 2007.	0 records pre 2000 2 records post 2000
Moths	<i>Chesias legatella</i>	Streak	1 record within 2km	Record is located 345m north-west of the site, dated 2005.	0 records pre 2000 1 record post 2000
Moths	<i>Spilosoma lubricipeda</i>	White Ermine	23 records within 2km	All records are beyond 250m of the site. Closest of which is 18 records located 345m north-west of the site, dated between 2003 and 2011.	0 records pre 2000 23 records post 2000
Mollusc	<i>Potamopyrgus antipodarum</i>	Mud Snail	1 record within 2km	Record is located 1.9km south-west of the site, dated 1988.	1 record pre 2000 0 records post 2000
Flowering Plant	<i>Scleranthus annuus</i>	Annual Knawel	1 record within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 0 records post 2000
Flowering Plant	<i>Hyacinthoides non-scripta</i>	Bluebell	2 records within 2km	The only known location record is 1.2km south-west of the site, dated 1999.	2 records pre 2000 0 records post 2000
Flowering Plant	<i>Centaurea cyanus</i>	Cornflower	2 records within 2km	Exact location unknown – within 2km of the site.	0 records pre 2000 2 records post 2000
Flowering Plant	<i>Stellaria palustris</i>	Marsh Stitchwort	2 records within 2km	Exact location unknown – within 2km of the site.	2 records pre 2000 0 records post 2000
Flowering Plant	<i>Torilis arvensis</i>	Spreading Hedge-parsley	1 records within 2km	Exact location unknown – within 2km of the site.	1 record pre 2000 0 records post 2000
Flowering Plant	<i>Oenanthe fistulosa</i>	Tubular Water-dropwort	12 records within 2km	All records beyond 250m of the site. Closest of which is located 825m south-west of the site, dated 2006.	9 record pre 2000 3 records post 2000



APPENDIX G: LOCAL PLANNING POLICY

Policy Reference	Key Policy Text
Central Lincolnshire Local Plan (Adopted April 2017)	
Policy LP19: Renewable Energy Proposals	<p>Proposals for non-wind renewable technology will be assessed on their merits, with the impacts, both individual and cumulative, considered against the benefits of the scheme, taking account of the following:</p> <p>The surrounding landscape and townscape;</p> <ul style="list-style-type: none"> • Heritage assets; • Ecology and diversity; • Residential and visual amenity; • Safety, including ensuring no adverse highway impact; • MoD operations, including having no unacceptable impact on the operation of aircraft • movement or operational radar; and • Agricultural Land Classification (including a presumption against photovoltaic solar farm proposals on the best and most versatile agricultural land). <p>Proposals will be supported where the benefit of the development outweighs the harm caused and it is demonstrated that any harm will be mitigated as far as is reasonably possible.</p> <p>Renewable energy proposals which will directly benefit a local community, have the support of the local community and / or are targeted at residents experiencing fuel poverty, will be particularly supported.</p>
Policy LP20: Green Infrastructure Network	<p>The Central Lincolnshire Authorities will aim to maintain and improve the green infrastructure network in Central Lincolnshire by enhancing, creating and managing multifunctional green space within and around settlements that are well connected to each other and the wider countryside.</p> <p>Development proposals which are consistent with and help deliver the opportunities, priorities and initiatives identified in the latest Central Lincolnshire Green Infrastructure Study and Biodiversity Opportunity Mapping Study, will be supported. Proposals that cause loss or harm to this network will not be permitted unless the need for and benefits of the development demonstrably outweigh any adverse impacts. Where adverse impacts on green infrastructure are unavoidable, development will only be permitted if suitable mitigation measures for the network are provided.</p> <p>Development proposals should ensure that existing and new green infrastructure is considered and integrated into the scheme design from the outset. Where new green infrastructure is proposed, the design should maximise the delivery of ecosystem services and support healthy and active lifestyles.</p> <p>Development proposals must protect the linear features of the green infrastructure network that provide connectivity between green infrastructure assets, including public rights of way, bridleways, cycleways and waterways, and take opportunities to improve such features.</p> <p>Development will be expected to make contributions proportionate to their scale towards the establishment, enhancement and on-going management of green infrastructure by contributing to the development of the strategic green infrastructure network within Central Lincolnshire, in line with guidance set out in LP12.</p>
Policy LP21: Biodiversity and Geodiversity	<p>All development should:</p> <ul style="list-style-type: none"> • protect, manage and enhance the network of habitats, species and sites of international, national and local importance (statutory and non-statutory), including sites that meet the criteria for selection as a Local Site; • minimise impacts on biodiversity and geodiversity; and • seek to deliver a net gain in biodiversity and geodiversity. <p>Development proposals that will have an adverse impact on a European Site or cause significant harm to a Site of Special Scientific Interest, located within or outside Central Lincolnshire, will not be permitted, in accordance with the NPPF.</p>



Policy Reference	Key Policy Text
	<p>Planning permission will be refused for development resulting in the loss, deterioration or fragmentation of irreplaceable habitats, including ancient woodland and aged or veteran trees, unless the need for, and benefits of, the development in that location clearly outweigh the loss or harm.</p> <p>Proposals for major development should adopt an ecosystem services approach, and for large scale major development schemes (such as Sustainable Urban Extensions) also a landscape scale approach, to biodiversity and geodiversity protection and enhancement identified in the Central Lincolnshire Biodiversity Opportunity Mapping Study.</p> <p>Development proposals should create new habitats, and links between habitats, in line with Biodiversity Opportunity Mapping evidence to maintain a network of wildlife sites and corridors to minimise habitat fragmentation and provide opportunities for species to respond and adapt to climate change. Development should seek to preserve, restore and re-create priority habitats, ecological networks and the protection and recovery of priority species set out in the Lincolnshire Biodiversity Action Plan and Geodiversity Action Plan.</p> <p>Where development is within a Nature Improvement Area (NIA), it should contribute to the aims and aspirations of the NIA.</p> <p>Development proposals should ensure opportunities are taken to retain, protect and enhance biodiversity and geodiversity features proportionate to their scale, through site layout, design of new buildings and proposals for existing buildings.</p> <p>Mitigation</p> <p>Any development which could have an adverse effect on sites with designated features and / or protected species, either individually or cumulatively, will require an assessment as required by the relevant legislation or national planning guidance.</p> <p>Where any potential adverse effects to the biodiversity or geodiversity value of designated sites are identified, the proposal will not normally be permitted. Development proposals will only be supported if the benefits of the development clearly outweigh the harm to the habitat and/or species.</p> <p>In exceptional circumstances, where adverse impacts are demonstrated to be unavoidable, developers will be required to ensure that impacts are appropriately mitigated, with compensation measures towards loss of habitat used only as a last resort where there is no alternative. Where any mitigation and compensation measures are required, they should be in place before development activities start that may disturb protected or important habitats and species.</p>
<p>Policy LP22: Green Wedges</p>	<p>Green Wedges, as identified on the Policies Map, have been identified to fulfil one or more of the following functions and policy aims:</p> <ul style="list-style-type: none"> • Prevention of the physical merging of settlements, preserving their separate identity, local character and historic character; • Creation of a multi-functional 'green lung' to offer communities a direct and continuous link to the open countryside beyond the urban area; • Provision of an accessible recreational resource, with both formal and informal opportunities, close to where people live, where public access is maximised without compromising the integrity of the Green Wedge; • Conservation and enhancement of local wildlife and protection of links between wildlife sites to support wildlife corridors. <p>Within the Green Wedges planning permission will not be granted for any form of development, including changes of use, unless:</p> <ol style="list-style-type: none"> a) it can be demonstrated that the development is not contrary or detrimental to the above functions and aims; or b) it is essential for the proposed development to be located within the Green Wedge, and the benefits of which override the potential impact on the Green Wedge. <p>Development proposals within a Green Wedge will be expected to have regard to:</p> <ol style="list-style-type: none"> c) the need to retain the open and undeveloped character of the Green Wedge, physical separation between settlements, historic environment character and green infrastructure value;



Policy Reference	Key Policy Text
	<p>d) the maintenance and enhancement of the network of footpaths, cycleways and bridleways, and their links to the countryside, to retain and enhance public access, where appropriate to the role and function of the Green Wedge;</p> <p>e) opportunities to improve the quality and function of green infrastructure within the Green Wedge with regard to the Central Lincolnshire Green Infrastructure network and Biodiversity Opportunity Mapping.</p> <p>Development proposals adjacent to the Green Wedges will be expected to demonstrate that:</p> <p>f) they do not adversely impact on the function of the Green Wedge, taking into account scale, siting, design, materials and landscape treatment;</p> <p>g) They have considered linkages to and enhancements of the adjacent Green Wedge.</p>
<p>Policy LP23: Local Green Space and other Important Open Space</p>	<p>An area identified as a Local Green Space on the Policies Map will be protected from development in line with the NPPF, which rules out development on these sites other than in very special circumstances.</p> <p>An area identified as an Important Open Space on the Policies Map is safeguarded from development unless it can be demonstrated that:</p> <p>a) In the case of publicly accessible open space, there is an identified over provision of that particular type of open space in the community area and the site is not required for alternative recreational uses or suitable alternative open space can be provided on a replacement site or by enhancing existing open space serving the community area; and</p> <p>b) In the case of all Important Open Spaces, there are no significant detrimental impacts on the character and appearance of the surrounding area, ecology and any heritage assets.</p>
<p>Central Lincolnshire Local Plan Review – Consultation Draft (June 2021)</p>	
<p>Policy S13: Renewable Energy</p>	<p>The Central Lincolnshire Joint Strategic Planning Committee is committed to supporting the transition to a net zero carbon future and will seek to maximise appropriately located renewable energy generated in Central Lincolnshire (such energy likely being wind and solar based).</p> <p>Proposals for renewable energy schemes, including ancillary development, will be supported where the direct, indirect, individual and cumulative impacts on the following considerations are, or will be made, acceptable:</p> <p>i. As a result of its scale, siting or design, the impacts on the following issues are satisfactorily addressed: landscape character; visual amenity; biodiversity; geodiversity; flood risk; townscape; historic assets; and highway safety...</p> <p>Testing compliance with part (i) above will be via applicable policies elsewhere in a development plan document for the area (i.e. this Local Plan; a Neighbourhood Plan, if one exists; any applicable policies in a Minerals or Waste Local Plan; and any further guidance set out in a Supplementary Planning Document).</p> <p>For all matters in (i)-(iii), the applicable local planning authority may commission its own independent assessment of the proposals, to ensure it is satisfied what the degree of harm may be and whether reasonable mitigation opportunities are being taken.</p> <p>Where significant adverse effects are concluded by the local planning authority following consideration of the above assessment(s), such effects will be weighed against the wider environmental, economic, social and community benefits provided by the proposal. In this regard, and as part of the planning balance, significant additional weight in favour of the proposal will arise for any proposal which is community-led for the benefit of that community.</p> <p>In areas that have been designated for their national importance, as identified in the National Planning Policy Framework, renewable energy infrastructure will only be permitted where it can be demonstrated that it would be appropriate in scale, located in areas that do not contribute positively to the objectives of the designation, is sympathetically designed and includes any necessary mitigation measures.</p> <p>Additional matters for solar based energy proposals Proposals for solar thermal or photovoltaics panels to be installed on existing property will be under a presumption in favour of permission unless there is clear and demonstrable significant harm arising.</p> <p>Proposals for ground based photovoltaics, including commercial large scale proposals, will be under a presumption in favour unless:</p> <ul style="list-style-type: none"> • there is clear and demonstrable significant harm arising; or



Policy Reference	Key Policy Text
	<ul style="list-style-type: none"> the proposal is (following a site specific soil assessment) to take place on Best and Most Versatile (BMV) agricultural land, unless such land is peat based and the proposal is part of a wider scheme to protect or enhance the carbon sink of such land; or the land is allocated for another purpose in this Local Plan or other statutory based document (such as a nature recovery strategy or a Local Transport Plan), and the proposal is not compatible with such other allocation. <p>Decommissioning renewable energy infrastructure</p> <p>Permitted proposals will be subject to a condition that will require the facility to be removed and the site fully restored to its original condition (or as near as reasonably practical to its original condition) within one year of that facility becoming non-operational.</p>
Policy S58: Green Infrastructure Network	<p>The Central Lincolnshire Authorities will safeguard green infrastructure in Central Lincolnshire from inappropriate development and work actively with partners to maintain and improve the quantity, quality, accessibility and management of the green infrastructure network.</p> <p>Proposals that cause loss or harm to the green infrastructure network will not be supported unless the need for and benefits of the development demonstrably outweigh any adverse impacts. Where adverse impacts on green infrastructure are unavoidable, development will only be supported if suitable mitigation measures for the network are provided.</p> <p>Development proposals should ensure that existing and new green infrastructure is considered and integrated into the scheme design from the outset. Where new green infrastructure is proposed, the design and layout should take opportunities to incorporate a range of green infrastructure to maximise the delivery of multi-functionality and ecosystem services, support climate change adaptation and encourage healthy and active lifestyles.</p> <p>Development proposals must protect the linear features of the green infrastructure network that provide connectivity between green infrastructure assets, including public rights of way, bridleways, cycleways and waterways, and take opportunities to improve and expand such features.</p> <p>Development will be expected to make a contribution proportionate to their scale towards the establishment, enhancement and on-going management of green infrastructure by contributing to the development of the strategic green infrastructure network within Central Lincolnshire, in accordance with the Developer Contributions SPD.</p>
Policy S59: Protecting Biodiversity and Geodiversity	<p>All development should:</p> <ol style="list-style-type: none"> protect, manage and enhance the ecological network of habitats, species and sites of international, national and local importance (statutory and non-statutory), including sites that meet the criteria for selection as a Local Site; minimise impacts on biodiversity and features of geodiversity value; deliver measurable and proportionate net gains in biodiversity; and protect and enhance the aquatic environment within or adjoining the site, including water quality and habitat. <p>Part One: Designated Sites</p> <p>The following hierarchy of sites will apply in the consideration of development proposals:</p> <ol style="list-style-type: none"> International Sites <p>The highest level of protection will be afforded to internationally protected sites. Development proposals that will have an adverse impact on the integrity of such areas, will not be supported other than in exceptional circumstances, in accordance with the NPPF.</p> <p>Development proposals that are likely to result in a significant adverse effect, either alone or in combination, on any internationally designated site, must satisfy the requirements of the Habitats Regulations (or any superseding similar UK legislation). Development requiring Appropriate Assessment will only be allowed where it can be determined, taking into account mitigation, that the proposal would not result in significant adverse effects on the site's integrity.</p> <ol style="list-style-type: none"> National Sites (NNRs and SSSIs as shown on the Policies Map)



Policy Reference	Key Policy Text
	<p>Development proposals should avoid impact on these nationally protected sites. Development proposals within or outside a national site, likely to have an adverse effect, either individually or in combination with other developments, will not normally be supported unless the benefits of the development, at this site clearly outweigh both the adverse impacts on the features of the site and any adverse impacts on the wider network of nationally protected sites.</p> <p>3. Irreplaceable Habitats</p> <p>Planning permission will be refused for development resulting in the loss, deterioration or fragmentation of irreplaceable habitats, including ancient woodland and aged or veteran trees, unless there are wholly exceptional reasons and a suitable compensation strategy will be delivered.</p> <p>4. Local Sites (LNR, LWS and LGS as shown on the Policies Map)</p> <p>Development likely to have an adverse effect on locally designated sites, their features or their function as part of the ecological network, will only be supported where the need and benefits of the development clearly outweigh the loss, and the coherence of the local ecological network is maintained. Where significant harm cannot be avoided, the mitigation hierarchy should be followed.</p> <p>Part Two: Species and Habitats of Principal Importance</p> <p>All development proposals will be considered in the context of the relevant Local Authority's duty to promote the protection and recovery of priority species and habitats.</p> <p>Development should seek to preserve, restore and re-create priority habitats, ecological networks and the protection and recovery of priority species set out in the Natural Environment and Rural Communities Act 2006, Lincolnshire Biodiversity Action Plan, Lincolnshire Geodiversity Strategy and Local Nature Recovery Strategy.</p> <p>Where adverse impacts are likely, development will only be supported where the need for and benefits of the development clearly outweigh these impacts. In such cases, appropriate mitigation or compensatory measures will be required.</p> <p>Part Three: Mitigation of Potential Adverse Impacts</p> <p>Development should avoid adverse impact on existing biodiversity and geodiversity features as a first principle, in line with the mitigation hierarchy. Where adverse impacts are unavoidable they must be adequately and proportionately mitigated. If full mitigation cannot be provided, compensation will be required as a last resort where there is no alternative.</p> <p>Development will only be supported where the proposed measures for mitigation and/or compensation along with details of net gain are acceptable to the Local Planning Authority in terms of design and location, and are secured for the lifetime of the development with appropriate funding mechanisms that are capable of being secured by condition and/or legal agreement.</p> <p>If significant harm to biodiversity resulting from development cannot be avoided, adequately mitigated, or, as a last resort, compensated for, then planning permission will be refused.</p>
<p>Policy S60: Biodiversity Opportunity and Delivering Measurable Net Gains</p>	<p>Following application of the mitigation hierarchy, development proposals should ensure opportunities are taken to retain, protect and enhance biodiversity and geodiversity features proportionate to their scale, through site layout, design of new buildings and proposals for existing buildings.</p> <p>Development proposals should create new habitats, and links between habitats, in line with Central Lincolnshire Biodiversity Opportunity and Green Infrastructure Mapping evidence, the biodiversity opportunity area principles set out in Appendix 4 to this Plan and the Local Nature Recovery Strategy, to maintain a network of wildlife sites and corridors, to minimise habitat fragmentation and provide opportunities for species to respond and adapt to climate change.</p> <p>Proposals for major and large scale development should seek to deliver wider environmental net gains where feasible.</p> <p>All development proposals must deliver, as a minimum, a 10% measurable biodiversity net gain attributable to the development. The net gain for biodiversity should be calculated using DEFRA's biodiversity metric.</p> <p>Appendix 4: Principles for Development within Biodiversity</p> <p>Opportunity Areas</p> <p>The following guidance provides a set of development principles which should be used when considering site allocations and determining planning applications in the context of the Central Lincolnshire Biodiversity</p>



Policy Reference	Key Policy Text
	<p>Opportunity Mapping (BOM) and the ecological network it alludes to. These principles are to be used in conjunction with policy S60 within this Local Plan. Ecological networks are key to creating a more robust natural environment which will be resilient to future pressures²⁵. They will play an integral role in the creation of Nature Recovery Networks and likely act as the basis of any local work towards a national strategy, for example Local Nature Recovery Strategies.</p> <p>Central Lincolnshire Biodiversity Opportunity Mapping Categories</p> <p>Dark Green: Ecological network - high quality</p> <p>Consists of Priority habitat, these are the core areas of an ecological network and are of high value in terms of distinctiveness. These may require management to either maintain or improve their current condition.</p> <p>Light Green: Ecological network - opportunity for management</p> <p>These areas are not currently Priority habitat, but are important for biodiversity and the functionality of the ecological network of which they are part. They provide an opportunity for their quality to be improved through management, with positive results for biodiversity.</p> <p>Dark Brown: Opportunity for creation - more joined up</p> <p>These are not currently part of an ecological network, but provide opportunities to connect together two or more ecological networks through habitat creation.</p> <p>Light Brown: Opportunity for creation</p> <p>These areas are not currently part of an ecological network, but provide opportunities for increasing the size of an ecological network through habitat creation. Guidance regarding site allocations and planning permission applications in a</p> <p>Biodiversity Opportunity Mapping context.</p> <p>Biodiversity opportunity mapping developed by the Greater Lincolnshire Nature Partnership highlights both the existing ecological network and where the best opportunities lie for improvement in regards to the extent of habitat in the network, the condition or distinctiveness of said habitat and overall connectivity of the network. All policy and decisions should take into account the impact of development to these networks and where possible avoid permitting proposals which may negatively affect the existing network. Where this is not possible, or where development is planned on areas identified as an opportunity for creation, principles should call for quality design which will protect and enhance the existing network.</p> <p>Biodiversity net gain should prioritise onsite habitat creation and management over offsite. Where land earmarked for development contains, either partially or entirely, any areas highlighted by the BOM, these should be seen as opportunities to contribute to onsite biodiversity net gain requirements in a way that will also conserve, restore and enhance ecological connectivity. However, it should be recognised that Ecological network - opportunity for management areas and Opportunity for creation areas identified by the BOM, which are not part of a development area, are well placed as locations for habitat creation or management. Doing so contributes towards any required offsite biodiversity net gain commitments for development. Additionally, habitat created in an ecologically desirable location or in an area identified for biodiversity by a local strategy are valued more highly by Defra's biodiversity net gain metric. Any sites recognised by the BOM which apply to be included on the register of biodiversity gain sites should be given due regard in planning for their importance to enhancing ecological networks.</p> <p>Notes on Development Principles</p> <p>For the purpose of ecological networks "habitat creation" refers to semi natural or natural habitats. Any habitat created should fit with the existing ecological network and be either the same habitat type or related habitat. A related habitat refers to habitats often found in association as part of a dynamic complex. Ecological advice should be sought in the preservation and enhancement of ecological networks and achievement of biodiversity net gain.</p> <p>Development Principles</p> <p>Where allocated sites or sites submitted for planning permission contain or overlap with any Ecological network – high quality area, the following principles should apply:</p> <ol style="list-style-type: none"> 1. High quality ecological network areas consist of Priority habitat and contain the most valuable habitats. It should not be built on and should be buffered against impacts of development. Where development is permitted on land containing areas of high quality ecological network, the development layout should use the principles of the Mitigation Hierarchy and be designed in such a way as to avoid damage to these areas.



Policy Reference	Key Policy Text
	<p>2. High quality ecological network areas should be recognised as a potential opportunity to achieve biodiversity net gain requirements by improving condition through sensitive management. Where allocated sites or sites submitted for planning permission contain or overlap with any</p> <p>Ecological network – opportunity for management area, the following development principles should apply:</p> <ol style="list-style-type: none"> Proposals should avoid development on Ecological network – opportunity for management areas where possible. Where this is not possible, the development layout should ensure that connectivity of the network is maintained. This can be achieved through quality design, for example by leaving strategically important habitat in place to create wildlife corridors or the use of green/brown roofing to act as stepping stones between larger areas of habitat; or through the effective creation of new habitat as part of a landscaping scheme which allows for the migration and dispersal of species. Proposals should fulfil onsite net gain requirements through creation and sensitive management of habitats, in a way that will enhance the ecological network either by ensuring connectivity or improving condition. <p>Where allocated sites or sites submitted for planning permission contain or overlap with any mapped Opportunity for creation areas, the following development principles should apply:</p> <ol style="list-style-type: none"> Where development takes place on Opportunity for creation areas, applications should include information clearly demonstrating how opportunities to maintain or enhance the ecological network (in regards to the extent of habitat in the network, the condition or distinctiveness of said habitat) and overall connectivity in the network, have or will be taken. It should include aspects of quality design; for example, by leaving strategically important habitat in place where possible to create wildlife corridors or the use of green/brown roofing to act as stepping stones between larger areas of habitat. It should also take any opportunities for effective habitat creation as part of a landscaping scheme which ensures connectivity between habitats for the species which utilise them. Proposals should prioritise any Opportunity for creation areas within the development site for habitat creation. This will ensure that requirements for both biodiversity net gain and the enhancement of ecological networks are achieved in an effective way. Habitat creation onsite should maximise the potential for the ecological network in regards to: the extent of habitat in the network, the condition or distinctiveness of said habitat and the overall connectivity of the network. Additionally, habitat created onsite in an ecologically desirable location or in an area identified by a local strategy, are valued more highly by Defra's biodiversity net gain metric.
<p>Policy S65: Trees, Woodland and Hedgerows</p>	<p>Development proposals should be prepared based on the overriding principle that:</p> <ul style="list-style-type: none"> the existing tree and woodland cover is maintained, improved and expanded; and opportunities for expanding woodland are actively considered, and implemented where practical and appropriate to do so. <p>Existing Trees and Woodland</p> <p>Planning permission will only be granted if the proposal provides evidence that it has been subject to adequate consideration of the impact of the development on any existing trees and woodland found on-site (and off-site, if there are any trees near the site, with 'near' defined as the distance comprising 12 times the stem diameter of the off-site tree). If any trees exist on or near the development site, 'adequate consideration' is likely to mean the completion of a British Standard 5837 Tree Survey and, if applicable, an Arboricultural Method Statement.</p> <p>Where the proposal will result in the loss or deterioration of:</p> <ol style="list-style-type: none"> ancient woodland; and/or the loss of aged or veteran trees found outside ancient woodland, <p>permission will be refused, unless and on an exceptional basis the need for, and benefits of, the development in that location clearly outweigh the loss.</p> <p>Where the proposal will result in the loss or deterioration of a tree protected by a Tree Preservation Order or a tree within a Conservation Area, then permission will be refused unless:</p> <ol style="list-style-type: none"> there is no net loss of amenity value which arises as a result of the development; or the need for, and benefits of, the development in that location clearly outweigh the loss.



Policy Reference	Key Policy Text														
	<p>Where the proposal will result in the loss of any other tree or woodland not covered by the above, then the Council will expect the proposal to retain those trees that make a significant contribution to the landscape or biodiversity value of the area, provided this can be done without compromising the achievement of good design for the site.</p> <p>Mitigating for loss of Trees and Woodland</p> <p>Where it is appropriate for higher value tree(s) (category A or B trees (BS5837)) and/or woodland to be lost as part of a development proposal, then appropriate mitigation, via compensatory tree planting, will be required. Such tree planting should be on-site wherever possible and should:</p> <ul style="list-style-type: none"> e) take all opportunities to meet the five Tree Planting Principles (see supporting text); and f) unless demonstrably impractical or inappropriate, provide the following specific quantity of compensatory trees: <table border="1" data-bbox="555 658 1066 1189"> <thead> <tr> <th data-bbox="555 658 820 813">Trunk diameter(mm) at 1.5m above ground of tree lost to development</th> <th data-bbox="820 658 1066 813">Number of replacement trees required, per tree lost*</th> </tr> </thead> <tbody> <tr> <td data-bbox="555 813 820 875">75-200</td> <td data-bbox="820 813 1066 875">1</td> </tr> <tr> <td data-bbox="555 875 820 938">210-400</td> <td data-bbox="820 875 1066 938">4</td> </tr> <tr> <td data-bbox="555 938 820 1001">410-600</td> <td data-bbox="820 938 1066 1001">6</td> </tr> <tr> <td data-bbox="555 1001 820 1064">610-800</td> <td data-bbox="820 1001 1066 1064">9</td> </tr> <tr> <td data-bbox="555 1064 820 1126">810-1000</td> <td data-bbox="820 1064 1066 1126">10</td> </tr> <tr> <td data-bbox="555 1126 820 1189">1000+</td> <td data-bbox="820 1126 1066 1189">11</td> </tr> </tbody> </table> <p>* replacement based on selected standards 10/12 cm girth at 1m</p> <p>New Trees and Woodland</p> <p>Where appropriate and practical, opportunities for new tree planting should be explored as part of all development proposals (in addition to, if applicable, any necessary compensatory tree provision). Where new trees are proposed, they should be done so on the basis of the five Tree Planting Principles. Proposals which fail to provide practical opportunities for new tree planting will be refused.</p> <p>Planting schemes should include provision to replace any plant failures within five years after the date of planting. Planting of trees must be considered in the context of wider plans for nature recovery which seeks to increase biodiversity and green infrastructure generally, not simply planting of trees, and protecting / enhancing soils, particularly peat soils. Tree planting should only be carried out in appropriate locations that will not impact on existing ecology or opportunities to create alternative habitats that could deliver better enhancements for people and wildlife, including carbon storage. Where woodland habitat creation is appropriate, consideration should be given to the economic and ecological benefits that can be achieved through natural regeneration. Any tree planting should use native and local provenance tree species suitable for the location.</p> <p>Management and Maintenance</p> <p>In instances where new trees and/or woodlands are proposed, it may be necessary for the council to require appropriate developer contributions to be provided, to ensure provision is made for appropriate management and maintenance of the new trees and/or woodland.</p> <p>Hedgerows</p> <p>Proposals for new development will be expected to retain existing hedgerows where appropriate and integrate them fully into the design having regard to their management requirements.</p> <p>Proposals for new development will not be supported that would result in the loss of hedges of high landscape, heritage, amenity or biodiversity value unless the need for, and benefits of, the development clearly outweigh the loss and this loss can be clearly demonstrated to be unavoidable.</p>	Trunk diameter(mm) at 1.5m above ground of tree lost to development	Number of replacement trees required, per tree lost*	75-200	1	210-400	4	410-600	6	610-800	9	810-1000	10	1000+	11
Trunk diameter(mm) at 1.5m above ground of tree lost to development	Number of replacement trees required, per tree lost*														
75-200	1														
210-400	4														
410-600	6														
610-800	9														
810-1000	10														
1000+	11														



Policy Reference	Key Policy Text
	Development requiring the loss of a hedgerow protected under The Hedgerow Regulations will only be supported where it would allow for a substantially improved overall approach to the design and landscaping of the development that would outweigh the loss of the hedgerow. Where any hedges are lost, suitable replacement planting or restoration of existing hedges, will be required within the



APPENDIX H – PHASE 1 HABITATS MAPS

A3 maps supplied as a separate volume:

West Burton 1

West Burton 2

West Burton 3



Clarkson and Woods Ltd.

Overbrook Business Centre,
Poolbridge Road, Blackford,
Somerset BS28 4PA

t: 01934 712500

e: info@clarksonwoods.co.uk

