

# Bramford to Twinstead Reinforcement

Volume 6: Environmental Information

Document 6.3.7.1.1: ES Appendix 7.1 – Annex  
A Habitats Baseline UKHab Descriptions

Final Issue A  
April 2023

Planning Inspectorate Reference: EN020002

Infrastructure Planning (Applications: Prescribed Forms and Procedure) Regulations 2009 Regulation 5(2)(a)

nationalgrid



Page intentionally blank

# Contents

---

<b>Annex A: Habitats Baseline UKHab Descriptions</b>	<b>1</b>
1.1    UK Habitat Classification (UKHab) Polygon Descriptions	1
1.2    Line of Trees Linear Habitat Descriptions	71
1.3    Hedgerow Descriptions	75

---

# Annex A: Habitats Baseline UKHab Descriptions

## 1.1 UK Habitat Classification (UKHab) Polygon Descriptions

1.1.1 Habitat classification is shown in Figure 7.1.4 (application document 6.4) for habitat areas as part of the UKHab mapping for the full project and the descriptions listed in Table A-1.

Table A-1 – UKHab Polygon Descriptions

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	A_1021	c1 – Arable and horticulture	-	-	Arable field margin used for vehicle access	-
AB	A_1070	c1c - Cereal Crops	-	-	Arable	-
AB	A_1076	c1c - Cereal Crops	-	-	Arable	-
AB	A_1078	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1079	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1080	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1103	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1191	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1197	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1198	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1361	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1364 / W_1061	r1 – Standing open water and canals	-	Habitat of Principal Importance (HPI)	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1369	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1487	u1c - Artificial unvegetated, unsealed surface	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1503	c1c - Cereal crops	-	-	Arable	-
AB	A_1504	c1c - Cereal Crops	-	-	Arable	-
AB	A_1509	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	A_1514	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_1542	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_401	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_402	c1c - Cereal Crops	-	-	Arable	-
AB	A_406	c1c - Cereal Crops	-	-	Arable	-
AB	A_407	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_409	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_413	c1c - Cereal Crops	-	-	Arable	-
AB	A_448	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_454	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_456	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_460	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_461	c1c - Cereal Crops	-	-	Arable	-
AB	A_529	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_535	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	W_1060	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
AB	W_1064	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
AB	W_1065	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
AB	W_1066	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	W_1069	r2a – Rivers (priority habitat)	-	HPI	<p>Belstead Brook.</p> <p>Steep banked stream approximately 5m wide with partly vegetated banks. Trees along bank including ash. No aquatic vegetation, slow flow, approximately 30cm deep.</p> <p>See Aquatic Baseline Report (<a href="#">application document 6.3.7.3</a>)</p>	-
AB	W_1070	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
AB	A_548	c1 - Arable and horticulture	-	-	Arable	-
AB	A_578	c1c - Cereal Crops	-	-	Arable	-
AB	A_580	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_586	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_587	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_647	c1c - Cereal Crops	-	-	Arable	-
AB	A_745	c1c - Cereal Crops	-	-	Arable	-
AB	A_748	u1 - Built-up areas and gardens	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_824	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_922	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_923	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	A_940	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_1041	w1f7 - Other Lowland mixed deciduous woodland	-	-	<p>Area is 10m strip to north-east of pylon lines. Predominately dense blackthorn interspersed with some taller trees such as ash and silver birch to 10 metres. Some ancient woodland indicators. At the south-west end where the Order Limits extends out due to the pylon the woodland more typical of Hintlesham with pedunculate oak (<i>Quercus robur</i>), hazel coppice and significant silver birch. However, this area is notably younger than the surrounding woodland perhaps due to felling during pylon construction.</p>	<p>Silver birch (<i>Betula pendula</i>) F, hazel (<i>Corylus avellana</i>) F, field maple (<i>Acer campestre</i>) O, ash (<i>Fraxinus excelsior</i>) O, currant family (<i>Ribes</i> sp.) R, pedunculate oak (<i>Quercus robur</i>) A, dog's mercury (<i>Mercurialis perennis</i>) F, enchanter's-nightshade (<i>Circaea lutetiana</i>) R, wood avens (<i>Geum urbanum</i>) O, tufted hair-grass (<i>Deschampsia cespitosa</i>) F, bramble (<i>Rubus fruticosus</i>) F, false brome (<i>Brachypodium sylvaticum</i>) F, common male fern (<i>Dryopteris filix-mas</i>) R, wood-sedge (<i>Carex sylvatica</i>) O, cowslip (<i>Primula veris</i>) R, bugle (<i>Ajuga reptans</i>) R, common dog-violet (<i>Viola riviniana</i>) R, germander speedwell (<i>Veronica chamaedrys</i>) R, wood dock (<i>Rumex sanguineus</i>) R, wood speedwell (<i>Veronica montana</i>) R</p>

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_1054	g4 - Modified grassland	-	-	Sheep grazed, probably reseeded recently. Few herbaceous plants.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, cock's-foot ( <i>Dactylis glomerata</i> ) A, Timothy ( <i>Phleum pratense</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) O, red fescue ( <i>Festuca rubra</i> ) R
AB	H_A_1056	h3h - Mixed scrub	-	-	Open scrubby woodland on the edge of an old, more mature pedunculate oak ( <i>Quercus robur</i> ) dominated woodland. Regrowth of trees with frequent open areas.	Bramble ( <i>Rubus fruticosus</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) F, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, silver birch ( <i>Betula pendula</i> ) O, tufted hair-grass ( <i>Deschampsia cespitosa</i> ) O, groundsel ( <i>Senecio vulgaris</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) F
AB	AB_T_906	h3h - Mixed scrub	-	-	Open scrubby woodland on the edge of an old, more mature pedunculate oak ( <i>Quercus robur</i> ) dominated woodland. Regrowth of trees with frequent open areas.	Bramble ( <i>Rubus fruticosus</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) F, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, silver birch ( <i>Betula pendula</i> ) O, tufted hair-grass ( <i>Deschampsia cespitosa</i> ) O, groundsel ( <i>Senecio vulgaris</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) F
AB	H_A_1058	c1c5 - Winter stubble	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_1059	g4 - Modified grassland	-	-	Vehicle track lines but not surfaced.	Ribwort plantain ( <i>Plantago lanceolata</i> ), bristly ox-tongue ( <i>Picris echioides</i> ), perennial rye-grass ( <i>Lolium perenne</i> )
AB	H_A_1060	c1c - Cereal crops	-	-	Arable	-
AB	H_A_1063	g4 - Modified grassland	-	-	Mown grass verge at side of arable field, some fungi present.	Field bindweed ( <i>Convolvulus arvensis</i> ), hogweed ( <i>Heracleum sphondylium</i> ), common sorrel ( <i>Rumex acetosa</i> ), common nettle ( <i>Urtica dioica</i> ), dandelion ( <i>Taraxacum agg.</i> ), thistle ( <i>Cirsium sp.</i> ), meadow buttercup ( <i>Ranunculus acris</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ),
AB	H_A_1076	g4 - Modified grassland	-	-	Arable field edge, regularly mown grassland. Low sward height. Approx. 2-3m wide.	dandelion ( <i>Taraxacum agg.</i> ), meadow grass ( <i>Poa sp.</i> ), yarrow ( <i>Achillea millefolium</i> ), cock's-foot ( <i>Dactylis glomerata</i> )
AB	H_A_1087	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_1123	g4 - Modified grassland	-	-	Sheep grazed probably reseeded recently. Forb poor. Only surveyed from gate hence low species count.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, cock's-foot ( <i>Dactylis glomerata</i> ) A, Timothy ( <i>Phleum pratense</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) O, red fescue ( <i>Festuca rubra</i> ) R
AB	H_A_1124	c1a - Arable field margins	-	HPI	Cut margin potentially used for access. Scattered trees of hawthorn and Salix sp. may be relic of old hedgerow. Ditch present on eastern side separating arable fields. Not wet throughout year. Dog's mercury in ditch.	Hawthorn ( <i>Crataegus monogyna</i> ), willow ( <i>Salix sp.</i> ), bristly ox-tongue ( <i>Picris echioides</i> ), dove's-foot crane's-bill ( <i>Geranium molle</i> ), common ragwort ( <i>Senecio jacobaea</i> ) LA, curled dock ( <i>Rumex crispus</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), hogweed ( <i>Heracleum sphondylium</i> ), primrose ( <i>Primula vulgaris</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), red fescue ( <i>Festuca rubra</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), trefoil ( <i>Trifolium sp.</i> ), white dead-nettle ( <i>Lamium album</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), soft-rush ( <i>Juncus effusus</i> ), cowslip ( <i>Primula veris</i> ), groundsel ( <i>Senecio vulgaris</i> )
AB	H_A_1127	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_636	u1 - Built-up areas and gardens	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_728	c1a - Arable field margins	HPI		Strip of land left as arable field margin.	Perennial rye-grass ( <i>Lolium perenne</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ), hogweed ( <i>Heracleum sphondylium</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) F, cowslip ( <i>Primula veris</i> ), groundsel ( <i>Senecio vulgaris</i> ), field bindweed ( <i>Convolvulus arvensis</i> ) R, black medick ( <i>Medicago lupulina</i> ) R, spear thistle ( <i>Cirsium vulgare</i> )
AB	H_A_731	c1 - Arable and horticulture	-	-	Arable	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_732	g3c - Other neutral grassland			Arable field margin grassland, could be field margin as looks like it is managed for wildlife however is larger than 12m.	Cock's-foot ( <i>Dactylis glomerata</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) O, Creeping buttercup ( <i>Ranunculus repens</i> ) O, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) R, meadow foxtail ( <i>Alopecurus pratensis</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) O
AB	H_A_736	c1a - Arable field margins	-	HPI	-	Creeping bent ( <i>Agrostis stolonifera</i> ), common bent ( <i>Agrostis capillaris</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), common couch grass ( <i>Elytrigia repens</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), daisy ( <i>Bellis perennis</i> ), common ragwort ( <i>Senecio jacobaea</i> ), common knapweed ( <i>Centaurea nigra</i> ), common vetch ( <i>Vicia sativa</i> )
AB	H_A_737	c1 - Arable and horticulture	-	-	Arable field	-
AB	H_A_738	u1b - Developed land; sealed surface	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_740	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_742	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_744	u1c - Artificial unvegetated, unsealed surface	-	-	Gravel track with early colonising species growing through. Largely unvegetated. Public footpath and bridleway.	Daisy ( <i>Bellis perennis</i> ), dandelion ( <i>Taraxacum agg</i> )
AB	H_A_745	h - Heathland and shrub	-	-	Rows of planted shrubs, not all successful. Tree guards present on all. Young plants mainly with small area of semi-mature field maple.	Privet ( <i>Ligustrum vulgare</i> ), rose ( <i>Rosa sp.</i> ) dogwood ( <i>Cornus sanguinea</i> ), blackthorn ( <i>Prunus spinosa</i> ), field maple ( <i>Acer campestre</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), bramble ( <i>Rubus fruticosus</i> ), lords and ladies ( <i>Arum maculatum</i> ), broad-leaved dock ( <i>Rumex obtusifolius</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), curled dock ( <i>Rumex crispus</i> ), dandelion ( <i>Taraxacum agg.</i> ), wild teasel ( <i>Dipsacus fullonum</i> ), cinquefoil ( <i>Potentilla sp.</i> )
AB	H_A_747	h3h - Mixed scrub	Plantation	-	Rows of planted shrubs, not all successful. Tree guards present on all. Young plants mainly with small area of semi-mature field maple.	Privet ( <i>Ligustrum vulgare</i> ), rose ( <i>Rosa sp.</i> ) dogwood ( <i>Cornus sanguinea</i> ), blackthorn ( <i>Prunus spinosa</i> ), field maple ( <i>Acer campestre</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), bramble ( <i>Rubus fruticosus</i> ), lords and ladies ( <i>Arum maculatum</i> ), broad-leaved dock ( <i>Rumex obtusifolius</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), curled dock ( <i>Rumex crispus</i> ), dandelion ( <i>Taraxacum agg.</i> ), wild teasel ( <i>Dipsacus fullonum</i> ), cinquefoil ( <i>Potentilla sp.</i> )
AB	H_A_748	u1b - Developed land; sealed surface	-	-	Road	-
AB	H_A_749	h3h - Mixed scrub	-	-	Planted scrubby species as found in adjacent area. Honeybees kept in fenced off area.	Rose ( <i>Rosa sp.</i> ), dogwood ( <i>Cornus sanguinea</i> ), blackthorn ( <i>Prunus spinosa</i> ), cowslip ( <i>Primula veris</i> ), rosemary ( <i>Rosmarinus officinalis</i> ), selfheal ( <i>Prunella vulgaris</i> ), wild teasel ( <i>Dipsacus fullonum</i> ), crested dog's-tail ( <i>Cynosurus cristatus</i> ), yarrow ( <i>Achillea millefolium</i> ), buttercup ( <i>Ranunculus sp.</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), crane's-bill ( <i>Geranium sp.</i> )
AB	H_A_750	g4 - Modified grassland	-	-	Mowed grassy strip on other side of fence. Herb rich but low diversity.	Perennial rye-grass ( <i>Lolium perenne</i> ), bristly oxtongue ( <i>Picris echioides</i> ) ribwort plantain ( <i>Plantago lanceolata</i> ), white clover ( <i>Trifolium repens</i> )
AB	H_A_751	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_752	h3h - Mixed scrub	-	-	Mount of scrub	Buddleia ( <i>Buddleja sp.</i> ) R, sycamore ( <i>Acer pseudoplatanus</i> ) O, bramble ( <i>Rubus fruticosus</i> ) A, elder ( <i>Sambucus nigra</i> ) R, common nettle ( <i>Urtica dioica</i> ) D, cow parsley ( <i>Anthriscus sylvestris</i> ) O, hemlock ( <i>Conium maculatum</i> ) F, red dead-nettle ( <i>Lamium purpureum</i> ) O

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_753	g4 - Modified grassland	-	-	Not arable. Field boundary as used as dump for horse dung and other soil. Flora same as nearby modified grassland but with groundsel and Columbine. Almost all perennial rye-grass.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, groundsel ( <i>Senecio vulgaris</i> ), columbine ( <i>Aquilegia vulgaris</i> ) R
AB	H_A_754	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_755	h3h - Mixed scrub	-	-	Likely historic pond present that is now completely overgrown with no water present. Pond area now contains almost solely elder scrub, encircled by trees including mature pedunculate oak ( <i>Quercus robur</i> ). Some planting of conifers on house side.	Elder ( <i>Sambucus nigra</i> ), field maple ( <i>Acer campestre</i> ), pedunculate oak ( <i>Quercus robur</i> ), English elm ( <i>Ulmus procera</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), burdock ( <i>Arctium sp.</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), perennial rye-grass ( <i>Lolium perenne</i> ) A
AB	H_A_756	g4 - Modified grassland	-	-	Transitioning between perennial rye-grass on golf course to species poor grassland near arable field. Only diversity is on areas of dumped soil and detritus where ruderal species are growing.	Perennial rye-grass ( <i>Lolium perenne</i> ), lesser stitchwort ( <i>Stellaria graminea</i> ), common nettle ( <i>Urtica dioica</i> ) A, burdock ( <i>Arctium sp.</i> ), groundsel ( <i>Senecio vulgaris</i> ), white dead-nettle ( <i>Lamium album</i> ), ground ivy ( <i>Glechoma hederacea</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), hemlock ( <i>Conium maculatum</i> )
AB	H_A_758	c1 - Arable and horticulture	-	-	Arable crop including narrow strip around edge used for access.	-
AB	H_A_759	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_760	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_761	w1f7 - Other Lowland mixed deciduous woodland	-	-	Area to the east of the public footpath, different composition, planted later in 17th century than other side which lacks the dominant canopy layer and is dominated by typical scrub species growing to young trees. Mature wild cherry. Some indication of planting of non-native trees historically. Some management indicated by old brash piles. Hazel not coppiced to extent of main woodland. Mature pedunculate oaks are present on eastern side of footpath with some other indicator species. Though these indicator species are common even in cities it is interesting to see the difference in ground flora composition and may mean the woodland to east of the trees is secondary. Mature horse chestnut likely planted on boundary to north-eastern extent. Mature Scots pine.	Pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) D, hazel ( <i>Corylus avellana</i> ) O, horse chestnut ( <i>Aesculus hippocastanum</i> ), ash ( <i>Fraxinus excelsior</i> ), wild cherry ( <i>Prunus avium</i> ) LA, willow ( <i>Salix sp.</i> ) O, hornbeam ( <i>Carpinus betulus</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) D, common nettle ( <i>Urtica dioica</i> ) A, lords and ladies ( <i>Arum maculatum</i> ) F, wood anemone ( <i>Anemone nemorosa</i> ) R, common dog-violet ( <i>Viola riviniana</i> ) R, primrose ( <i>Primula vulgaris</i> ), lesser celandine ( <i>Ranunculus ficaria</i> ), wood avens ( <i>Geum urbanum</i> ), herb-Robert ( <i>Geranium robertianum</i> ), Scots pine ( <i>Pinus sylvestris</i> ) O
AB	H_A_763	g3c - Other neutral grassland	-	-	Field margin not meeting definition as used as a track, not for biodiversity benefit.	Cut-leaved crane's-bill ( <i>Geranium dissectum</i> ), lady's-mantle ( <i>Alchemilla vulgaris</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), creeping bent ( <i>Agrostis stolonifera</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), shepherd's-purse ( <i>Capsella bursa-pastoris</i> ), white dead-nettle ( <i>Lamium album</i> ), dandelion ( <i>Taraxacum agg.</i> ), common whitlowgrass ( <i>Erophila verna agg.</i> ), primrose ( <i>Primula vulgaris</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ), groundsel ( <i>Senecio vulgaris</i> ), common knapweed ( <i>Centaurea nigra</i> ), cowslip ( <i>Primula veris</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_764	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H_A_765	r1a - Eutrophic standing waters	-	HPI	Pond in arable field. Lacks suitable connectivity to nearby habitats or woodland. Large mature likely white willow ( <i>Salix alba</i> ).	White willow ( <i>Salix alba</i> ), bulrush ( <i>Typha latifolia</i> ) A, pendulous sedge ( <i>Carex pendula</i> ), soft-rush ( <i>Juncus effusus</i> ), hard-rush ( <i>Juncus inflexus</i> ), celery-leaved buttercup ( <i>Ranunculus sceleratus</i> ), pondweed ( <i>Potamogeton</i> sp.)
AB	H_A_766	c1 - Arable and horticulture	-	-	Arable field margin surrounding pond. Cleavers locally abundant to west of pond.	Cock's-foot ( <i>Dactylis glomerata</i> ) A, common nettle ( <i>Urtica dioica</i> ) A, perennial rye-grass ( <i>Lolium perenne</i> ) A, red dead-nettle ( <i>Lamium purpureum</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) R, burdock ( <i>Arctium</i> sp.) R, hawthorn ( <i>Crataegus monogyna</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) F, cleavers ( <i>Galium aparine</i> ) LA, Yorkshire fog ( <i>Holcus lanatus</i> ) R
AB	H_A_767	w1f7 - Other Lowland mixed deciduous woodland	-	-	Enclosed by ditches. Similar to PoAWS3 to southwest in structure with semi-mature and couple of mature pedunculate oaks. Young lime, potentially coppiced. Some evidence of cut trees, including sycamore, which have started to grow back. Clearing where power line tower is but woodland composition not significantly different. Elder and bramble growing in clearing. Same condition as nearby PoAWS.	Pedunculate oak ( <i>Quercus robur</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), hornbeam ( <i>Carpinus betulus</i> ), horse chestnut ( <i>Aesculus hippocastanum</i> ), lime ( <i>Tilia</i> sp.) hawthorn ( <i>Crataegus monogyna</i> ), elm ( <i>Ulmus</i> sp.), holly ( <i>Ilex aquifolium</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) D, common dog-violet ( <i>Viola riviniana</i> ) R, lords and ladies ( <i>Arum maculatum</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) O, common nettle ( <i>Urtica dioica</i> ) LA, garlic mustard ( <i>Alliaria petiolata</i> ) O, ivy ( <i>Hedera helix</i> ) D, cleavers ( <i>Galium aparine</i> ), hedge bedstraw ( <i>Galium mollugo</i> ), cowslip ( <i>Primula veris</i> ), elder ( <i>Sambucus nigra</i> ), bramble ( <i>Rubus fruticosus</i> ) A
AB	H_A_769	w1f7 - Other Lowland mixed deciduous woodland	-	-	Area of young and semi mature trees. Canopy of coppiced trees with a few conifers. On northern border of woodland, a line of hazel and hawthorn is present, indicating old hedgerow. Common nettle and cleaver locally abundant and bramble frequent in woodland clearing where woodland goes east. Dry ditch along north of woodland, east to west. Eastern section of woodland separated by access track and ditch running north to south, small amount of water no flow.	Lime ( <i>Tilia</i> sp.), pedunculate oak ( <i>Quercus robur</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), red oak ( <i>Quercus rubra</i> ) R, hornbeam ( <i>Carpinus betulus</i> ) R, wild cherry ( <i>Prunus avium</i> ) F, Scots pine ( <i>Pinus sylvestris</i> ) R, Leyland cypress ( <i>Cupressus x leylandii</i> ) R, holly ( <i>Ilex aquifolium</i> ) O, hazel ( <i>Corylus avellana</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, dog's mercury ( <i>Mercurialis perennis</i> ) A, wood anemone ( <i>Anemone nemorosa</i> ) F, wood avens ( <i>Geum urbanum</i> ) R, lords and ladies ( <i>Arum maculatum</i> ) R, lesser celandine ( <i>Ranunculus ficaria</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) R, common chickweed ( <i>Stellaria media</i> ) R, false brome ( <i>Brachypodium sylvaticum</i> ) R, cleavers ( <i>Galium aparine</i> ) LA, common nettle ( <i>Urtica dioica</i> ) LA, bramble ( <i>Rubus fruticosus</i> ) F
AB	H_A_770	c1 - Arable and horticulture	-	-	Arable field.	-
AB	H_A_771	c1a - Arable field margins	-	HPI	Very short grass in northern section. Southern section longer.	Cock's-foot ( <i>Dactylis glomerata</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), red fescue ( <i>Festuca rubra</i> ) yellow oat-grass ( <i>Trisetum flavescens</i> ), Creeping bent ( <i>Agrostis stolonifera</i> ), bristly ox-tongue ( <i>Picris echioides</i> ), spear thistle ( <i>Cirsium vulgare</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), forget-me-not ( <i>Myosotis</i> sp.) R, curled dock ( <i>Rumex crispus</i> ), field wood-rush ( <i>Luzula campestris</i> ), goat's-beard ( <i>Tragopogon pratensis</i> ), wood dock ( <i>Rumex sanguineus</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_772	w1f7 - Other Lowland mixed deciduous woodland	-	-	This section of woodland separated by a ditch is much younger and planted more recently than the confirmed ancient woodland areas, the ground flora composition much less diverse and species poor. Canopy is predominantly mature pedunculate oak and semi-mature ash, remnants of Scots pine plantation still present in areas. Common nettle mostly present along public right of way (PRoW). Internal ditches through area, along areas of PRoW.	Scots pine ( <i>Pinus sylvestris</i> ) LF, pedunculate oak ( <i>Quercus robur</i> ), ash ( <i>Fraxinus excelsior</i> ), downy birch ( <i>Betula pubescens</i> ) F, silver birch ( <i>Betula pendula</i> ) F, European larch ( <i>Larix decidua</i> ) R, hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) A, spindle ( <i>Euonymus europaeus</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, field maple ( <i>Acer campestre</i> ) F, Midland hawthorn ( <i>Crataegus laevigata</i> ) R, hybrid hawthorn ( <i>Crataegus x media</i> ) R, common dog-violet ( <i>Viola riviniana</i> ), primrose ( <i>Primula vulgaris</i> ) R, lords and ladies ( <i>Arum maculatum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) D, false brome ( <i>Brachypodium sylvaticum</i> ) R, wild garlic ( <i>Allium ursinum</i> ), wood spurge ( <i>Euphorbia amygdaloides</i> ), barren strawberry ( <i>Fragaria sterilis</i> ), wood avens ( <i>Geum urbanum</i> ), wood forget-me-not ( <i>Myosotis sylvatica</i> ) F, common male fern ( <i>Dryopteris filix-mas</i> ), enchanter's-nightshade ( <i>Circaea lutetiana</i> ) R, wood millet ( <i>Milium effusum</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, herb-Paris ( <i>Paris quadrifolia</i> ) LO, common nettle ( <i>Urtica dioica</i> ) LA
AB	H_A_775	g3c - Other neutral grassland	Scattered trees	-	Grassland meadow, well managed with varying sward heights.	Perennial rye-grass ( <i>Lolium perenne</i> ) F, Timothy ( <i>Phleum pratense</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, tufted hair-grass ( <i>Deschampsia cespitosa</i> ) O, thistle ( <i>Cirsium</i> sp.) R, dandelion ( <i>Taraxacum</i> agg.) O, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) LF, ribwort plantain ( <i>Plantago lanceolata</i> ) O, marsh-mallow ( <i>Althaea officinalis</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) O, horsetail ( <i>Equisetum</i> sp.) R, silver birch ( <i>Betula pendula</i> ), alder ( <i>Alnus glutinosa</i> )
AB	H_A_776	w1f7 - Other Lowland mixed deciduous woodland	-	-	Area of ancient woodland pedunculate oak ( <i>Quercus robur</i> ) and ash, some ash in poor health, field maple, downy birch, and lime. Rides present within woodland species rich.	Pedunculate oak ( <i>Quercus robur</i> ) A, ash ( <i>Fraxinus excelsior</i> ) A, field maple ( <i>Acer campestre</i> ) F, downy birch ( <i>Betula pubescens</i> ) O, large leaved-lime ( <i>Tilia platyphyllos</i> ) R, hazel ( <i>Corylus avellana</i> ) A, spindle ( <i>Euonymus europaeus</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) A, bluebell ( <i>Hyacinthoides non-scripta</i> ) F, wood anemone ( <i>Anemone nemorosa</i> ) A, wild garlic ( <i>Allium ursinum</i> ) F, wood speedwell ( <i>Veronica montana</i> ) R, yellow archangel ( <i>Lamiastrum galeobdolon</i> ) O, wood forget-me-not ( <i>Myosotis sylvatica</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) O, lords and ladies ( <i>Arum maculatum</i> ) R, primrose ( <i>Primula vulgaris</i> ) O, lesser celandine ( <i>Ranunculus ficaria</i> ) R, green-flowered helleborine ( <i>Epipactis phyllanthes</i> ) LF, wood millet ( <i>Milium effusum</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) R, barren strawberry ( <i>Fragaria sterilis</i> ) R, wood spurge ( <i>Euphorbia amygdaloides</i> ) LF, common dog-violet ( <i>Viola riviniana</i> ) O, Creeping soft-grass ( <i>Holcus mollis</i> ) R, broad buckler fern ( <i>Dryopteris dilatata</i> ) R, bugle ( <i>Ajuga reptans</i> ) R, remote sedge ( <i>Carex remota</i> ) O, wild strawberry ( <i>Fragaria vesca</i> ) R, meadowsweet ( <i>Filipendula ulmaria</i> ) R, pendulous sedge ( <i>Carex pendula</i> ) R, hairy sedge ( <i>Carex hirta</i> ) R, wood avens ( <i>Geum urbanum</i> ) R, tufted hair-grass ( <i>Deschampsia cespitosa</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) R, yellow pimpernel ( <i>Lysimachia nemorum</i> ) R, wood-sorrel ( <i>Oxalis acetosella</i> ) LF, common figwort ( <i>Scrophularia nodosa</i> ) R, cuckooflower ( <i>Cardamine pratensis</i> ) R, man orchid ( <i>Orchis anthropophora</i> ) R.
AB	H_A_779	g3c - Other neutral grassland	-	-	Grassland	Creeping soft-grass ( <i>Holcus mollis</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) A, Creeping buttercup ( <i>Ranunculus repens</i> ) O, groundsel ( <i>Senecio vulgaris</i> ) F, sheep's sorrel ( <i>Rumex acetosella</i> ) O, thistle ( <i>Cirsium</i> sp.) R, black medick ( <i>Medicago lupulina</i> ) A, dandelion ( <i>Taraxacum</i> agg.) R, selfheal ( <i>Prunella vulgaris</i> ) O
AB	H_A_785	c1 - Arable and horticulture	-	-	Arable field	-
AB	AB_T_855	C1 – Arable and horticulture	-	-	Arable field	-
AB	H_A_788	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_791	c1 - Arable and horticulture	-	-	Arable	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_795	c1 - Arable and horticulture	-	-	Arable field.	-
AB	H_A_796	c1a8 - Game bird mix strips and corners	-	-	Corn crop used for game cover. Pheasant feeders present.	-
AB	H_A_802	g3c - Other neutral grassland	-	-	Not considered to be left for arable field margin biodiversity, just left for access for shooting and to ditches. Couple of scattered ash along field boundary.	Common knapweed ( <i>Centaurea nigra</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), cowslip ( <i>Primula veris</i> ), goat's-rue ( <i>Galega officinalis</i> ), bramble ( <i>Rubus fruticosus</i> ) LA, hedge bedstraw ( <i>Galium mollugo</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), ash ( <i>Fraxinus excelsior</i> )
AB	H_A_803	h3h - Mixed scrub	Anthills		Potentially planted for game cover. Dominated by willow likely goat willow, with hazel and blackthorn seedlings. Very dense in places. Ant hills present though grassland lacks tufts. Artificial game shelter present.	False oat grass ( <i>Arrhenatherum elatius</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), bramble ( <i>Rubus fruticosus</i> ) LA, common knapweed ( <i>Centaurea nigra</i> ), dog-rose ( <i>Rosa canina</i> agg.), silverweed ( <i>Potentilla anserina</i> ), creeping cinquefoil ( <i>Potentilla recta</i> )
AB	H_A_804	c1 - Arable and horticulture	-	-	Ploughed arable land	-
AB	H_A_805	g3c - Other neutral grassland	-	-	Field that may be grazed occasionally but no animals present at time of survey.	Rough meadow-grass ( <i>Poa trivialis</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), meadow foxtail ( <i>Alopecurus pratensis</i> ), cowslip ( <i>Primula veris</i> ), common nettle ( <i>Urtica dioica</i> ), buttercup ( <i>Ranunculus</i> sp.), dandelion ( <i>Taraxacum</i> agg.), vetch ( <i>Vicia</i> sp.), ribwort plantain ( <i>Plantago lanceolata</i> ), white clover ( <i>Trifolium repens</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> )
AB	H_A_806	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_807	f2b - Purple moor grass and rush pastures	Rushes dominant	HPI	Rushes Dominant. Area of pastureland with high water table dominated by soft rush and ground flora by brooklime, with scattered scrub grey willow.	Brooklime ( <i>Veronica beccabunga</i> ) A, soft-rush ( <i>Juncus effusus</i> ) D, willowherb ( <i>Epilobium</i> sp.) A, Yorkshire fog ( <i>Holcus lanatus</i> ) A, Creeping buttercup ( <i>Ranunculus repens</i> ) F, mind-your-own-business ( <i>Soleirolia soleirolii</i> ) A, sedge ( <i>Carex</i> sp.) F, sphagnum moss ( <i>Sphagnum</i> spp.) O,
AB	H_A_808	c1 - Arable and horticulture	-	-	Arable.	-
AB	H_A_817	g - Grassland	-	-	Horse grazed paddock, no access	N/A – Not surveyed
AB	H_A_818	g3c - Other neutral grassland	Horse grazed	-	Grassland strip along field boundary and horse grazed. Fields	Meadow-grass ( <i>Poa</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), common nettle ( <i>Urtica dioica</i> ), brome ( <i>Bromopsis</i> sp.)
AB	H_A_819	w1f7 - Other Lowland mixed deciduous woodland	-	-	Canopy - predominantly mature pedunculate oak ( <i>Quercus robur</i> ) and semi mature ash. Under storey - hazel, hawthorn, silver birch, spindle, Blackthorn, sycamore. Internal ditches through area, along areas of PRoW and footpaths.	Pedunculate oak ( <i>Quercus robur</i> ), ash ( <i>Fraxinus excelsior</i> ), hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), silver birch ( <i>Betula pendula</i> ), spindle ( <i>Euonymus europaeus</i> ), blackthorn ( <i>Prunus spinosa</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), common dog-violet ( <i>Viola riviniana</i> ), primrose ( <i>Primula vulgaris</i> ), lords and ladies ( <i>Arum maculatum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), wild garlic ( <i>Allium ursinum</i> ), wood spurge ( <i>Euphorbia amygdaloides</i> ), barren strawberry ( <i>Fragaria sterilis</i> ), wood avens ( <i>Geum urbanum</i> )
AB	H_A_824	g3c - Other neutral grassland	-	-	Grassland strip along field boundary. Short sward height.	Ribwort plantain ( <i>Plantago lanceolata</i> ), yarrow ( <i>Achillea millefolium</i> ), dandelion ( <i>Taraxacum</i> agg.), meadow-grass ( <i>Poa</i> sp.)
AB	H_A_825	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_826	g3c - Other neutral grassland	Tall herb		Grassland strip that hasn't been used for vehicle access. Nettle dominated.	Common nettle ( <i>Urtica dioica</i> ), white dead-nettle ( <i>Lamium album</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) A, spear thistle ( <i>Cirsium vulgare</i> ) F, willowherb ( <i>Epilobium</i> sp.) O, perennial rye-grass ( <i>Lolium perenne</i> ) D, cowslip ( <i>Primula veris</i> ) O, rose ( <i>Rosa</i> sp.) O, bristly ox-tongue ( <i>Picris echioides</i> ) F, vetch ( <i>Vicia</i> sp.) F, daffodil ( <i>Narcissus</i> sp.) O

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_827	g3c - Other neutral grassland	Horse grazed	-	Horse grazed paddocks, divided by semi-permanent electric fencing. Subject to varied grazing pressures throughout the year. Handful of trees towards centre of fields.	Red dead-nettle ( <i>Lamium purpureum</i> ), yarrow ( <i>Achillea millefolium</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), dandelion ( <i>Taraxacum</i> agg.), common ragwort ( <i>Senecio jacobaea</i> ), bristly oxtongue ( <i>Picris echioides</i> ), dock ( <i>Rumex</i> sp.), ribwort plantain ( <i>Plantago lanceolata</i> ), hogweed ( <i>Heracleum sphondylium</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), white clover ( <i>Trifolium repens</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), pedunculate oak ( <i>Quercus robur</i> ), buddleia ( <i>Buddleja</i> sp.), silver birch ( <i>Betula pendula</i> ), evergreen oak ( <i>Quercus ilex</i> )
AB	H_A_829	g3c - Other neutral grassland	-	-	Grassland adjacent to horse-grazed fields that doesn't appear to be grazed at any time of the year. Sward height more varied.	Red dead-nettle ( <i>Lamium purpureum</i> ), yarrow ( <i>Achillea millefolium</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), dandelion ( <i>Taraxacum</i> agg.), common ragwort ( <i>Senecio jacobaea</i> ), bristly oxtongue ( <i>Picris echioides</i> ), dock ( <i>Rumex</i> sp.), ribwort plantain ( <i>Plantago lanceolata</i> ), hogweed ( <i>Heracleum sphondylium</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), white clover ( <i>Trifolium repens</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), false brome ( <i>Brachypodium sylvaticum</i> )
AB	H_A_833	c1 - Arable and horticulture	-	-	Arable field boundaries, used for vehicle access.	-
AB	H_A_835	g4 - Modified grassland	-	-	Large, mowed field, perhaps grazed at times. Likely yearly cut.	Perennial rye-grass ( <i>Lolium perenne</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), daisy ( <i>Bellis perennis</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), cowslip ( <i>Primula veris</i> )
AB	H_A_836	g4 - Modified grassland	-	-	-	White dead-nettle ( <i>Lamium album</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), spear thistle ( <i>Cirsium vulgare</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), common knapweed ( <i>Centaurea nigra</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), wild lettuce ( <i>Lactuca</i> sp.)
AB	H_A_837	g3c - Other neutral grassland	Ruderal/ephemeral	-	Disturbed grassland edge along boundary fence.	Common nettle ( <i>Urtica dioica</i> ), dock ( <i>Rumex</i> sp.), cleavers ( <i>Galium aparine</i> ), cow parsley ( <i>Anthriscus sylvestris</i> )
AB	H_A_839	u1b - Developed land; sealed surface	-	-	Road	-
AB	H_A_841	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_842	g3c - Other neutral grassland	-	-	Wider grassland strip near to pylon. Overgrown underneath pylon with some small elder, periwinkle, nettle, cleavers, ragwort, spear thistle and cow parsley at base. Remainder grassland.	Elder ( <i>Sambucus nigra</i> ), periwinkle ( <i>Vinca</i> sp.), common nettle ( <i>Urtica dioica</i> ), spear thistle ( <i>Cirsium vulgare</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), common ragwort ( <i>Senecio jacobaea</i> ), bristly ox-tongue ( <i>Picris echioides</i> ), dandelion ( <i>Taraxacum</i> agg.), ribwort plantain ( <i>Plantago lanceolata</i> ), cock's-foot ( <i>Dactylis glomerata</i> )
AB	H_A_843	c1 - Arable and horticulture	-	-	Ploughed field.	-
AB	H_A_844	g4 - Modified grassland	Horse grazed	-	Horse grazed field - no access.	N/A - Not Surveyed
AB	H_A_846	c1 - Arable and horticulture	-	-	Ploughed field.	-
AB	H_A_848	c1 - Arable and horticulture	-	-	Arable crops	-
AB	H_A_850	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_853	g3c - Other neutral grassland	Scattered scrub and tall herb	-	Grassland at top of ditch bank and adjacent to road. Highly disturbed.	Common nettle ( <i>Urtica dioica</i> ), willowherb ( <i>Epilobium</i> sp.), bramble ( <i>Rubus fruticosus</i> ), dock ( <i>Rumex</i> sp.), lords and ladies ( <i>Arum maculatum</i> ), cleavers ( <i>Galium aparine</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), rough meadow-grass ( <i>Poa trivialis</i> )
AB	H_A_854	h3d - Bramble scrub	-	-	Small area of scrub where hedgerow once was. Bare ground.	Bramble ( <i>Rubus fruticosus</i> ) D, common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), cow parsley ( <i>Anthriscus sylvestris</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_856	h3d - Bramble scrub	-	-	Small parcel of dense bramble scrub at field corner. Dry ditch running through.	Bramble ( <i>Rubus fruticosus</i> ), rose ( <i>Rosa sp.</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> )
AB	H_A_857	c1 - Arable and horticulture	-	-	Arable field. Boundary used for vehicle access. More herbs nearer to dry ditch	Common nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), dog's mercury ( <i>Mercurialis perennis</i> )
AB	H_A_859	c1a - Arable field margins	-	HPI	-	Common nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), dandelion ( <i>Taraxacum agg.</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), cleavers ( <i>Galium aparine</i> ), curled dock ( <i>Rumex crispus</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), meadowsweet ( <i>Filipendula ulmaria</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), cowslip ( <i>Primula veris</i> )
AB	H_A_860	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_861	c1 - Arable and horticulture	-	-	-	Arable field and narrow margin
AB	H_A_864	c1a - Arable field margins	-	HPI	Heavily cut making grass identification difficult.	Red fescue ( <i>Festuca rubra</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), common bent ( <i>Agrostis capillaris</i> ), common couch grass ( <i>Elytrigia repens</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), curled dock ( <i>Rumex crispus</i> ), hogweed ( <i>Heracleum sphondylium</i> ), common nettle ( <i>Urtica dioica</i> ) LA, creeping cinquefoil ( <i>Potentilla reptans</i> ), cat's-ear ( <i>Hypochoeris radicata</i> ), burdock ( <i>Arctium sp.</i> ), creeping thistle ( <i>Cirsium arvense</i> ), willowherb ( <i>Epilobium sp.</i> ), daisy ( <i>Bellis perennis</i> ), common knapweed ( <i>Centaurea nigra</i> ), spear thistle ( <i>Cirsium vulgare</i> ), germander speedwell ( <i>Veronica chamaedrys</i> )
AB	H_A_866	r2a - Rivers (priority habitat)	-	HPI	Belstead Brook See Aquatic Baseline Report ( <a href="#">application document 6.3.7.3</a> )	-
AB	H_A_867	g3c - Other neutral grassland	-	-	Field margin not meeting definition as use as track not for biodiversity benefit. Part shaded by woodland edge. Mammals passing through.	Common knapweed ( <i>Centaurea nigra</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), dandelion ( <i>Taraxacum agg.</i> ), dock ( <i>Rumex sp.</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), bristly oxtongue ( <i>Picris echioides</i> ), common nettle ( <i>Urtica dioica</i> ), lords and ladies ( <i>Arum maculatum</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), primrose ( <i>Primula vulgaris</i> ), spear thistle ( <i>Cirsium vulgare</i> )
AB	H_A_868	c1a - Arable field margins	-	HPI	Relic hedgerow and scattered trees present. Not meeting hedge definition. Semi-mature to mature pedunculate oak ( <i>Quercus robur</i> ), blackthorn, dog rose, field maple and dog wood. Ditch separating arable field. Not likely yearly wet.	Pedunculate oak ( <i>Quercus robur</i> ), blackthorn ( <i>Prunus spinosa</i> ), dog-rose ( <i>Rosa canina</i> ), field maple ( <i>Acer campestre</i> ), dogwood ( <i>Cornus sanguinea</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), annual meadow-grass ( <i>Poa annua</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), creeping thistle ( <i>Cirsium arvense</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), common bent ( <i>Agrostis capillaris</i> ), shepherd's-purse ( <i>Capsella bursa-pastoris</i> ), white clover ( <i>Trifolium repens</i> ), common knapweed ( <i>Centaurea nigra</i> ), herb-Robert ( <i>Geranium robertianum</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), willowherb ( <i>Epilobium sp.</i> ), cowslip ( <i>Primula veris</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), curled dock ( <i>Rumex crispus</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), groundsel ( <i>Senecio vulgaris</i> ), hedge bedstraw ( <i>Galium mollugo</i> )
AB	H_A_869	c1a - Arable field margins	-	HPI	Grassland dominated by perennial rye-grass. Ditch containing cock's foot and false brome is central of margin area.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, Yorkshire fog ( <i>Holcus lanatus</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) O, cleavers ( <i>Galium aparine</i> ) LA, common nettle ( <i>Urtica dioica</i> ) LA, cowslip ( <i>Primula veris</i> ) R
AB	H_A_870	c1 - Arable and horticulture	-	-	Arable crop	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_871	h3h - Mixed scrub	-	-	Hawthorn scrub with no bramble in corner of arable field. Dense patch of bramble with young hawthorn in open areas.	Hawthorn ( <i>Crataegus monogyna</i> ), bramble ( <i>Rubus fruticosus</i> ), hogweed ( <i>Heracleum sphondylium</i> ) O, cowslip ( <i>Primula veris</i> ) F, common knapweed ( <i>Centaurea nigra</i> ) A, dandelion ( <i>Taraxacum</i> agg.) O, spear thistle ( <i>Cirsium vulgare</i> ) F, horsetail ( <i>Equisetum</i> sp.) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, dock ( <i>Rumex</i> sp.) O, cock's-foot ( <i>Dactylis glomerata</i> ) A, cinquefoil ( <i>Potentilla</i> sp.) F, willow ( <i>Salix</i> sp.) R
AB	H_A_882	w1d - Wet woodland	-	HPI	Wet woodland edge with open areas containing bulrush. Willow dominant including young and mature trees. Hawthorn and elder at higher elevations along the bank. Scattered bramble. Occasional mosses.	Willow ( <i>Salix</i> sp.), bulrush ( <i>Typha</i> sp.), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ), bramble ( <i>Rubus fruticosus</i> ), common nettle ( <i>Urtica dioica</i> ) A, horsetail ( <i>Equisetum</i> sp.) O, sedge ( <i>Carex</i> sp.) O, ground ivy ( <i>Glechoma hederacea</i> ) A, lords and ladies ( <i>Arum maculatum</i> ) O, dandelion ( <i>Taraxacum</i> agg.) O, ivy ( <i>Hedera helix</i> ssp. <i>helix</i> ) O, lesser celandine ( <i>Ranunculus ficaria</i> ) A, bramble ( <i>Rubus fruticosus</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) F, meadowsweet ( <i>Filipendula ulmaria</i> ) R, mint ( <i>Mentha</i> sp.) F
AB	H_A_883	c1 - Arable and horticulture	-	-	Arable field boundaries, used for vehicle access.	-
AB	H_A_884	c1 - Arable and horticulture	-	-	Arable field boundaries, used for vehicle access.	-
AB	H_A_885	w2c - Other coniferous woodland	-	-	Leyland Cypress woodland with small, shed present. Ground largely bare. Some cherry saplings and field maple on edge but dominated by leylandii.	Leyland cypress ( <i>Cupressus x leylandii</i> ), wild cherry ( <i>Prunus avium</i> ), field maple ( <i>Acer campestre</i> )
AB	H_A_887	u1b - Developed land; sealed surface	-	-	Road/driveway	-
AB	H_A_888	u1b - Developed land; sealed surface	-	-	Road	-
AB	H_A_889	g4 - Modified grassland	-	-	Approx. 9m wide strip of grassland running around edge of arable field. Very short sward height with clear signs of vehicle usage. Lots of bare ground where vehicles have tracked over.	Dandelion ( <i>Taraxacum</i> agg.) F, ribwort plantain ( <i>Plantago lanceolata</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) D, white clover ( <i>Trifolium repens</i> ) F, rough meadow-grass ( <i>Poa trivialis</i> ) O
AB	H_A_890	g3c - Other neutral grassland	Disused quarry	-	Deep excavation with exposed ground and gravel amongst vegetation. Short sward height. Perhaps previously used as a quarry. Abundance of moss at lowest point suggest periodic inundation. Abundant mosses.	Germander speedwell ( <i>Veronica chamaedrys</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, dandelion ( <i>Taraxacum</i> agg.) O, daisy ( <i>Bellis perennis</i> ) O, yarrow ( <i>Achillea millefolium</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ), bent grass ( <i>Agrostis</i> sp.) A, red dead-nettle ( <i>Lamium purpureum</i> ) O, mouse-ear-hawkweed ( <i>Pilosella officinarum</i> ) F
AB	H_A_891	g3c - Other neutral grassland	-	-	Grassland associated with dry ditch. Willow herb in ditch. Not managed unlike adjacent strip.	Cow parsley ( <i>Anthriscus sylvestris</i> ), white dead-nettle ( <i>Lamium album</i> ), common nettle ( <i>Urtica dioica</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), hogweed ( <i>Heracleum sphondylium</i> ), dock ( <i>Rumex</i> sp.), field mustard ( <i>Sinapis arvensis</i> ), dandelion ( <i>Taraxacum</i> agg.), Timothy ( <i>Phleum pratense</i> ), greater burdock ( <i>Arctium lappa</i> ), stitchwort ( <i>Stellaria</i> sp.)
AB	H_A_892	c1 - Arable and horticulture	-	-	Ploughed field. Including narrow field margin	-
AB	H_A_893	g4 - Modified grassland	-	-	Perennial rye-grass dominated grassland with short sward height.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, white clover ( <i>Trifolium repens</i> ) A, dandelion ( <i>Taraxacum</i> agg.) A, ribwort plantain ( <i>Plantago lanceolata</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, daisy ( <i>Bellis perennis</i> ) F, bristly oxtongue ( <i>Picris echioides</i> ) O, chickweed ( <i>Stellaria</i> sp.) A, common ragwort ( <i>Senecio jacobaea</i> ) R
AB	H_A_894	c1 - Arable and horticulture	-	-	Ploughed field. Inc narrow field margin	-
AB	H_A_895	c1 - Arable and horticulture	-	-	Arable field	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_896	g4 - Modified grassland	-	-	Approx. 9m wide strip of grassland running around edge of arable field. Very short sward height with clear signs of vehicle usage.	Dandelion ( <i>Taraxacum</i> agg.) F, ribwort plantain ( <i>Plantago lanceolata</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) D, white clover ( <i>Trifolium repens</i> ) F, rough meadow-grass ( <i>Poa trivialis</i> ) O, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) O.
AB	H_A_897	w1d - Wet woodland	-	HPI	Woodland dominated by <i>Salix</i> species, including <i>Salix caprea</i> . Alder is frequent also. Hazel at edge of wood on bank with elder. Areas of vehicle damage and evidence of clay pigeon shooting in woodland.	Willow ( <i>Salix</i> sp.) D, goat willow ( <i>Salix caprea</i> ) D, alder ( <i>Alnus glutinosa</i> ) F, hazel ( <i>Corylus avellana</i> ), elder ( <i>Sambucus nigra</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) A, common nettle ( <i>Urtica dioica</i> ) A, lords and ladies ( <i>Arum maculatum</i> ) F, horsetail ( <i>Equisetum</i> sp.) O, opposite-leaved golden-saxifrage ( <i>Chrysosplenium oppositifolium</i> ) O, mint ( <i>Mentha</i> sp.) O, cleavers ( <i>Galium aparine</i> ) F, moschatel ( <i>Adoxa moschatellina</i> ) O, lesser celandine ( <i>Ranunculus ficaria</i> ) R, sedge ( <i>Carex</i> sp.)
AB	H_A_898	h3h - Mixed scrub	-	-	Hawthorn, blackthorn and rose scrub with some young pedunculate oak ( <i>Quercus robur</i> ) and larger cypress trees.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), rose ( <i>Rosa</i> sp.), pedunculate oak ( <i>Quercus robur</i> ), cypress ( <i>Cupressus</i> sp.)
AB	H_A_899	h3d - Bramble scrub	Scattered trees. Scattered rushes.	-	Scattered willow and rushes within scrub. Wetter towards west near woodland and drier in east where rush presence reduced. Ground flora very similar to that of adjacent woodland.	Willow ( <i>Salix</i> sp.) F, rush ( <i>Juncus</i> sp.), bramble ( <i>Rubus fruticosus</i> ) D, lesser celandine ( <i>Ranunculus ficaria</i> ) A, ground ivy ( <i>Glechoma hederacea</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) F, common nettle ( <i>Urtica dioica</i> ) F, dock ( <i>Rumex</i> sp.) F, sedge ( <i>Carex</i> sp.) F, horsetail ( <i>Equisetum</i> sp.) F
AB	H_A_900	c1 - Arable and horticulture	-	-	Ploughed field.	-
AB	H_A_901	g4 - Modified grassland	-	-	Perennial rye-grass dominated grassland with short sward height.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, white clover ( <i>Trifolium repens</i> ) A, dandelion ( <i>Taraxacum</i> agg.) A, ribwort plantain ( <i>Plantago lanceolata</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, daisy ( <i>Bellis perennis</i> ) F, bristly oxtongue ( <i>Picris echioides</i> ) O, chickweed ( <i>Stellaria</i> sp.) A, yarrow ( <i>Achillea millefolium</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) F, dock ( <i>Rumex</i> sp.) O, bramble ( <i>Rubus fruticosus</i> ) R, horsetail ( <i>Equisetum</i> sp.) R
AB	H_A_902	g3c - Other neutral grassland	-	-	Small grassland patch adjacent to arable field.	Yorkshire fog ( <i>Holcus lanatus</i> ) A, perennial rye-grass ( <i>Lolium perenne</i> ) F, dandelion ( <i>Taraxacum</i> agg.) F, red dead-nettle ( <i>Lamium purpureum</i> ) O, bristly oxtongue ( <i>Picris echioides</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, white dead-nettle ( <i>Lamium album</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) F, red campion ( <i>Silene dioica</i> ) R, cleavers ( <i>Galium aparine</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) A, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) F.
AB	H_A_904	g4 - Modified grassland	-	-	Grassland with low sward height, accessed by vehicles. Areas previously covered with scrub have been cleared.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, ground ivy ( <i>Glechoma hederacea</i> ) A, dandelion ( <i>Taraxacum</i> agg.) A, daisy ( <i>Bellis perennis</i> ) A, hogweed ( <i>Heracleum sphondylium</i> ) F, thistle ( <i>Cirsium</i> sp.) F, ribwort plantain ( <i>Plantago lanceolata</i> ) A
AB	H_A_905	g4 - Modified grassland	-	-	Approx. 9m wide strip of grassland running around edge of arable field. Very short sward height with clear signs of vehicle usage.	Dandelion ( <i>Taraxacum</i> agg.) F, ribwort plantain ( <i>Plantago lanceolata</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) D, white clover ( <i>Trifolium repens</i> ) F, rough meadow-grass ( <i>Poa trivialis</i> ) O, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) O.
AB	H_A_906	g3c - Other neutral grassland	Tall herb	-	Unmanaged grassland border, adjacent to woodland.	Common nettle ( <i>Urtica dioica</i> ) D, garlic mustard ( <i>Alliaria petiolata</i> ) O, cleavers ( <i>Galium aparine</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) A, dandelion ( <i>Taraxacum</i> agg.) O, cow parsley ( <i>Anthriscus sylvestris</i> ) F, bramble ( <i>Rubus fruticosus</i> ) O, white dead-nettle ( <i>Lamium album</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) O, daffodil ( <i>Narcissus</i> sp.) R, bluebell ( <i>Hyacinthoides non-scripta</i> ) R, crane's-bill ( <i>Geranium</i> sp.) F, dock ( <i>Rumex</i> sp.) F,

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_907	g3c - Other neutral grassland	Scattered trees.	-	Tussocks Grassland present on south facing slope. Scattered hawthorn and field maple. No signs of regular management, grazing or vehicle access.	Cock's-foot ( <i>Dactylis glomerata</i> ) A, meadow foxtail ( <i>Alopecurus pratensis</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) F, common vetch ( <i>Vicia sativa</i> ) F, red dead-nettle ( <i>Lamium purpureum</i> ) O, common nettle ( <i>Urtica dioica</i> ) O, yarrow ( <i>Achillea millefolium</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, white dead-nettle ( <i>Lamium album</i> ) O, dock ( <i>Rumex</i> sp.) O, meadow-grass ( <i>Poa</i> sp.) A, cinquefoil ( <i>Potentilla</i> sp.) F, hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> )
AB	H_A_915	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_916	g4 - Modified grassland	-	-	Does not reach plant threshold per m <sup>2</sup> , potentially area of left arable.	Creeping buttercup ( <i>Ranunculus repens</i> ) A, perennial rye-grass ( <i>Lolium perenne</i> ) D, red dead-nettle ( <i>Lamium purpureum</i> ) F, white dead-nettle ( <i>Lamium album</i> ) O, chickweed ( <i>Stellaria</i> sp.) O, dock ( <i>Rumex</i> sp.) R, spear thistle ( <i>Cirsium vulgare</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, common bent ( <i>Agrostis capillaris</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ) O, white clover ( <i>Trifolium repens</i> ) O
AB	H_A_920	c1a - Arable field margins	-	HPI	False brome by woodland margin.	Yorkshire fog ( <i>Holcus lanatus</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), common couch grass ( <i>Elytrigia repens</i> ), red fescue ( <i>Festuca rubra</i> ), common knapweed ( <i>Centaurea nigra</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), creeping thistle ( <i>Cirsium arvense</i> )
AB	H_A_921	g4 - Modified grassland	-	-	Large, mowed field, perhaps grazed at times.	Perennial rye-grass ( <i>Lolium perenne</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), daisy ( <i>Bellis perennis</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), cowslip ( <i>Primula vulgaris</i> )
AB	H_A_923	g4 - Modified grassland	Active management. Track.	-	Grassland track leading along woodland and adjacent areas. Short sward with signs of extensive use by heavy machinery. Large areas of bare ground with some herbaceous species. Areas of completely cleared bramble scrub and brash piles burnt. Small areas of hay storage too.	Common nettle ( <i>Urtica dioica</i> ), daisy ( <i>Bellis perennis</i> ), lords and ladies ( <i>Arum maculatum</i> ), cleavers ( <i>Galium aparine</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), dock ( <i>Rumex</i> sp.), meadow-grass ( <i>Poa</i> sp.), white clover ( <i>Trifolium repens</i> ), spear thistle ( <i>Cirsium vulgare</i> )
AB	H_A_925	g4 - Modified grassland	-	-	Cattle or horse grazed grassland with low sward height.	Cock's-foot ( <i>Dactylis glomerata</i> ) D, crested dog's-tail ( <i>Cynosurus cristatus</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) A, cowslip ( <i>Primula veris</i> ) F, daisy ( <i>Bellis perennis</i> ) O, meadow-grass ( <i>Poa</i> sp.) F, primrose ( <i>Primula vulgaris</i> ) O, dandelion ( <i>Taraxacum</i> agg.) O, creeping buttercup ( <i>Ranunculus repens</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) O, meadow foxtail ( <i>Alopecurus pratensis</i> ) O
AB	H_A_926	c1 - Arable and horticulture	-	-	Arable fields ploughed and sown. Includes margins used for vehicle access.	-
AB	H_A_928	c1 - Arable and horticulture	-	-	Arable fields ploughed and sown. Includes margins used for vehicle access.	-
AB	H_A_929	c1 - Arable and horticulture	-	-	Arable	-
AB	H_A_934	g4 - Modified grassland	-	-	Field possibly used for grazing or in rotation with others.	Cock's-foot ( <i>Dactylis glomerata</i> ), crested dog's-tail ( <i>Cynosurus cristatus</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), dandelion ( <i>Taraxacum</i> agg.), meadow-grass ( <i>Poa</i> sp.), St John's-wort ( <i>Hypericum</i> sp.)

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H_A_935	w1f7 - Other Lowland mixed deciduous woodland	-	-	Belt of Woodland which has grown around a central track PRoW. Ditches on either side separating from arable. Potentially old hedgerow once present either side but this is outgrown to extent of no return to stock proof and has a width on either side better classified as woodland than line of trees. Fully mature pedunculate oak ( <i>Quercus robur</i> ) and field maple trees on either side with some evident if historic coppicing. Semi-mature elm present. Ground flora has ancient woodland indicators but high proportion of ruderal and species potentially indicative of nutrient enrichment.	Pedunculate oak ( <i>Quercus robur</i> ), field maple ( <i>Acer campestre</i> ), hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), holly ( <i>Ilex aquifolium</i> ), elder ( <i>Sambucus nigra</i> ), elm ( <i>Ulmus</i> sp.), dog-rose ( <i>Rosa canina</i> ), yew ( <i>Taxus baccata</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) A, primrose ( <i>Primula vulgaris</i> ) O, common dog-violet ( <i>Viola riviniana</i> ) R, red campion ( <i>Silene dioica</i> ) R, herb-Robert ( <i>Geranium robertianum</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) A, cleavers ( <i>Galium aparine</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) F, broad-leaved dock ( <i>Rumex obtusifolius</i> ), curled dock ( <i>Rumex crispus</i> ), bramble ( <i>Rubus fruticosus</i> ) LA, wood-sedge ( <i>Carex sylvatica</i> ) O, wood avens ( <i>Geum urbanum</i> ) O, wild strawberry ( <i>Fragaria vesca</i> ) R, Goldilocks buttercup ( <i>Ranunculus auricomus</i> ) LO, early purple orchid ( <i>Orchis mascula</i> ).
AB	H_A_936	c1 - Arable and horticulture	-	-	Arable fields including margins used for vehicle access. Margins are narrow with frequent cowslip.	-
AB	H_A_937	c1 - Arable and horticulture	-	-	Confirmed aerial assessment	-
AB	H_A_938	g4 - Modified grassland	Cattle grazed	-	Not accessed because of cattle but assume same as adjacent grassland. Cattle grazed grassland with low sward height.	Cock's-foot ( <i>Dactylis glomerata</i> ) D, crested dog's-tail ( <i>Cynosurus cristatus</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) A, cowslip ( <i>Primula veris</i> ) F, daisy ( <i>Bellis perennis</i> ) O, meadow-grass ( <i>Poa</i> sp.) F, primrose ( <i>Primula vulgaris</i> ) O, dandelion ( <i>Taraxacum</i> agg) O, Creeping buttercup ( <i>Ranunculus repens</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) O, meadow foxtail ( <i>Alopecurus pratensis</i> ) O
AB	H_A_939	g4 - Modified grassland	Active management. Track.	-	Grassland track leading along woodland and field edge. Short sward with signs of extensive use by heavy machinery. Large areas of bare ground with some herbaceous species.	Common nettle ( <i>Urtica dioica</i> ), daisy ( <i>Bellis perennis</i> ), lords and ladies ( <i>Arum maculatum</i> ), cleavers ( <i>Galium aparine</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), dock ( <i>Rumex</i> sp.), meadow-grass ( <i>Poa</i> sp.), white clover ( <i>Trifolium repens</i> ), spear thistle ( <i>Cirsium vulgare</i> )
AB	H_A_940	c1 - Arable and horticulture	-	-	Arable crops	-
AB	H_A_809	f2 - Fen marsh and swamp	Rushes	HPI	-	Dominated by <i>Typha</i> sp. think is <i>Typha domagensis</i> . Scattered willow seedlings ( <i>Salix</i> sp.). Ground flora - creeping buttercup ( <i>Ranunculus repens</i> ), spear thistle ( <i>Cirsium vulgare</i> ), sphagnum sp ( <i>Sphagnum</i> spp.), marsh marigold R ( <i>Caltha palustris</i> ), willowherb O ( <i>Epilobium</i> sp.), dock sp. ( <i>Rumex</i> sp.).
AB	H_A_944	w1d - Wet woodland	-	HPI	Small area of young seedling grey willow and alder, including some semi-mature alder.	Grey willow ( <i>Salix cinerea</i> ) D, alder ( <i>Alnus glutinosa</i> ) F, sphagnum moss ( <i>Sphagnum</i> spp.) A, spear thistle ( <i>Cirsium vulgare</i> ), water mint ( <i>Mentha aquatica</i> ) O, meadowsweet ( <i>Filipendula ulmaria</i> ) R, willowherb ( <i>Epilobium</i> sp.) R, Yorkshire fog ( <i>Holcus lanatus</i> ) F, soft-rush ( <i>Juncus effusus</i> ) F, common nettle ( <i>Urtica dioica</i> ) O
AB	H3_1005	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1006	c1c - Cereal Crops	-	-	Arable crop	-
AB	H3_1012	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1013	w1 - Broadleaved mixed and yew woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	H3_1027	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1051	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1052	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1053	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1055	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1066	g4 – Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1100	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1104	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1105	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1107	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1108	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_1111	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_972	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_973	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_974	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_976	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_977	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_994	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	H3_995	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	HA3_460	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	HA3_839	h - Heathland and shrub	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	HA3_897	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	HA3_938	w1g - Other woodland; broadleaved	-		Plantation of white poplar	White poplar ( <i>Populus alba</i> )
AB	HA3_949	w1d - Wet woodland	-	HPI	-	Pedunculate oak ( <i>Quercus robur</i> ), goat willow ( <i>Salix caprea</i> ) D, hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) F, elder ( <i>Sambucus nigra</i> ) F, common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), red campion ( <i>Silene dioica</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), green alkanet ( <i>Pentaglottis sempervirens</i> ), sow-thistle ( <i>Sonchus</i> sp.)
AB	HA3_952	g - Grassland	-	-	Perennial rye-grass dominated grassland with short sward height.	White clover ( <i>Trifolium repens</i> ) A, dandelion ( <i>Taraxacum</i> agg.), ribwort plantain ( <i>Plantago lanceolata</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, daisy ( <i>Bellis perennis</i> ) F, bristly ox-tongue ( <i>Picris echioides</i> ) O, chickweed ( <i>Stellaria</i> sp.) A, common ragwort ( <i>Senecio jacobaea</i> ) R
AB	HL_252	g4 - Modified grassland	-	-	Sheep grazed probably reseeded recently. Dominated by grasses, lack of herbaceous plants. Surveyed from gate therefore low species count due to lack of access.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, cock's-foot ( <i>Dactylis glomerata</i> ) A, Timothy ( <i>Phleum pratense</i> ) O, false oat-grass ( <i>Arrhenatherum elatius</i> ) O, red fescue ( <i>Festuca rubra</i> ) R
AB	HL_258	g4 - Modified grassland	-	-	Amenity grass track	-
AB	HL_262	w1f - Lowland mixed deciduous woodland	-	HPI	-	Elder ( <i>Sambucus nigra</i> ), alder ( <i>Alnus glutinosa</i> ), blackthorn ( <i>Prunus spinosa</i> ), willow ( <i>Salix</i> sp.), ash ( <i>Fraxinus excelsior</i> ), field maple ( <i>Acer campestre</i> ), honeysuckle ( <i>Lonicera periclymenum</i> ), common nettle ( <i>Urtica dioica</i> ), ground ivy ( <i>Glechoma hederacea</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), hedge woundwort ( <i>Stachys sylvatica</i> ), old man's beard ( <i>Clematis vitalba</i> ), cleavers ( <i>Galium aparine</i> ), bramble ( <i>Rubus fruticosus</i> ), pedunculate oak ( <i>Quercus robur</i> ), hazel ( <i>Corylus avellana</i> ), rose ( <i>Rosa</i> sp.), wood avens ( <i>Geum urbanum</i> ), creeping thistle ( <i>Cirsium arvense</i> ), hogweed ( <i>Heracleum sphondylium</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), false brome ( <i>Brachypodium sylvaticum</i> ),
AB	HL_263	c1c5 - Winter stubble	-	-	Arable	-
AB	HL_264	g4 - Modified grassland	-	-	Species poor, good reptile habitat.	Broad-leaved dock ( <i>Rumex obtusifolius</i> ), prickly sow thistle ( <i>Sonchus asper</i> ), meadow foxtail ( <i>Alopecurus pratensis</i> ), bristly ox-tongue ( <i>Picris echioides</i> ), thistle ( <i>Cirsium</i> sp.),
AB	HL_265	g3c - Other neutral grassland	-	-	Field boundary of ruderal species. Poplar saplings.	Common nettle ( <i>Urtica dioica</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), spear thistle ( <i>Cirsium vulgare</i> ), curled dock ( <i>Rumex crispus</i> ), poplar ( <i>Populus</i> sp.)
AB	HL_269	c1 - Arable and horticulture	-	-	Arable	-
AB	HL_270	c1c6 - Game bird mix fields	Ruderal / Ephemeral	-	Possibly semi-managed due to prevalence of ruderal species.	Sunflower ( <i>Helianthus annua</i> ), millet ( <i>Panicum miliaceum</i> ), creeping thistle ( <i>Cirsium arvense</i> ), burdock ( <i>Arctium</i> sp.)
AB	HL_272	g4 - Modified grassland	-	-	Amenity grass short mown track, approx. 10m wide.	White clover ( <i>Trifolium repens</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), dock ( <i>Rumex</i> sp.)
AB	HL_273	c1d - Non-cereal crops	-	-	Non-cereal crops - brassica/buckwheat mix.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	HL_274	w1g - Other woodland; broadleaved	-	-	-	White poplar ( <i>Populus alba</i> ), ash ( <i>Fraxinus excelsior</i> ), burdock ( <i>Arctium sp.</i> ), elder ( <i>Sambucus nigra</i> ), blackthorn ( <i>Prunus spinosa</i> ), common nettle ( <i>Urtica dioica</i> ), ground ivy ( <i>Glechoma hederacea</i> ), spear thistle ( <i>Cirsium vulgare</i> ), alder ( <i>Alnus glutinosa</i> ), old man's beard ( <i>Clematis vitalba</i> ), bramble ( <i>Rubus fruticosus</i> ), field maple ( <i>Acer campestre</i> ), hawthorn ( <i>Crataegus monogyna</i> ), rose ( <i>Rosa sp.</i> )
AB	HL_276	g3c - Other neutral grassland	-	-	Review habitat type. Field boundary of ruderal species and some ash saplings. Cock's-foot, horsetail, and cow parsley most frequent.	Common nettle ( <i>Urtica dioica</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), spear thistle ( <i>Cirsium vulgare</i> ), common knapweed ( <i>Centaurea nigra</i> ), ground ivy ( <i>Glechoma hederacea</i> ), hogweed ( <i>Heracleum sphondylium</i> ), ash ( <i>Fraxinus excelsior</i> ), bristly oxtongue ( <i>Picris echioides</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), annual meadow grass ( <i>Poa annua</i> ), horsetail ( <i>Equisetum sp.</i> ), narrow-leaved ragwort ( <i>Senecio inaequidens</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), bramble ( <i>Rubus fruticosus</i> ), agrimony ( <i>Agrimonia eupatoria</i> )
AB	HL_278	g3c - Other neutral grassland	-	-	Field boundary of ruderal species, sloping to stream.	Common nettle ( <i>Urtica dioica</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), spear thistle ( <i>Cirsium vulgare</i> ), common knapweed ( <i>Centaurea nigra</i> ), ground ivy ( <i>Glechoma hederacea</i> ), hogweed ( <i>Heracleum sphondylium</i> ), curled dock ( <i>Rumex crispus</i> ), clematis ( <i>Clematis sp.</i> ), hedge bindweed ( <i>Calystegia sepium</i> ), cleavers ( <i>Galium aparine</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), smooth bedstraw ( <i>Cruciata laevipes</i> )
AB	HL_279	g4 - Modified grassland	Sheep grazed	-	Grazed grassland with sheep present.	Bristly oxtongue ( <i>Picris echioides</i> ), white clover ( <i>Trifolium repens</i> ), creeping thistle ( <i>Cirsium arvense</i> ), perennial rye-grass ( <i>Lolium perenne</i> )
AB	HL_282	g3c - Other neutral grassland	Ruderal and ephemeral land.	-	Ruderal and ephemeral land	Common nettle ( <i>Urtica dioica</i> ), thistle ( <i>Cirsium sp.</i> ), burdock ( <i>Arctium sp.</i> ), common couch grass ( <i>Elytrigia repens</i> ), common knapweed ( <i>Centaurea nigra</i> ), bristly oxtongue ( <i>Picris echioides</i> ), prickly sow thistle ( <i>Sonchus asper</i> ), narrow-leaved ragwort ( <i>Senecio inaequidens</i> )
AB	HL_340	g3c - Other neutral grassland	Scattered scrub. Ruderal and ephemeral land.	-	Grassy bank with tall ruderal vegetation and scattered scrub. Discarded materials on bank, including plastic piping, pallets, and containers.	Common nettle ( <i>Urtica dioica</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ), thistle ( <i>Cirsium sp.</i> ), pedunculate oak ( <i>Quercus robur</i> ), hazel ( <i>Corylus avellana</i> )
AB	HL_341	h3d - Bramble scrub	-	-	Bramble scrub along steep bank at gravel track edge. Largely dry ditch at base of steep bank. Southern edge has more scattered trees.	Horsetail ( <i>Equisetum sp.</i> ) F, common nettle ( <i>Urtica dioica</i> ) F, bindweed (Convolvulaceae sp.), F, elder ( <i>Sambucus nigra</i> ), blackthorn ( <i>Prunus spinosa</i> ), willow ( <i>Salix sp.</i> ), nightshade ( <i>Solanum sp.</i> ), willowherb ( <i>Epilobium sp.</i> ), sedges ( <i>Carex spp.</i> )
AB	HL_342	g4 - Modified grassland	-	-	Grass strip at edge of arable field approximately 1 metre wide. Recently mown. Not surveyed due to no access to arable land parcel.	-
AB	HL_343	u1c - Artificial unvegetated, unsealed surface	-	-	Gravel track and car park with areas used for storage of wood chip and logs. Log piles and wood chip support small mammals. Includes large spoil mound with some logs discarded on top. Lots of bare exposed ground with early colonising plants.	Common nettle ( <i>Urtica dioica</i> ), buddleia ( <i>Buddleja sp.</i> ), winter-cress ( <i>Barbarea vulgaris</i> ), thistle ( <i>Cirsium sp.</i> )
AB	HL_344	r1 - Standing open water and canals	Fishing lake	HPI	Little no aquatic vegetation, water lilies in tree shade. Used by birds such as moor hens and mallards.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	HL_345	g4 - Modified grassland	-	-	Mowed grassland path. Steep grassy bank alongside gravel track. Regularly used. Fungi species present.	Creeping buttercup ( <i>Ranunculus repens</i> ), daisy ( <i>Bellis perennis</i> ), creeping cinquefoil ( <i>Potentilla reptans</i> ), prickly sow thistle ( <i>Sonchus asper</i> ), bristly oxtongue ( <i>Picris echioides</i> ), smooth sow-thistle ( <i>Sonchus oleraceus</i> ), yarrow ( <i>Achillea millefolium</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), greater plantain ( <i>Plantago major</i> ), spotted medick ( <i>Medicago arabica</i> ), common couch grass ( <i>Elytrigia repens</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), bent ( <i>Agrostis</i> sp.)
AB	HL_346	u1c - Artificial unvegetated, unsealed surface	-	-	Gravel track used by cars	-
AB	HL_347	g4 - Modified grassland	-	-	Infrequently managed grassland patch, scattered white willows.	White willow ( <i>Salix alba</i> ), ground ivy ( <i>Glechoma hederacea</i> ), thistle ( <i>Cirsium</i> sp.), cinquefoil ( <i>Potentilla</i> sp.), broad-leaved dock ( <i>Rumex obtusifolius</i> ), dock ( <i>Rumex</i> sp.) creeping buttercup ( <i>Ranunculus repens</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), common ragwort ( <i>Senecio jacobaea</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), Narrow leaved dock species ( <i>Rumex</i> sp.)
AB	HL_348	g4 - Modified grassland	-	-	Mowed grass bank.	Ribwort plantain ( <i>Plantago lanceolata</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), dandelion ( <i>Taraxacum</i> agg.), bristly oxtongue ( <i>Picris echioides</i> )
AB	HL_349	g4 - Modified grassland	-	-	Mowed grassland path. Several clusters of different fungi species.	Ribwort plantain ( <i>Plantago lanceolata</i> ), Creeping buttercup ( <i>Ranunculus repens</i> ), daisy ( <i>Bellis perennis</i> ), creeping cinquefoil ( <i>Potentilla reptans</i> ), bristly oxtongue ( <i>Picris echioides</i> ), yarrow ( <i>Achillea millefolium</i> ), smooth sow-thistle ( <i>Sonchus oleraceus</i> ), prickly sow thistle ( <i>Sonchus asper</i> ), common ragwort ( <i>Senecio jacobaea</i> ), greater plantain ( <i>Plantago major</i> ), spotted medick ( <i>Medicago arabica</i> ), common couch grass ( <i>Elytrigia repens</i> ), bent ( <i>Agrostis</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ).
AB	W_1072	r2a – Rivers (priority habitat)		HPI	Belstead Brook. See Aquatic Baseline Report ( <a href="#">application document 6.3.7.3</a> )	-
AB	HL_351	c1 - Arable and horticulture	-	-	Arable field recently ploughed and sown, not surveyed due to lack of access.	-
AB	HL_353	h3a - Blackthorn scrub	-	-	Scattered blackthorn scrub alongside grazed field. Southern boundary comprises large overgrown spoil mound.	Common nettle ( <i>Urtica dioica</i> ), docks ( <i>Rumex</i> sp.), thistles ( <i>Cirsium</i> sp.)
AB	HL_354	u1b - Developed land; sealed surface	Residential	-	New build house, including driveway.	-
AB	HL_355	g3c - Other neutral grassland	-	-	Occasionally mown grassland adjacent to chicken coop and stables.	Cock's-foot ( <i>Dactylis glomerata</i> ), crested dog's-tail ( <i>Cynosurus cristatus</i> ), Timothy ( <i>Phleum pratense</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), dock ( <i>Rumex</i> sp.), clover ( <i>Trifolium</i> sp.), buttercup ( <i>Ranunculus</i> sp.)
AB	HL_356	h3h - Mixed scrub	-	-	Thin strip of scrubby vegetation at top of steep bank. Low level with gaps. Grades into grassland. Single ash specimen.	Hawthorn ( <i>Crataegus monogyna</i> ), rose ( <i>Rosa</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), bramble ( <i>Rubus fruticosus</i> ), ash ( <i>Fraxinus excelsior</i> ), old man's beard ( <i>Clematis vitalba</i> ), field maple ( <i>Acer campestre</i> ), dogwood ( <i>Cornus sanguinea</i> ), common nettle ( <i>Urtica dioica</i> )
AB	HL_357	c1 - Arable and horticulture	-	-	Recently ploughed arable field	-
AB	HL_358	u1b - Developed land; sealed surface	-	-	Road	-
AB	HL_359	u1b5 - Buildings	-	-	Collection of stables, sheds, and storage.	-
AB	HL_360	u1b - Developed land; sealed surface	-	-	Gravel track and parking locations.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
AB	HL_362	g4 - Modified grassland	Horse grazed	-	Low sward height, grazed by horses.	Perennial rye-grass ( <i>Lolium perenne</i> ), Italian rye-grass ( <i>Lolium multiflorum</i> ), clover ( <i>Trifolium</i> sp.), crested dog's-tail ( <i>Cynosurus cristatus</i> ), buttercup ( <i>Ranunculus</i> sp.), greater plantain ( <i>Plantago major</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), thistle ( <i>Cirsium</i> sp.), selfheal ( <i>Prunella vulgaris</i> ), geranium ( <i>Geranium</i> sp.), black medick ( <i>Medicago lupulina</i> )
AB	HL_363	g3c - Other neutral grassland	Horse grazed	-	Currently a not grazed horse paddock divided by electric fencing. Large pylon present within, with bramble scrub at base.	Cock's-foot ( <i>Dactylis glomerata</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), bent grass ( <i>Agrostis</i> sp.), buttercup ( <i>Ranunculus</i> sp.), clover ( <i>Trifolium</i> sp.), ribwort plantain ( <i>Plantago lanceolata</i> ), common ragwort ( <i>Senecio jacobaea</i> ), thistle ( <i>Cirsium</i> sp.), false oat grass ( <i>Arrhenatherum elatius</i> ), self-heal ( <i>Prunella vulgaris</i> ), spear thistle ( <i>Cirsium vulgare</i> ), ground ivy ( <i>Glechoma hederacea</i> ), hawkbit ( <i>Leontodon</i> sp.), sheep's fescue ( <i>Festuca</i> sp.), black medick ( <i>Medicago lupulina</i> ), common knotgrass ( <i>Polygonum aviculare</i> ), speedwell ( <i>Veronica</i> sp.)
AB	HL_364	g3c - Other neutral grassland	Ruderal and ephemeral land	-	Large spoil mound with some manure and hay dumped on top. common nettle, thistle and dock dominate.	Common nettle ( <i>Urtica dioica</i> ), thistle ( <i>Cirsium</i> sp.), dock ( <i>Rumex</i> sp.), hogweed ( <i>Heracleum sphondylium</i> )
AB	HL_383	g4 - Modified grassland	-	-	Arable field edge, regularly mown grassland. Approx. 2-3m wide. Low sward height.	Cock's-foot ( <i>Dactylis glomerata</i> ), dandelion ( <i>Taraxacum</i> agg.), meadow grass ( <i>Poa</i> sp.), yarrow ( <i>Achillea millefolium</i> )
AB	HL_384	u1d - Suburban/ mosaic of developed/ natural surface	Garden	-	Gardens, no access. Garden back onto dry ditch. Variety of garden types and levels of management. Continuous line of trees, scrub and other vegetation including some conifers and non-native ornamental species. Not hedgerow.	-
AB	HL_385	g4 - Modified grassland	-	-	Arable field edge, regularly mown grassland. Low sward height. Approx. 2-3m wide.	Dandelion ( <i>Taraxacum</i> agg.), meadow grass ( <i>Poa</i> sp.), yarrow ( <i>Achillea millefolium</i> ), cock's-foot ( <i>Dactylis glomerata</i> )
AB	HL_386	g3c - Other neutral grassland	Scattered trees	-	Grassland strip adjacent to arable and dry ditch. Scattered trees predominantly to southern side.	Common nettle ( <i>Urtica dioica</i> ) LA, common couch grass ( <i>Elytrigia repens</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), bramble ( <i>Rubus fruticosus</i> ), dock ( <i>Rumex</i> sp.), bindweed ( <i>Convolvulaceae</i> ), meadow grass ( <i>Poa</i> sp.), buttercup ( <i>Ranunculus</i> sp.), creeping cinquefoil ( <i>Potentilla reptans</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), hogweed ( <i>Heracleum sphondylium</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), wood avens ( <i>Geum urbanum</i> )
AB	HL_387	c1 - Arable and horticulture	-	-	Arable field	-
AB	HL_388	c1 - Arable and horticulture	-	-	Arable field	-
AB	HL_389	c1 - Arable and horticulture	-	-	Arable field	-
AB	HL_390	c1 - Arable and horticulture	-	-	Arable field	-
AB	HL_391	g4 - Modified grassland	-	-	Arable field edge, regularly mown grassland. Low sward height. Approx. 2-3m wide.	Dandelion ( <i>Taraxacum</i> agg.), meadow grass ( <i>Poa</i> sp.), yarrow ( <i>Achillea millefolium</i> ), cock's-foot ( <i>Dactylis glomerata</i> )
AB	HL_403	w1g - Other woodland; broadleaved	-	-	Pedunculate oak ( <i>Quercus robur</i> ) dominated broad leaved woodland. Evidence of historic coppicing. Ground flora similar to scrub with 25% bare earth	Pedunculate oak ( <i>Quercus robur</i> ) D, blackthorn ( <i>Prunus spinosa</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ), English elm ( <i>Ulmus procera</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) LA to south
AB	HL_427	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery	-
AB	HL_430	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
C	A_142	c1c - Cereal Crops	-	-	Arable	-
C	H_A_782	w1f7 - Other Lowland mixed deciduous woodland	-	-	-	Sycamore ( <i>Acer pseudoplatanus</i> ) F, sweet chestnut ( <i>Castanea sativa</i> ) A, beech ( <i>Fagus sylvatica</i> ) A, pine ( <i>Pinus</i> sp.) O, elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ), holly ( <i>Ilex aquifolium</i> ), common nettle ( <i>Urtica dioica</i> ) D, garlic mustard ( <i>Alliaria petiolata</i> ), red campion ( <i>Silene dioica</i> ), cleavers ( <i>Galium aparine</i> ), bramble ( <i>Rubus fruticosus</i> )
C	H_A_792	w1g - Other woodland; broadleaved	-	-	Small woodland strip on bank that is used extensively by badgers. Some large mature pedunculate oak ( <i>Quercus robur</i> ) with elder, hawthorn and blackthorn interspersed. Lots of leaf litter.	Pedunculate oak ( <i>Quercus robur</i> ), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), common nettle ( <i>Urtica dioica</i> ), lords and ladies ( <i>Arum maculatum</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), bramble ( <i>Rubus fruticosus</i> ), dog-rose ( <i>Rosa canina</i> ),
C	H_A_793	h3d - Bramble scrub	-	-	Strip of bramble scrub at base of bank. Other side of dog walking field fence.	Bramble ( <i>Rubus fruticosus</i> )
C	H_A_794	w1h - Other woodland; mixed	-	-	Woodland	Scot's pine ( <i>Pinus sylvestris</i> ), beech ( <i>Fagus sylvatica</i> ), sweet chestnut ( <i>Castanea sativa</i> ), pedunculate oak ( <i>Quercus robur</i> ), hawthorn ( <i>Crataegus monogyna</i> ) O, elder ( <i>Sambucus nigra</i> ) O, common nettle ( <i>Urtica dioica</i> )
C	H_A_823	g3c - Other neutral grassland	Ruderal/ephemeral	-	Area beyond dog walking facility and alongside river. Aquatic/bank side vegetation forms narrow strip that is described in H_A_831. Small patch of reeds nearer to river but still containing grassland species.	Common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), spear thistle ( <i>Cirsium vulgare</i> ), willowherb ( <i>Epilobium</i> sp.), wild teasel ( <i>Dipsacus fullonum</i> ), dock ( <i>Rumex</i> sp.), red dead-nettle ( <i>Lamium purpureum</i> )
C	H_A_831	r2a - Rivers (priority habitat)	HPI		River Brett.  Not condition assessed. Gentle meander through arable landscape. 3-4m wide, slow to moderate flow. Bank side vegetation include reeds and bulrushes, some trees also. Patchy along both sides however.	Reeds ( <i>Phragmites</i> sp.), bulrush ( <i>Typha</i> sp.)
					See Aquatic Baseline Report ( <a href="#">application document 6.3.7.3</a> )	
C	H_A_847	u1 - Built-up areas and gardens	Garden	-	Not accessed.	N/A - Not Surveyed
C	HA3_907	g- grassland		-	Assessed via aerial imagery.	
C	H_A_849	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
C	H_A_852	f - Wetland	-	-	Pond, sedge dominant with some rush species. Waterlogged but probably not permanent.	Sedge ( <i>Carex</i> sp.), bulrush ( <i>Typha</i> sp.)
C	H_A_914	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
C	H_A_922	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
C	H_A_924	g4 - Modified grassland	-	-	Modified grassland used as dog run facility. Fenced off. Grass dominant with lots of buttercup.	creeping buttercup ( <i>Ranunculus repens</i> ), red dead-nettle ( <i>Lamium purpureum</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
C	H_A_945	g3c - Other neutral grassland	Scattered rushes. Coastal and floodplain grazing marsh.	HPI (Secondary code)	Grassland adjacent to river. Slopes gradually down to the water becoming increasingly herb rich further down the slope. Areas of periodic inundation throughout with occasional juncus species.	Yorkshire fog ( <i>Holcus lanatus</i> ), meadow-grass ( <i>Poa</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ), dandelion ( <i>Taraxacum</i> agg.), daisy ( <i>Bellis perennis</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), white dead-nettle ( <i>Lamium album</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), common nettle ( <i>Urtica dioica</i> ), spear thistle ( <i>Cirsium vulgare</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), common ragwort ( <i>Senecio jacobaea</i> ), chickweed ( <i>Stellaria</i> sp.), dove's-foot crane's-bill ( <i>Geranium molle</i> )
C	H3_930	c1e - Intensive orchards	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
C	HA3_808	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
C	HA3_809	u1 - Built-up areas and gardens	-	-	Access tracks and sealed surfaces such as tarmac.	-
D	A_141	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	A_47	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	A_55	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	H_A_773	w1h - Other woodland; mixed	-	-	Seems to have once been a Scots pine plantation. Area for rearing game birds exists. Broadleaved species now make up at East 50% of woodland. Watercourse at W of woodland, slightly wetter however no significant change in species composition.	Scot's pine ( <i>Pinus sylvestris</i> ), ash ( <i>Fraxinus excelsior</i> ) O, beech ( <i>Fagus sylvatica</i> ) F, willow ( <i>Salix</i> sp.) R, hazel ( <i>Corylus avellana</i> ) O, elder ( <i>Sambucus nigra</i> ) O, common nettle ( <i>Urtica dioica</i> ) D, creeping buttercup ( <i>Ranunculus repens</i> ) A, red campion ( <i>Silene dioica</i> ) O, dock ( <i>Rumex</i> sp.) O, dog's mercury ( <i>Mercurialis perennis</i> ) O.
D	H_A_774	g4 - Modified grassland	-	-	Horse paddock, grazed grassland.	White clover ( <i>Trifolium repens</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) O, thistle ( <i>Cirsium</i> sp.) R, ribwort plantain ( <i>Plantago lanceolata</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) O.
D	H_A_778	g4 - Modified grassland	-	-	Horse paddock, grazed grassland.	White clover ( <i>Trifolium repens</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) O, thistle ( <i>Cirsium</i> sp.) R, ribwort plantain ( <i>Plantago lanceolata</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) O.
D	H_A_780	g4 - Modified grassland	-	-	Horse paddock, grazed grassland.	White clover ( <i>Trifolium repens</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) O, thistle ( <i>Cirsium</i> sp.) R, ribwort plantain ( <i>Plantago lanceolata</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) O.
D	H_A_781	w1g7 - Other broadleaved woodland types	-	-	Used for game bird rearing. Understorey contains sycamore saplings.	Beech ( <i>Fagus sylvatica</i> ) D, sweet chestnut ( <i>Castanea sativa</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) R, ash ( <i>Fraxinus excelsior</i> ) R, elder ( <i>Sambucus nigra</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), hawthorn ( <i>Crataegus monogyna</i> ), common nettle ( <i>Urtica dioica</i> )
D	H_A_784	w1f7 - Other Lowland mixed deciduous woodland	-	-	Woodland area with planted cricket bat willow, with wet ditch North to South, no flow, almost dry. Mature oak, lime in understorey. Pond sedge occasional where ditch is. No very mature trees.	Cricket bat willow ( <i>Salix alba</i> 'Caerulea'), alder ( <i>Alnus glutinosa</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) R, hazel ( <i>Corylus avellana</i> ) R, lime ( <i>Tilia</i> sp.) O, elder ( <i>Sambucus nigra</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, common nettle ( <i>Urtica dioica</i> ) D, red campion ( <i>Silene dioica</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) F, cleavers ( <i>Galium aparine</i> ) F, pond sedge ( <i>Carex</i> sp.) LO

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
D	W_1038	r2a – Rivers (priority habitat)	N/A	HPI	River Box See Aquatic Baseline Report ( <a href="#">application document 6.3.7.3</a> )	-
D	W_1043	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
D	W_1047	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
D	W_1048	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
D	W_1058	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
D	H_A_799	g4 - Modified grassland	-	-	-	Cock's-foot ( <i>Dactylis glomerata</i> ), bramble ( <i>Rubus fruticosus</i> ), dandelion ( <i>Taraxacum agg.</i> ), clover ( <i>Trifolium sp.</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), buttercup ( <i>Ranunculus sp.</i> )
D	H_A_801	c1c - Cereal crops	-	-	-	Arable.
D	H_A_812	c1d8 - Other non-cereal crops	-	-	Italian rye grass planted. No till.	-
D	H_A_855	c1a8 - Game bird mix strips and corners	-	-	Arable	-
D	H_A_872	g4 - Modified grassland	-	-	Heavily grazed patch of grassland bordered by arable field on one side. Currently used as paddock for horses. Edges of field dominated by common nettle. Surveyed from fence, unable to gain full access due to horses.	Common nettle ( <i>Urtica dioica</i> ), spear thistle ( <i>Cirsium vulgare</i> ), common ragwort ( <i>Senecio jacobaea</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), dandelion ( <i>Taraxacum agg.</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), bristly oxtongue ( <i>Picris echioides</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), white clover ( <i>Trifolium repens</i> ).
D	H_A_927	c1 - Arable and horticulture	-	-	Indian Buckwheat / cabbage. Young crop	-
D	H_A_953	c1c - Cereal crops	-	-	Arable field of wheat	-
D	H_A_813	g3c – Other neutral grassland	-	-	Area of unmown/ungrazed grassland bordered by arable fields.	Cock's-foot ( <i>Dactylus glomerata</i> ), dandelion ( <i>Taraxacum sp.</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), ragwort ( <i>Senecio jacobaea</i> ), bramble ( <i>Rubus fruticosus</i> ), dog rose ( <i>Rosa canina</i> ), primrose ( <i>Primula vulgaris</i> ), white clover ( <i>Trifolium repens</i> ), creeping thistle ( <i>Cirsium arvense</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), bristly oxtongue ( <i>Picris echioides</i> ), dove's-foot cranes-bill ( <i>Geranium molle</i> ), common vetch ( <i>Vicia sativa</i> ).
D	H3_1115	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	H3_920	c1c - Cereal crops	-	-	Arable	-
D	H3_923	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HA3_807	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HA3_810	u1b - Developed land; sealed surface	-	-	Hard standing	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
D	HA3_819	h3d - Bramble scrub	-	-	Dense bramble on edge of grassland and woodland, large part has been cut back.	Bramble ( <i>Rubus fruticosus</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), ground ivy ( <i>Glechoma hederacea</i> ), elder ( <i>Sambucus nigra</i> )
D	HA3_822	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HA3_824	w1 - Broadleaved mixed and yew woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	W_1046	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HL_151	c1d8 - Other non-cereal crops	-	-	Sugar beet, 2.5m strip surrounding field as buffer zone with arable weeds.	False oat grass ( <i>Arrhenatherum elatius</i> ), creeping thistle ( <i>Cirsium arvense</i> ) D, wild lettuce ( <i>Lactuca</i> sp.), common nettle ( <i>Urtica dioica</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), mayweed ( <i>Tripleurospermum inodorum</i> )
D	HL_247	g1c - Bracken	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HL_248	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	W_1045	r1 – standing open water and canals	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HL_249	w1g - Other woodland; broadleaved	-	-	Only surveyed corner of woodland as no agreed access to land parcel. Hazel coppice.	wood avens ( <i>Geum urbanum</i> ), wild cherry ( <i>Prunus avium</i> ), sedge ( <i>Carex</i> sp.), hedge woundwort ( <i>Stachys sylvatica</i> ), creeping thistle ( <i>Cirsium arvense</i> ), hazel ( <i>Corylus avellana</i> ), ash ( <i>Fraxinus excelsior</i> ), bramble ( <i>Rubus fruticosus</i> ), wood speedwell ( <i>Veronica montana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), elder ( <i>Sambucus nigra</i> ), field maple ( <i>Acer campestre</i> )
D	HL_251	g1c - Bracken	-	-	Bracken either side of dry ditch	Bracken ( <i>Pteridium aquilinum</i> ) D
D	HL_255	w1f - Lowland mixed deciduous woodland	-	HPI	Southern boundary including blackthorn and the other understory species. Bare ground/ leaf litter.	Pedunculate ( <i>Quercus robur</i> ), wild cherry ( <i>Prunus avium</i> ), willow ( <i>Salix</i> sp.), hazel ( <i>Corylus avellana</i> ), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ), dog-rose ( <i>Rosa canina</i> ), black bryony ( <i>Tamus communis</i> ), common nettle ( <i>Urtica dioica</i> ), wood speedwell ( <i>Veronica montana</i> ), bluebell ( <i>Hyacinthoides non-scripta</i> ), wood avens ( <i>Geum urbanum</i> ), cleavers ( <i>Galium aparine</i> ), common sorrel ( <i>Rumex acetosa</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), ivy ( <i>Hedera helix</i> ), ground ivy ( <i>Glechoma hederacea</i> ), wood dock ( <i>Rumex sanguineus</i> ), dock ( <i>Rumex</i> sp.), common feather-moss ( <i>Kindbergia praelonga</i> ), blackthorn ( <i>Prunus spinosa</i> ), snapping bonnet ( <i>Mycena vitilis</i> )
D	W_1044	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-
D	HL_281	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HL_378	c1a - Arable field margins	-	HPI	Thin grassland strip adjacent to g4. No obvious management or disturbance. Oak saplings. Low-distinctiveness grassland. Scattered bramble scrub.	Dock ( <i>Rumex</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), pedunculate oak ( <i>Quercus robur</i> ), bramble ( <i>Rubus fruticosus</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), crested dog's-tail ( <i>Cynosurus cristatus</i> ), bent grass ( <i>Agrostis</i> sp.), meadow-grass ( <i>Poa</i> sp.), common couch grass ( <i>Elytrigia repens</i> ), Timothy ( <i>Phleum pratense</i> ), old man's beard ( <i>Clematis vitalba</i> ), bramble ( <i>Rubus fruticosus</i> )
D	HL_423	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
D	HL_424	h3h - Mixed scrub	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	HL_60	h3h - Mixed scrub	-	-	Scrub on top of artificial mound.	Rose ( <i>Rosa</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), silver birch ( <i>Betula pendula</i> )
D	HL_62	h3h - Mixed scrub	-	-	Dense scrub dominated by similar species to scattered scrub nearby but abundant bramble. Limited gaps.	Bramble ( <i>Rubus fruticosus</i> ) A
D	HL_63	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Previous quarry activity. Successional species. Spoil piles with some industrial bricks and other artificial materials within spoil. Banks are steep with more scrub species. Grey willow saplings.	Wild teasel ( <i>Dipsacus fullonum</i> ) A, selfheal ( <i>Prunella vulgaris</i> ) O, creeping cinquefoil ( <i>Potentilla reptans</i> ) F, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) F, bristly ox-tongue ( <i>Picris echioides</i> ) F, bramble ( <i>Rubus fruticosus</i> ) R, common knapweed ( <i>Centaurea nigra</i> ) R, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) O, burdock ( <i>Arctium</i> sp.) R, common centaury ( <i>Centaurium erythraea</i> ) LA, blackthorn ( <i>Prunus spinosa</i> ) R, grey willow ( <i>Salix cinerea</i> ) R, gorse ( <i>Ulex europaeus</i> ) R, red-stemmed feather-moss ( <i>Pleurozium schreberi</i> ), perforate St John's-wort ( <i>Hypericum perforatum</i> ) O, hard-rush ( <i>Juncus inflexus</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, yellow pimpernel ( <i>Lysimachia nemorum</i> ) O, smooth tare ( <i>Vicia tetrasperma</i> ) R, Osier ( <i>Salix viminalis</i> ) R, goat willow ( <i>Salix caprea</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) R, carline thistle ( <i>Carlina vulgaris</i> ) R, great horsetail ( <i>Equisetum telmateia</i> ) LA
D	HL_64	w1g7 - Other broadleaved woodland types	Secondary woodland	-	Silver birch dominant. Two age ranges. Older approx. 20 years on slope as heading north with younger age range between slope and pathway. To eastern extent a mound with overgrown hawthorn line separating from habitat to east.	Common nettle ( <i>Urtica dioica</i> ) LA, ground ivy ( <i>Glechoma hederacea</i> ) F, bramble ( <i>Rubus fruticosus</i> ) R, cleavers ( <i>Galium aparine</i> ) R, wood forget-me-not ( <i>Myosotis sylvatica</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, common male fern ( <i>Dryopteris filix-mas</i> ) R, red campion ( <i>Silene dioica</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, creeping soft-grass ( <i>Holcus mollis</i> ) O
D	HL_65	h3 - Dense scrub	-	-	Dense scrub with abundant bramble. Limited gaps	Bramble ( <i>Rubus fructosus</i> ) A.
D	HL_66	r1b - Mesotrophic lakes	-	HPI	Bulrush frequently present on bankside	Pondweed ( <i>Potamogeton</i> sp.) O, bulrush ( <i>Typha</i> sp.) F, soft-rush ( <i>Juncus effusus</i> ) O, hard-rush ( <i>Juncus inflexus</i> ) O, water forget-me-not ( <i>Myosotis scorpioides</i> ) O, water mint ( <i>Mentha aquatica</i> ) O, great willowherb ( <i>Epilobium hirsutum</i> ) R, common club-rush ( <i>Schoenoplectus lacustris</i> ) R, water-cress ( <i>Rorippa nasturtium-aquaticum</i> ) R, clustered dock ( <i>Rumex conglomeratus</i> ) R, broad-leaved pondweed ( <i>Potamogeton natans</i> )
D	HL_67	w1f7 - Other Lowland mixed deciduous woodland	-	-	Very dense preventing access. Young trees with bramble and bracken covering most gaps. Old dry ditch to western aspect used by quarry with older mature trees of grey and white willow ( <i>Salix alba</i> ).	Bracken ( <i>Pteridium aquilinum</i> ), bramble ( <i>Rubus fruticosus</i> ), white willow ( <i>Salix alba</i> ), grey willow ( <i>Salix cinerea</i> )
D	HL_68	w1g - Other woodland; broadleaved	-	-	Likely boundary features which though lack of management have grown into woodland. Liverworts rare.	Wild cherry ( <i>Prunus avium</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) M, sycamore ( <i>Acer pseudoplatanus</i> ) M, hawthorn ( <i>Crataegus monogyna</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, hazel ( <i>Corylus avellana</i> ) C, common nettle ( <i>Urtica dioica</i> ) F, common male fern ( <i>Dryopteris filix-mas</i> ), red campion ( <i>Silene dioica</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) S, chickweed ( <i>Stellaria</i> sp.) R, field maple ( <i>Acer campestre</i> ), dogwood ( <i>Cornus sanguinea</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
D	HL_69	g3c - Other neutral grassland	-	-	-	Yorkshire fog ( <i>Holcus lanatus</i> ), creeping bent ( <i>Agrostis stolonifera</i> ), white clover ( <i>Trifolium repens</i> ), smooth tare ( <i>Vicia tetrasperma</i> ), common bent ( <i>Agrostis capillaris</i> ), bramble ( <i>Rubus fruticosus</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ), lesser stitchwort ( <i>Stellaria graminea</i> ), common centaury ( <i>Centaurium erythraea</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ), curled dock ( <i>Rumex crispus</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), common ragwort ( <i>Senecio jacobaea</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), selfheal ( <i>Prunella vulgaris</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), lady's bedstraw ( <i>Galium verum</i> ), field horsetail ( <i>Equisetum arvense</i> ).
D	W_1054	r1 - Standing open water and canals	-	HPI	Old large pond from quarry activities. Likely silt lake. Dry to western aspect and getting wetter to west, but lacking water. Not priority. Margin of Salix and silver birch up to 10m from pond but narrows. Dryer areas red feather moss.	Silver birch ( <i>Betula pendula</i> ), willows ( <i>Salix spp.</i> ), lesser bulrush ( <i>Typha angustifolia</i> ) A, bulrush ( <i>Typha latifolia</i> ) F, stonecrop ( <i>Crassula sp.</i> ) A, red-stemmed feather-moss ( <i>Pleurozium schreberi</i> ) O
D	HL_71	h3d - Bramble scrub	-	-	Very dense bramble.	Bramble ( <i>Rubus fruticosus</i> ) A
D	HL_72	h3d - Bramble scrub	-	-	-	Bramble ( <i>Rubus fruticosus</i> ) A, rose ( <i>Rosa sp.</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, gorse ( <i>Ulex europaeus</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) S
D	HL_73	g1c - Bracken	-	-	Bracken smothering all vegetation and scrub if present.	Bracken ( <i>Pteridium aquilinum</i> ) D
D	HL_74	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Areas of bare ground from quarry activity which is under succession. Lichens developing. Non-native panicle willowherb ( <i>Epilobium brachycarpum</i> ) present.	Common centaury ( <i>Centaurium erythraea</i> ) F, creeping cinquefoil ( <i>Potentilla reptans</i> ) F, perforate St John's-wort ( <i>Hypericum perforatum</i> ) F, selfheal ( <i>Prunella vulgaris</i> ) O, American willowherb ( <i>Epilobium ciliatum</i> ), panicle willowherb ( <i>Epilobium brachycarpum</i> ) F, creeping buttercup ( <i>Ranunculus repens</i> ) O, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) O, mouse-ear-hawkweed ( <i>Pilosella officinarum</i> ), dark mullein ( <i>Verbascum nigrum</i> ) R
D	HL_75	f2f - Other swamps	-	-	Dominated by bulrush	Bulrush ( <i>Typha sp.</i> ) D
D	HL_76	w1d - Wet woodland	-	-	Old area for quarry water which is under succession to woodland. Contains stream flowing to pond to west, standing water in centre where open glades dominated by aquatic species and horsetails. Bulrush locally abundant near stream. Abundant mosses in wetter areas. Young willow species dominant in woodland aspect.	Water forget-me-not ( <i>Myosotis scorpioides</i> ) O, marsh thistle ( <i>Cirsium palustre</i> ) F, water mint ( <i>Mentha aquatica</i> ) F, common club-rush ( <i>Schoenoplectus lacustris</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, bulrush ( <i>Typha sp.</i> ) LA, willow ( <i>Salix sp.</i> ) LD, common nettle ( <i>Urtica dioica</i> ) O, great horsetail ( <i>Equisetum telmateia</i> ) O, cleavers ( <i>Galium aparine</i> ) R, common feather-moss ( <i>Kindbergia praelonga</i> ) A
D	HL_77	g1a6 - Other lowland dry acid grassland	Rabbit grazed.	HPI	Patches of lichens.	Common sorrel ( <i>Rumex acetosa</i> ) (Rumex acetosa) F, heath speedwell ( <i>Veronica officinalis</i> ) O, heath bedstraw ( <i>Galium saxatile</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) R, yarrow ( <i>Achillea millefolium</i> ) O, lesser stitchwort ( <i>Stellaria graminea</i> ) R, common bent ( <i>Agrostis capillaris</i> ), red fescue ( <i>Festuca rubra</i> ), mouse-ear-hawkweed ( <i>Pilosella officinarum</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ), creeping bent ( <i>Agrostis stolonifera</i> ), common ragwort ( <i>Senecio jacobaea</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
D	HL_79	w1f - Lowland mixed deciduous woodland	-	HPI	Secondary woodland, on dryer banks and upper areas around old silt lake or quarry area. Two age classes within silver birch canopy.	Silver birch ( <i>Betula pendula</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) R, wood avens ( <i>Geum urbanum</i> ) R, bramble ( <i>Rubus fruticosus</i> ) A, germander speedwell ( <i>Veronica chamaedrys</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) O, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ), common centaury ( <i>Centaurium erythraea</i> ) R, wood forget-me-not ( <i>Myosotis sylvatica</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) O, bent grass ( <i>Agrostis</i> sp.) O, common nettle ( <i>Urtica dioica</i> ) LA, common feather-moss ( <i>Kindbergia praelonga</i> ) F, ( <i>Physcomitrium</i> sp.) moss A, scarlet pimpernel ( <i>Anagallis arvensis</i> ) R.
D	HL_80	u1a - Open Mosaic Habitats on Previously Developed Land	Scattered scrub	HPI	Areas of bare ground from quarry activity which is under succession. Grassland, includes areas of scattered scrub.	Willows ( <i>Salix</i> spp.), bramble ( <i>Rubus fruticosus</i> ), silver birch ( <i>Betula pendula</i> ), common ragwort ( <i>Senecio jacobaea</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) F, creeping cinquefoil ( <i>Potentilla reptans</i> ) F, wild teasel ( <i>Dipsacus fullonum</i> ), perforate St John's-wort ( <i>Hypericum perforatum</i> ) F, hard-rush ( <i>Juncus inflexus</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) O, American willowherb ( <i>Epilobium ciliatum</i> ), smooth hawk's-beard ( <i>Crepis capillaris</i> ) f, creeping buttercup ( <i>Ranunculus repens</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) O, hairy sedge ( <i>Carex hirta</i> ) LF, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) LA, dark mullein ( <i>Verbascum nigrum</i> ) R, great mullein ( <i>Verbascum thapsus</i> ) R, evening-primrose ( <i>Oenothera</i> sp.), willowherb ( <i>Epilobium</i> sp.) F, water figwort ( <i>Scrophularia auriculata</i> ) R, bristly haircap ( <i>Polytrichum piliferum</i> ) A, garden cress ( <i>Lepidium campestre</i> ) F
E	A_1017	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_1427	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_185	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_191	c1e - Intensive orchards	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_49	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_49	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_71	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_73	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	A_955	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
E	H_A_1040	g4 - Modified grassland	Cattle grazed	-	Cattle grazed grassland.	Ribwort plantain ( <i>Plantago lanceolata</i> ) O, dandelion ( <i>Taraxacum</i> agg.) R, cock's-foot ( <i>Dactylis glomerata</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) F, red fescue ( <i>Festuca rubra</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) F, common bent ( <i>Agrostis capillaris</i> ) F, cat's-ear ( <i>Hypochaeris radicata</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, common sorrel ( <i>Rumex acetosa</i> ) R, yarrow ( <i>Achillea millefolium</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, hairy sedge ( <i>Carex hirta</i> ) R, downy oat-grass ( <i>Avenula pubescens</i> ) O, meadow crane's-bill ( <i>Geranium pratense</i> ) O, white clover ( <i>Trifolium repens</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, meadow buttercup ( <i>Ranunculus acris</i> ) O, common knapweed ( <i>Centaurea nigra</i> ) R, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, dove's-foot crane's-bill ( <i>Geranium molle</i> ) R,
E	H_A_1050	c1e - Intensive orchards	-	-	Intensive commercial apple orchard	-
E	H_A_1051	c1e - Intensive orchards	-	-	Commercial apple orchards	-
E	H_A_798	w1f7 - Other Lowland mixed deciduous woodland	-	-	Woodland bordered by arable field to north and grassland with plantation trees to south. Woodland becomes increasingly scrubby towards west with blackthorn and hawthorn dominant.	Ground ivy ( <i>Glechoma hederacea</i> ), common nettle ( <i>Urtica dioica</i> ), lords and ladies ( <i>Arum maculatum</i> ), cleavers ( <i>Galium aparine</i> ), bluebell ( <i>Hyacinthoides non-scripta</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), lesser burdock ( <i>Arctium</i> sp.), dog's mercury ( <i>Mercurialis perennis</i> ), bugle ( <i>Ajuga reptans</i> ), ivy ( <i>Hedera helix</i> ), dandelion ( <i>Taraxacum</i> agg.), hazel ( <i>Corylus avellana</i> ), elder ( <i>Sambucus nigra</i> ), pedunculate oak ( <i>Quercus robur</i> ), field maple ( <i>Acer campestre</i> ), white willow ( <i>Salix alba</i> ), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), moschatel ( <i>Adoxa moschatellina</i> )
E	H_A_878	g3c - Other neutral grassland	Plantation	-	Grassland with white willow ( <i>Salix alba</i> ) plantation. Two ditches run through habitat creating wetter areas. Wetter areas wood sedge more common.	Broad-leaved dock ( <i>Rumex obtusifolius</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), dove's-foot crane's-bill ( <i>Geranium molle</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), meadowsweet ( <i>Filipendula ulmaria</i> ), wood-sedge ( <i>Carex sylvatica</i> ), common nettle ( <i>Urtica dioica</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), dandelion ( <i>Taraxacum</i> agg.), yellow iris ( <i>Iris pseudacorus</i> ), marsh marigold ( <i>Caltha palustris</i> ), lords and ladies ( <i>Arum maculatum</i> )
E	H_A_954	c1d - Non-cereal crops	-	-	Arable field of rapeseed. Access track/arable field margin approximately 7 metres across.	-
E	E_T_867	w - Woodland and forest	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	H_A_963	c1c - Cereal crops	-	-	Linseed	-
E	H_A_964	c1c - Cereal crops	-	-	Wheat, no hedge on eastern boundary by road	-
E	H_A_965	c1d - Non-cereal crops	-	-	Rape seed	-
E	H3_1135	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	H3_914	c1c - Cereal crops	-	-	Arable	-
E	H3_914	c1c - Cereal crops	-	-	Arable	-
E	H3_915	c1c - Cereal crops	-	-	Arable	-
E	H3_931	c1e - Intensive orchards	-	-	Arable	-
E	H3_932	w1f7 - Other Lowland mixed deciduous woodland	-	-	Arable	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
E	HA3_950	w1d - Wet woodland	-	HPI	Alder wet woodland, wet under foot with ditch running northern border of woodland. Southern edge of woodland is large brash piles.	Alder ( <i>Alnus glutinosa</i> ) (d), hybrid poplar ( <i>Populus x</i> ) (O), willow ( <i>Salix sp.</i> ) (O), pedunculate oak ( <i>Quercus robur</i> ) (r), wild cherry ( <i>Prunus avium</i> ) (o), elder ( <i>Sambucus nigra</i> ) (d), hazel ( <i>Corylus avellana</i> ) (o), hawthorn ( <i>Crataegus monogyna</i> ) (R), common nettle ( <i>Urtica dioica</i> ) (d), cleavers ( <i>Galium aparine</i> ) (A), brooklime ( <i>Veronica beccabunga</i> ) (o), crane's-bill ( <i>Geranium sp.</i> ) (r), herb-Robert ( <i>Geranium robertianum</i> ) (r), creeping buttercup ( <i>Ranunculus repens</i> ) (f), soft-rush ( <i>Juncus effusus</i> ) (R), yellow pimpernel ( <i>Lysimachia nemorum</i> ) (r).
E	HL_13	c1c - Cereal crops	-	-	Linseed	-
E	HL_222	f2f - Other swamps	Rushes dominant	-	Dense patches of wetland arising within pasture from occasional springs. Dominated by soft rush.	Soft-rush ( <i>Juncus effusus</i> ) D, common nettle ( <i>Urtica dioica</i> ) O, common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) R, wood dock ( <i>Rumex sanguineus</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) O, marsh thistle ( <i>Cirsium palustre</i> ) O, hard-rush ( <i>Juncus inflexus</i> ) R, hairy sedge ( <i>Carex hirta</i> ) R, great willowherb ( <i>Epilobium hirsutum</i> ) R, broad-leaved willowherb ( <i>Epilobium montanum</i> ) R
E	HL_223	g4 - Modified grassland	-	-	Cattle grazed grassland.	Ribwort plantain ( <i>Plantago lanceolata</i> ) O, dandelion ( <i>Taraxacum agg.</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) F, red fescue ( <i>Festuca rubra</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) F, common bent ( <i>Agrostis capillaris</i> ) F, cat's-ear ( <i>Hypochaeris radicata</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, common sorrel ( <i>Rumex acetosa</i> ) R, yarrow ( <i>Achillea millefolium</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, hairy sedge ( <i>Carex hirta</i> ) R, downy oat-grass ( <i>Avenula pubescens</i> ) O, meadow crane's-bill ( <i>Geranium pratense</i> ) O, white clover ( <i>Trifolium repens</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, meadow buttercup ( <i>Ranunculus acris</i> ) O, common knapweed ( <i>Centaurea nigra</i> ) R, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, dove's-foot crane's-bill ( <i>Geranium molle</i> ) R,
E	E_T_869	c1 – Arable and horticulture	-	-	UKHab category derived from aerial imagery/2012 Phase 1 survey.	-
E	HL_224	r1 – Standing open water and canals	-	HPI	This is not a woodland but a ring of trees. A shallow stream fed pond heavily shaded by young, pollarded alder ( <i>Alnus glutinosa</i> ) trees on the bank. No aquatic macrophytes evident	Alder ( <i>Alnus glutinosa</i> ) D
E	W_1037	r1 – Standing open water and canals	-	HPI	Pond with shallow standing water fed by at least two streams.	Bulrush ( <i>Typha sp.</i> ) D.
E	HL_245	c1e - Intensive orchards	-	-	Intensive commercial apple orchard	-
E	HL_246	c1e - Intensive orchards	-	-	Commercial apple orchards	-
E	HL_26	g1a6 - Other lowland dry acid grassland	-	HPI	-	Sheep's sorrel ( <i>Rumex acetosella</i> ) F to LA, sand spurrey ( <i>Spergularia rubra</i> ) O, parsley-piert ( <i>Aphanes arvensis</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) R, common bent ( <i>Agrostis capillaris</i> ) A, heath bedstraw ( <i>Galium saxatile</i> ) R, sticky mouse-ear ( <i>Cerastium glomeratum</i> ) R, shepherd's-purse ( <i>Capsella bursa-pastoris</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) O, lesser trefoil ( <i>Trifolium dubium</i> ) O, annual meadow-grass ( <i>Poa annua</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) F, ribwort plantain ( <i>Plantago lanceolata</i> ), cat's-ear ( <i>Hypochaeris radicata</i> ), squirreltail fescue ( <i>Vulpia bromoides</i> ) R, greater plantain ( <i>Plantago major</i> ), thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R
E	HL_27	h3e - Gorse scrub	-	-	Horse scrub, very dense	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
E	F_T_872	C1 – Arable and horticulture	-	-	UKHab category derived from aerial imagery/ 2012 Phase 1 survey.	-
E	HL_292	c1 - Arable and horticulture	-	-	Arable field recently ploughed.	-
E	HL_32	c1c - Cereal crops	-	-	Barley	-
E	HL_366	c1a - Arable field margins	-	HPI	Grassland track around arable field for vehicular access.	Ribwort plantain ( <i>Plantago lanceolata</i> ), dock ( <i>Rumex</i> sp.), perennial rye-grass ( <i>Lolium perenne</i> ), meadow-grass ( <i>Poa</i> sp.).
E	HL_367	c1a - Arable field margins	-	HPI	Grassland grading into unkept arable field. Large patches of nettle with encroaching bramble and old man's beard. Oak saplings.	Common nettle ( <i>Urtica dioica</i> ), bramble ( <i>Rubus fruticosus</i> ), old man's beard ( <i>Clematis vitalba</i> ), thistle ( <i>Cirsium</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ), common couch grass ( <i>Elytrigia repens</i> ) grass, bent grass ( <i>Agrostis</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ), pedunculate oak ( <i>Quercus robur</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), blackthorn ( <i>Prunus spinosa</i> ) sapling, scarlet pimpernel ( <i>Anagallis arvensis</i> )
E	HL_369	c1 - Arable and horticulture	-	-	Arable	-
E	HL_372	c1a - Arable field margins	-	HPI	Arable field boundaries often include farm tracks. Grassland track around arable field for vehicular access.	Ribwort plantain ( <i>Plantago lanceolata</i> ), dock ( <i>Rumex</i> sp.), perennial rye-grass ( <i>Lolium perenne</i> ), meadow-grass ( <i>Poa</i> sp.)
E	HL_374	c1a - Arable field margins	-	HPI	Grassland strip along hedgerow of slightly different composition than grassland on track. Scattered bracken present.	Bracken ( <i>Pteridium aquilinum</i> ), common couch grass ( <i>Elytrigia repens</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), Timothy ( <i>Phleum pratense</i> ), bent grass ( <i>Agrostis</i> sp.), false oat grass ( <i>Arrhenatherum elatius</i> ), dock ( <i>Rumex</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), common nettle ( <i>Urtica dioica</i> ), thistle ( <i>Cirsium</i> sp.)
E	HL_375	c1 - Arable and horticulture	-	-	Arable crops	-
E	HL_376	c1a - Arable field margins	-	HPI	Grassland strip adjacent to track. One small oak.	Cock's-foot ( <i>Dactylis glomerata</i> ), common couch grass ( <i>Elytrigia repens</i> ), Timothy ( <i>Phleum pratense</i> ), bent grass ( <i>Agrostis</i> sp.), ribwort plantain ( <i>Plantago lanceolata</i> ), meadow-grass ( <i>Poa</i> sp.), spear thistle ( <i>Cirsium vulgare</i> ), crested dog's-tail ( <i>Cynosurus cristatus</i> ), pedunculate oak ( <i>Quercus robur</i> ).
E	HL_377	c1a - Arable field margins	-	HPI	Grassland along edge of track. Small oak.	Thistle ( <i>Cirsium</i> sp.), bent grass ( <i>Agrostis</i> sp.), perennial rye-grass ( <i>Lolium perenne</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), common couch grass ( <i>Elytrigia repens</i> ), crested dog's-tail ( <i>Cynosurus cristatus</i> ), pedunculate oak ( <i>Quercus robur</i> )
E	HL_379	c1 - Arable and horticulture	-	-	Arable	-
E	HL_382	c1a - Arable field margins	-	HPI	Mowed grassland strip along arable field boundary. Used for vehicular access. Same as adjacent grassland but managed differently.	Dock ( <i>Rumex</i> sp.), broad-leaved plantain ( <i>Plantago major</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), thistle ( <i>Cirsium</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ), dandelion ( <i>Taraxacum</i> agg.), common nettle ( <i>Urtica dioica</i> ), rough meadow-grass ( <i>Poa trivialis</i> ).
E	HL_39	c1d - Non-cereal crops	-	-	Rape seed	-
E	HL_39	c1d - Non-cereal crops	-	-	Rape seed	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
E	HL_41	w1f7 - Other Lowland mixed deciduous woodland	-	-	Areas of leaf litter. Hedgerow separated from highway. Canopy sweet chestnut, cherry, ash, field maple. Understorey includes oak, English elm. Some standing and fallen deadwood but not much.	Cow parsley ( <i>Anthriscus sylvestris</i> ) A, common nettle ( <i>Urtica dioica</i> ) F, burdock ( <i>Arctium</i> sp.) O, garlic mustard ( <i>Alliaria petiolata</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) O to LA, ground ivy ( <i>Glechoma hederacea</i> ) O, greater stitchwort ( <i>Stellaria holostea</i> ) R, white dead-nettle ( <i>Lamium album</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) R, bluebell hybrid ( <i>Hyacinthoides</i> x) R, pignut ( <i>Conopodium majus</i> ) F, rough meadow-grass ( <i>Poa trivialis</i> ) LA, sweet chestnut ( <i>Castanea sativa</i> ) F, wild cherry ( <i>Prunus avium</i> ) R, ash ( <i>Fraxinus excelsior</i> ) Y, field maple ( <i>Acer campestre</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) R, English elm ( <i>Ulmus procera</i> ) f, blackthorn ( <i>Prunus spinosa</i> ) R, holly ( <i>Ilex aquifolium</i> ) R,
E	HL_42	c1d - Non-cereal crops	-	-	Possible potato	-
E	HL_437	w1d - Wet woodland	-	HPI	A wet woodland which has not been accessed internally due to no access, which may be w1d5.	-
E	HL_43a	w1f - Lowland mixed deciduous woodland	-	HPI	Deadwood including large fallen cherry and standing deadwood. Canopy oak, mature multi-stem Hornbeam, mature ash, mature wild cherry. Large areas of leaf litter. Topography is down a short but very steep hill from west to east	Wild cherry ( <i>Prunus avium</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) A, ash ( <i>Fraxinus excelsior</i> ) O, hornbeam ( <i>Carpinus betulus</i> ), hawthorn ( <i>Crataegus monogyna</i> ) R, holly ( <i>Ilex aquifolium</i> ) O, elder ( <i>Sambucus nigra</i> ) R, bluebell ( <i>Hyacinthoides non-scripta</i> ) LA, garlic mustard ( <i>Alliaria petiolata</i> ), common mouse-ear ( <i>Cerastium fontanum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), bracken ( <i>Pteridium aquilinum</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) LA, red campion ( <i>Silene dioica</i> ) R, yellow archangel ( <i>Lamiastrum galeobdolon</i> ) R.
E	HL_441	g - Grassland	-	-	Priority habitat floodplain grassland	-
E	HL_45	g3c - Other neutral grassland	-	-	Eastern end nettles and brambles and bracken to southern side along hedgerow.	Common nettle ( <i>Urtica dioica</i> ) A, bramble ( <i>Rubus fruticosus</i> ) LA, bracken ( <i>Pteridium aquilinum</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) F to LA, common knapweed ( <i>Centaurea nigra</i> ) LA, germander speedwell ( <i>Veronica chamaedrys</i> ) O, lesser stitchwort ( <i>Stellaria graminea</i> ) O, yarrow ( <i>Achillea millefolium</i> ) O, smooth meadow-grass ( <i>Poa pratensis</i> ) R, sticky mouse-ear ( <i>Cerastium glomeratum</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) O, cleavers ( <i>Galium aparine</i> ) O, red campion ( <i>Silene dioica</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) R, white dead-nettle ( <i>Lamium album</i> ) R, rough meadow-grass ( <i>Poa trivialis</i> ) O, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ), red fescue ( <i>Festuca rubra</i> )
F	A_205	c1c - Cereal Crops	-	-	Arable	-
F	A_236	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H_A_1004	g3c - Other neutral grassland	Horse grazed	-	Informed will have a hay cut in summer and horse grazed in winter. Patches have higher bare earth % maybe from over grazing. Ant hills present. Lacking more than 10/11 species per m2.	Autumn hawkbit ( <i>Leontodon autumnalis</i> ) O, marsh sow-thistle ( <i>Sonchus asper</i> ) R, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) R, greater plantain ( <i>Plantago major</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) A, soft brome ( <i>Bromus hordeaceus</i> ) O, selfheal ( <i>Prunella vulgaris</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) F, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, common bent ( <i>Agrostis capillaris</i> ) A, common mouse-ear ( <i>Cerastium fontanum</i> ) O, dandelion ( <i>Taraxacum</i> agg.) F, white clover ( <i>Trifolium repens</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) F, hedgerow crane's-bill ( <i>Geranium pyrenaicum</i> ) O, Timothy ( <i>Phleum pratense</i> ) O, daisy ( <i>Bellis perennis</i> ) O, bristly oxtongue ( <i>Picris echioides</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) R, smooth tare ( <i>Vicia tetrasperma</i> ) R, lesser trefoil ( <i>Trifolium dubium</i> ) O, greater bird's-foot-trefoil ( <i>Lotus pedunculatus</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) LA, red fescue ( <i>Festuca rubra</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) F, mouse-ear-hawkweed ( <i>Pilosella officinarum</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) O, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R, field madder ( <i>Sherardia arvensis</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	H_A_1005	g4 - Modified grassland	-	-	Less than 9 sp per m <sup>2</sup> . Short horse grazed pasture likely sown. Road through field of gravel with young trees planted.	Yorkshire fog ( <i>Holcus lanatus</i> ) A, perennial rye-grass ( <i>Lolium perenne</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) R, white clover ( <i>Trifolium repens</i> ) F, yarrow ( <i>Achillea millefolium</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, field maple ( <i>Acer campestre</i> ), beech ( <i>Fagus sylvatica</i> ), smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R.
F	H_A_1007	g4 - Modified grassland	Plantation, non-native, scattered trees	-	Native and non-native mix, less than 20 years in age. Turkey oak present.	Turkey oak ( <i>Quercus cerris</i> )
F	H_A_1038	c1c - Cereal crops	-	-	A ploughed wheat field with a 10-metre margin around the edge of the field	-
F	H_A_1043	c1c - Cereal crops	-	-	Field of winter wheat	-
F	H_A_1044a	w1d5 - Alder woodland on floodplains (H91E0)	-	Subset of HPI Annex 1 Habitat	The accessibility of the wood is extremely limited and was surveyed around the perimeter at two points where entry was possible. Therefore, accurate percentage cover of willow and alder is not possible. Some large, coppiced alder present with open areas and mixed deciduous trees and shrubs including some large ash. A small stream runs through the woodland. Pheasant rearing.	Common nettle ( <i>Urtica dioica</i> ) D, hogweed ( <i>Heracleum sphondylium</i> ) O, eagle fern ( <i>Pteridium aquilinum</i> ) A, elder ( <i>Sambucus nigra</i> ) R, willow ( <i>Salix</i> sp.) F, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, bulrush ( <i>Typha latifolia</i> ) R, wild angelica ( <i>Angelica sylvestris</i> ) O, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) O, ash ( <i>Fraxinus excelsior</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) R, woody nightshade ( <i>Solanum dulcamara</i> ) R, opposite-leaved golden-saxifrage ( <i>Chrysosplenium oppositifolium</i> ) R, fool's parsley ( <i>Aethusa cynapium</i> ) R, red campion ( <i>Silene dioica</i> ) R, curled dock ( <i>Rumex crispus</i> ) R, field maple ( <i>Acer campestre</i> ) O, alder ( <i>Alnus glutinosa</i> ) A,
F	H_A_1044b	w1d5 - Alder woodland on floodplains (H91E0)	-	Subset of HPI Annex 1 Habitat	The accessibility of the wood is extremely limited and was surveyed around the perimeter at two points where entry was possible. Therefore, accurate percentage cover of willow and alder is not possible. Some large, coppiced alder present with open areas and mixed deciduous trees and shrubs including some large ash. A small stream runs through the woodland. Pheasant rearing.	Common nettle ( <i>Urtica dioica</i> ) D, hogweed ( <i>Heracleum sphondylium</i> ) O, eagle fern ( <i>Pteridium aquilinum</i> ) A, elder ( <i>Sambucus nigra</i> ) R, willow ( <i>Salix</i> sp.) F, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, bulrush ( <i>Typha latifolia</i> ) R, wild angelica ( <i>Angelica sylvestris</i> ) O, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) O, ash ( <i>Fraxinus excelsior</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) R, woody nightshade ( <i>Solanum dulcamara</i> ) R, opposite-leaved golden-saxifrage ( <i>Chrysosplenium oppositifolium</i> ) R, fool's parsley ( <i>Aethusa cynapium</i> ) R, red campion ( <i>Silene dioica</i> ) R, curled dock ( <i>Rumex crispus</i> ) R, field maple ( <i>Acer campestre</i> ) O, alder ( <i>Alnus glutinosa</i> ) A,
F	W_1029	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ 2012 Phase 1 surveys.	-
F	H_A_1048	c1c - Cereal crops	-	-	Field of winter wheat	-
F	H_A_1062	g4 - Modified grassland	-	-	-	Oxeye daisy ( <i>Leucanthemum vulgare</i> ), common mallow ( <i>Malva sylvestris</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), St. John's wort ( <i>Hypericum</i> sp.), perennial rye-grass ( <i>Lolium perenne</i> )
F	H_A_1067	g3c - Other neutral grassland	-	-	Mowed grassland in between arable fields. Herb rich with low award height. No obvious use. Oak saplings.	Ribwort plantain ( <i>Plantago lanceolata</i> ) A, bittercress ( <i>Cardamine</i> sp.) F, dandelion ( <i>Taraxacum</i> agg.) O, common ragwort ( <i>Senecio jacobaea</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R, hogweed ( <i>Heracleum sphondylium</i> ) O, figwort ( <i>Scrophularia</i> sp.) R, common centaury ( <i>Centaurium erythraea</i> ) R, speedwell ( <i>Veronica</i> sp.). R. cock's-foot ( <i>Dactylis glomerata</i> ) A, meadow buttercup ( <i>Ranunculus acris</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) dock ( <i>Rumex</i> sp.) O, bramble ( <i>Rubus fruticosus</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) A, false oat grass ( <i>Arrhenatherum elatius</i> ) F.
F	H_A_1068	c1 - Arable and horticulture	-	-	Arable field recently plotted.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	H_A_1069	g3c - Other neutral grassland	Scattered scrub	-	Neutral grassland with dense common nettle. Grassland with evidence of past disturbance.	Common nettle ( <i>Urtica dioica</i> ) A, field bindweed ( <i>Convolvulus arvensis</i> ) F, cleavers ( <i>Galium aparine</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) R, dock ( <i>Rumex</i> sp.) O, false oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, perennial rye-grass ( <i>Lolium perenne</i> ) F hogweed ( <i>Heracleum sphondylium</i> ) O, soft brome ( <i>Bromus mollis</i> ) O.
F	H_A_1071	c1 - Arable and horticulture	-	-	Arable	-
F	H_A_1072	g3c - Other neutral grassland	Scattered scrub	-	Grassland with evidence of past disturbance. Mix of tall ruderal and bramble scrub vegetation.	Bramble ( <i>Rubus fruticosus</i> ) A, thistle ( <i>Cirsium</i> sp.) O, creeping buttercup ( <i>Ranunculus repens</i> ) O, common sorrel ( <i>Rumex acetosa</i> ) F, common nettle ( <i>Urtica dioica</i> ) A, cleavers ( <i>Galium aparine</i> ) O, mugwort ( <i>Artemisia vulgaris</i> ) O, elder ( <i>Sambucus nigra</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) O, dock ( <i>Rumex</i> sp.) O, ground ivy ( <i>Glechoma hederacea</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, common mallow ( <i>Malva sylvestris</i> ) O, wild teasel ( <i>Dipsacus fullonum</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) O, wild cherry ( <i>Prunus avium</i> ) R.
F	H_A_789	u1c - Artificial unvegetated, unsealed surface	-	-	Stable yard, gravel	-
F	H_A_815	g3c8 - Holcus-Juncus neutral grassland	-	-	Believed to be g3c8 which has deteriorated. Some alder saplings. Presence of abundant marsh thistle, rare yellow archangel and frequent cock's-foot taken as evidence of deterioration so poor condition.	Alder ( <i>Alnus glutinosa</i> ), greater bird's-foot-trefoil ( <i>Lotus pedunculatus</i> ), common chickweed ( <i>Stellaria media</i> ), water mint ( <i>Mentha aquatica</i> ), soft-rush ( <i>Juncus effusus</i> ) D, hard-rush ( <i>Juncus inflexus</i> ) D, Yorkshire fog ( <i>Holcus lanatus</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), cock's-foot ( <i>Dactylis glomerata</i> ) F, yellow archangel ( <i>Lamiastrum galeobdolon</i> ) R, marsh thistle ( <i>Cirsium palustre</i> ) A
F	H_A_832	c1d8 - Other non-cereal crops	-	-	Arable	-
F	H_A_851	g3c - Other neutral grassland	-	-	-	Common nettle ( <i>Urtica dioica</i> ), mullein ( <i>Verbascum</i> sp.), germander speedwell ( <i>Veronica chamaedrys</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), mugwort ( <i>Artemisia vulgaris</i> ), lesser celandine ( <i>Ranunculus ficaria</i> ), bluebell ( <i>Hyacinthoides non-scripta</i> ), docks ( <i>Rumex</i> spp.), dandelion ( <i>Taraxacum</i> agg.), red dead-nettle ( <i>Lamium purpureum</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), red campion ( <i>Silene dioica</i> ), cress ( <i>Arabidopsis</i> sp.), shining crane's-bill ( <i>Geranium lucidum</i> ), greater plantain ( <i>Plantago major</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), cut-leaved crane's-bill ( <i>Geranium dissectum</i> ), lords and ladies ( <i>Arum maculatum</i> ), common knapweed ( <i>Centaurea nigra</i> ), burdock ( <i>Arctium</i> sp.), groundsel ( <i>Senecio vulgaris</i> ), hedge woundwort ( <i>Stachys sylvatica</i> ), thistle ( <i>Cirsium</i> sp.), scabious species.
F	H_A_858	g4 - Modified grassland	-	-	Planted tree saplings, max. 50cm tall around field margins. Grass mown	Dandelion ( <i>Taraxacum</i> agg.) s, groundsel ( <i>Senecio vulgaris</i> ), common mallow ( <i>Malva sylvestris</i> ).
F	H_A_919	u1d - Suburban/ mosaic of developed/ natural surface	-	-	Partly vegetated gravel track.	Oxeye daisy ( <i>Leucanthemum vulgare</i> ), common knapweed ( <i>Centaurea nigra</i> ), common mallow ( <i>Malva sylvestris</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), daisy ( <i>Bellis perennis</i> ), common nettle ( <i>Urtica dioica</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), common ragwort ( <i>Senecio jacobaea</i> ).
F	H_A_958	g4 - Modified grassland	-	-	Possibly grazed annually, some tussocks, limited sward diversity or forbs.	Wood-sedge ( <i>Carex sylvatica</i> ), Yorkshire fog ( <i>Holcus lanatus</i> )
F	H_A_999	w1f - Lowland mixed deciduous woodland	-	HPI	Hazel coppice with occasional mature oak. Large amount of bare ground.	Hazel ( <i>Corylus avellana</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, elder ( <i>Sambucus nigra</i> ) R, common nettle ( <i>Urtica dioica</i> ) A, wood avens ( <i>Geum urbanum</i> ) R, great horsetail ( <i>Equisetum telmateia</i> ) R.
F	H3_1089	c1c - Cereal Crops	-	-	Arable	-
F	H3_1116	c1c - Cereal Crops	-	-	Arable	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	H3_1118	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_1119	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_1121	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_1124	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_913	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_936	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_937	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_941	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	H3_943	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	HA3_860	f2f - Other swamps	-	-	Swamp that does not meet other UKHab definitions. High proportion of ruderal species including and willow herb. Sedges and bulrush making other components	Common nettle ( <i>Urtica dioica</i> )
F	HA3_957	g - Grassland	Plantation, non-native, scattered trees	-	Native and non-native mix, less than 20 years in age. Turkey oak present.	Turkey oak ( <i>Quercus cerris</i> )
F	HA3_958	g - Grassland	Plantation, non-native, scattered trees	-	Native and non-native mix, less than 20 years in age. Turkey oak present.	Turkey oak ( <i>Quercus cerris</i> )
F	HL_126	w1d - Wet woodland	-	HPI	Brook running as land boundary through area of wet woodland. As head east topography slowly increases. Stream lacks vegetation. Over mature hazel coppice to western aspect. Just where it starts to get a little dryer.	Great horsetail ( <i>Equisetum telmateia</i> ) F, lesser pond-sedge ( <i>Carex acutiformis</i> ) O, common nettle ( <i>Urtica dioica</i> ) F, pendulous sedge ( <i>Carex pendula</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) F, herb-Robert ( <i>Geranium robertianum</i> ) R, reed sweet-grass ( <i>Glyceria maxima</i> ) O, grey willow ( <i>Salix cinerea</i> ) A, willow ( <i>Salix</i> sp.) A, alder ( <i>Alnus glutinosa</i> ) A, tufted hair-grass ( <i>Deschampsia cespitosa</i> ) R, remote sedge ( <i>Carex remota</i> ) R, wild angelica ( <i>Angelica sylvestris</i> ) R, Indian balsam ( <i>Impatiens glandulifera</i> ) R, hazel ( <i>Corylus avellana</i> ) C, hawthorn ( <i>Crataegus monogyna</i> ) F, soft-rush ( <i>Juncus effusus</i> ) R, water forget-me-not ( <i>Myosotis scorpioides</i> ) R, toad rush ( <i>Juncus bufonius</i> ) R, brooklime ( <i>Veronica beccabunga</i> ) R, meadow vetchling ( <i>Lathyrus pratensis</i> ) O, rough meadow-grass ( <i>Poa trivialis</i> ) O, silver birch ( <i>Betula pendula</i> ) R, creeping bent ( <i>Agrostis stolonifera</i> ) R, common hemp-nettle ( <i>Galeopsis tetrahit</i> ) O, bugle ( <i>Ajuga reptans</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	HL_137	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Land impacted by activities of scrap yard to north. Left to become vegetated, it has a bund surrounding the area from the woodland. Further dumped materials present. Note that two <i>Polypogon monspeliensis</i> plants were found. This is a potentially rare plant away from the coast, but it can be introduced.	Bristly oxtongue ( <i>Picris echioides</i> ) R, creeping cinquefoil ( <i>Potentilla reptans</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) R, common bent ( <i>Agrostis capillaris</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) O, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) F, wood dock ( <i>Rumex sanguineus</i> ) R, greater plantain ( <i>Plantago major</i> ) R, ribbed melilot ( <i>Melilotus officinalis</i> ) R, goat's-rue ( <i>Galega officinalis</i> ) (intro) A, rosebay willowherb ( <i>Chamerion angustifolium</i> ) R, broad-leaved willowherb ( <i>Epilobium montanum</i> ) R, pendulous sedge ( <i>Carex pendula</i> ) R, yarrow ( <i>Achillea millefolium</i> ) R, common nettle ( <i>Urtica dioica</i> ) O, black medick ( <i>Medicago lupulina</i> ) R, wild teasel ( <i>Dipsacus fullonum</i> ) R, hard-rush ( <i>Juncus inflexus</i> ) R, white clover ( <i>Trifolium repens</i> ) O, common vetch ( <i>Vicia sativa</i> ) R, verbena ( <i>Verbena sp.</i> ) (intro) O, creeping buttercup ( <i>Ranunculus repens</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) R, creeping bent ( <i>Agrostis stolonifera</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) O, great mullein ( <i>Verbascum thapsus</i> ) R, annual beard-grass ( <i>Polypogon monspeliensis</i> ) R
F	HL_141	g3c5 - Arrhenatherum neutral grassland	Traditional orchard	-	Only management is cut grass between orchard strips. Shrubs and trees growing over and killing orchard in places. Was an intensive orchard, that could have been abandoned approx. 30 years ago. Some trees already dead and some dying. As UKHab only mention intensive this polygon will access the grassland. Fits best g3c5 but is not a perfect fit, scrub coming through to western section.	Apple ( <i>Malus</i> var) D, bramble ( <i>Rubus fruticosus</i> ) A, corn parsley ( <i>Petroselinum segetum</i> ) F, smooth tare ( <i>Vicia tetrasperma</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, common bent ( <i>Agrostis capillaris</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) O, Italian alder ( <i>Alnus cordata</i> ) (intro) A, pedunculate oak ( <i>Quercus robur</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, marsh thistle ( <i>Cirsium palustre</i> ) R, tufted hair-grass ( <i>Deschampsia caespitosa</i> ) R, Timothy ( <i>Phleum pratense</i> ) O, white clover ( <i>Trifolium repens</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) O, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, Italian alder ( <i>Alnus cordata</i> ) F
F	HL_142	g4 - Modified grassland	Horse grazed.	-	Informed to have a hay cut one to twice a year. Lacking more than 8/9 sp per m2.	Timothy ( <i>Phleum pratense</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) A, common bent ( <i>Agrostis capillaris</i> ) A, creeping thistle ( <i>Cirsium arvense</i> ) O, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) R, common mouse-ear ( <i>Cerastium fontanum</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) O, red fescue ( <i>Festuca rubra</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) R, bristly ox-tongue ( <i>Picris echioides</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, dandelion ( <i>Taraxacum agg.</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R
F	HL_143	w1g7 - Other broadleaved woodland types	Plantation	-	No evidence of recent felling. Uniform age ranged. On a bank that rises up to the south by 1m every 3m. Log piles from ride where wire goes through. Bare ground flora dominant.	Sweet chestnut ( <i>Castanea sativa</i> ) D, silver birch ( <i>Betula pendula</i> ) R, ash ( <i>Fraxinus excelsior</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, holly ( <i>Ilex aquifolium</i> ) S, common nettle ( <i>Urtica dioica</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) O, common bent ( <i>Agrostis capillaris</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) R, wood avens ( <i>Geum urbanum</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) F
F	HL_144	h3d - Bramble scrub	-	-	A G3c grassland that has transitioned to bramble scrub. Starting to have saplings of woody species. Owner informed area to have been cleared of trees approx., 5 years ago. Grassland present as glades.	Tufted hair-grass ( <i>Deschampsia cespitosa</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) R, wood dock ( <i>Rumex sanguineus</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) O, bramble ( <i>Rubus fruticosus</i> ) A, common nettle ( <i>Urtica dioica</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, corn parsley ( <i>Petroselinum segetum</i> ) R, marsh thistle ( <i>Cirsium palustre</i> ) F, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) R, smooth tare ( <i>Vicia tetrasperma</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	HL_147	w1g - Other woodland; broadleaved	-	-	Different ages of trees planted. Hybrid black poplars almost in a going through marking the eastern boundary. These being of intermediate age. Rest of woodland to west planted, young. Divided into different species groups in sections e.g., hazel.	Hybrid black poplar ( <i>Populus deltoides x nigra</i> ) F, eagle fern ( <i>Pteridium aquilinum</i> ) R, common nettle ( <i>Urtica dioica</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) F, dog's mercury ( <i>Mercurialis perennis</i> ) R, elder ( <i>Sambucus nigra</i> ) R, silver birch ( <i>Betula pendula</i> ) F, ash ( <i>Fraxinus excelsior</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) O, common bent ( <i>Agrostis capillaris</i> ) A, dog-rose ( <i>Rosa canina</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) O, wild cherry ( <i>Prunus avium</i> ) F, field bindweed ( <i>Convolvulus arvensis</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) F, hazel ( <i>Corylus avellana</i> ) F, willow ( <i>Salix</i> sp.) R, cat's-ear ( <i>Hypochaeris radicata</i> ) O, hairy tare ( <i>Vicia hirsuta</i> ) R, fox and cubs ( <i>Pilosella aurantiaca</i> ) R
F	HL_149	g3c - Other neutral grassland	-	-	Possibly left ungrazed or cut this year as lacks tussocks of grass, presence of moderate ant hills and vegetation growing densely and wrapping around gates to field. Lacks tussocks grassland and short sword height Av. 15-20 cm to 30cm to south. This combined with lack of <i>Arrhenatherum elatius</i> means it must be getting cut. Young trees planted ash x3 on eastern side. Possibly a sown mixture.	Red fescue ( <i>Festuca rubra</i> ) A, common ragwort ( <i>Senecio jacobaea</i> ) F, lesser trefoil ( <i>Trifolium dubium</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) F, ribwort plantain ( <i>Plantago lanceolata</i> ) A, smooth tare ( <i>Vicia tetrasperma</i> ) O, dandelion ( <i>Taraxacum</i> agg.) F, common bent ( <i>Agrostis capillaris</i> ) F, hairy tare ( <i>Vicia hirsuta</i> ) LA, white clover ( <i>Trifolium repens</i> ) O, field bindweed ( <i>Convolvulus arvensis</i> ) F, curled dock ( <i>Rumex crispus</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) A, white campion ( <i>Silene latifolia</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) R, Agrimony ( <i>Agrimonia eupatoria</i> ) R, Timothy ( <i>Phleum pratense</i> ) ,R spear thistle ( <i>Cirsium vulgaris</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) O, goat's-beard ( <i>Tragopogon pratensis</i> agg) R, wild carrot ( <i>Daucus carota</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) R, lesser burdock ( <i>Arctium minus</i> ) R, downy oat-grass ( <i>Avenula pubescens</i> ) R
F	HL_230	c1c - Cereal crops	-	-	Arable field crop not currently planted.	-
F	HL_231	c1c - Cereal crops	-	-	Field of winter wheat	-
F	HL_233	c1c - Cereal crops	-	-	Arable field	-
F	HL_234	c1c5 - Winter stubble	-	-	Arable field of winter stubble.	-
F	HL_235	c1d8 - Other non-cereal crops	-	-	-	-
F	F_T_880	c1 - Arable and horticulture	-	-	Arable field.	-
F	F_T_878	w1g7 - Other broadleaved woodland types	Plantation	-	Primarily a hornbeam plantation interspersed with other species and scrub. Very little ground flora. No ancient woodland indicators, suggesting that the area has been arable for some time. Area approximately 40 years old. Flailed on arable margin.	Hornbeam ( <i>Carpinus betulus</i> ) D, ash ( <i>Fraxinus excelsior</i> ) O, oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) LO, blackthorn ( <i>Prunus spinosa</i> ) LO, dogrose ( <i>Rosa canina</i> ) R, nettle ( <i>Urtica dioica</i> ) O.
F	F_T_879	c1 - Arable and horticulture	-	-	Arable field.	-
F	HL_237	c1a5 - Arable margins sown with tussocky grasses	-	-	Adjacent to a pheasant rearing station. Rabbit activity. Margin about 15 metres wide. Probably sown with red fescue seed mix. 15 species. Grass dominated, less than 8 species per square metre.	Red fescue ( <i>Festuca rubra</i> ) D, pedunculate oak ( <i>Quercus robur</i> ) R, Timothy ( <i>Phleum pratense</i> ) O, crested dog's-tail ( <i>Cynosurus cristatus</i> ) R, common bent ( <i>Agrostis capillaris</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) O, white clover ( <i>Trifolium repens</i> ) O, dandelion ( <i>Taraxacum</i> agg.) O, cat's-ear ( <i>Hypochaeris radicata</i> ) R, smooth tare ( <i>Vicia tetrasperma</i> ) R, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) R, lesser trefoil ( <i>Trifolium dubium</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	HL_238	g3c6 - Lolium-Cynosurus neutral grassland	-	-	Area of land left unploughed by farmer because of subsidence and safety.	Cock's-foot ( <i>Dactylis glomerata</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, Timothy ( <i>Phleum pratense</i> ) O, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, red fescue ( <i>Festuca rubra</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, common centaury ( <i>Centaurium erythraea</i> ) O, lesser trefoil ( <i>Trifolium dubium</i> ) O, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) R, Tall fescue ( <i>Schedonardus arundinaceus</i> ) F, oxeye daisy ( <i>Leucanthemum vulgare</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) R, common bent ( <i>Agrostis capillaris</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) R, dandelion ( <i>Taraxacum agg.</i> ) R, smooth tare ( <i>Vicia tetrasperma</i> ) LA, cow parsley ( <i>Anthriscus sylvestris</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, Agrimony ( <i>Agrimonia eupatoria</i> ) R, blue fleabane ( <i>Erigeron acer</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, gorse ( <i>Ulex europaeus</i> ) R, common mouse-ear ( <i>Cerastium fontanum</i> ) R, alder ( <i>Alnus glutinosa</i> ) R, Italian alder ( <i>Alnus cordata</i> )
F	HL_239	c1c5 - Winter stubble	-	-	Arable	-
F	HL_244	w1f7 - Other Lowland mixed deciduous woodland	-	-	Ring of mixed deciduous tree surrounding a freshwater pond constrained by an arable field. One mature oak and significant willow, often fallen. Semi mature ash frequent. Steep bank down to pond approximately 2.5 metres deep. Ground vegetation sparse due to heavy shading. Some ancient woodland indicators with some nettle and ivy.	Ash ( <i>Fraxinus excelsior</i> ) A, willow ( <i>Salix sp.</i> ) A, hazel ( <i>Corylus avellana</i> ) F, elder ( <i>Sambucus nigra</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, wild cherry ( <i>Prunus avium</i> ) R, field maple ( <i>Acer campestre</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) R, common nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) R, ivy ( <i>Hedera helix</i> ) F, wood dock ( <i>Rumex sanguineus</i> ) R, bramble ( <i>Rubus fruticosus</i> ) O, woody nightshade ( <i>Solanum dulcamara</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) R, corn parsley ( <i>Petroselinum segetum</i> ) R
F	W_1034	- r1 – Standing open water and canals	-	HPI	-	-
F	HL_285	w1f7 - Other Lowland mixed deciduous woodland	-	-	Updated using v3.0. Small woodland parcel surrounded by arable grassland. Set within depression perhaps previously filled with water. Extensive rabbit/other mammal burrows. Pedunculate oak are mature, other trees young.	Pedunculate oak ( <i>Quercus robur</i> ) D, elder ( <i>Sambucus nigra</i> ) O, wild cherry ( <i>Prunus avium</i> ) R, willow ( <i>Salix sp.</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) O, common nettle ( <i>Urtica dioica</i> ) D, ivy ( <i>Hedera helix</i> ) R.
F	W_1036	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
F	HL_288	g3c - Other neutral grassland	Scattered scrub	-	Grassland with evidence of past disturbance. Mix of tall ruderal and bramble scrub vegetation.	Bramble ( <i>Rubus fruticosus</i> ) A, thistle ( <i>Cirsium sp.</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) O, common sorrel ( <i>Rumex acetosa</i> ) F, common nettle ( <i>Urtica dioica</i> ) A, willowherb ( <i>Epilobium sp.</i> ) A, spear thistle ( <i>Cirsium vulgare</i> ) O, sedge ( <i>Carex sp.</i> ) O, cleavers ( <i>Galium aparine</i> ) O, mugwort ( <i>Artemisia vulgaris</i> ) O, heath speedwell ( <i>Veronica officinalis</i> ) O, elder ( <i>Sambucus nigra</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) O, dock ( <i>Rumex sp.</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, common mallow ( <i>Malva sylvestris</i> ) O, wild teasel ( <i>Dipsacus fullonum</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) F.

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
F	HL_291	g3c - Other neutral grassland	-	-	Mowed grassland in between arable fields. Herb rich with low award height. No obvious use. Oak saplings.	Ribwort plantain ( <i>Plantago lanceolata</i> ) A, bittercress ( <i>Cardamine</i> sp.) F, dandelion ( <i>Taraxacum</i> agg.) O, common ragwort ( <i>Senecio jacobaea</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R, hogweed ( <i>Heracleum sphondylium</i> ) O, figwort ( <i>Scrophularia</i> sp.) R, common centaury ( <i>Centaurium erythraea</i> ) R, speedwell ( <i>Veronica</i> sp.) R, cock's-foot ( <i>Dactylis glomerata</i> ) A, meadow buttercup ( <i>Ranunculus acris</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ), dock ( <i>Rumex</i> sp.) O, bramble ( <i>Rubus fruticosus</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) A, false oat grass ( <i>Arrhenatherum elatius</i> ) F.
F	HL_295	w1f - Lowland mixed deciduous woodland	-	HPI	Updated using v3.0. Woodland on grassland edge moving into private garden. Not fully accessed.	Crack willow ( <i>Salix fragilis</i> ) R, elder ( <i>Sambucus nigra</i> ) O, holly ( <i>Ilex aquifolium</i> ) A, lime ( <i>Tilia</i> sp.) O, pedunculate oak ( <i>Quercus robur</i> ) F, sycamore ( <i>Acer pseudoplatanus</i> ) R, willow ( <i>Salix</i> sp.) O, wild cherry ( <i>Prunus avium</i> ) R.
F	HL_298	c1 - Arable and horticulture	-	-	Arable field with one mature oak (target noted). Pylons within field that have some longer vegetation at base.	Pedunculate oak ( <i>Quercus robur</i> )
F	HL_302	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
F	HL_435	g4 - Modified grassland	Cattle grazed.		Grassland not meeting enough species per m <sup>2</sup> to be neutral grassland.	White clover ( <i>Trifolium repens</i> ) D, perennial rye-grass ( <i>Lolium perenne</i> ), red fescue ( <i>Festuca rubra</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), creeping buttercup ( <i>Ranunculus repens</i> )
G	A_1147	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1216	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1257	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1265	g1a - Lowland dry acid grassland	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1267	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1356	g1a - Lowland dry acid grassland	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1377	r2a – Rivers (priority habitat)	-	HPI	River Stour UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1394	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1398	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1436a	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HA3_898	h3h – mixed scrub	-	-	Tall mixed scrub that was once a hedgerow that has grown into an area of dense scrub with a dry ditch running though.	-
G	A_1699	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1708	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1710	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1787	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1800	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_291	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_293	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_294	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_311	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_319	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_333	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_376	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_1296	w1f7 – Other lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_385	h3h - Mixed scrub	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_386	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_392	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_742	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	A_868	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_1012	w1h5 - Other woodland; mixed; mainly broadleaved	Plantation	-	Woodland planted in rows (plantation), no understory, trees less than 50yrs old. Canopy still open, no age structure. Herb layer dominated by common nettles with some elder (photo included to show reference of herb layer height). pedunculate oak dominated with sweet chestnut and occasional wild cherry. Field maple cut down within woodland. Ancient woodland indicators present. Condition assessment attached separately - score of 23 gives poor condition.	Elder ( <i>Sambucus nigra</i> ) O, wild cherry ( <i>Prunus avium</i> ) O, field maple ( <i>Acer campestre</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) D, sweet chestnut ( <i>Castanea sativa</i> ) O, elder ( <i>Sambucus nigra</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, walnut ( <i>Juglans regia</i> ) R, lime ( <i>Tilia x europaea</i> ) R, common nettle ( <i>Urtica dioica</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, bramble ( <i>Rubus fruticosus</i> ) A
G	H_A_1014	c1c - Cereal crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H_A_1016	w1f7 - Other Lowland mixed deciduous woodland	-	-	Various tree species present including medium sized oaks. Mainly dominated by field maple. Oak seedlings at the bottom. Links to woodland. Some areas too difficult to survey due to overgrown shrub layer. Shrub layer mainly consisting of dog's mercury and common nettles. Dead badger found also. 15m gap between two parts of green lane woodland for vehicle entrance. Polygon of green lane extends further north than the buffer and connects to vegetation in gardens to the north. Condition assessment: At least 2 ages present (moderate), no herbivore damage (good), no invasive species (good), 5 or more native species (good), >80% of canopy trees and understory is native (good), no open space (poor), woodland regeneration is good, tree health is good, ancient woodland flora indicators present x4 (good), 2 or more stories present across plot (moderate), no veteran tree present (poor), not much dead wood - although couldn't be assessed in all areas (poor), woodland enrichment (moderate). Good condition.	Pedunculate oak ( <i>Quercus robur</i> ), field maple ( <i>Acer campestre</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), common nettle ( <i>Urtica dioica</i> )
G	H_A_1017	c1c - Cereal crops	-	-	More gently sloping crop field bordered by grassland, houses, and hedgerow trees.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_1018	g3c - Other neutral grassland	-	-	Grassy area next to woodland and planted maize used for game purposes. Species present in both fields are the same/very similar. Grassland was being cut at the time of surveying. Red kite present in area.	Perforate St John's-wort ( <i>Hypericum perforatum</i> ) O, cat's-ear ( <i>Hypochaeris radicata</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, creeping bent ( <i>Agrostis stolonifera</i> ) F, bramble ( <i>Rubus fruticosus</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) A, common ragwort ( <i>Senecio jacobaea</i> ) F, common bent ( <i>Agrostis capillaris</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, Agrimony ( <i>Agrimonia eupatoria</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) R, downy oat-grass ( <i>Avenula pubescens</i> ) O, common nettle ( <i>Urtica dioica</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, common sorrel ( <i>Rumex acetosa</i> ) R, yellow oat-grass ( <i>Trisetum flavescens</i> ) R, common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) F, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) F, wild carrot ( <i>Daucus carota</i> ) F, red clover ( <i>Trifolium pratense</i> ) O, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) F, smooth tare ( <i>Vicia tetrasperma</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) F, bristly oxtongue ( <i>Picris echioides</i> ) R, oxeye daisy ( <i>Leucanthemum vulgare</i> ) A, goat's-beard ( <i>Tragopogon pratensis</i> ) O, blue fleabane ( <i>Erigeron acer</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R, yellow-wort ( <i>Blackstonia perfoliata</i> ) R, common calamint ( <i>Calamintha ascendens</i> ) R, old man's beard ( <i>Clematis vitalba</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, dandelion ( <i>Taraxacum agg.</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R,
G	H_A_1020	h3h - Mixed scrub	-	-	Over 5m wide, bordering two fields. Numerous dead elms, elm/elder, wormwood etc. Steep bank from one field to another. Passed 2/5 on condition assessment = poor condition. May have been an elm hedge at one point, elm still present but many dead. Unmanaged, very tall vegetation.	Elm ( <i>Ulmus</i> sp.), elder ( <i>Sambucus nigra</i> ), wormwood ( <i>Artemisia absinthium</i> )
G	H_A_1021	c1c - Cereal crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H_A_1025	c1c5 - Winter stubble	-	-	Arable	-
G	H_A_1027	c1c6 - Game bird mix fields	-	-	Arable	-
G	H_A_1028	c1c6 - Game bird mix fields	-	-	Maize for partridges	-
G	H_A_1029	w1f7 - Other Lowland mixed deciduous woodland	-	-	A strip of elm-dominated woodland spanning a three-metre-deep ditch between five and ten metres wide. Elms multi-stemmed 15-20m tall, small amount of dieback. 12 native woody species. Ditch very dry with greater than 50% cover bare ground.	Elm ( <i>Ulmus</i> sp.) D, field maple ( <i>Acer campestre</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) R, wild cherry ( <i>Prunus avium</i> ) R, spindle ( <i>Euonymus europaeus</i> ) R, ash ( <i>Fraxinus excelsior</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, elder ( <i>Sambucus nigra</i> ) R, willow ( <i>Salix</i> sp.) R, dogwood ( <i>Cornus sanguinea</i> ) O, hazel ( <i>Corylus avellana</i> ) R, common nettle ( <i>Urtica dioica</i> ) F, ivy ( <i>Hedera helix</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) O, wood dock ( <i>Rumex sanguineus</i> ) R, old man's beard ( <i>Clematis vitalba</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) F, wood avens ( <i>Geum urbanum</i> ) R
G	H_A_1034	c1c - Cereal crops	-	-	Ploughed field with no fringe of vegetation, with young self-set trees.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_1036	f2f - Other swamps	Tall herb. Unmanaged.		Fen priority habitat transitioning to alder woodland. Open wet woodland mosaics with alder saplings to 3 metres tall bounded by tall willow to 25 metres and oak and hazel trees. Frequent open areas with abundant Bracken, nettle, and giant horsetail. Adjacent to a stream. One 200 square stand of lesser pond-sedge. 14/4/22 MS and BF found Moschatel growing next to stream and wild garlic ancient woodland indicators.	Pedunculate oak ( <i>Quercus robur</i> ) F, hazel ( <i>Corylus avellana</i> ) F, alder ( <i>Alnus glutinosa</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) O, willow ( <i>Salix</i> sp.) F, marsh thistle ( <i>Cirsium palustre</i> ) F, eagle fern ( <i>Pteridium aquilinum</i> ) F, common nettle ( <i>Urtica dioica</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) O, giant fescue ( <i>Festuca gigantea</i> ) R, hogweed ( <i>Heracleum sphondylium</i> ) O, great horsetail ( <i>Equisetum telmateia</i> ) A, great willowherb ( <i>Epilobium hirsutum</i> ) F, angelica ( <i>Angelica sylvestris</i> ) O, hedge woundwort ( <i>Stachys sylvatica</i> ) R, hard-rush ( <i>Juncus inflexus</i> ) R, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) O, woody nightshade ( <i>Solanum dulcamara</i> ) R, meadowsweet ( <i>Filipendula ulmaria</i> ) O, lesser pond-sedge ( <i>Carex acutiformis</i> ) LA, moschatel ( <i>Adoxa moschatellina</i> )
G	H_A_1037	w1h5 - Other woodland; mixed; mainly broadleaved	-	-	A 30-year-old plantation with three mature trees used as pheasant feeding station, ground flora scarce no ancient woodland indicators, bisected by farm track with several neophytes.	Ash ( <i>Fraxinus excelsior</i> ) A, wild cherry ( <i>Prunus avium</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) A, sweet chestnut ( <i>Castanea sativa</i> ) F, cherry laurel ( <i>Prunus laurocerasus</i> ) O, white poplar ( <i>Populus alba</i> ) R, bramble ( <i>Rubus fruticosus</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) F, field maple ( <i>Acer campestre</i> ) R, hedge woundwort ( <i>Stachys sylvatica</i> ) R, common nettle ( <i>Urtica dioica</i> ) F
G	H_A_1039	g4 - Modified grassland	-	-	The field has been mown and sheep are grazing on the field, predominately grass with occasional herbaceous plants.	Cock's-foot ( <i>Dactylis glomerata</i> ) O, red fescue ( <i>Festuca rubra</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) F, yarrow ( <i>Achillea millefolium</i> ) O, common bent ( <i>Agrostis capillaris</i> ) A, sheep's sorrel ( <i>Rumex acetosella</i> ) R, rough meadow-grass ( <i>Poa trivialis</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, dandelion ( <i>Taraxacum</i> agg.) O
G	H_A_1073	w1g7 - Other broadleaved woodland types	-	-	Plantation woodland parcel. Sparse understory with no shrub layer. Small patches of bramble ( <i>Rubus fruticosus</i> )	Pedunculate oak ( <i>Quercus robur</i> ) F, ash ( <i>Fraxinus excelsior</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, wild cherry ( <i>Prunus avium</i> ) O, sweet chestnut ( <i>Castanea sativa</i> ) R, hazel ( <i>Corylus avellana</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, field maple ( <i>Acer campestre</i> ) R, bramble ( <i>Rubus fruticosus</i> ), ground ivy ( <i>Glechoma hederacea</i> ), common nettle ( <i>Urtica dioica</i> ), wood speedwell ( <i>Veronica montana</i> )
G	H_A_1077	c1c7 - Other cereal crops	Scattered trees		Oak within field	Pedunculate oak ( <i>Quercus robur</i> )
G	H_A_1078	g4 - Modified grassland	-	-	Farmland historically used for grazing / pigs which has been left abandoned. Composition uniform. Not reaching enough species to be other neutral grassland. Common lizards present according to owner.	Cock's-foot ( <i>Dactylis glomerata</i> ) D, common nettle ( <i>Urtica dioica</i> ), bramble ( <i>Rubus fruticosus</i> ) O, cinquefoil ( <i>Potentilla</i> sp.), creeping bent ( <i>Agrostis stolonifera</i> ), curled dock ( <i>Rumex crispus</i> ), yellow oat-grass ( <i>Trisetum flavescens</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), common knapweed ( <i>Centaurea nigra</i> ).
G	H_A_1079	g4 - Modified grassland	Horse grazed	-	Short grazing land. Access not possible due to horses.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, cinquefoil ( <i>Potentilla</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ), O, curled dock ( <i>Rumex crispus</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ).

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_1126	g1a6 - Other lowland dry acid grassland	Dry. Sward type mosaic. Other grazed. Scattered scrub.	HPI	A mosaic of grassland types, predominantly G1a6 with some G4. Patches of bramble and nettle, management probably none but heavy rabbit grazing and burrowing.	False oat grass ( <i>Arrhenatherum elatius</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, common bent ( <i>Agrostis capillaris</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) F, common ragwort ( <i>Senecio jacobaea</i> ) F, musk thistle ( <i>Carduus nutans</i> ) R, perforate St John's-wort ( <i>Hypericum perforatum</i> ) F, hop trefoil ( <i>Trifolium campestre</i> ) R, sheep's sorrel ( <i>Rumex acetosella</i> ) O, common mouse-ear ( <i>Cerastium fontanum</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) O, dandelion ( <i>Taraxacum</i> agg.) R, creeping thistle ( <i>Cirsium arvense</i> ) F, bramble ( <i>Rubus fruticosus</i> ) F, common nettle ( <i>Urtica dioica</i> ) F, dog-rose ( <i>Rosa canina</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) O, small-flowered crane's-bill ( <i>Geranium pusillum</i> ) R, yarrow ( <i>Achillea millefolium</i> ) O, hairy sedge ( <i>Carex hirta</i> ) R, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R, red fescue ( <i>Festuca rubra</i> ) F, sheep's fescue ( <i>Festuca ovina</i> ) O, yellow oat-grass ( <i>Trisetum flavescens</i> ) O
G	H_A_1130	g3c - Other neutral grassland	-	-	Habitat same as polygon/field west. Mature pedunculate oak tree within field.	Pedunculate oak ( <i>Quercus robur</i> )
G	H_A_1131	w1d - Wet woodland	-	HPI	Goat and crack willow dominant south-east of stream.	Pedunculate oak ( <i>Quercus robur</i> ) (A), alder ( <i>Alnus glutinosa</i> ) (A), hazel ( <i>Corylus avellana</i> ), horse chestnut ( <i>Aesculus hippocastanum</i> ) (R), crack willow ( <i>Salix fragilis</i> ), goat willow ( <i>Salix caprea</i> ), hawthorn ( <i>Crataegus monogyna</i> ) (D), holly ( <i>Ilex aquifolium</i> ) (R), ground ivy ( <i>Glechoma hederacea</i> ) (O), wild garlic ( <i>Allium ursinum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) (R), common nettle ( <i>Urtica dioica</i> ) (D), cleavers ( <i>Galium aparine</i> ) (R), wood avens ( <i>Geum urbanum</i> ) (O), red campion ( <i>Silene dioica</i> ) (R), herb-Robert ( <i>Geranium robertianum</i> ) (R).
G	H_A_729	g3c - Other neutral grassland	Scattered scrub.		Grassland area, some areas of bare ground where rabbit warrens present and heavily grazed by rabbit however more than 8 species present in 1m.	Cock's-foot ( <i>Dactylis glomerata</i> ) A, Yorkshire fog ( <i>Holcus lanatus</i> ) A, false brome ( <i>Brachypodium sylvaticum</i> ) O, field bindweed ( <i>Convolvulus arvensis</i> ) R, groundsel ( <i>Senecio vulgaris</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, cranesbill ( <i>Geranium</i> sp.) R, common nettle ( <i>Urtica dioica</i> ) LF, spear thistle ( <i>Cirsium vulgare</i> ) R, dandelion ( <i>Taraxacum</i> agg.) O, ribwort plantain ( <i>Plantago lanceolata</i> ) R, cinquefoil ( <i>Potentilla</i> sp.) F.
G	H_A_733	g4 - Modified grassland	-	-	Heavily grazed by rabbits, few species present therefore, does not fit criteria g3.	-
G	H_A_734	g3c - Other neutral grassland	Scattered scrub		Grassland area, some areas of bare ground where rabbit warrens present and heavily grazed by rabbit however majority sward height above 20cm. Alder scrub at bottom of hill next to stream, scattered scrub of bramble and blackthorn.	Alder ( <i>Alnus glutinosa</i> ), bramble ( <i>Rubus fruticosus</i> ), blackthorn ( <i>Prunus spinosa</i> ), cock's-foot ( <i>Dactylis glomerata</i> ) A, Yorkshire fog ( <i>Holcus lanatus</i> ) A, false brome ( <i>Brachypodium sylvaticum</i> ) O, field bindweed ( <i>Convolvulus arvensis</i> ) R, groundsel ( <i>Senecio vulgaris</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, cranesbill ( <i>Geranium</i> sp.) R, common nettle ( <i>Urtica dioica</i> ) LF, spear thistle ( <i>Cirsium vulgare</i> ) R, dandelion ( <i>Taraxacum</i> agg.) O, ribwort plantain ( <i>Plantago lanceolata</i> ) R, cinquefoil ( <i>Potentilla</i> sp.) F.
G	H_A_739	h3h - Mixed scrub	-	-	Area of mixed scrub, some areas denser than others dominated by bramble and young pedunculate oak trees.	Bramble ( <i>Rubus fruticosus</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O
G	H_A_757	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Land left from farming activities species composition same as adjacent u1a habitat	-
G	H_A_762	c1 - Arable and horticulture	-	-	Bare earth	-
G	H_A_768	g3c - Other neutral grassland	-	-	-	Yorkshire fog ( <i>Holcus lanatus</i> ) (D), perennial rye-grass ( <i>Lolium perenne</i> ) (O), cock's-foot ( <i>Dactylis glomerata</i> ) (D), creeping buttercup ( <i>Ranunculus repens</i> ) (O), common nettle ( <i>Urtica dioica</i> ) (O), tufted hair-grass ( <i>Deschampsia cespitosa</i> ) (A), thistle ( <i>Cirsium</i> sp.) (O), false oat grass ( <i>Arrhenatherum elatius</i> ) (A)

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_787	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Car park area and access route. Large stone 5-7cm. Loosely vegetated. Buddleia and oak saplings.	Sticky mouse-ear ( <i>Cerastium glomeratum</i> ), yarrow ( <i>Achillea millefolium</i> ), creeping cinquefoil ( <i>Potentilla reptans</i> ), common stork's-bill ( <i>Erodium cicutarium</i> ), hedgerow crane's-bill ( <i>Geranium pyrenaicum</i> ), early forget-me-not ( <i>Myosotis ramosissima</i> ), perforate St John's-wort ( <i>Hypericum perforatum</i> ), daisy ( <i>Bellis perennis</i> ), red fescue ( <i>Festuca rubra</i> ), sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), buddleia ( <i>Buddleja</i> sp.), pedunculate oak ( <i>Quercus robur</i> ), common ragwort ( <i>Senecio jacobaea</i> ), common hogweed ( <i>Heracleum sphondylium</i> ),
G	H_A_797	w1f7 - Other Lowland mixed deciduous woodland	-	-	Oak dominated woodland, more than 75% presence. Northern edge of woodland has dry ditch with old line of trees present that have become part of woodland with hazel coppice, oaks here mature.	Pedunculate oak ( <i>Quercus robur</i> ), ground ivy ( <i>Glechoma hederacea</i> ) (O), bramble ( <i>Rubus fruticosus</i> ) (O), common dog-violet ( <i>Viola riviniana</i> ) (O), primrose ( <i>Primula vulgaris</i> ) (R), dog's mercury ( <i>Mercurialis perennis</i> ) (A), common nettle ( <i>Urtica dioica</i> ) (F), currant ( <i>Ribes</i> sp.) (O), cleavers ( <i>Galium aparine</i> ) (F), wild garlic ( <i>Allium ursinum</i> ) (R), holly ( <i>Ilex aquifolium</i> ) (R), pendulous sedge ( <i>Carex pendula</i> ) (R), dock ( <i>Rumex</i> sp.) (O), bluebell ( <i>Hyacinthoides non-scripta</i> ) (R), lesser celandine ( <i>Ranunculus ficaria</i> ) (A), lords and ladies ( <i>Arum maculatum</i> ) (O), dandelion ( <i>Taraxacum</i> agg.) (R), hazel ( <i>Corylus avellana</i> ), holly ( <i>Ilex aquifolium</i> ), hazel ( <i>Corylus avellana</i> ), blackthorn ( <i>Prunus spinosa</i> ), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ).
G	H_A_811	c1b - Temporary grass and clover leys	-	-	C1b7 stewardship scheme winter bird mix AB9. No till, drilled into soil. Planted 2018	-
G	H_A_812	c1d8 - Other non-cereal crops	-	-	Italian rye grass planted. No till.	-
G	H_A_814	w1f7 - Other Lowland mixed deciduous woodland	-	-	Plantation. Evenly aged. North of EWT nature reserve. Approx. 30 years age. Single storey. Sections used for educational purposes including tents and tree felling. Lack of ground cover dominant.	Alder ( <i>Alnus glutinosa</i> ) LA, hazel ( <i>Corylus avellana</i> ) C, wild cherry ( <i>Prunus avium</i> ), pedunculate oak ( <i>Quercus robur</i> ), blackthorn ( <i>Prunus spinosa</i> ), silver birch ( <i>Betula pendula</i> ), common nettle ( <i>Urtica dioica</i> ) LA, ground ivy ( <i>Glechoma hederacea</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), bramble ( <i>Rubus fruticosus</i> ).
G	H_A_815	g3c8 - Holcus-Juncus neutral grassland	-	-	Believed to be g3c8 which has deteriorated. Some alder saplings. Presence of abundant marsh thistle, rare yellow archangel and frequent cock's-foot taken as evidence of deterioration so poor condition.	Alder ( <i>Alnus glutinosa</i> ), greater bird's-foot-trefoil ( <i>Lotus pedunculatus</i> ), common chickweed ( <i>Stellaria media</i> ), water mint ( <i>Mentha aquatica</i> ), soft-rush ( <i>Juncus effusus</i> ) D, hard-rush ( <i>Juncus inflexus</i> ) D, Yorkshire fog ( <i>Holcus lanatus</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), cock's-foot ( <i>Dactylis glomerata</i> ) F, yellow archangel ( <i>Lamiastrum galeobdolon</i> ) R, marsh thistle ( <i>Cirsium palustre</i> ) A
G	H_A_820	g3c8 - Holcus-Juncus neutral grassland	-	-	Gap in centre of plantation under power line where there are no trees, ground flora similar to that of plantation.	Yorkshire fog ( <i>Holcus lanatus</i> ) A, hard-rush ( <i>Juncus inflexus</i> ) F, angelica ( <i>Angelica sylvestris</i> ) O, jointed rush ( <i>Juncus articulatus</i> ) R, common nettle ( <i>Urtica dioica</i> ) O, marsh thistle ( <i>Cirsium palustre</i> ) R, common hogweed ( <i>Heracleum sphondylium</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, square-stalked St John's-wort ( <i>Hypericum tetrapterum</i> ) R, water mint ( <i>Mentha aquatica</i> ) R, curled dock ( <i>Rumex crispus</i> ) O, soft-rush ( <i>Juncus effusus</i> ) R, fleabane ( <i>Erigeron</i> sp.) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, common fleabane ( <i>Pulicaria dysenterica</i> ), common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) R.
G	H_A_834	u1b - Developed land; sealed surface	-	-	Hard-standing, done from aerial.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_873	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Rocky gravel substrate with varied flora present. Largely young hazel. Abundant bramble towards edges. Mixture of open gravel ground, small, vegetated patches, emergent vegetation, and young hazel.	Hazel ( <i>Corylus avellana</i> ), buddleia ( <i>Buddleja</i> sp.) (O), willow ( <i>Salix</i> sp.) (O), pendulous sedge ( <i>Carex pendula</i> ) (O), spear thistle ( <i>Cirsium vulgare</i> ) (F), ground ivy ( <i>Glechoma hederacea</i> ) (F), speedwell ( <i>Veronica</i> sp.) (O), crane's-bill ( <i>Geranium</i> sp.) (O), forget-me-not ( <i>Myosotis</i> sp.) (O), great mullein ( <i>Verbascum thapsus</i> ) (R), common nettle ( <i>Urtica dioica</i> ) (O), groundsel ( <i>Senecio vulgaris</i> ) (O), bramble ( <i>Rubus fruticosus</i> ) (A), dandelion ( <i>Taraxacum</i> agg.) (R), bristly ox-tongue ( <i>Picris echioides</i> ) (O), willow herb ( <i>Epilobium</i> sp.) (O), cudweed sp. ( <i>Filago</i> sp.) O, bonfire moss ( <i>Funaria hygrometrica</i> ) O, wild teasel ( <i>Dipsacus fullonum</i> ) (R), alder ( <i>Alnus glutinosa</i> ) (R).
G	H_A_879	h3b - Hazel scrub	-	-	Hazel coppice with most trees similar age and height. Open areas with more bramble and nettle. Ground flora suggests previous cleared woodland. Willow increasingly present towards the east where ground is softer and sedge more common. Still hazel coppice, however. Small area of rushes amongst scrub but no open water.	Hazel ( <i>Corylus avellana</i> ), lords and ladies ( <i>Arum maculatum</i> ) (O), common nettle ( <i>Urtica dioica</i> ) (D), cleavers ( <i>Galium aparine</i> ) (F), red dead-nettle ( <i>Lamium purpureum</i> ) (F), bramble ( <i>Rubus fruticosus</i> ), (O), hogweed ( <i>Heracleum sphondylium</i> ) (R), dock ( <i>Rumex</i> sp.) (O), wild garlic ( <i>Allium ursinum</i> ) (F), rush ( <i>Juncus</i> sp.) (O), willow ( <i>Salix</i> sp.) O
G	H_A_880	g3c - Other neutral grassland	Tall herb.	-	Heavily disturbed grassland sloping down to river. Dominated by common nettle with damp areas containing horsetail and rush species. Wild garlic and dog's mercury both nearer to woodland edge. Some discarded wood clippings and logs.	Common nettle ( <i>Urtica dioica</i> ) D, elder ( <i>Sambucus nigra</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) (F), wild garlic ( <i>Allium ursinum</i> ) (O), red dead-nettle ( <i>Lamium purpureum</i> ), white dead-nettle ( <i>Lamium album</i> ) (O), lords and ladies ( <i>Arum maculatum</i> ) (O), spear thistle ( <i>Cirsium vulgare</i> ) (O), burdock ( <i>Arctium</i> sp.) (F), willow herb ( <i>Epilobium</i> sp.) (A), cleavers ( <i>Galium aparine</i> ) (R), hop ( <i>Humulus lupulus</i> ) (R), ground ivy ( <i>Glechoma hederacea</i> ) (O), sycamore ( <i>Acer pseudoplatanus</i> ) (R), sedge ( <i>Carex</i> sp.) (R)
G	H_A_881	r2a – Rivers (priority habitat)	-	HPI	Small stream running though woodland. See Aquatic Baseline Report ( <a href="#">application document 6.3.7.3</a> )	-
G	H_A_910	w1g - Other woodland; broadleaved	Plantation		Cricket willow plantation, areas to the south closer to stream is more wet.	Creeping buttercup ( <i>Ranunculus repens</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), marsh thistle ( <i>Cirsium palustre</i> ), false brome ( <i>Brachypodium sylvaticum</i> ), soft-rush ( <i>Juncus effusus</i> ), water mint ( <i>Mentha aquatica</i> ) O, ragged-robin ( <i>Lychnis flos-cuculi</i> ) O.
G	H_A_911	w1f - Lowland mixed deciduous woodland	-	HPI	Ash and oak woodland. No signs of flushing or species to indicate that is ground water dependent terrestrial ecosystem. Ancient woodland PoAWS10.	Ash ( <i>Fraxinus excelsior</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) A, sweet chestnut ( <i>Castanea sativa</i> ) R, wild cherry ( <i>Prunus avium</i> ) LO, elm ( <i>Ulmus</i> sp.) R, field maple ( <i>Acer campestre</i> ) C LF, hornbeam ( <i>Carpinus betulus</i> ) LO, hazel ( <i>Corylus avellana</i> ) C A, holly ( <i>Ilex aquifolium</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, bracken ( <i>Pteridium aquilinum</i> ) F, bramble ( <i>Rubus fruticosus</i> ) F, bluebell ( <i>Hyacinthoides non-scripta</i> ) A, wild garlic ( <i>Allium ursinum</i> ) A, three-nerved sandwort ( <i>Moehringia trinervia</i> ) R, garlic mustard ( <i>Alliaria petiolata</i> ) LF, red campion ( <i>Silene dioica</i> ) R, wood avens ( <i>Geum urbanum</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) LA, ground ivy ( <i>Glechoma hederacea</i> ).
G	H_A_913	w1d - Wet woodland	-	HPI	Ash woodland adjacent to pond, some mature ash in healthy good condition.	Ash ( <i>Fraxinus excelsior</i> ), alder ( <i>Alnus glutinosa</i> ) O, hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), elder ( <i>Sambucus nigra</i> ), common nettle ( <i>Urtica dioica</i> ) A, dog's mercury ( <i>Mercurialis perennis</i> ) A, ground ivy ( <i>Glechoma hederacea</i> ) F, lords and ladies ( <i>Arum maculatum</i> ) R, lesser celandine ( <i>Ranunculus ficaria</i> ) LO, three-nerved sandwort ( <i>Moehringia trinervia</i> ) O, bugle ( <i>Ajuga reptans</i> ) R, great horsetail ( <i>Equisetum telmateia</i> ) LA, field-rose ( <i>Rosa arvensis</i> ).

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_933	g4 - Modified grassland	-	-	Field of heavily grazed grassland.	Perennial rye-grass ( <i>Lolium perenne</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, Yorkshire fog ( <i>Holcus lanatus</i> ) O, field wood-rush ( <i>Luzula campestris</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, forget-me-not ( <i>Myosotis</i> sp.) R, creeping thistle ( <i>Cirsium arvense</i> ) R, bluebell ( <i>Hyacinthoides non-scripta</i> ) R, dandelion ( <i>Taraxacum</i> agg.) R
G	H_A_941	g4 - Modified grassland	Scattered trees.	-	Fenced strip of unmanaged land. Scattered holly and oak trees.	Common nettle ( <i>Urtica dioica</i> ) A, white dead-nettle ( <i>Lamium album</i> ) F, red dead-nettle ( <i>Lamium purpureum</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) O, red fescue ( <i>Festuca rubra</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) A, dandelion ( <i>Taraxacum</i> agg.) R, creeping thistle ( <i>Cirsium arvense</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, holly ( <i>Ilex aquifolium</i> ), pedunculate oak ( <i>Quercus robur</i> )
G	H_A_942	g4 - Modified grassland	-	-	Field of heavily grazed grassland with small areas of bracken.	Bracken ( <i>Pteridium aquilinum</i> ), perennial rye-grass ( <i>Lolium perenne</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, Yorkshire fog ( <i>Holcus lanatus</i> ) O, field wood-rush ( <i>Luzula campestris</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, forget-me-not ( <i>Myosotis</i> sp.) R, creeping thistle ( <i>Cirsium arvense</i> ) R, bluebell ( <i>Hyacinthoides non-scripta</i> ) R, dandelion ( <i>Taraxacum</i> agg.) R
G	H_A_943	h3h - Mixed scrub	-	-	Mixed scrub containing some larger oak trees and bed of bramble and nettle.	Hawthorn ( <i>Crataegus monogyna</i> ), pedunculate oak ( <i>Quercus robur</i> ), willow ( <i>Salix</i> sp.), bramble ( <i>Rubus fruticosus</i> ), ground ivy ( <i>Glechoma hederacea</i> ), ash ( <i>Fraxinus excelsior</i> ), common sorrel ( <i>Rumex acetosa</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), common nettle ( <i>Urtica dioica</i> )
G	H_A_947	w1d - Wet woodland	-	HPI	Incredibly dense. Access limited due to boggy nature. Ground flora dominated by nutrient enrichment species, mainly nettle, grass, some willowherb. Low parts willow, banks have alder and some hazel with scrub species like hawthorn. Willow and alder dominant. Historic alder coppiced in north-west section. Dry areas only at current pylon.	Common nettle ( <i>Urtica dioica</i> ), alder ( <i>Alnus glutinosa</i> ) D, willow ( <i>Salix</i> sp.) D, willowherb ( <i>Epilobium</i> sp.), hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), water mint ( <i>Mentha aquatica</i> ), large bitter-cress ( <i>Cardamine amara</i> ), hard-rush ( <i>Juncus inflexus</i> ), horsetail ( <i>Equisetum</i> sp.) cleavers ( <i>Galium aparine</i> ), marsh thistle ( <i>Cirsium palustre</i> )
G	W_1032 / H_A_948	r1a6 - Other eutrophic standing waters	-	HPI	Potentially used for fishing. Old peer and other rotten wooden structures on banks. Old small boats. But evident something gone in and out on southern bank. Trees overhang on northern and eastern aspect. Lacks any kind of aquatic vegetation apart water mint. Pair herons present.	Water mint ( <i>Mentha aquatica</i> ) R
G	W_1030	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
G	W_1033	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
G	H_A_949	h3h - Mixed scrub	-	-	Approx. 50m bramble along track running through to waterbody. Dense nettle.	Bramble ( <i>Rubus fruticosus</i> ), common nettle ( <i>Urtica dioica</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), creeping thistle ( <i>Cirsium arvense</i> ), dandelion ( <i>Taraxacum</i> agg.), dead-nettle ( <i>Lamium</i> sp.), spear thistle ( <i>Cirsium vulgare</i> ), sallow willow ( <i>Salix viminalis</i> ), SM ash ( <i>Fraxinus excelsior</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_950	c1b - Temporary grass and clover leys	-	-	Meant to contain notable species such as corncockle, corn marigolds, cornflower. AB8. C1b7. Track contains some overlap though slightly different. Good variety of species. Left over oil-seed rape.	Red campion ( <i>Silene dioica</i> ), greater bird's-foot-trefoil ( <i>Lotus pedunculatus</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), rape ( <i>Brassica napus</i> ), red fescue ( <i>Festuca rubra</i> ), lesser trefoil ( <i>Trifolium dubium</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), white clover ( <i>Trifolium repens</i> ), cut-leaved crane's-bill ( <i>Geranium dissectum</i> )
G	H_A_952	c1b - Temporary grass and clover leys	-	-	Meant to contain notable species such as corncockle, corn marigolds, cornflower. AB8. C1b7. Track contains some overlap though slightly different. Good variety of species. Left over oil-seed rape.	Red campion ( <i>Silene dioica</i> ), greater bird's-foot-trefoil ( <i>Lotus pedunculatus</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), rape ( <i>Brassica napus</i> ), red fescue ( <i>Festuca rubra</i> ), lesser trefoil ( <i>Trifolium dubium</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), white clover ( <i>Trifolium repens</i> ), cut-leaved crane's-bill ( <i>Geranium dissectum</i> )
G	H_A_955	g4 - Modified grassland	-	-	Access by horse field not taken as horses present. Assessment done of agave to northern bank. Likely cut at least once a year. Likely kept as horse pasture.	Meadow foxtail ( <i>Alopecurus pratensis</i> ), cock's-foot ( <i>Dactylis glomerata</i> ) D, Yorkshire fog ( <i>Holcus lanatus</i> ), curled dock ( <i>Rumex crispus</i> ), creeping thistle ( <i>Cirsium arvense</i> ), creeping buttercup ( <i>Ranunculus repens</i> )
G	H_A_957	w1g7 - Other broadleaved woodland types	-	-	Mixed species woodland, some wet ground. Small clearing where trees have been planted. Previously coppiced hazel.	Hazel ( <i>Corylus avellana</i> ) (A) , sycamore ( <i>Acer pseudoplatanus</i> ) (O), hawthorn ( <i>Crataegus monogyna</i> ) (O), elder ( <i>Sambucus nigra</i> ) (O), alder ( <i>Alnus glutinosa</i> ) (F), willow ( <i>Salix</i> sp.) (O), common nettle ( <i>Urtica dioica</i> ) (D), cleavers ( <i>Galium aparine</i> ) (O), lesser celandine ( <i>Ranunculus ficaria</i> ) (O), moschatel ( <i>Adoxa moschatellina</i> ) (F), lords and ladies ( <i>Arum maculatum</i> ) (F), dog's mercury ( <i>Mercurialis perennis</i> ) (F), elder ( <i>Sambucus nigra</i> ) (O), horsetail ( <i>Equisetum</i> sp.) (R), ground ivy ( <i>Glechoma hederacea</i> ) (O), sycamore ( <i>Acer pseudoplatanus</i> ) (O), garlic mustard ( <i>Alliaria petiolata</i> ) (O).
G	H_A_961	c1b - Temporary grass and clover leys	-	-	Arable land that for some reason has been left alone after being ploughed. Frequent common grasses and herbs with 30 to 50 % bare earth. Likely for sheep grazing.	Perennial rye-grass ( <i>Lolium perenne</i> ), fat-hen ( <i>Chenopodium album</i> ) LA, , shepherd's-purse ( <i>Capsella bursa-pastoris</i> ) O, scented mayweed ( <i>Matricaria recutita</i> ), common couch grass ( <i>Elytrigia repens</i> ), dandelion ( <i>Taraxacum</i> agg.), common field speedwell ( <i>Veronica persica</i> ), creeping thistle ( <i>Cirsium arvense</i> ), scentless mayweed ( <i>Tripleurospermum inodorum</i> ), white clover ( <i>Trifolium repens</i> ), groundsel ( <i>Senecio vulgaris</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), common chickweed ( <i>Stellaria media</i> ), common mouse-ear ( <i>Cerastium fontanum</i> ), small nettle ( <i>Urtica urens</i> ), scarlet pimpernel ( <i>Anagallis arvensis</i> )
G	H_A_962	g4 - Modified grassland	-	-	Field used for arable practices.	Perennial rye-grass ( <i>Lolium perenne</i> ) A, common poppy ( <i>Papaver rhoeas</i> ) R, common chickweed ( <i>Stellaria media</i> ), prickly sow thistle ( <i>Sonchus asper</i> ), spear thistle ( <i>Cirsium vulgare</i> ), hedge mustard ( <i>Sisymbrium officinale</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), bur chervil ( <i>Anthriscus caucalis</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), scentless mayweed ( <i>Tripleurospermum inodorum</i> ), scented mayweed ( <i>Matricaria recutita</i> ), wall speedwell ( <i>Veronica arvensis</i> ) R, common chickweed ( <i>Stellaria media</i> ), meadow fescue ( <i>Festuca pratensis</i> ) F, Italian ryegrass ( <i>Lolium multiflorum</i> ) F, white campion ( <i>Silene latifolia</i> ) R, meadow thistle ( <i>Cirsium dissectum</i> ), Timothy ( <i>Phleum pratense</i> ), red campion ( <i>Silene dioica</i> ), borage ( <i>Borago officinalis</i> ), common fumitory ( <i>Fumaria officinalis</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_969	w1f7 - Other Lowland mixed deciduous woodland	-	-	Ash canopy. Goat willow on steep banks forming 20m canopy layer. Old path in wood with banks from east to west. Topography odd with steep areas and valley which haven't seen in local area. Southern section has a stream, dry during survey, from east to west with mature wych elms and standing deadwood in banks. Update. Have been informed by local residents this area was quarried for aggregate for American WW2 airfield	Wild strawberry ( <i>Fragaria vesca</i> ) LA, false brome ( <i>Brachypodium sylvaticum</i> ) F, wood avens ( <i>Geum urbanum</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) O, primrose ( <i>Primula vulgaris</i> ) F, creeping buttercup ( <i>Ranunculus repens</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) R, wood speedwell ( <i>Veronica montana</i> ) R, wood doc ( <i>Rumex sanguineus</i> ) R, ash ( <i>Fraxinus excelsior</i> ) O, goat willow ( <i>Salix caprea</i> ) A, field maple ( <i>Acer campestre</i> ), spindle ( <i>Euonymus europaeus</i> ), hawthorn ( <i>Crataegus monogyna</i> ), dogwood ( <i>Cornus sanguinea</i> ) F, rose ( <i>Rosa sp</i> ), wych elm ( <i>Ulmus glabra</i> ) R.
G	H_A_971	g3c - Other neutral grassland	Cattle grazed		Pasture, cattle in field, dominated by grasses over 90% of composition. Young blackthorn by tower. 15/06 - checked for updated Order Limits and extended habitat.	Black medick ( <i>Medicago lupulina</i> ) O, oxeye daisy ( <i>Leucanthemum vulgare</i> ) F, selfheal ( <i>Prunella vulgaris</i> ), field bindweed ( <i>Convolvulus arvensis</i> ), common ragwort ( <i>Senecio jacobaea</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) O, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, red fescue ( <i>Festuca rubra</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ), Timothy ( <i>Phleum pratense</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, false brome ( <i>Brachypodium sylvaticum</i> ) R, agrimony ( <i>Agrimonia eupatoria</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) R, common bent ( <i>Agrostis capillaris</i> ), creeping bent ( <i>Agrostis stolonifera</i> ) O, goat's-beard ( <i>Tragopogon pratensis</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) R, yellow oat-grass ( <i>Trisetum flavescens</i> ) R, lady's bedstraw ( <i>Galium verum</i> ) R, lesser trefoil ( <i>Trifolium dubium</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) R, bramble ( <i>Rubus fruticosus</i> ) and wild teasel ( <i>Dipsacus fullonum</i> ), blackthorn ( <i>Prunus spinosa</i> ) Y, perforate St John's-wort ( <i>Hypericum perforatum</i> ) R, common nettle ( <i>Urtica dioica</i> ) R, small-flowered crane's-bill ( <i>Geranium pusillum</i> ) R, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) R, dandelion ( <i>Taraxacum agg.</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) R barley ( <i>Hordeum sp.</i> )
G	H_A_972	u1a - Open Mosaic Habitats on Previously Developed Land	-	HPI	Land which has been left by arable activities. Maybe due to stewardship for skylarks. Main body dominated by creeping thistle and bristly oxtongue. With a thin strip shooting off to north-east which has volunteer arable plants including wheat and rape.	Creeping thistle ( <i>Cirsium arvense</i> ) D, bristly oxtongue ( <i>Picris echioides</i> ) D, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) R, common poppy ( <i>Papaver rhoeas</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) R, curled dock ( <i>Rumex crispus</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) R
G	H_A_973	g3c - Other neutral grassland	-	-	Ant hills present of ok size. Part of skylark stewardship. Indicators of calcareous tendencies - rough hawkbit and wild carrot. About 10% bare earth. Chicory locally abundant to southern aspect. Area overall does not reach priority habitat, but it has pockets which are species rich. Looks to be established within last 10 years, with management could be priority habitat.	Rough hawkbit ( <i>Leontodon hispidus</i> ) R, wild carrot ( <i>Daucus carota</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) O, red fescue ( <i>Festuca rubra</i> ) O, yellow oat-grass ( <i>Trisetum flavescens</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, hop trefoil ( <i>Trifolium campestre</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) O, creeping cinquefoil ( <i>Potentilla reptans</i> ) O, oxeye daisy ( <i>Leucanthemum vulgare</i> ) O, creeping bent ( <i>Agrostis stolonifera</i> ), common bent ( <i>Agrostis capillaris</i> ) O, daisy ( <i>Bellis perennis</i> ) O, red clover ( <i>Trifolium pratense</i> ) O, smooth meadow-grass ( <i>Poa pratensis</i> ) R, common knapweed ( <i>Centaurea nigra</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) LA, pyramidal orchid ( <i>Anacamptis pyramidalis</i> ), chicory ( <i>Cichorium intybus</i> ) LA, common centaury ( <i>Centaurium erythraea</i> ) O, common agrimony ( <i>Agrimonia eupatoria</i> ) O.
G	H_A_974	g4 - Modified grassland	-	-	Modified grassland of Italian rye	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_975	w1d - Wet woodland	-	HPI	Continuation of wet woodland. Canopy mature crack willows and ash. Quite an open canopy. Understory of over-mature hazel. Ground very wet and lack of management apart from were used for pheasants. Ditch by grassland to north which water drains into. Mature specimen coppice ash ( <i>Fraxinus excelsior</i> ) with veteran features present.	Marsh thistle ( <i>Cirsium palustre</i> ) R, hedge woundwort ( <i>Stachys sylvatica</i> ) R, great horsetail ( <i>Equisetum telmateia</i> ) O, common nettle ( <i>Urtica dioica</i> ) F, dog's mercury ( <i>Mercurialis perennis</i> ) O, wood avens ( <i>Geum urbanum</i> ) R, water mint ( <i>Mentha aquatica</i> ) O, agrimony ( <i>Agrimonia eupatoria</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) F, opposite-leaved golden-saxifrage ( <i>Chrysosplenium oppositifolium</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) LA, ash ( <i>Fraxinus excelsior</i> ) F, crack-willow ( <i>Salix fragilis</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) F, hazel ( <i>Corylus avellana</i> ) C O, hawthorn ( <i>Crataegus monogyna</i> ) O, grey willow ( <i>Salix cinerea</i> ) LA.
G	H_A_976	g1c - Bracken	Scattered bracken	-	-	Bracken ( <i>Pteridium aquilinum</i> )
G	H_A_977	w1g7 - Other broadleaved woodland types	Plantation	-	Plantation, evenly aged and placed. Canopy oak, Norway spruce, silver birch, sweet chestnut. Two glades present in centre. Previous farm use likely pheasants keeping.	Pedunculate oak ( <i>Quercus robur</i> ) A, Norway spruce ( <i>Picea abies</i> ) F, silver birch ( <i>Betula pendula</i> ) O, sweet chestnut ( <i>Castanea sativa</i> ) R, field maple ( <i>Acer campestre</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, rose ( <i>Rosa sp.</i> ) R.
G	H_A_978	g4 - Modified grassland	-	-	Recently mown, leaving cuttings on the lawn. Likely to be cut repeatedly as part of garden management. Overall lacks diversity. Low distinctiveness.	Yorkshire fog ( <i>Holcus lanatus</i> ) A, rough meadow-grass ( <i>Poa trivialis</i> ) F, creeping buttercup ( <i>Ranunculus repens</i> ) F, creeping bent ( <i>Agrostis stolonifera</i> ) F.
G	H_A_979	g4 - Modified grassland	Cattle grazed	-	Bull in field so did not enter. Bare ground damage up to 10%.	False oat grass ( <i>Arrhenatherum elatius</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) F, white clover ( <i>Trifolium repens</i> ) F, common nettle ( <i>Urtica dioica</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ).
G	H_A_980	w1d - Wet woodland	-	HPI	Low topographical area, boggy under foot. Area looks to have been left for a while with no obvious management or pathways apart from one bridge but that looks rotting and old. Very dense vegetation.	Enchanter's-nightshade ( <i>Circaeа lutetiana</i> ) R, common nettle ( <i>Urtica dioica</i> ) A, dog's mercury ( <i>Mercurialis perennis</i> ) O, great horsetail ( <i>Equisetum telmateia</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, yellow iris ( <i>Iris pseudacorus</i> ) R, false brome ( <i>Brachypodium sylvaticum</i> ) R, red campion ( <i>Silene dioica</i> ) R, hart's-tongue thyme-moss ( <i>Plagiomnium undulatum</i> ) R, elder ( <i>Sambucus nigra</i> ) R, alder ( <i>Alnus glutinosa</i> ) O, grey willow ( <i>Salix cinerea</i> ) A, variegated yellow archangel ( <i>Lamiastrum galeobdolon</i> subsp. <i>argentatum</i> ) O
G	H_A_981	g3c5 - Arrhenatherum neutral grassland	-	-	More bramble scrub present on southern part of grassland. Topography goes uphill towards south. Hard rush in wet seepage with marsh thistle, water mint and soft rush.	False oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) O, great horsetail ( <i>Equisetum telmateia</i> ) O becoming A to east, smooth tare ( <i>Vicia tetrasperma</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) O, hemp-nettle ( <i>Galeopsis sp.</i> ) R, wood dock ( <i>Rumex sanguineus</i> ) R, hard-rush ( <i>Juncus inflexus</i> ) R, soft-rush ( <i>Juncus effusus</i> ), marsh thistle ( <i>Cirsium palustre</i> ), water mint ( <i>Mentha aquatica</i> ) R, cleavers ( <i>Galium aparine</i> ) R, hedge woundwort ( <i>Stachys sylvatica</i> ) R, common fleabane ( <i>Pulicaria dysenterica</i> ) R, lesser stitchwort ( <i>Stellaria graminea</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) O, creeping cinquefoil ( <i>Potentilla reptans</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, common bent ( <i>Agrostis capillaris</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, red fescue ( <i>Festuca rubra</i> ) O, yarrow ( <i>Achillea millefolium</i> ) R, upright hedge-parsley ( <i>Torilis japonica</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, common nettle ( <i>Urtica dioica</i> ) O, bramble ( <i>Rubus fruticosus</i> )
G	H_A_982	g1c - Bracken	Scattered bracken	-	-	Bracken ( <i>Pteridium aquilinum</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_983		Anthills. Woodland open space.	-	Ant hills present.	Common centaury ( <i>Centaurium erythraea</i> ) O becoming LA to west, red-stemmed feather-moss ( <i>Pleurozium schreberi</i> ) O, creeping cinquefoil ( <i>Potentilla reptans</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) LA, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) O, common mouse-ear ( <i>Cerastium fontanum</i> ) R, scarlet pimpernel ( <i>Anagallis arvensis</i> ) R, dove's-foot crane's-bill ( <i>Geranium molle</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) R, creeping bent ( <i>Agrostis stolonifera</i> ) O
G	H_A_984	w1g7 - Other broadleaved woodland types	-	-	Woodland on steep topography heading down approx. 10m from south to north. Possible boundary feature as mature pedunculate oak, coppice hazel and different species present.	Pedunculate oak ( <i>Quercus robur</i> ) M, wild cherry ( <i>Prunus avium</i> ) M O R, ash ( <i>Fraxinus excelsior</i> ) M O, sweet chestnut ( <i>Castanea sativa</i> ) R, wych elm ( <i>Ulmus glabra</i> ) R, holly ( <i>Ilex aquifolium</i> ) M O, elder ( <i>Sambucus nigra</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R, hazel ( <i>Corylus avellana</i> ) C F, English elm ( <i>Ulmus procera</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, common nettle ( <i>Urtica dioica</i> ) LA, ground ivy ( <i>Glechoma hederacea</i> ) LA, dog's mercury ( <i>Mercurialis perennis</i> ) O, primrose ( <i>Primula vulgaris</i> ) O, variegated yellow archangel ( <i>Lamiastrum galeobdolon</i> subsp. <i>argentatum</i> ) LA.
G	H_A_985	g3c - Other neutral grassland	-	-	Current managed leaving 5 strips of vegetation maybe under stewardship. Strip all the way round along hedges.	Common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) O, common fleabane ( <i>Pulicaria dysenterica</i> ) O, white clover ( <i>Trifolium repens</i> ) O, red clover ( <i>Trifolium pratense</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) A, common bent ( <i>Agrostis capillaris</i> ) O, bristly ox-tongue ( <i>Picris echioides</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) O, cat's-ear ( <i>Hypochaeris radicata</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) O, wood dock ( <i>Rumex sanguineus</i> ) R, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) R, rough hawkbit ( <i>Leontodon hispidus</i> ) R, smooth tare ( <i>Vicia tetrasperma</i> ) R, curled dock ( <i>Rumex crispus</i> ) R, scarlet pimpernel ( <i>Anagallis arvensis</i> ) R, hedge mustard ( <i>Sisymbrium officinale</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R
G	H_A_986	g1c - Bracken	Scattered bracken	-		Bracken ( <i>Pteridium aquilinum</i> )
G	H_A_987	g3c - Other neutral grassland	-	-	Blackthorn saplings. Indian balsam present north-western corner.	False oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, smooth tare ( <i>Vicia tetrasperma</i> ) F, great horsetail ( <i>Equisetum telmateia</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) O, angelica ( <i>Angelica sylvestris</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) R, creeping bent ( <i>Agrostis stolonifera</i> ) O, common bent ( <i>Agrostis capillaris</i> ) O, rough meadow-grass ( <i>Poa trivialis</i> ) O, smooth meadow-grass ( <i>Poa pratensis</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, greater stitchwort ( <i>Stellaria holostea</i> ) R, common nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) R, upright hedge-parsley ( <i>Torilis japonica</i> ) R, tufted vetch ( <i>Vicia cracca</i> ) O, lesser stitchwort ( <i>Stellaria graminea</i> ) R, Indian balsam ( <i>Impatiens glandulifera</i> )
G	H_A_988	g3c5 - Arrhenatherum neutral grassland	Other grazed	-	Rabbit grazed with bare soil patches. On top of hill heading down northwards.	False oat grass ( <i>Arrhenatherum elatius</i> ) A, common ragwort ( <i>Senecio jacobaea</i> ) LA, grey sedge ( <i>Carex divulsa</i> subsp. <i>divulsa</i> ) O, wild marjoram ( <i>Origanum vulgare</i> ) LA, St John's-wort ( <i>Hypericum sp.</i> ) F, white campion ( <i>Silene latifolia</i> ) R, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) LA, musk thistle ( <i>Carduus nutans</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) O, fine-leaved sandwort ( <i>Minuartia hybrida</i> ) R, common chickweed ( <i>Stellaria media</i> ) R, common bent ( <i>Agrostis capillaris</i> ) O, tufted hair grass ( <i>Deschampsia cespitosa</i> ).

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_989	g3c - Other neutral grassland	Scattered bracken	-	-	False oat grass ( <i>Arrhenatherum elatius</i> ) A, angelica ( <i>Angelica sylvestris</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, meadow vetchling ( <i>Lathyrus pratensis</i> ) F, tare ( <i>Vicia</i> sp.), lesser stitchwort ( <i>Stellaria graminea</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) R, marsh thistle ( <i>Cirsium palustre</i> ) R, common nettle ( <i>Urtica dioica</i> ) O, hairy tare ( <i>Vicia hirsuta</i> ) O, common fleabane ( <i>Pulicaria dysenterica</i> ) O, hard-rush ( <i>Juncus inflexus</i> ) R, great horsetail ( <i>Equisetum telmateia</i> ) O, wood dock ( <i>Rumex sanguineus</i> ) R, hoary willowherb ( <i>Epilobium parviflorum</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) R
G	H_A_990	w1d - Wet woodland	-	HPI	Indian balsam present in small stream on western aspect flowing north to glade in woodland, slow flow less than 3cm depth. Stream northern side flowing west to east. Further ditch to south side. Woodland bit to south possible spring but access limited. Embankment with dryer habitat to south boundary of grassland and western boundary including walnut understory coppiced hazel. Brooklime in slow stream. Common bent and primrose on dry banks.	Alder ( <i>Alnus glutinosa</i> ) O M, Indian balsam ( <i>Impatiens glandulifera</i> ), walnut ( <i>Juglans regia</i> ) R, grey willow ( <i>Salix cinerea</i> ) O, goat willow ( <i>Salix caprea</i> ), pedunculate oak ( <i>Quercus robur</i> ), alder ( <i>Alnus glutinosa</i> ), grey willow ( <i>Salix cinerea</i> ), crab apple ( <i>Malus sylvestris</i> ) R, hazel ( <i>Corylus avellana</i> ) C O, field bindweed ( <i>Convolvulus arvensis</i> ), marsh thistle ( <i>Cirsium palustre</i> ), hemp-agrimony ( <i>Eupatorium cannabinum</i> ), water mint ( <i>Mentha aquatica</i> ) F, great horsetail ( <i>Equisetum telmateia</i> ) O, hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), holly ( <i>Ilex aquifolium</i> ) R, water forget-me-not ( <i>Myosotis scorpioides</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ) O, soft-rush ( <i>Juncus effusus</i> ) R, yellow iris ( <i>Iris pseudacorus</i> ) R, reed sweet-grass ( <i>Glyceria maxima</i> ) O, brooklime ( <i>Veronica beccabunga</i> ) O, distant sedge ( <i>Carex distans</i> ) LA, creeping buttercup ( <i>Ranunculus repens</i> ) O, herb-Robert ( <i>Geranium robertianum</i> ) R, common nettle ( <i>Urtica dioica</i> ) F, jointed rush ( <i>Juncus articulatus</i> ) O, primrose ( <i>Primula vulgaris</i> ) O, cleavers ( <i>Galium aparine</i> ) R, red campion ( <i>Silene dioica</i> ) R, hedge woundwort ( <i>Stachys sylvatica</i> ), ground ivy ( <i>Glechoma hederacea</i> ) O, common bent ( <i>Agrostis capillaris</i> ) F, viola ( <i>Viola</i> sp.) R, cock's-foot ( <i>Dactylis glomerata</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, hoary willowherb ( <i>Epilobium parviflorum</i> ) R
G	H_A_991	r1a6 - Other eutrophic standing waters	-	HPI	Some limitations algae and duckweed. Grey willow on northern side. Could be fed from possible springs.	Soft-rush ( <i>Juncus effusus</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) O, water-cress ( <i>Rorippa nasturtium-aquaticum</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) R, water mint ( <i>Mentha aquatica</i> ) O, great horsetail ( <i>Equisetum telmateia</i> ) O, hemp-agrimony ( <i>Eupatorium cannabinum</i> ) R, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) F.
G	H_A_992	g3c6 - Lolium-Cynosurus neutral grassland	Cattle grazed	-	Common cudweed present. Cattle grazed. Mosaic between g3c6 in higher areas with G4 in wetter rushes. Dominated g3c6, transitioning to G4 where density stops grazing.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, common ragwort ( <i>Senecio jacobaea</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) F, greater plantain ( <i>Plantago major</i> ) O, crested dog's-tail ( <i>Cynosurus cristatus</i> ) A, downy oat-grass ( <i>Avenula pubescens</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) A, common bent ( <i>Agrostis capillaris</i> ) A, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, dandelion ( <i>Taraxacum</i> agg.) R, common stork's-bill ( <i>Erodium cicutarium</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) R, yellow oat-grass ( <i>Trisetum flavescens</i> ) F, cat's-ear ( <i>Hypochaeris radicata</i> ) O, Timothy ( <i>Phleum pratense</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, hop trefoil ( <i>Trifolium campestre</i> ) R, red fescue ( <i>Festuca rubra</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) R, bugloss ( <i>Anchusa arvensis</i> ) R, hare's-foot clover ( <i>Trifolium arvense</i> ) R, goat's-beard ( <i>Tragopogon pratensis</i> ) R, soft brome ( <i>Bromus mollis</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_993	g3c6 - Lolium-Cynosurus neutral grassland	Cattle grazed	-	-	White clover ( <i>Trifolium repens</i> ) F, wood dock ( <i>Rumex sanguineus</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) F, ribwort plantain ( <i>Plantago lanceolata</i> ) F, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) A, crested dog's-tail ( <i>Cynosurus cristatus</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) O, common mouse-ear ( <i>Cerastium fontanum</i> ) O, selfheal ( <i>Prunella vulgaris</i> ) O, common centaury ( <i>Centaurium erythraea</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) F, meadow buttercup ( <i>Ranunculus acris</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, pale Persicaria ( <i>Polygonum lapathifolia</i> ) O, dandelion ( <i>Taraxacum agg.</i> ) R, marsh thistle ( <i>Cirsium palustre</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, common bent ( <i>Agrostis capillaris</i> ) F, common fleabane ( <i>Pulicaria dysenterica</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) R, bristly ox-tongue ( <i>Picris echioides</i> ) R, procumbent pearlwort ( <i>Sagina procumbens</i> ) R, Agrimony ( <i>Agrimonia eupatoria</i> ) R, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) R, lady's bedstraw ( <i>Galium verum</i> ) R
G	H_A_994	g4 - Modified grassland	Sheep grazed	-	Lacking 9 species per average m <sup>2</sup> . Young bramble starting to emerge. Variety species found across area, but majority are of rare distribution.	Creeping thistle ( <i>Cirsium arvense</i> ) R, greater plantain ( <i>Plantago major</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) A, ribwort plantain ( <i>Plantago lanceolata</i> ) A, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, white clover ( <i>Trifolium repens</i> ) F, common nettle ( <i>Urtica dioica</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) F, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, common bent ( <i>Agrostis capillaris</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, soft brome ( <i>Bromus mollis</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, wood dock ( <i>Rumex sanguineus</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, common mouse-ear ( <i>Cerastium fontanum</i> ) O, cat's-ear ( <i>Hypochaeris radicata</i> ) F, black medick ( <i>Medicago lupulina</i> ) R, bramble ( <i>Rubus fruticosus</i> ) F, dandelion ( <i>Taraxacum agg.</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) F, meadow buttercup ( <i>Ranunculus acris</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) R, smooth tare ( <i>Vicia tetrasperma</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R
G	H_A_995	w1f7 - Other Lowland mixed deciduous woodland	Earth bank	-	Mature oak canopy. Evidence of ditch present in wood on either side which could be old boundary feature.	Pedunculate oak ( <i>Quercus robur</i> ) M O, ash ( <i>Fraxinus excelsior</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, rose ( <i>Rosa sp.</i> ) R, elder ( <i>Sambucus nigra</i> ) O, English elm ( <i>Ulmus procera</i> ) R Y, hawthorn ( <i>Crataegus monogyna</i> ) R.
G	H_A_996	g4 - Modified grassland	Veteran trees. Sheep grazed. Ruderal. Improved ground.	-	Three veteran fruit trees. If left unmanned would be dominated by nettles. Lack richness per average m <sup>2</sup> . Historic used could be for chickens due to hen house present.	Common nettle ( <i>Urtica dioica</i> ) A, broad-leaved dock ( <i>Rumex obtusifolius</i> ) O, herb-Robert ( <i>Geranium robertianum</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, hedge woundwort ( <i>Stachys sylvatica</i> ) R, wood avens ( <i>Geum urbanum</i> ) R, common knotgrass ( <i>Polygonum aviculare</i> ) R, garlic mustard ( <i>Alliaria petiolata</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, lords and ladies ( <i>Arum maculatum</i> ) R, greater plantain ( <i>Plantago major</i> ) O, rough meadow-grass ( <i>Poa trivialis</i> ) F, creeping buttercup ( <i>Ranunculus repens</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) R, shepherd's-purse ( <i>Capsella bursa-pastoris</i> ) R, fat-hen ( <i>Chenopodium album</i> ) R, lesser burdock ( <i>Actium minus</i> ) R, hedge mustard ( <i>Sisymbrium officinale</i> ) R, eagle fern ( <i>Pteridium aquilinum</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, common field speedwell ( <i>Veronica persica</i> ) R, field madder ( <i>Sherardia arvensis</i> ) R, pineapple weed ( <i>Matricaria discoidea</i> ) R, dandelion ( <i>Taraxacum agg.</i> ) R, bramble ( <i>Rubus fruticosus</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H_A_997	g4 - Modified grassland	Horse grazed	-	Horse paddock. Species poor. Species poor.	Cock's-foot ( <i>Dactylis glomerata</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) A, yarrow ( <i>Achillea millefolium</i> ) F, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, white clover ( <i>Trifolium repens</i> ) A, greater plantain ( <i>Plantago major</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) O, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, common nettle ( <i>Urtica dioica</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) A, Timothy ( <i>Phleum pratense</i> ) F, bristly oxtongue ( <i>Picris echioides</i> ) R
G	H3_1026	c1c - Cereal Crops	-	-	Arable	-
G	H3_1043	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1068	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1075	g1a - Lowland dry acid grassland	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1079	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1081	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1090	c1c - Cereal Crops	-	-	Arable	-
G	H3_1091	c1c - Cereal Crops	-	-	Arable	-
G	H3_1124	c1c - Cereal Crops	-	-	Arable	-
G	H3_1126	c1c - Cereal Crops	-	-	Arable	-
G	H3_1141	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1141	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1142	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1144	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_1149	c1c - Cereal Crops	-	-	Arable	-
G	H3_943	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_945	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_947	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_950	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	H3_953	g3c - Other neutral grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_954	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_955	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	H3_959	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_853	c1c6 - Game bird mix fields	-	-	This is maize probably planted for pheasants, supported by the presence of bird feeders at the top of the field.	-
G	HA3_867	H – heathland scrub	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_907	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_908	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_909	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_910	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_911	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_913	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_914	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HA3_915	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_108	w1d - Wet woodland	Veteran tree	HPI	Ash die-back on mature trees. More ancient woodland indicators to western side with certain trees. Woodland along stream, mature trees and canopy cover limiting light to ground. Lots of bear earth and where vegetation present dominated by nettles. Veteran oak. Lacks wet woodland ground flora.	Ash ( <i>Fraxinus excelsior</i> ), common nettle ( <i>Urtica dioica</i> ) D, grey willow ( <i>Salix cinerea</i> ), willow ( <i>Salix</i> sp.), alder ( <i>Alnus glutinosa</i> ), hazel ( <i>Corylus avellana</i> ), field maple ( <i>Acer campestre</i> ), blackthorn ( <i>Prunus spinosa</i> ), holly ( <i>Ilex aquifolium</i> ), ash ( <i>Fraxinus excelsior</i> ), dogwood ( <i>Cornus sanguinea</i> ), pedunculate oak ( <i>Quercus robur</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ), common nettle ( <i>Urtica dioica</i> ) A, false brome ( <i>Brachypodium sylvaticum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), hedge woundwort ( <i>Stachys sylvatica</i> ), great horsetail ( <i>Equisetum telmateia</i> ), laburnum ( <i>Laburnum anagyroides</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_115	g3c - Other neutral grassland	Anthills	-	16 tall herb community abundant to make grass occasional. 140 Ant hills present. High forbs cover. Land left by farmer as it is often too wet for grazing. Mentioned rare orchids present but none seen during survey, he did not mention which ones. Likely to be wet throughout the year.	Hairy tare ( <i>Vicia hirsuta</i> ) D, pale Persicaria ( <i>Polygonum lapathifolia</i> ) A, creeping thistle ( <i>Cirsium arvense</i> ) F, marsh thistle ( <i>Cirsium palustre</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) F, common fleabane ( <i>Pulicaria dysenterica</i> ) F, hairy sedge ( <i>Carex hirta</i> ) R, white clover ( <i>Trifolium repens</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) O, red barista ( <i>Odontites verna</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, soft-rush ( <i>Juncus effusus</i> ) O, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) R wood dock ( <i>Rumex sanguineus</i> ) R crested dog's-tail ( <i>Cynosurus cristatus</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) F, wild carrot ( <i>Daucus carota</i> ) F common centaury ( <i>Centaurium erythraea</i> ) R, Agrimony ( <i>Agrimonia eupatoria</i> ) R, black medick ( <i>Medicago lupulina</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) R, meadow buttercup ( <i>Ranunculus acris</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) F, perennial sow-thistle ( <i>Sonchus arvensis</i> ) R, goat's-beard ( <i>Tragopogon pratensis</i> ) R, creeping cinquefoil ( <i>Potentilla reptans</i> ) R, perforate St John's-wort ( <i>Hypericum perforatum</i> ) R, common knapweed ( <i>Centaurea nigra</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R
G	HL_116	g3c6 - Lolium-Cynosurus neutral grassland	Cattle grazed.	-	Mosaic between Lolium-Cynosurus neutral grassland in higher areas with improved grassland in wetter rushes. Dominated by Lolium-Cynosurus neutral grassland, transitioning to improved grassland where dense stops grazing.	Perennial rye-grass ( <i>Lolium perenne</i> ) D, common ragwort ( <i>Senecio jacobaea</i> ) F, spear thistle ( <i>Cirsium vulgare</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) F, greater plantain ( <i>Plantago major</i> ) O, crested dog's-tail ( <i>Cynosurus cristatus</i> ) A, downy oat-grass ( <i>Avenula pubescens</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) A, common bent ( <i>Agrostis capillaris</i> ) A, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O dandelion ( <i>Taraxacum agg.</i> ) R, common stork's-bill ( <i>Erodium cicutarium</i> ) R, common cudweed ( <i>Filago vulgaris</i> ) R, yellow oat-grass ( <i>Trisetum flavescens</i> ) F, cat's-ear ( <i>Hypochaeris radicata</i> ) O, Timothy ( <i>Phleum pratense</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, hop trefoil ( <i>Trifolium campestre</i> ) R, red fescue ( <i>Festuca rubra</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) R, bugloss ( <i>Anchusa arvensis</i> ) R, hare's-foot clover ( <i>Trifolium arvensis</i> ) R, goat's-beard ( <i>Tragopogon pratensis</i> ) R, soft brome ( <i>Bromus mollis</i> ) R
G	G_T_1147	w – woodland and forest	-	-	Woodland either side of the road, UKHab derived from aerial imagery.	-
G	G_T_1148	w – woodland and forest	-	-	Woodland either side of the road, UKHab derived from aerial imagery.	-
G	HL_117	g3c6 - Lolium-Cynosurus neutral grassland	Cattle grazed	-	-	White clover ( <i>Trifolium repens</i> ) F, wood dock ( <i>Rumex sanguineus</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) F, ribwort plantain ( <i>Plantago lanceolata</i> ) F, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) A, crested dog's-tail ( <i>Cynosurus cristatus</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) O, common mouse-ear ( <i>Cerastium fontanum</i> ) O, selfheal ( <i>Prunella vulgaris</i> ) O, common centaury ( <i>Centaurium erythraea</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) F, meadow buttercup ( <i>Ranunculus acris</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, pale Persicaria ( <i>Polygonum lapathifolia</i> ) O, dandelion ( <i>Taraxacum agg.</i> ) R, marsh thistle ( <i>Cirsium palustre</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, common bent ( <i>Agrostis capillaris</i> ) F, common fleabane ( <i>Pulicaria dysenterica</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) R, procumbent pearlwort ( <i>Sagina procumbens</i> ) R, Agrimony ( <i>Agrimonia eupatoria</i> ) R, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) R lady's bedstraw ( <i>Galium verum</i> ) R
G	HL_118	w1g7 - Other broadleaved woodland types	Plantation	-	Young alder of similar age, used for shelter for sheep. Lots of bare ground.	Alder ( <i>Alnus glutinosa</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), common nettle ( <i>Urtica dioica</i> ), Yorkshire fog ( <i>Holcus lanatus</i> )

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_120	h3a - Blackthorn scrub	Earth bank	-	Hazel and blackthorn where old hedgerow could have once been. Indicators of bank where hedge used to go through but completely overtaken by blackthorn. Ground flora mainly bare. Lots of dead wood.	Blackthorn ( <i>Prunus spinosa</i> ) D, hazel ( <i>Corylus avellana</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, common nettle ( <i>Urtica dioica</i> ) R
G	HL_121	w1f7 - Other Lowland mixed deciduous woodland	Earth bank	-	Evidence of ditch present in wood on either side which could be old boundary feature.	Pedunculate oak ( <i>Quercus robur</i> ) M O, ash ( <i>Fraxinus excelsior</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, rose ( <i>Rosa sp.</i> ) R, elder ( <i>Sambucus nigra</i> ) O, English elm ( <i>Ulmus procera</i> ) R Y, hawthorn ( <i>Crataegus monogyna</i> ) R.
G	HL_123	w1g7 - Other broadleaved woodland types	Plantation	-	Young alder of similar age, used for shelter for sheep. Lots of bare ground. Lot of bare ground. Some recently cut and replanted.	Alder ( <i>Alnus glutinosa</i> ), perennial rye-grass ( <i>Lolium perenne</i> ), common nettle ( <i>Urtica dioica</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), bramble ( <i>Rubus fruticosus</i> ), hawthorn ( <i>Crataegus monogyna</i> )
G	HL_124	g4 - Modified grassland	Horse grazed	-	Horse paddock. Species poor.	Cock's-foot ( <i>Dactylis glomerata</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) A, yarrow ( <i>Achillea millefolium</i> ) F, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R, creeping buttercup ( <i>Ranunculus repens</i> ) O, white clover ( <i>Trifolium repens</i> ) A, greater plantain ( <i>Plantago major</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) O, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, common nettle ( <i>Urtica dioica</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) A, Timothy ( <i>Phleum pratense</i> ) F, bristly oxtongue ( <i>Picris echioides</i> ) R
G	HL_129	h3h - Mixed scrub	-	-	Young scrub likely kept low due to pylon. Very dense, no obvious ground flora as no light. All similar age	Blackthorn ( <i>Prunus spinosa</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R, elm ( <i>Ulmus sp.</i> )
G	HL_130	g1c - Bracken	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_132	g4 - Modified grassland	Cattle grazed	-	Field done by hedge line as bull in field.	Smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) A, common bent ( <i>Agrostis capillaris</i> ) A, common centaury ( <i>Centaurium erythraea</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) A, false oat grass ( <i>Arrhenatherum elatius</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O,
G	HL_135	g3c6 - <i>Lolium-Cynosurus</i> neutral grassland	-	-	Very similar to the field to west however has been cut to approx. 10-20 cm. Similar species present, but slightly less per m <sup>2</sup> but still above 9 per m <sup>2</sup> . Abundances on dominant species the same. Sandy well drained soil.	Smooth hawk's-beard ( <i>Crepis capillaris</i> ) F, common cudweed ( <i>Filago vulgaris</i> ) F, yellow oat-grass ( <i>Trisetum flavescens</i> ) F, common mouse-ear ( <i>Cerastium fontanum</i> ) R, crested dog's-tail ( <i>Cynosurus cristatus</i> ) F, white clover ( <i>Trifolium repens</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) F, common ragwort ( <i>Senecio jacobaea</i> ) F, hop trefoil ( <i>Trifolium campestre</i> ) R, common bent ( <i>Agrostis capillaris</i> ) A, Timothy ( <i>Phleum pratense</i> ) O, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) O, Yorkshire fog ( <i>Holcus lanatus</i> ) A, spear thistle ( <i>Cirsium vulgare</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) O, selfheal ( <i>Prunella vulgaris</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) R, grass ( <i>Vulpia sp.</i> ) R, red fescue ( <i>Festuca rubra</i> ) R, sweet vernal-grass ( <i>Anthoxanthum odoratum</i> ) R, common nettle ( <i>Urtica dioica</i> ) R, common calamint ( <i>Calamintha ascendens</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, fairy flax ( <i>Linum catharticum</i> ) O musk thistle ( <i>Carduus nutans</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_138	g3c6 - Lolium-Cynosurus neutral grassland	Anthills present	-	Informed by farmer to have medieval ploughing marks in field. Likely grazed intermittently but no evidence of this being so recently. Paths have been cut through field and rest likely cut as absence of tufts etc. Topography varies with valleys going down near 10-15m approx. but looks to be uniform in species abundance with false oat grass and nettle becoming more common. Small ant hills present. Cudweed present.	Smooth hawk's-beard ( <i>Crepis capillaris</i> ) F, common cudweed ( <i>Filago vulgaris</i> ) O, yellow oat-grass ( <i>Trisetum flavescens</i> ) F, common mouse-ear ( <i>Cerastium fontanum</i> ) R, crested dog's-tail ( <i>Cynosurus cristatus</i> ) F, white clover ( <i>Trifolium repens</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) F, common ragwort ( <i>Senecio jacobaea</i> ) F, hop trefoil ( <i>Trifolium campestre</i> ) R, common bent ( <i>Agrostis capillaris</i> ) A, Timothy ( <i>Phleum pratense</i> ) O, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, downy oat-grass ( <i>Avenula pubescens</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) O, yarrow ( <i>Achillea millefolium</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) A, spear thistle ( <i>Cirsium vulgare</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) O, selfheal ( <i>Prunella vulgaris</i> ) O, dove's-foot crane's-bill ( <i>Geranium molle</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) O, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) R, perennial rye-grass ( <i>Lolium perenne</i> ) R, grass ( <i>Vulpia</i> sp.) R, creeping buttercup ( <i>Ranunculus repens</i> ) R, hairy tare ( <i>Vicia hirsuta</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) O
G	HL_139	h3h - Mixed scrub	-	-	Likely cut for power lines. Planted in places. Nettle abundant in areas where scrub not present.	Pedunculate oak ( <i>Quercus robur</i> ) O Y, elm ( <i>Ulmus</i> sp.) O Y, blackthorn ( <i>Prunus spinosa</i> ) R, elder ( <i>Sambucus nigra</i> ) R, dog-rose ( <i>Rosa canina</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, hazel ( <i>Corylus avellana</i> ) O, bramble ( <i>Rubus fruticosus</i> ) F, common nettle ( <i>Urtica dioica</i> ) A, wild marjoram ( <i>Origanum vulgare</i> ) R, dove's-foot crane's-bill ( <i>Geranium molle</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, false oat grass ( <i>Arrhenatherum elatius</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) LA, common cudweed ( <i>Filago vulgaris</i> ) O, common bent ( <i>Agrostis capillaris</i> ), wood dock ( <i>Rumex sanguineus</i> ) R, common mouse-ear ( <i>Cerastium fontanum</i> ) R
G	HL_152	c1d8 - Other non-cereal crops	-	-	Hilly sloping field from the east down to west. Arable crop of peas. Arable weeds. Native species present at the west edge of field.	Field pansy ( <i>Viola arvensis</i> ) O, black-bindweed ( <i>Fallopia convolvulus</i> ) O, common mallow ( <i>Malva sylvestris</i> ) R, common poppy ( <i>Papaver rhoeas</i> ) R, common field speedwell ( <i>Veronica persica</i> ), white campion ( <i>Silene latifolia</i> ) R, knotgrass ( <i>Polygonum aviculare</i> ) O, curled dock ( <i>Rumex crispus</i> ), prickly lettuce ( <i>Lactuca serriola</i> ) R, great willowherb ( <i>Epilobium hirsutum</i> ) F, toad rush ( <i>Juncus bufonius</i> ) R, scarlet pimpernel ( <i>Anagallis arvensis</i> ) R, slender meadow foxtail ( <i>Alopecurus myosuroides</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) O, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) F, cockspur grass ( <i>Echinochloa crus-galli</i> ) R, greater plantain ( <i>Plantago major</i> ) O, curled dock ( <i>Rumex crispus</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) F, knotgrass ( <i>Polygonum aviculare</i> ) O, marsh sow-thistle ( <i>Sonchus asper</i> ) O, rough meadow-grass ( <i>Poa trivialis</i> ) O, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, field horsetail ( <i>Equisetum arvense</i> ) F, prickly lettuce ( <i>Lactuca serriola</i> ) R, red clover ( <i>Trifolium pratense</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, compact rush ( <i>Juncus glomeratus</i> ) O, smooth tare ( <i>Vicia tetrasperma</i> ) O, common cudweed ( <i>Filago vulgaris</i> ) R, bugloss ( <i>Anchusa arvensis</i> ) R,
G	HL_153	c1a7 - Arable margins cultivated annually with an annual flora	-	-	Rutted area, 10% bare ground. Mixture of grasses and herbaceous plants. Arable weeds dominated by horsetail in some areas and cudweed present in NE corner.	Field horsetail ( <i>Equisetum arvense</i> ) D, Common cudweed ( <i>Filago vulgaris</i> ) O.

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_154	w1h5 - Other woodland; mixed; mainly broadleaved	Plantation.	-	Woodland planted in rows (plantation), no understory, trees less than 50yrs old. Canopy still open, no age structure. Herb layer dominated by common nettles with some elder (photo included to show reference of herb layer height). pedunculate oak dominated with sweet chestnut and occasional wild cherry. Field maple cut down within woodland. Ancient woodland indicators present. Condition assessment attached separately - score of 23 gives poor condition.	Elder ( <i>Sambucus nigra</i> ) O, wild cherry ( <i>Prunus avium</i> ) O, field maple ( <i>Acer campestre</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) D, sweet chestnut ( <i>Castanea sativa</i> ) O, elder ( <i>Sambucus nigra</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, walnut ( <i>Juglans regia</i> ) R, lime ( <i>Tilia x europaea</i> ) R, common nettle ( <i>Urtica dioica</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, bramble ( <i>Rubus fruticosus</i> ) A
G	HL_157	g3c - Other neutral grassland	-	-	Over 8 species per metre squared. Long grass area. Only passed two out of five assessments = poor condition.	Whorled mint ( <i>Mentha x verticillata</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) F, bramble ( <i>Rubus fruticosus</i> ) R, prickly lettuce ( <i>Lactuca serriola</i> ) R, perennial sow-thistle ( <i>Sonchus arvensis</i> ) O, old man's beard ( <i>Clematis vitalba</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) O, crested dog's-tail ( <i>Cynosurus cristatus</i> ) R, common knapweed ( <i>Centaurea nigra</i> ) O, spear thistle ( <i>Cirsium vulgare</i> ) R, goat's-beard ( <i>Tragopogon pratensis</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) F, wild carrot ( <i>Daucus carota</i> ) R, Timothy ( <i>Phleum pratense</i> ) O.
G	HL_158	c1c6 - Game bird mix fields	-	-	This is maize probably planted for pheasants, supported by the presence of bird feeders at the top of the field.	-
G	HL_161	g3c5 - Arrhenatherum neutral grassland	-	-	Hilly, sloping grass land area next to ploughed land and maize areas, also bordering minty woodland. Cannot be easily classified. Relatively species rich. Moderate condition from assessment	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_162	w1f7 - Other Lowland mixed deciduous woodland	-	-	Various tree species present including medium sized oaks. Mainly dominated by field maple. Oak seedlings at the bottom. Links to woodland. Some areas too difficult to survey due to overgrown shrub layer. Dead badger found also. 15m gap between two parts of green lane woodland for vehicle entrance. Polygon of green lane extends further north than the buffer and connects to vegetation in gardens north. Condition assessment: At least 2 ages present (moderate), no herbivore damage (good), no invasive species (good), 5 or more native species (good), >80% of canopy trees and understory is native (good), no open space (poor), woodland regeneration is good, tree health is good, ancient woodland flora indicators present x4 (good), 2 or more stories present across plot (moderate), no veteran tree present (poor), not much dead wood - although couldn't be assessed in all areas (poor), woodland enrichment (moderate). Good condition.	Pedunculate oak ( <i>Quercus robur</i> ), field maple ( <i>Acer campestre</i> ) D, dog's mercury ( <i>Mercurialis perennis</i> ), common nettle ( <i>Urtica dioica</i> )
G	HL_163	c1c - Cereal crops	-	-	More gently sloping crop field bordered by grassland, houses and hedgerow trees.	-
G	HL_164	w1g - Other woodland; broadleaved	-	-	Woodland starts at bend of river away from willow field and borders the field to the west. Unable to get into woodland due to wire, and T59 plantation of single age group, no understory, no mature trees, herb layer dominated by nettles, approx. 40yrs old ash. Field maple is outside land boundary.	Elder ( <i>Sambucus nigra</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, willow ( <i>Salix</i> sp.) A, blackthorn ( <i>Prunus spinosa</i> ) O, hazel ( <i>Corylus avellana</i> ) O, common nettle ( <i>Urtica dioica</i> ), ash ( <i>Fraxinus excelsior</i> ), wild cherry ( <i>Prunus avium</i> ), hawthorn ( <i>Crataegus monogyna</i> ) O, field maple ( <i>Acer campestre</i> ) O
G	HL_165	w1h5 - Other woodland; mixed; mainly broadleaved	-	-	Part-open plantation of mixed species. Disused badger sett outside of the buffer. Moderate condition based on assessment.	Pedunculate oak ( <i>Quercus robur</i> ) A, sweet chestnut ( <i>Castanea sativa</i> ) F, wild cherry ( <i>Prunus avium</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, hazel ( <i>Corylus avellana</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, small-leaved lime ( <i>Tilia cordata</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, blue fleabane ( <i>Erigeron acer</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) D, Yorkshire fog ( <i>Holcus lanatus</i> ) A, false oat grass ( <i>Arrhenatherum elatius</i> ) O, black horehound ( <i>Ballota nigra</i> ) R, white campion ( <i>Silene latifolia</i> ) R, wild carrot ( <i>Daucus carota</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, perforate St John's-wort ( <i>Hypericum perforatum</i> ) R, common calamint ( <i>Calamintha ascendens</i> ) F, ploughman's-spikenard ( <i>Inula conyzae</i> ) R

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_166	g3c - Other neutral grassland	-	-	Grassy area next to woodland and planted maize used for game purposes. Species present in both fields are the same/very similar. Grassland was being cut at the time of surveying. Red kite present in area.	Perforate St John's-wort ( <i>Hypericum perforatum</i> ) O, cat's-ear ( <i>Hypochaeris radicata</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, creeping bent ( <i>Agrostis stolonifera</i> ) F, bramble ( <i>Rubus fruticosus</i> ) R, field bindweed ( <i>Convolvulus arvensis</i> ) R, Cock's-foot ( <i>Dactylis glomerata</i> ) A, common ragwort ( <i>Senecio jacobaea</i> ) F, common bent ( <i>Agrostis capillaris</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O, Agrimony ( <i>Agrimonia eupatoria</i> ) R, hop trefoil ( <i>Trifolium campestre</i> ) R, downy oat-grass ( <i>Avenula pubescens</i> ) O, common nettle ( <i>Urtica dioica</i> ) R, smooth hawk's-beard ( <i>Crepis capillaris</i> ) R, common sorrel ( <i>Rumex acetosa</i> ) R, yellow oat-grass ( <i>Trisetum flavescens</i> ) R, common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) F, scentless mayweed ( <i>Tripleurospermum inodorum</i> ) F, wild carrot ( <i>Daucus carota</i> ) F, red clover ( <i>Trifolium pratense</i> ) O, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) F, smooth tare ( <i>Vicia tetrasperma</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) F, bristly oxtongue ( <i>Picris echioides</i> ) R, oxeye daisy ( <i>Leucanthemum vulgare</i> ) A, goat's-beard ( <i>Tragopogon pratensis</i> ) O, blue fleabane ( <i>Erigeron acer</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R, yellow-wort ( <i>Blackstonia perfoliata</i> ) R, common calamint ( <i>Calamintha ascendens</i> ) R, old man's beard ( <i>Clematis vitalba</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, dandelion ( <i>Taraxacum agg.</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R,
G	HL_167	c1c6 - Game bird mix fields	-	-	This is maize probably planted for pheasants, supported by the presence of bird feeders at the top of the field.	-
G	HL_171	h3h - Mixed scrub	-	-	Over 5m wide, bordering two fields. Numerous dead elm, elm/elder, wormwood etc. Steep bank from one field to another. Passed 2/5 on condition assessment = poor condition. May have been an elm hedge at one point, elm still present but many dead. Unmanaged, very tall vegetation.	Elm ( <i>Ulmus sp.</i> ), elder ( <i>Sambucus nigra</i> ), wormwood ( <i>Artemisia absinthium</i> )
G	HL_174	c1c - Cereal crops	-	-	Hilly sloping wheat crop field. Ploughed recently, stubble ploughed in. Surrounded by maize and grassland.	-
G	HL_186	c1c6 - Game bird mix fields	-	-	Arable	-
G	HL_189	c1c6 - Game bird mix fields	-	-	Cockspur and millet planted for turtle dove habitat	-
G	HL_190	g3c5 - Arrhenatherum neutral grassland	-	-	Cut in swathes for partridge cover uncut to 1.5m high.	Cock's-foot ( <i>Dactylis glomerata</i> ) A, Agrimony ( <i>Agrimonia eupatoria</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, wood dock ( <i>Rumex sanguineus</i> ) O, old man's beard ( <i>Clematis vitalba</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) O, bramble ( <i>Rubus fruticosus</i> ) R, bristly oxtongue ( <i>Picris echioides</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) O, wild carrot ( <i>Daucus carota</i> ) F, blue fleabane ( <i>Erigeron acer</i> ) R, cat's-ear ( <i>Hypochaeris radicata</i> ) R, false brome ( <i>Brachypodium sylvaticum</i> ) O, common nettle ( <i>Urtica dioica</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) O, yellow-wort ( <i>Blackstonia perfoliata</i> ) O, oxeye daisy ( <i>Leucanthemum vulgare</i> ) F, dandelion ( <i>Taraxacum agg.</i> ) O, wild basil ( <i>Clinopodium vulgare</i> ) F, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, lesser burdock ( <i>Arctium minus</i> ) R, perforate St John's-wort ( <i>Hypericum perforatum</i> ) R, Yorkshire fog ( <i>Holcus lanatus</i> ) O
G	HL_191	c1c6 - Game bird mix fields	-	-	Maize for partridges	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_194	c1c6 - Game bird mix fields	-	-	Maize for game bird cover and food	-
G	HL_196	w1f7 - Other Lowland mixed deciduous woodland	-	-	Heavily used for game bird rearing. Native trees include field maple, hornbeam and oak. Significant disturbance of herb layer with large tracts of bare ground in upper drier areas and invasive bracken in lower areas, no regeneration seen, lower, wetter area with abundant by <i>Salix</i> species.	Field maple ( <i>Acer campestre</i> ) A, dogwood ( <i>Cornus sanguinea</i> ) F, wild cherry ( <i>Prunus avium</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) O, willow ( <i>Salix</i> sp.) O, hazel ( <i>Corylus avellana</i> ) O, elder ( <i>Sambucus nigra</i> ) O, hornbeam ( <i>Carpinus betulus</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, ash ( <i>Fraxinus excelsior</i> ) O, common nettle ( <i>Urtica dioica</i> ) A, eagle fern ( <i>Pteridium aquilinum</i> ) F, great horsetail ( <i>Equisetum telmateia</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) F, wood dock ( <i>Rumex sanguineus</i> ) O, meadow vetchling ( <i>Lathyrus pratensis</i> ) R, false oat grass ( <i>Arrhenatherum elatius</i> ) R, cock's-foot ( <i>Dactylis glomerata</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, Agrimony ( <i>Agrimonia eupatoria</i> ) R, broad-leaved dock ( <i>Rumex obtusifolius</i> ) R, creeping bent ( <i>Agrostis stolonifera</i> ) O, perennial rye-grass ( <i>Lolium perenne</i> ) O, wild angelica ( <i>Angelica sylvestris</i> ) R, false brome ( <i>Brachypodium sylvaticum</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) F
G	HL_197	w1f - Lowland mixed deciduous woodland	-	HPI	Broad-leaved woodland dominated by twenty-metre-high ash with six maturing oaks closest to pylon on the line of old track which probably had two hedges and outer edge is dominated by blackthorn. Very steep sloping in places stand of mature field maple with badger set possibly disused. Understorey hazel and hawthorn with evidence of hawthorn and holly new growth. On western boundary of polygon is open area dominated by willow and elder with high cover of common nettles, creeping thistle and some giant horsetail. Herb layer of Jazzy wood dominated by dog's mercury with abundant false wood brome.	Ash ( <i>Fraxinus excelsior</i> ), pedunculate oak ( <i>Quercus robur</i> ), blackthorn ( <i>Prunus spinosa</i> ) O, field maple ( <i>Acer campestre</i> ) A, holly ( <i>Ilex aquifolium</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) A, elder ( <i>Sambucus nigra</i> ) R, common nettle ( <i>Urtica dioica</i> ) D, creeping thistle ( <i>Cirsium arvense</i> ), dog's mercury ( <i>Mercurialis perennis</i> ) D, pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) D, hazel ( <i>Corylus avellana</i> ) F, dog-rose ( <i>Rosa canina</i> ) R, apple ( <i>Malus</i> sp.) R, aspen ( <i>Populus tremula</i> ) R, willow ( <i>Salix</i> sp.), elm ( <i>Ulmus</i> sp.), ground ivy ( <i>Glechoma hederacea</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) O, nipplewort ( <i>Lapsana communis</i> ) R, bramble ( <i>Rubus fruticosus</i> ) O, hop ( <i>Humulus lupulus</i> ) O, lesser burdock ( <i>Arctium minus</i> ) R, great horsetail ( <i>Equisetum telmateia</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) O, wood avens ( <i>Geum urbanum</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) A
G	HL_2	g3c - Other neutral grassland	Anthills	-	Likely used for historic arable practices. Most of glade is species poor. Lots of black ants, low mounds ant hills.	Squirreltail fescue ( <i>Vulpia bromoides</i> ), perennial rye-grass ( <i>Lolium perenne</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F-LA, cock's-foot ( <i>Dactylis glomerata</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), common mouse-ear ( <i>Cerastium fontanum</i> ), field madder ( <i>Sherardia arvensis</i> ), hop trefoil ( <i>Trifolium campestre</i> ), soft brome ( <i>Bromus hordeaceus</i> ) F, cat's-ear ( <i>Hypocharis radicata</i> ), dove's-foot crane's-bill ( <i>Geranium molle</i> ), common ragwort ( <i>Senecio jacobaea</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), white clover ( <i>Trifolium repens</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), sticky mouse-ear ( <i>Cerastium glomeratum</i> ), thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ), false brome ( <i>Brachypodium sylvaticum</i> ) F, possible common cudweed ( <i>Filago vulgaris</i> ), annual meadow-grass ( <i>Poa annua</i> ), bugloss ( <i>Anchusa arvensis</i> ), spear thistle ( <i>Cirsium vulgare</i> ), strawberry clover ( <i>Trifolium fragiferum</i> ) LF, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ), common centaury ( <i>Centaurium erythraea</i> ), dandelion ( <i>Taraxacum</i> agg.), common bent ( <i>Agrostis capillaris</i> ).
G	HL_205	c1c - Cereal crops	-	-	A ploughed field, was a wheat crop with no margins	-
G	HL_210	w1h5 - Other woodland; mixed; mainly broadleaved	-	-	A 30-year-old plantation with three mature trees used as pheasant feeding station, ground flora scarce no ancient woodland indicators, bisected by farm track with several neophytes.	Ash ( <i>Fraxinus excelsior</i> ) A, wild cherry ( <i>Prunus avium</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) A, sweet chestnut ( <i>Castanea sativa</i> ) F, cherry laurel ( <i>Prunus laurocerasus</i> ) O, white poplar ( <i>Populus alba</i> ) R, bramble ( <i>Rubus fruticosus</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) F, field maple ( <i>Acer campestre</i> ) R, hedge woundwort ( <i>Stachys sylvatica</i> ) R, common nettle ( <i>Urtica dioica</i> ) F

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_212	c1c - Cereal crops	-	-	A ploughed wheat field with a 10m margin around the edge of the field	-
G	HL_213	g3c - Other neutral grassland	Sward type mosaic	-	Rolling hills with steep slopes, the higher parts often heavily rabbit grazed with significant exposed sandy soil with sheep's sorrel and cudweed. Species more than 10 per square metre, transitioning in lower parts to more typical G4 grassland with longer sward dominated by Yorkshire fog, cock's foot and bent and fewer herbaceous plants.	Sheep's fescue ( <i>Festuca ovina</i> ) O, sheep's sorrel ( <i>Rumex acetosella</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) F, common cudweed ( <i>Filago vulgaris</i> ) O, bramble ( <i>Rubus fruticosus</i> ) R, red fescue ( <i>Festuca rubra</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, creeping bent ( <i>Agrostis stolonifera</i> ) F, common centaury ( <i>Centaurium erythraea</i> ) R, perforate St John's-wort ( <i>Hypericum perforatum</i> ) R, small-flowered crane's-bill ( <i>Geranium pusillum</i> ) R, common stork's-bill ( <i>Erodium cicutarium</i> ) R, mouse-ear-hawkweed ( <i>Pilosella officinarum</i> ) R
G	HL_215	g4 - Modified grassland	-	-	The field has been mown and sheep are grazing on the field, predominately grass with occasional herbaceous plants.	Cock's-foot ( <i>Dactylis glomerata</i> ) O, red fescue ( <i>Festuca rubra</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, common ragwort ( <i>Senecio jacobaea</i> ) F, yarrow ( <i>Achillea millefolium</i> ) O, common bent ( <i>Agrostis capillaris</i> ) A, sheep's sorrel ( <i>Rumex acetosella</i> ) R, rough meadow-grass ( <i>Poa trivialis</i> ) F, germander speedwell ( <i>Veronica chamaedrys</i> ) O, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, creeping thistle ( <i>Cirsium arvense</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, ground ivy ( <i>Glechoma hederacea</i> ) O, dandelion ( <i>Taraxacum agg.</i> ) O
G	HL_217	g1a6 - Other lowland dry acid grassland	Dry. Sward type mosaic. Other grazed. Scattered scrub.	HPI	A mosaic of grassland types, predominantly G1a6 with some G4. Patches of bramble and nettle, management probably none but heavy rabbit grazing and burrowing.	False oat grass ( <i>Arrhenatherum elatius</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, smooth hawk's-beard ( <i>Crepis capillaris</i> ) O, common bent ( <i>Agrostis capillaris</i> ) F, cock's-foot ( <i>Dactylis glomerata</i> ) F, common ragwort ( <i>Senecio jacobaea</i> ) F, musk thistle ( <i>Carduus nutans</i> ) R, perforate St John's-wort ( <i>Hypericum perforatum</i> ) F, hop trefoil ( <i>Trifolium campestre</i> ) R, sheep's sorrel ( <i>Rumex acetosella</i> ) O, common mouse-ear ( <i>Cerastium fontanum</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R, germander speedwell ( <i>Veronica chamaedrys</i> ) O, dandelion ( <i>Taraxacum agg.</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) F, bramble ( <i>Rubus fruticosus</i> ) F, common nettle ( <i>Urtica dioica</i> ) F, dog-rose ( <i>Rosa canina</i> ) F, ground ivy ( <i>Glechoma hederacea</i> ) O, small-flowered crane's-bill ( <i>Geranium pusillum</i> ) R, yarrow ( <i>Achillea millefolium</i> ) O, hairy sedge ( <i>Carex hirta</i> ) R, thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ) R, red fescue ( <i>Festuca rubra</i> ) F, sheep's fescue ( <i>Festuca ovina</i> ) O, yellow oat-grass ( <i>Trisetum flavescens</i> ) O
G	HL_304	g4 - Modified grassland	-	-	Willow plantation, some willow dead. Passes 6/7 criteria in assessment = good condition.	False oat grass ( <i>Arrhenatherum elatius</i> ) A, cock's-foot ( <i>Dactylis glomerata</i> ) A, common bent ( <i>Agrostis capillaris</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) F, common cudweed ( <i>Filago vulgaris</i> ) R
G	G_T_881	w1g – Other woodland; broadleaved	Plantation	-	Young plantation woodland, trees approximately 7m tall.	Willow ( <i>Salix sp.</i> ) D
G	HL_306	w1g - Other woodland; broadleaved	-	-	Woodland starts at bend of river away from willow field and borders the field to the west. Unable to get into woodland due to wire, and T59 plantation of single age group, no understory, no mature trees, herb layer dominated by nettles. Approx. 40yrs old ash, wild cherry, hawthorn and field maple. field maple is outside land boundary.	Elder ( <i>Sambucus nigra</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, willow ( <i>Salix sp.</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) O, hazel ( <i>Corylus avellana</i> ) O, field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ), common nettle ( <i>Urtica dioica</i> ) D, wild cherry ( <i>Prunus avium</i> )
G	HL_308	g4 - Modified grassland	-	-	Grassland with low sward height, perhaps grazed or managed.	Meadow grass ( <i>Poa sp.</i> ) A, common nettle ( <i>Urtica dioica</i> ) O, dock ( <i>Rumex sp.</i> ) O, thistle ( <i>Cirsium sp.</i> ) O, dandelion ( <i>Taraxacum agg.</i> ) R.

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_335	w1g - Other woodland; broadleaved	Plantation	-	Planted group of trees of similar age with Arrhenatherum dominated grassland understorey, as adjacent. Occasional scattered bramble scrub. W1g 36, lowland broadleaved woodland, plantation. Tree guards present on younger trees.	Apple ( <i>Malus</i> sp.), field maple ( <i>Acer campestre</i> ), hawthorn ( <i>Crataegus monogyna</i> ), pedunculate oak ( <i>Quercus robur</i> ), hazel ( <i>Corylus avellana</i> ), silver birch ( <i>Betula pendula</i> ), hornbeam ( <i>Carpinus betulus</i> ), rose ( <i>Rosa</i> sp.), bramble ( <i>Rubus fruticosus</i> ) O
G	HL_336	h3d - Bramble scrub	-	-	Bramble scrub border to woodland.	Bramble ( <i>Rubus fruticosus</i> ), common nettle ( <i>Urtica dioica</i> ) F, horsetail ( <i>Equisetum</i> sp.), bindweed ( <i>Convolvulus</i> sp), rose ( <i>Rosa</i> sp.)
G	HL_352	c1a - Arable field margins	-	HPI	-	Agrimony ( <i>Agrimonia eupatoria</i> ) O, common knapweed ( <i>Centaurea nigra</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) F, red barista ( <i>Odontites verna</i> ) F, curled dock ( <i>Rumex crispus</i> ) O, upright hedge-parsley ( <i>Torilis japonica</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, black medick ( <i>Medicago lupulina</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) R, common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, Autumn hawkbit ( <i>Leontodon autumnalis</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, white clover ( <i>Trifolium repens</i> ) R, corn parsley ( <i>Petroselinum segetum</i> ) R
G	HL_352a	c1a - Arable field margins	-	HPI	-	Agrimony ( <i>Agrimonia eupatoria</i> ) O, common knapweed ( <i>Centaurea nigra</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) F, red barista ( <i>Odontites verna</i> ) F, curled dock ( <i>Rumex crispus</i> ) O, upright hedge-parsley ( <i>Torilis japonica</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) R, ribwort plantain ( <i>Plantago lanceolata</i> ) A, creeping buttercup ( <i>Ranunculus repens</i> ) F, Yorkshire fog ( <i>Holcus lanatus</i> ) O, black medick ( <i>Medicago lupulina</i> ) R, creeping thistle ( <i>Cirsium arvense</i> ) R, common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) R, autumn hawkbit ( <i>Leontodon autumnalis</i> ) R, common centaury ( <i>Centaurium erythraea</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, white clover ( <i>Trifolium repens</i> ) R, corn parsley ( <i>Petroselinum segetum</i> ) R
G	HL_392	c1c7 - Other cereal crops	Scattered trees		Scattered oak through field	Pedunculate oak ( <i>Quercus robur</i> )
G	A_1436a	g4 - Modified grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_397	h3h - Mixed scrub	-	-	Young scrub	Hawthorn ( <i>Crataegus monogyna</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) O, elm ( <i>Ulmus</i> sp.) R, pedunculate oak ( <i>Quercus robur</i> ) R, spindle ( <i>Euonymus europaeus</i> ) R
G	HL_398	g4 - Modified grassland	-	-	Farmland historically used for grazing / pigs which has been left abandoned. Composition uniform. Not reaching enough species to be other neutral grassland. Common lizards present according to owner.	Cock's-foot ( <i>Dactylis glomerata</i> ) D, common nettle ( <i>Urtica dioica</i> ), bramble ( <i>Rubus fruticosus</i> ) O, cinquefoil ( <i>Potentilla</i> sp.), creeping bent ( <i>Agrostis stolonifera</i> ), curled dock ( <i>Rumex crispus</i> ), yellow oat-grass ( <i>Trisetum flavescens</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), common bird's-foot-trefoil ( <i>Lotus corniculatus</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), common knapweed ( <i>Centaurea nigra</i> ).
G	HL_406	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_409	r1 - Standing open water and canals	-	HPI	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_411	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_413	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_416	g - Grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_417	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_418	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_420	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
G	HL_7	c1b - Temporary grass and clover leys	-	-	Arable land that for some reason has been left alone after being ploughed. Frequent common grasses and herbs with 30 to 50 % bare earth. Likely for sheep grazing.	Perennial rye-grass ( <i>Lolium perenne</i> ), fat-hen ( <i>Chenopodium album</i> ) LA, shepherd's-purse ( <i>Capsella bursa-pastoris</i> ) O, scented mayweed ( <i>Matricaria recutita</i> ), common couch grass ( <i>Elytrigia repens</i> ), dandelion ( <i>Taraxacum</i> agg.), common field speedwell ( <i>Veronica persica</i> ), creeping thistle ( <i>Cirsium arvense</i> ), scentless mayweed ( <i>Tripleurospermum inodorum</i> ), white clover ( <i>Trifolium repens</i> ), groundsel ( <i>Senecio vulgaris</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), common chickweed ( <i>Stellaria media</i> ), common mouse-ear ( <i>Cerastium fontanum</i> ), small nettle ( <i>Urtica urens</i> ), scarlet pimpernel ( <i>Anagallis arvensis</i> )
G	HL_8	g4 - Modified grassland	-	-	Field used for arable practices.	Perennial rye-grass ( <i>Lolium perenne</i> ) A, common poppy ( <i>Papaver rhoeas</i> ) R, common chickweed ( <i>Stellaria media</i> ), prickly sow thistle ( <i>Sonchus asper</i> ), spear thistle ( <i>Cirsium vulgare</i> ), hedge mustard ( <i>Sisymbrium officinale</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), bur chervil ( <i>Anthriscus caucalis</i> ), rough meadow-grass ( <i>Poa trivialis</i> ), scentless mayweed ( <i>Tripleurospermum inodorum</i> ), scented mayweed ( <i>Matricaria recutita</i> ), wall speedwell ( <i>Veronica arvensis</i> ) R, chickweed ( <i>Stellaria</i> sp.), meadow fescue ( <i>Festuca pratensis</i> ) F, Italian ryegrass ( <i>Lolium multiflorum</i> ) F, white campion ( <i>Silene latifolia</i> ) R, meadow thistle ( <i>Cirsium dissectum</i> ), Timothy ( <i>Phleum pratense</i> ), red campion ( <i>Silene dioica</i> ), borage ( <i>Borago officinalis</i> ), common fumitory ( <i>Fumaria officinalis</i> )
G	HL_81	h3h - Mixed scrub	-	-	Cut area of woodland directly below current line. Likely to stop trees touching line.	Dogwood ( <i>Cornus sanguinea</i> ) F, goat willow ( <i>Salix caprea</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, English elm ( <i>Ulmus procera</i> ) O, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R
G	HL_82	w - Woodland and forest	-	-	Some evidence of internal ditch and mound indicating old boundary feature, some apparently disused pheasant pens or trapping. Some tree markers and dormouse boxes about, from old work with the line perhaps, or us. Dying mature elm. Goat willow coppice. Hazel coppice. Ground steep sided valleys, with depression 10m deep.	Dog's mercury ( <i>Mercurialis perennis</i> ) F, false brome ( <i>Brachypodium sylvaticum</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, wood avens ( <i>Geum urbanum</i> ) O, wood dock ( <i>Rumex sanguineus</i> ) R, rough meadow-grass ( <i>Poa trivialis</i> ) R, Primrose F, M pedunculate oak ( <i>Quercus robur</i> ) R, elm ( <i>Ulmus</i> sp.), goat willow ( <i>Salix caprea</i> ) MS R, hazel ( <i>Corylus avellana</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) F, dogwood ( <i>Cornus sanguinea</i> ) R
G	W_1017	r1 – Standing open water and canals	-	HPI	Pond that is dry seasonally. UKHab category derived from aerial imagery.	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
G	HL_86	w1f7 - Other Lowland mixed deciduous woodland	-	-	Ash canopy. Goat willow on steep banks forming 20m canopy layer. Old path in wood with banks from east to west. Topography odd with steep areas and valley which haven't seen in local area. Southern section has a stream, dry during survey, from east to west with mature wych elms and standing deadwood in banks. Update. Have been informed by local residents this area was quarried for aggregate for American WW2 airfield	Wild strawberry ( <i>Fragaria vesca</i> ) LA, false brome ( <i>Brachypodium sylvaticum</i> ) F, wood avens ( <i>Geum urbanum</i> ) R, selfheal ( <i>Prunella vulgaris</i> ) O, primrose ( <i>Primula vulgaris</i> ) F, creeping buttercup ( <i>Ranunculus repens</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, germander speedwell ( <i>Veronica chamaedrys</i> ) R, wood speedwell ( <i>Veronica montana</i> ) R, wood doc ( <i>Rumex sanguineus</i> ) R, ash ( <i>Fraxinus excelsior</i> ) O, goat willow ( <i>Salix caprea</i> ) A, field maple ( <i>Acer campestre</i> ), spindle ( <i>Euonymus europaeus</i> ), hawthorn ( <i>Crataegus monogyna</i> ), dogwood ( <i>Cornus sanguinea</i> ) F, rose ( <i>Rosa sp</i> ), wych elm ( <i>Ulmus glabra</i> ) R.
G	W_1018	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
G	W_1020	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
G	W_1035	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
G	HL_91	g3c - Other neutral grassland	-	-	Ant hills present of ok size. Part of skylark stewardship. Indicators of calcareous tendencies - rough hawkbit and wild carrot. About 10% bare earth. Chicory locally abundant to southern aspect. Area overall does not reach priority habitat, but it has pockets which are species rich. Looks to be established within last 10 years, with management could be priority habitat.	Rough hawkbit ( <i>Leontodon hispidus</i> ) R, wild carrot ( <i>Daucus carota</i> ) F, false oat grass ( <i>Arrhenatherum elatius</i> ) F, crested dog's-tail ( <i>Cynosurus cristatus</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) O, red fescue ( <i>Festuca rubra</i> ) O, yellow oat-grass ( <i>Trisetum flavescens</i> ) O, ribwort plantain ( <i>Plantago lanceolata</i> ) O, hop trefoil ( <i>Trifolium campestre</i> ) R, common ragwort ( <i>Senecio jacobaea</i> ) O, creeping cinquefoil ( <i>Potentilla reptans</i> ) O, oxeye daisy ( <i>Leucanthemum vulgare</i> ) O, creeping bent ( <i>Agrostis stolonifera</i> ), common bent ( <i>Agrostis capillaris</i> ) O, daisy ( <i>Bellis perennis</i> ) O, red clover ( <i>Trifolium pratense</i> ) O, smooth meadow-grass ( <i>Poa pratensis</i> ) R, common knapweed ( <i>Centaurea nigra</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) LA, pyramidal orchids ( <i>Anacamptis pyramidalis</i> ), chicory ( <i>Cichorium intybus</i> ) LA, common centaury ( <i>Centaurium erythraea</i> ) O, common agrimony ( <i>Agrimonia eupatoria</i> ) O, bristly haircap ( <i>Polytrichum piliferum</i> ) O.
G	HL_93	g4 - Modified grassland	-	-	Modified grassland of Italian rye - c1b5	
H	H_T_887	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_T_889	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_T_894	g3c – Other neutral grassland	-	-	Marginal grassland approx. 4-6m wide adjacent to woodland and dry ditch and arable field.	Cock's-foot ( <i>Dactylis glomerata</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), meadow grass ( <i>Poa annua</i> ), bracken ( <i>Pteridium</i> ) encroaching from woodland, buttercup sp. ( <i>Ranunculus sp.</i> ), runes sp., Yorkshire fog ( <i>Holcus lanatus</i> ), spear thistle ( <i>Cirsium vulgare</i> ), selfheal ( <i>Prunella vulgaris</i> ), birch saplings ( <i>Betula sp.</i> ), rush sp. ( <i>Juncus sp.</i> ), common cudweed ( <i>Filago vulgaris</i> ), violet sp. ( <i>Viola sp.</i> ), oak saplings ( <i>Quercus sp.</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), ragwort ( <i>Jacobaea vulgaris</i> ), dandelion sp. ( <i>Taraxacum sp.</i> )
H	A_1445	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_T_895	c1 – Arable and horticulture	-	-	Arable field	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
H	H_T_897	w1f7 - Other Lowland mixed deciduous woodland	-	-	Butler's Wood ancient woodland, oak ( <i>Quercus sp.</i> ), and hazel ( <i>Corylus avellana</i> ) dominated, silver birch ( <i>Betula pendula</i> ) present but failing in areas - rotten birch and fallen trees observed more than in Waldegrave woodland. Evidence of management present, man-made log and brash piles from coppicing, anecdotal evidence from landowner of coppicing undertaken and oak starting to out compete birch - both species replanted after WW2, numerous ancient woodland indicators present. Bracken ( <i>Pteridium</i> ) present in localised patches. Diverse age and height structure but not high amount of scrub layer - hazel abundant but other than that just occasional holly ( <i>Ilex</i> ) and bramble ( <i>Rubus fruticosus</i> ).	-
H	W_1023	r1 – Standing open water and canals	-	HPI	UKHab category derived from aerial imagery/ Phase 1 habitat survey.	-
H	A_1447	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_1776	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_1778	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_1780	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_1782	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_1783	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_673	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_705	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	A_716	w1f7 - Other Lowland mixed deciduous woodland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
H	H_A_1092	w1g - Other woodland; broadleaved	-	-	Small woodland parcel, mature plantation. Notes from 27th April - badgers. Ground flora indicative of ancient woodland. Ditch on woodland boundary with mature pedunculate oak and dead stumps.	Burdock ( <i>Arctium</i> sp.), thyme-leaved speedwell ( <i>Veronica serpyllifolia</i> ), pedunculate oak ( <i>Quercus robur</i> ) D, field maple ( <i>Acer campestre</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) O, elder ( <i>Sambucus nigra</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, native bluebell ( <i>Hyacinthoides non-scripta</i> ) D, greater stitchwort ( <i>Stellaria holostea</i> ) A, red campion ( <i>Silene dioica</i> ) A, lesser celandine ( <i>Ranunculus ficaria</i> ) A, lords and ladies ( <i>Arum maculatum</i> ) A, barren strawberry ( <i>Fragaria sterilis</i> ) O, pedunculate oak ( <i>Quercus robur</i> ), bramble ( <i>Rubus fruticosus</i> ) LA, white dead-nettle ( <i>Lamium album</i> ), lesser stitchwort ( <i>Stellaria graminea</i> ), common chickweed ( <i>Stellaria media</i> ), common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ).
H	H_A_1094	c1 - Arable and horticulture	-	-	Arable crops	-
H	H_A_1095	c1 - Arable and horticulture	-	-	Arable	-
H	A_1454	g- grassland	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_A_1096	c1 - Arable and horticulture	-	-	Arable field with some sunflowers. Radish species growing as crop.	-
H	H_A_1100	c1 - Arable and horticulture	-	-	Arable	-
H	H_A_1101	c1 - Arable and horticulture	-	-	Arable	-
H	H_A_1102	c1e - Intensive orchards	-	-	Apple orchard	-
H	H_A_1104	g3c - Other neutral grassland	-	-	Rough grassland margin, over 1m high sward, false oat, cock's-foot grass dominated,	false oat grass ( <i>Arrhenatherum elatius</i> ) D, cock's-foot ( <i>Dactylis glomerata</i> ) D
H	H_A_1110	g3c - Other neutral grassland	-	-	False oat dominated road verge, low herbaceous coverage.	False oat grass ( <i>Arrhenatherum elatius</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), perennial rye-grass ( <i>Lolium perenne</i> ) O, soft brome ( <i>Bromus mollis</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) A, cleavers ( <i>Galium aparine</i> ) A, docks ( <i>Rumex</i> sp.), spear thistle ( <i>Cirsium vulgare</i> ), white dead-nettle ( <i>Lamium album</i> ) O
H	H_A_1111	w1f7 - Other Lowland mixed deciduous woodland	-	-	Ancient woodland. less signs of management practices compared to Butler's Wood, northern edge locally dominated by hazel, silver birch, western edge mix of pedunculate oak, aspen and porch	Pedunculate oak ( <i>Quercus robur</i> ), hazel ( <i>Corylus avellana</i> ), birch ( <i>Betula</i> sp.), poplar ( <i>Populus</i> sp.)
H	H_A_1114	g4 - Modified grassland	-	-	Small grassland path between arable fields. Regular mowing and vehicle access. G4 115 track.	Greater plantain ( <i>Plantago major</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) A, meadow-grass ( <i>Poa</i> sp.) A, ribwort plantain ( <i>Plantago lanceolata</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) O
H	H_A_1115	g3c - Other neutral grassland	-	-	Tall sward over 1.5m.	False oat grass ( <i>Arrhenatherum elatius</i> ), tufted hair-grass ( <i>Deschampsia cespitosa</i> ), thistle ( <i>Cirsium</i> sp.) LA, cow parsley ( <i>Anthriscus sylvestris</i> ), dock ( <i>Rumex</i> sp.)
H	H_A_1119	u1 - Built-up areas and gardens	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_A_1120	c1 - Arable and horticulture	-	-	Arable	-
H	H_A_1122	g3c - Other neutral grassland	-	-	Grassland normally used for horses - horses have not been on land in some time, grass is long sward over 1m, dominated by Yorkshire fog. Horses to be introduced to land in coming weeks (landowner)	Yorkshire fog ( <i>Holcus lanatus</i> ) D

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
H	H_A_1129	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_A_863	c1 - Arable and horticulture	-	-	Field of rapeseed	-
H	H_A_874	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H_A_875	w1d - Wet woodland	-	HPI	Disturbed area of woodland with main tree species hazel, elder, silver birch, grey willow. Ground flora dominated by common nettle. Ground wet with number of streams and large pond to the north of Order Limits. Occasional ivy, lords and ladies, cleavers, water mint, dog's mercury. Towards pylon woodland becomes increasingly scrub-like and disturbed in patches with main species blackthorn, hawthorn and hops.	Hazel ( <i>Corylus avellana</i> ), elder ( <i>Sambucus nigra</i> ), silver birch ( <i>Betula pendula</i> ), grey willow ( <i>Salix cinerea</i> ), common nettle ( <i>Urtica dioica</i> ) O, ivy ( <i>Hedera helix</i> ) O, lords and ladies ( <i>Arum maculatum</i> ) O, cleavers ( <i>Galium aparine</i> ) O, water mint ( <i>Mentha aquatica</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) O, blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), hop ( <i>Humulus lupulus</i> )
H	HA3_896	h3h – mixed scrub	-	-	Mixed scrub under pylon	-
H	H_A_876	g4 - Modified grassland	Scattered trees		Managed grassland possible grazed and cut. Large pedunculate oaks and pine trees planted sporadically.	Bramble ( <i>Rubus fruticosus</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), creeping thistle ( <i>Cirsium arvense</i> ), primrose ( <i>Primula vulgaris</i> ), white clover ( <i>Trifolium repens</i> ), common ragwort ( <i>Senecio jacobaea</i> ), speedwell ( <i>Veronica sp.</i> ), cuckooflower ( <i>Cardamine pratensis</i> ), wood-sedge ( <i>Carex sylvatica</i> ), dove's-foot crane's-bill ( <i>Geranium molle</i> ), pedunculate oak ( <i>Quercus robur</i> ), pine ( <i>Pinus sp.</i> ), garlic mustard ( <i>Alliaria petiolata</i> )
H	H3_1015	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H3_1020	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H3_1023	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H3_1093	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H3_1135	w1g - Other woodland; broadleaved	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	H3_1139	c1c - Cereal Crops	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	HL_11	h - Heathland and shrub	-	-	Line of semi mature to mature oaks, scrub underlayer.	Pedunculate oak ( <i>Quercus robur</i> )
H	HL_155	c1 - Arable and horticulture	-	-	Arable	-
H	HL_156	c1 - Arable and horticulture	-	-	Arable	-
H	HL_175	c1 - Arable and horticulture	-	-	Arable crops	-
H	HL_177	c1 - Arable and horticulture	-	-	Arable	-

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
H	HL_21	w1f7 - Other Lowland mixed deciduous woodland	-	-	Butler's Wood is an ancient woodland, pedunculate oak and hazel dominated, silver birch present but failing in areas - rotten birch and fallen trees observed more than in Waldegrave woodland. Evidence of management present, man-made log and brash (ash) piles from coppicing, anecdotal evidence from landowner of coppicing undertaken and pedunculate oak starting to out-compete birch - both species replanted after WW2, numerous ancient woodland indicators present. Bracken present in localised patches.	Pedunculate oak ( <i>Quercus robur</i> ) D, hazel ( <i>Corylus avellana</i> ) D, silver birch ( <i>Betula pendula</i> ), ash ( <i>Fraxinus excelsior</i> ), bracken ( <i>Pteridium aquilinum</i> )
H	HL_23	g3c - Other neutral grassland	-	-	False oat dominated road verge, low herbaceous coverage.	False oat grass ( <i>Arrhenatherum elatius</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), perennial rye-grass ( <i>Lolium perenne</i> ) O, soft brome ( <i>Bromus mollis</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) A, cleavers ( <i>Galium aparine</i> ) A, docks ( <i>Rumex</i> sp.), spear thistle ( <i>Cirsium vulgare</i> ), white dead-nettle ( <i>Lamium album</i> ) O
H	HL_25	g3c - Other neutral grassland	Horse grazed. Sheep grazed.	-	Grassland grazed by horses and sheep, northern half of field grazed to ground by animals, where horse fence splits small field in two (see line indicating split) grassland in southern half has higher sward and less evidence of grazing.	-
H	HL_311	c1 - Arable and horticulture	-	-	Arable	-
H	HL_313	g4 - Modified grassland	-	-	Small grassland path between arable fields. Regular mowing and vehicle access. G4 115 track.	Greater plantain ( <i>Plantago major</i> ) F, perennial rye-grass ( <i>Lolium perenne</i> ) A, meadow-grass ( <i>Poa</i> sp.) A, ribwort plantain ( <i>Plantago lanceolata</i> ) O, creeping buttercup ( <i>Ranunculus repens</i> ) O
H	HL_314	c1 - Arable and horticulture	-	-	Not harvested	-
H	HL_320	c1 - Arable and horticulture	-	-	No access but viewed from adjacent field.	-
H	HL_321	g3c - Other neutral grassland	-	-	Marginal grassland approx. 4-6m wide adjacent to woodland and dry ditch and arable field. Bracken encroaching from woodland, birch and oak saplings.	Cock's-foot ( <i>Dactylis glomerata</i> ), false oat grass ( <i>Arrhenatherum elatius</i> ), meadow grass ( <i>Poa</i> sp.), bracken ( <i>Pteridium aquilinum</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), spear thistle ( <i>Cirsium vulgare</i> ), selfheal ( <i>Prunella vulgaris</i> ), birch ( <i>Betula</i> sp.), rush ( <i>Juncus</i> sp.), common cudweed ( <i>Filago vulgaris</i> ), violet ( <i>Viola</i> sp.), pedunculate oak ( <i>Quercus robur</i> ), germander speedwell ( <i>Veronica chamaedrys</i> ), common ragwort ( <i>Senecio jacobaea</i> ), dandelion ( <i>Taraxacum</i> agg.) sp., dock ( <i>Rumex</i> sp.)
H	HL_322	g3c - Other neutral grassland	-	-	Marginal grassland approx. 4-6m wide adjacent to woodland and dry ditch and arable field	-
H	HL_327	c1 - Arable and horticulture	-	-	Arable	-
H	HL_330	w1g - Other woodland; broadleaved	-	-	Small woodland parcel, mature plantation.	Pedunculate oak ( <i>Quercus robur</i> ), field maple ( <i>Acer campestre</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) O, elder ( <i>Sambucus nigra</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O.
H	HL_331	g3c - Other neutral grassland	-	-	Tall sward over 1.5m.	False oat grass ( <i>Arrhenatherum elatius</i> ), tufted hair-grass ( <i>Deschampsia cespitosa</i> ), thistle ( <i>Cirsium</i> sp.) LA, cow parsley ( <i>Anthriscus sylvestris</i> ), dock ( <i>Rumex</i> sp.)

Section	Habitat ID	UKHab Category	Secondary Code	Status	Description	Species List with DAFOR
H	HL_339	w1g - Other woodland; broadleaved	-	-	Small woodland parcel where line of trees widens on arable field boundary.	Hawthorn ( <i>Crataegus monogyna</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) F, ash ( <i>Fraxinus excelsior</i> ) O, field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, elm ( <i>Ulmus</i> sp.) R
H	HL_406	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	HL_406	c1 - Arable and horticulture	-	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
H	HL_46	g3c - Other neutral grassland	-	-	Tall sward over 1.5m.	False oat grass ( <i>Arrhenatherum elatius</i> ), tufted hair-grass ( <i>Deschampsia cespitosa</i> ), thistle ( <i>Cirsium</i> sp.) LA, cow parsley ( <i>Anthriscus sylvestris</i> ), dock ( <i>Rumex</i> sp.)
H	HL_48	g4 - Modified grassland	-	-	Amenity grassland used by scout group, regularly mown, common species, evidence of badger in foraging holes and latrines	-
H	SH_01	g- Grassland	--	-	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

## 1.2 Line of Trees Linear Habitat Descriptions

1.2.1 Habitat classification is shown in Figure 7.1.5 (application document 6.4) for linear features as part of the UKHab mapping for the full project and the descriptions for line of trees (habitat type w1g6) listed in Table A-2.

Table A-2 – Line of Trees (Habitat Type w1g6) Linear Habitat Descriptions

Section	Habitat ID	Description	Species List with DAFOR
AB	LOT-AB-01	Line of trees associated with a dry ditch.	-
AB	LOT-AB-011	Line of trees with shrubs that was once a hedgerow.	Mature oak ( <i>Quercus robur</i> ), hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ) O, ivy ( <i>Hedera helix</i> ). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), bramble ( <i>Rubus fruticosus</i> ).
AB	LOT-AB-02	Line of trees that may have naturally grown along path.	Sycamore ( <i>Acer pseudoplatanus</i> ), lime ( <i>Ulmus</i> sp.), oak ( <i>Quercus robur</i> ), understorey elder ( <i>Sambucus nigra</i> ). Ground flora: nettle ( <i>Urtica dioica</i> ) D, cleavers ( <i>Galium aparine</i> ) D, hemlock ( <i>Conium maculatum</i> ) D.
AB	LOT-AB-03	Line of trees on the roadside of a pond.	Elm ( <i>Ulmus</i> sp.), oak ( <i>Quercus robur</i> ), goat willow ( <i>Salix caprea</i> ). Understorey: elder ( <i>Sambucus nigra</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), common nettle ( <i>Urtica dioica</i> ).
AB	LOT-AB-04	Line of ornamental trees.	-
AB	LOT-AB-05	Line of trees alongside gravel vehicle track, and wet ditch.	Ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), oak ( <i>Quercus robur</i> ), Italian poplar ( <i>Populus x canadensis</i> agg.).
AB	LOT-AB-06	Line of trees along PRoW.	Oak ( <i>Quercus robur</i> ), ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ). Ground flora: primrose ( <i>Primula vulgaris</i> ) F, wood anemone ( <i>Anemone nemorosa</i> ) A, bluebell ( <i>Hyacinthoides non-scripta</i> ) R, lords-and-ladies ( <i>Arum maculatum</i> ) R, honeysuckle ( <i>Lonicera periclymenum</i> ) LA, lesser celandine ( <i>Ranunculus ficaria</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) R, wood spurge ( <i>Euphorbia amygdaloides</i> ).
AB	LOT-AB-07	Line of trees along a footpath, with a ditch on the western aspect.	Oak ( <i>Quercus robur</i> ) D, semi mature ash ( <i>Fraxinus excelsior</i> ), silver birch ( <i>Betula pendula</i> ), young horse chestnut ( <i>Aesculus hippocastanum</i> ), hornbeam ( <i>Carpinus betulus</i> ). Ground flora: primrose ( <i>Primula vulgaris</i> ), bluebell ( <i>Hyacinthoides non-scripta</i> ), wood spurge ( <i>Euphorbia amygdaloides</i> ), greater stitchwort ( <i>Stellaria holostea</i> ).
AB	LOT-AB-08	Small line of semi mature and mature trees, associated with ditch.	Oak ( <i>Quercus robur</i> ), ash ( <i>Fraxinus excelsior</i> ).
AB	LOT-AB-09	Row of four trees along field edge, semi mature to mature.	Sycamore ( <i>Acer pseudoplatanus</i> ) D.
AB	LOT-AB-10	Small line of trees over a ditch.	Oak ( <i>Quercus robur</i> ), hazel ( <i>Corylus avellana</i> ), willow ( <i>Salix</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), ash ( <i>Fraxinus excelsior</i> ), field maple ( <i>Acer campestre</i> ).
AB	LOT-AB-11	Line of trees along dry ditch with mature trees nearer to the field edge and smaller saplings towards centre of the field.	Oak ( <i>Quercus robur</i> ), ash ( <i>Fraxinus excelsior</i> ), elm ( <i>Ulmus</i> sp.).
AB	LOT-AB-12	Line of trees at garden edge, taller than 12m in height.	Cypress ( <i>Chamaecyparis</i> sp.)
AB	LOT-AB-18	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-19	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-20	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-21	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-22	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-23	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

Section	Habitat ID	Description	Species List with DAFOR
AB	LOT-AB-24	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-25	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-26	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
AB	LOT-AB-27	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
C	LOT-C-01	Row of sparse trees along an arable field boundary with some large gaps, all young trees.	Poplar ( <i>Populus</i> sp.), and oak ( <i>Quercus robur</i> ).
D	LOT-D-01	Line of trees over 5m tall with open mosaic on either side. Dry ditch to southern aspect.	Goat willow ( <i>Salix caprea</i> ) and grey willow ( <i>Salix cinerea</i> ).
D	LOT-D-02	Line of trees on a top of a bank that heads down to the south.	Goat willow ( <i>Salix caprea</i> ), grey willow ( <i>Salix cinerea</i> ) and white willow ( <i>Salix alba</i> ).
D	LOT-D-03	Line of trees parallel to LOT-D-04.	Italian poplar ( <i>Populus x canadensis</i> agg.), sycamore ( <i>Acer pseudoplatanus</i> ), wild cherry ( <i>Prunus avium</i> ), sweet chestnut ( <i>Castanea sativa</i> ).
D	LOT-D-04	Line of trees parallel to LOT-D-03.	Italian poplar ( <i>Populus x canadensis</i> agg.), sycamore ( <i>Acer pseudoplatanus</i> ), wild cherry ( <i>Prunus avium</i> ), sweet chestnut ( <i>Castanea sativa</i> ).
D	LOT-D-05	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
D	LOT-D-06	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-
E	LOT-E-01	UKHab category derived from aerial imagery	-
E	LOT-E-02	UKHab category derived from aerial imagery	-
F	LOT-F-01	Line of trees less than 30 years old, spaced approximately 1.5m apart, 8m in height.	Italian alder ( <i>Alnus cordata</i> ).
F	LOT-F-02	Non-native planted tree.	Non-native hornbeam ( <i>Carpinus</i> sp.)
F	LOT-F-03	Line planted trees, that are all the same age. Evidence of historical pollarding.	Italian alder ( <i>Alnus cordata</i> ).
F	LOT-F-04	Line of trees that borders garden. The garden could not be access to land access constraints.	Oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, balsam poplar ( <i>Populus</i> sp.) R, horse chestnut ( <i>Aesculus hippocastanum</i> ) R, willow ( <i>Salix</i> sp.) R, rose ( <i>Rosa</i> sp.) R, birch ( <i>Betula</i> sp.) O, field maple ( <i>Acer campestre</i> ) R, semi mature pedunculate oak ( <i>Quercus robur</i> ).
F	LOT-F-05	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-
F	LOT-F-06	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-
F	LOT-F-07	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-
G	LOT-G-01	Line of six semi mature trees, 10m to 14m in height.	Willow ( <i>Salix</i> sp.)
G	LOT-G-02	Trees running along watercourse.	Willow ( <i>Salix</i> sp.) F, wild hops ( <i>Humulus lupulus</i> ), elder ( <i>Sambucus nigra</i> ) O, bramble ( <i>Rubus fruticosus</i> ) O, wych elm ( <i>Ulmus glabra</i> ) O, goat willow ( <i>Salix caprea</i> ) A, common nettle ( <i>Urtica dioica</i> ) F, apple ( <i>Malus sylvestris</i> ) R.
G	LOT-G-03	Line of trees along watercourse.	Willow ( <i>Salix</i> sp.)
G	LOT-G-04	Line of trees.	White willow ( <i>Salix alba</i> ) A.

Section	Habitat ID	Description	Species List with DAFOR
G	LOT-G-05	Semi mature line of trees.	Oak ( <i>Quercus robur</i> ) and small-leaved lime ( <i>Tilia cordata</i> ). Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ) A, Yorkshire fog ( <i>Holcus lanatus</i> ) A, perennial rye grass ( <i>Lolium perenne</i> ) F, white dead-nettle ( <i>Lamium album</i> ) O, nettle ( <i>Urtica dioica</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) F, foxtail ( <i>Alopecurus pratensis</i> ) R.
G	LOT-G-06	Two semi-mature oak trees with central eucalyptus.	Pedunculate oak ( <i>Quercus robur</i> ), eucalyptus ( <i>Eucalyptus</i> sp.)
G	LOT-G-07	An overgrown hedgerow turned into a line of trees. Along a road, on a bank and flailed at 4m high with the higher branches forming a tunnel across the road to touch the opposite hedge.	Elm ( <i>Ulmus</i> sp) D, dog-rose ( <i>Rosa canina</i> ) R, mature pedunculate oak ( <i>Quercus robur</i> ) O, holly ( <i>Ilex aquifolium</i> ) O, false-brome ( <i>Brachypodium sylvaticum</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) O, herb-Robert ( <i>Geranium robertianum</i> ) R.
G	LOT-G-08	Line of semi mature trees.	Elm ( <i>Ulmus</i> sp.)
G	LOT-G-09	Previously laid hedgerow along a sandy bank which has ground into a line of trees. The trees are multi-stemmed between 6m and 20m in height.	Spindle ( <i>Euonymus europaeus</i> ) R, Pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, elder ( <i>Sambucus nigra</i> ) R, elm ( <i>Ulmus</i> sp) F, Holly ( <i>Ilex aquifolium</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, field maple ( <i>Acer campestre</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) O, false-brome ( <i>Brachypodium sylvaticum</i> ) O
G	LOT-G-10	Coppiced overgrown hedge which has evolved into an overgrown line of trees with large gaps greater than 5m. Trees were in good health stems as large as 40cm dbh.	Elm ( <i>Ulmus</i> sp,) A, elder ( <i>Sambucus nigra</i> ) R.
G	LOT-G-11	Lime of trees either side of the road.	Elm ( <i>Ulmus</i> sp.)
G	LOT-G-12	Line of semi mature trees on the side of the road.	
G	LOT-G-13	Line of trees.	Field maple ( <i>Acer campestre</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) O, crack willow ( <i>Salix fragilis</i> ) O, ivy ( <i>Hedera helix</i> ) O, Hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, Holly ( <i>Ilex aquifolium</i> ) R, elder ( <i>Sambucus nigra</i> ) R, common nettle ( <i>Urtica dioica</i> ) A, dock ( <i>Rumex</i> sp.) O, thistle ( <i>Cirsium</i> sp.) O.
G	LOT-G-14	Non-native willow trees planted on northern side of ditch. Patchy in places where been cut down.	Willow ( <i>Salix</i> sp.)
G	LOT-G-15	Line of trees of similar ages on northern side of a wet ditch and PRoW. Evidence of historic pollarding and coppicing. Shrubs cut underneath.	Pedunculate oak ( <i>Quercus robur</i> ).
G	LOT-G-16	Line of trees on steep roadside bank. Small area of grassland between line of trees and arable.	English elm ( <i>Ulmus procera</i> ) D, holly ( <i>Ilex aquifolium</i> ) R, ivy ( <i>Hedera helix</i> ). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) A, cock's foot ( <i>Dactylis glomerata</i> ) A, creeping thistle ( <i>Cirsium arvense</i> ) R, ground ivy ( <i>Glechoma hederacea</i> ) F, nettle ( <i>Urtica dioica</i> ) F.
G	LOT-G-17	Line of trees.	White willow ( <i>Salix alba</i> ) A. Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ) A, dock ( <i>Rumex</i> sp.) O, white dead-nettle ( <i>Lamium album</i> ) O, common nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) F, thistle ( <i>Cirsium</i> sp.) O, foxtail ( <i>Alopecurus pratensis</i> ) O.
G	LOT-G-018	Line of trees along a stream, semi mature to mature trees.	Alder ( <i>Alnus glutinosa</i> ). Understorey: elder ( <i>Sambucus nigra</i> ) O. Ground flora: nettle ( <i>Urtica dioica</i> ) D, ramsons ( <i>Allium ursinum</i> ) O.
G	LOT-G-019	Line of trees planted approximately 20 years ago by the landowner.	Pedunculate oak trees ( <i>Quercus robur</i> )
G	LOT-G-020	Line of mature trees along road.	Elm ( <i>Ulmus</i> sp.). Ground flora: garlic mustard ( <i>Alliaria petiolata</i> ), nettle ( <i>Urtica dioica</i> ), false-brome ( <i>Brachypodium sylvaticum</i> )
G	LOT-G-021	Line of trees along a dry ditch with mature trees.	Mature oak ( <i>Quercus robur</i> ) F, field maple ( <i>Acer campestre</i> ) O, coppiced hazel (F), elder ( <i>Sambucus nigra</i> ) R, ash ( <i>Fraxinus excelsior</i> ) O, elm ( <i>Ulmus</i> sp.) F, lime ( <i>T. x europaea</i> ) (F). Ground flora: garlic mustard ( <i>Alliaria petiolata</i> ), cleavers ( <i>Galium aparine</i> ), nettle ( <i>Urtica dioica</i> ), bramble ( <i>Rubus fruticosus</i> ), creeping soft-grass ( <i>Holcus mollis</i> ), dock ( <i>Rumex</i> sp.) thistle ( <i>Cirsium</i> sp.)
G	LOT-G-22	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-

Section	Habitat ID	Description	Species List with DAFOR
H	LOT-H-01	Line of trees/over mature hedgerow running adjacent to track around the scout camp.	Oak ( <i>Quercus robur</i> ), bramble ( <i>Rubus fruticosus</i> ), blackthorn ( <i>Prunus spinosa</i> ), common hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ), rose ( <i>Rosa sp.</i> ), field maple ( <i>Acer campestre</i> ), ivy ( <i>hedera helix</i> ), yew ( <i>Taxus baccata</i> )
H	LOT-H-02	Line of trees/over mature hedgerow along a small and dry ditch.	Elder ( <i>Sambucus nigra</i> ), Oak ( <i>Quercus robur</i> ), bramble ( <i>Rubus fruticosus</i> ), rose ( <i>Rosa sp.</i> ), willow ( <i>Salix sp.</i> ), ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ), nettle ( <i>Urtica dioica</i> ), field maple ( <i>Acer campestre</i> ), elm ( <i>Ulmus sp.</i> )
H	LOT-H-03	Line of trees with extensive gaps. Ruderal ground flora.	Oak ( <i>Quercus robur</i> ), blackthorn ( <i>Prunus spinosa</i> ), holly ( <i>Ilex aquifolium</i> ), hazel ( <i>Corylus avellana</i> ). Ground flora: bramble ( <i>Rubus fruticosus</i> ), bracken ( <i>Pteridium aquilinum</i> ).
H	LOT-H-04	Broken canopy of line of six trees, with some understorey scrub.	-
H	LOT-H-05	Line of mature trees located along a grassy track and arable field. The arable field side has no grass margin. Parallel line of trees present opposite grassy track.	Ash ( <i>Fraxinus excelsior</i> ), field maple ( <i>Acer campestre</i> ), English elm ( <i>Ulmus procera</i> ), pedunculate oak ( <i>Quercus robur</i> ), blackthorn ( <i>Prunus spinosa</i> ), hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> ), ivy ( <i>Hedera helix</i> ), nettle ( <i>Urtica dioica</i> ).
H	LOT-H-06	Line of semi mature oak trees that is dominated with mature hawthorn, tall sward grass along margin of fields.	Semi mature oak ( <i>Quercus robur</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) D. Ground flora: nettle ( <i>Urtica dioica</i> ).
H	LOT-H-07	Short line of trees within dry ditch, acting as boundary to two arable fields. Dead trees present.	Elm ( <i>Ulmus sp.</i> ) D.
H	LOT-H-08	Planted line of poplar trees, all approximately 4m in height, well maintained.	Poplar ( <i>Populus sp.</i> )
H	LOT-H-09	Planted line of trees, heavily pollarded and managed for aesthetics.	Lime ( <i>Tilia sp.</i> )
H	LOT-H-10	Planted line of trees, heavily pollarded and managed for aesthetics.	Lime ( <i>Tilia sp.</i> )
H	LOT-H-12	UKHab category derived from aerial imagery and/or 2012 Phase 1 results	-

## 1.3 Hedgerow Descriptions

1.3.1 Habitat classification is shown in Figure 7.1.5 (application document 6.4) for linear features as part of the UKHab mapping for the full project and the descriptions for hedgerows listed in Table A-3.

Table A-3 – Hedgerow Descriptions

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-122	h2 - hedgerow	UKHab category derived from aerial imagery.	-
AB	H-AB-120	h2 - hedgerow	UKHab category derived from aerial imagery.	-
AB	H-AB-119	h2 - hedgerow	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-
AB	H-AB-118	h2 - hedgerow	UKHab category derived from aerial imagery and/or 2012 Phase 1 results.	-
AB	H-AB-117	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees - Associated with bank or ditch.	N/A. Not surveyed
AB	H-AB-115	h2a - hedgerow (priority habitat)	Shrubby, flailed, native hedgerow - associated with bank.	Hawthorn ( <i>Crataegus monogyna</i> ) D, oak ( <i>Quercus</i> sp.) O. Ground flora: bristly oxtongue ( <i>Picris echioides</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), common hogweed ( <i>Heracleum sphondylium</i> ), cleavers ( <i>Galium aparine</i> ), willowherb sp. ( <i>Epilobium</i> sp.), common nettle ( <i>Urtica dioica</i> ).
AB	H-AB-113	h2a - hedgerow (priority habitat)	Hedgerow with trees.	Pedunculate oak ( <i>Quercus robur</i> ), ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), bramble ( <i>Rubus</i> sp.), Ground flora cowslip ( <i>Primula veris</i> ) F, dog's mercury F, knapweed ( <i>Centaurea nigra</i> ) A, cow parsley ( <i>Anthriscus sylvestris</i> ) O.
AB	H-AB-112	h2a - hedgerow (priority habitat)	Hedgerow uniform in height, width, and species composition along length. Flailed and with a bank.	Hawthorn ( <i>Crataegus monogyna</i> ) D. Ground flora: common nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), common hogweed ( <i>Heracleum sphondylium</i> ), dandelion sp. ( <i>Taraxacum</i> sp.).
AB	H-AB-111	h2a - hedgerow (priority habitat)	Native hedgerow with trees, along a dry ditch.	Hawthorn ( <i>Crataegus monogyna</i> ) D, pedunculate oak ( <i>Quercus robur</i> ), sweet chestnut ( <i>Castanea sativa</i> ). Ground flora: cleavers ( <i>Galium aparine</i> ), common nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), dock sp. ( <i>Rumex</i> sp.), garlic mustard ( <i>Alliaria petiolata</i> ), bramble ( <i>Rubus</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ).
AB	H-AB-109	h2a - hedgerow (priority habitat)	Hedgerow adjacent to arable field boundary associated with dry ditch.	Blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), field maple ( <i>Acer campestre</i> ). Ground flora include cleavers ( <i>Galium aparine</i> ), dandelion sp. ( <i>Taraxacum</i> sp.), garlic mustard ( <i>Alliaria petiolata</i> ), bramble ( <i>Rubus</i> sp.), common nettle ( <i>Urtica dioica</i> ), thistle sp. ( <i>Cirsium</i> sp.), cow parsley ( <i>Anthriscus sylvestris</i> ), ivy ( <i>Hedera helix</i> ), buttercup sp. ( <i>Ranunculus</i> sp.), lesser celandine ( <i>Ranunculus ficaria</i> ), willowherb sp. ( <i>Epilobium</i> sp.), buttercup sp. ( <i>Ranunculus</i> sp.), bristly oxtongue ( <i>Helminthotheca echioides</i> ).
AB	H-AB-108	h2a - hedgerow (priority habitat)	Hedgerow with trees located between arable fields, with a ditch along its length. Wetter area where willow ( <i>Salix</i> sp.) present.	Elm sp. ( <i>Ulmus</i> sp.), semi mature ash ( <i>Fraxinus excelsior</i> ), dog-rose ( <i>Rosa canina</i> ), blackthorn ( <i>Prunus spinosa</i> ) LA, hawthorn ( <i>Crataegus monogyna</i> ) LA, spindle ( <i>Euonymus europaeus</i> ) R, sallow willow ( <i>Salix caprea</i> ) LA. Ground flora: wood anemone ( <i>Anemone nemorosa</i> ), hedge bedstraw ( <i>Galium mollugo</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), curled dock ( <i>Rumex crispus</i> ), cowslip ( <i>Primula veris</i> ).
AB	H-AB-104	h2a - hedgerow (priority habitat)	Hedgerow along highway and well cut. Ditch to south of hedgerow.	Elm sp. ( <i>Ulmus</i> sp.), hawthorn ( <i>Crataegus monogyna</i> ) and hazel ( <i>Corylus avellana</i> ). Nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ).
AB	H-AB-103	h2a - hedgerow (priority habitat)	Planted and flailed hedgerow.	Hawthorn ( <i>Crataegus monogyna</i> ), elm sp. ( <i>Ulmus</i> sp.). Ground flora: nettle ( <i>Urtica dioica</i> ) D, cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), dock sp. ( <i>Rumex</i> sp.), garlic mustard.
AB	H-AB-102	h2a - hedgerow (priority habitat)	Hedgerow with western hedgerow heavily flailed and managed with no trees.	Blackthorn ( <i>Prunus spinosa</i> ) A, ash ( <i>Fraxinus excelsior</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, sycamore ( <i>Acer pseudoplatanus</i> ) R, mature pedunculate oak ( <i>Quercus robur</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ) F, dog's mercury ( <i>Mercurialis perennis</i> ) A, cleavers ( <i>Galium aparine</i> ) O, primrose ( <i>Primula vulgaris</i> ) R.

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-101	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow with trees, along a dry ditch. Has been flailed.	Hazel ( <i>Corylus avellana</i> ) C, blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ), mature pedunculate oak ( <i>Quercus robur</i> ). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) A, lords-and-ladies ( <i>Arum maculatum</i> ) F, spurge laurel ( <i>Daphne laureola</i> ) R, false brome ( <i>Brachypodium sylvaticum</i> ) A, cleavers ( <i>Galium aparine</i> ) F, cut-leaved crane's-bill ( <i>Geranium dissectum</i> ).
AB	H-AB-099	h2a - hedgerow (priority habitat)	Flailed hedgerow adjacent to road and field. Entire hedgerow was not surveyed.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), oak sp. ( <i>Quercus</i> sp.), rose sp. ( <i>Rosa</i> sp.), ivy ( <i>Hedera helix</i> ). Ground flora: cowslip ( <i>Primula veris</i> ), cleavers ( <i>Galium aparine</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), nettle ( <i>Urtica dioica</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), bramble ( <i>Rubus</i> sp.), dock sp. ( <i>Rumex</i> sp.).
AB	H-AB-098	h2a - hedgerow (priority habitat)	Flailed hedgerow running parallel to another hedgerow.	Hawthorn ( <i>Crataegus monogyna</i> ), oak sp. ( <i>Quercus</i> sp.), ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ), rose sp. ( <i>Rosa</i> sp.). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), nettle ( <i>Urtica dioica</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), dock sp. ( <i>Rumex</i> sp.).
AB	H-AB-097	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees.	N/A. Not surveyed
AB	H-AB-096	h2a - hedgerow (priority habitat)	Native Hedgerow.	N/A. Not surveyed
AB	H-AB-095	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees.	N/A. Not surveyed
AB	H-AB-094	h2a - hedgerow (priority habitat)	Defunct hedgerow with several gaps and no longer stockproof. Short and overgrown.	Elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ). Ground flora: cleavers ( <i>Galium aparine</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), common nettle ( <i>Urtica dioica</i> ), lords-and-ladies ( <i>Arum maculatum</i> ).
AB	H-AB-093	h2a - hedgerow (priority habitat)	Hedgerow with trees, associated with bank and ditch.	Field maple ( <i>Acer campestre</i> ), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), elder ( <i>Sambucus nigra</i> ), rose sp. ( <i>Rosa</i> sp.), hazel ( <i>Corylus avellana</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), white dead-nettle ( <i>Lamium album</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), cowslip ( <i>Primula veris</i> ), stitchwort sp. ( <i>Stellaria</i> sp.), greater burdock ( <i>Arctium lappa</i> ).
AB	H-AB-092	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees - associated with bank or ditch.	N/A. Not surveyed
AB	H-AB-090	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
AB	H-AB-089	h2a - hedgerow (priority habitat)	Hedgerow with trees along a road. Gate towards southern end. Manure and hay dumped on eastern boundary.	Field maple ( <i>Acer campestre</i> ), spindle ( <i>Euonymus europaeus</i> ), ivy ( <i>Hedera helix</i> ), blackthorn ( <i>Prunus spinosa</i> ), old man's beard ( <i>Clematis vitalba</i> ), rose sp. ( <i>Rosa</i> sp.), bramble ( <i>Rubus</i> sp.), elder ( <i>Sambucus nigra</i> ), hazel ( <i>Corylus avellana</i> ), oak sp. ( <i>Quercus</i> sp.), dogwood ( <i>Cornus sanguinea</i> )
AB	H-AB-088	h2a - hedgerow (priority habitat)	Flailed hedgerow alongside a road and dry ditch.	Hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), ivy ( <i>Hedera helix</i> ), blackthorn ( <i>Prunus spinosa</i> ), rose sp. ( <i>Rosa</i> sp.), dogwood ( <i>Cornus sanguinea</i> ), apple sp. ( <i>Malus</i> sp.), ash ( <i>Fraxinus excelsior</i> ), black bryony ( <i>Tamus communis</i> ), oak sp. ( <i>Quercus</i> sp.)
AB	H-AB-087	h2a - hedgerow (priority habitat)	Hedgerow with trees along a dry ditch. Possibly outside of Order Limits.	Field maple ( <i>Acer campestre</i> ), ivy ( <i>Hedera helix</i> ), rose sp. ( <i>Rosa</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), bramble ( <i>Rubus</i> sp.), ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ), oak sp. ( <i>Quercus</i> sp.), dogwood ( <i>Cornus sanguinea</i> ), hazel ( <i>Corylus avellana</i> )

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-086	h2a - hedgerow (priority habitat)	Hedgerow with trees along dry ditch and grassland strip.	Ash ( <i>Fraxinus excelsior</i> ), bramble ( <i>Rubus sp.</i> ), rose sp. ( <i>Rosa sp.</i> ), field maple ( <i>Acer campestre</i> ), plum ( <i>Prunus domestica</i> ), hawthorn ( <i>Crataegus monogyna</i> ), goat willow ( <i>Salix caprea</i> ), oak sp. ( <i>Quercus sp.</i> ), blackthorn ( <i>Prunus spinosa</i> ).
AB	H-AB-085	h2a - hedgerow (priority habitat)	Hedgerow along a dry ditch and garden. Height 8-10m, width 2-3m, less than 30 years old.	Dogwood ( <i>Cornus sanguinea</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), rose sp. ( <i>Rosa sp.</i> ), bramble ( <i>Rubus sp.</i> )
AB	H-AB-084	h2a - hedgerow (priority habitat)	A managed hedgerow along garden and public footpath. Three trees along length, evenly spaced.	Hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), pedunculate oak ( <i>Quercus robur</i> ), blackthorn ( <i>Prunus spinosa</i> ), ash ( <i>Fraxinus excelsior</i> ), hazel ( <i>Corylus avellana</i> ), cranberry ( <i>Vaccinium myrtillus</i> ), rose sp. ( <i>Rosa sp.</i> )
AB	H-AB-083	h2a - hedgerow (priority habitat)	Native hedgerow with trees. Associated with bank or ditch. Hedgerow has been flailed.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), dogwood ( <i>Cornus sanguinea</i> ), young lime ( <i>Tilia x europaea</i> ). Ground flora: ivy ( <i>Hedera helix</i> ) D, cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), ground-ivy ( <i>Hedera helix</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), hogweed ( <i>Heracleum sphondylium</i> ).
AB	H-AB-082	h2a - hedgerow (priority habitat)	Hedgerow along a ditch, with large gaps.	Section 1: hazel ( <i>Corylus avellana</i> ), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), sweet chestnut ( <i>Castanea sativa</i> ). Section 2: hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), rose ( <i>Rosa sp.</i> ), field maple ( <i>Acer campestre</i> ). Ground flora: Primrose, nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), plantain sp. ( <i>Plantago sp.</i> ), dock sp. ( <i>Rumex sp.</i> ), cowslip ( <i>Primula veris</i> ), dandelion sp. ( <i>Taraxacum sp.</i> ).
AB	H-AB-079	h2a - hedgerow (priority habitat)	Defunct hedgerow uniform in height and species composition. Ground flora sparse with most comprising arable field edge. Gaps filled with bramble.	Hawthorn ( <i>Crataegus monogyna</i> ), bramble ( <i>Rubus sp.</i> ), willow sp. ( <i>Salix sp.</i> ), sycamore ( <i>Acer pseudoplatanus</i> ), blackthorn ( <i>Prunus spinosa</i> ). Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), plantain sp. ( <i>Plantago sp.</i> ), nettle ( <i>Urtica dioica</i> ), cowslip ( <i>Primula veris</i> ). Lots of bramble ( <i>Rubus sp.</i> ) in hedge gaps.
AB	H-AB-077	h2a - hedgerow (priority habitat)	Native species rich hedgerow.	N/A. Not surveyed
AB	H-AB-076	h2a - hedgerow (priority habitat)	Native species rich hedgerow.	N/A. Not surveyed
AB	H-AB-075	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-074	h2a - hedgerow (priority habitat)	Native hedgerow with trees.	N/A. Not surveyed
AB	H-AB-073	h2a - hedgerow (priority habitat)	Native hedgerow associated with a bank or ditch.	N/A. Not surveyed
AB	H-AB-071	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-069	h2a - hedgerow (priority habitat)	Native species rich hedgerow.	N/A. Not surveyed

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-068	h2a - hedgerow (priority habitat)	Native species rich, flailed hedgerow with trees.	Elm sp. ( <i>Ulmus</i> sp.) R, dogwood ( <i>Cornus sanguinea</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) A, field maple ( <i>Acer campestre</i> ) F, Persian ivy ( <i>Hedera colchica</i> ) F, ash ( <i>Fraxinus excelsior</i> ) R, hazel ( <i>Corylus avellana</i> ) O, dog-rose ( <i>Rosa canina</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) F. Ground flora: white bryony ( <i>Bryonia dioica</i> ) O, false oat-grass ( <i>Arrhenatherum elatius</i> ) A, perennial ryegrass ( <i>Lolium perenne</i> ) A, hedge parsley sp. ( <i>Torilis</i> sp.), dock sp. ( <i>Rumex</i> sp.) O, hogweed ( <i>Heracleum sphondylium</i> ) O, nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) O, greater burdock ( <i>Arctium lappa</i> ) R, nipplewort ( <i>Lapsana communis</i> ) R
AB	H-AB-067	h2a - hedgerow (priority habitat)	Hedgerow with trees. Northern section comprises more trees, much taller and less managed than the southern section. Southern section less species diversity with field maple and hazel dominating.	Field maple ( <i>Acer campestre</i> ) A, dogwood ( <i>Cornus sanguinea</i> ) O, hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, mature pedunculate oak ( <i>Quercus robur</i> ) O, mature sweet chestnut ( <i>Castanea sativa</i> ) R.
AB	H-AB-066	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow which has been flailed.	Dogwood ( <i>Cornus sanguinea</i> ), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ), pedunculate oak ( <i>Quercus robur</i> ). Ground flora: false oat-grass ( <i>Arrhenatherum elatius</i> ), perennial ryegrass ( <i>Lolium perenne</i> ), four-seeded vetch ( <i>Vicia tetrasperma</i> ), field sorrel ( <i>Rumex acetosa</i> ), Timothy ( <i>Phleum pratense</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), black medick ( <i>Medicago lupulina</i> ), cowslip ( <i>Primula veris</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), hogweed ( <i>Heracleum sphondylium</i> ), curled dock ( <i>Rumex crispus</i> ), mint sp. ( <i>Mentha</i> sp.)
AB	H-AB-065	h2a - hedgerow (priority habitat)	Historical hedgerow now bordering a power plant.	Blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), hazel ( <i>Corylus avellana</i> ), dogwood ( <i>Cornus sanguinea</i> ), dog-rose ( <i>Rosa canina</i> ). Ground flora: false oat-grass ( <i>Arrhenatherum elatius</i> ) A, perennial ryegrass ( <i>Lolium perenne</i> ) F, lords-and-ladies ( <i>Arum maculatum</i> ) R, hedge parsley sp. ( <i>Torilis</i> sp.) R, bracken ( <i>Pteridium aquilinum</i> ) R
AB	H-AB-064	h2a - hedgerow (priority habitat)	Hedgerow, uniform along the whole length. Continuation of planted scrub but as a thin strip, all in tree guards and young.	Blackthorn ( <i>Prunus spinosa</i> ), holly ( <i>Ilex aquifolium</i> ), rose ( <i>Rosa</i> sp.), hazel ( <i>Corylus avellana</i> ), dogwood ( <i>Cornus sanguinea</i> ).
AB	H-AB-063	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-062	h2a - hedgerow (priority habitat)	Thin hedgerow consists of continuous planted scrub. All young trees present in tree guards.	Blackthorn ( <i>Prunus spinosa</i> ), holly ( <i>Ilex aquifolium</i> ), rose sp. ( <i>Rosa</i> sp.), hazel ( <i>Corylus avellana</i> ), dogwood ( <i>Cornus sanguinea</i> ). Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), spear thistle ( <i>Cirsium vulgare</i> )
AB	H-AB-061	h2a - hedgerow (priority habitat)	Historical hedgerow next to bridleway.	Blackthorn ( <i>Prunus spinosa</i> ) A, hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) F, dogwood ( <i>Cornus sanguinea</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, dog-rose ( <i>Rosa canina</i> ) F, ash ( <i>Fraxinus excelsior</i> ) R, field maple ( <i>Acer campestre</i> ) F. Ground flora: nettle ( <i>Urtica dioica</i> ) R, spear thistle ( <i>Cirsium vulgare</i> ) O, lady's bedstraw ( <i>Galium verum</i> ) O, giant fescue ( <i>Schedonardus giganteus</i> ) R, false oat-grass ( <i>Arrhenatherum elatius</i> ) A, perennial ryegrass ( <i>Lolium perenne</i> ) A, wood false-brome ( <i>Brachypodium sylvaticum</i> ) F
AB	H-AB-060	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-059	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-058	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-057	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-056	h2a - hedgerow (priority habitat)	Native species rich hedgerow.	N/A. Not surveyed
AB	H-AB-055	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
AB	H-AB-054	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
AB	H-AB-053	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees	N/A. Not surveyed
AB	H-AB-051	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
AB	H-AB-050	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-049	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
AB	H-AB-048	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-046	h2a - hedgerow (priority habitat)	Hedgerow greatly improves to western part - fewer gaps, more trees, more species.	Ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), elm sp. ( <i>Ulmus</i> sp.), field rose sp. ( <i>Rosa</i> sp.), dog-rose ( <i>Rosa canina</i> ), field maple ( <i>Acer campestre</i> ), sycamore ( <i>Acer pseudoplatanus</i> ). Ground flora: hogweed ( <i>Heracleum sphondylium</i> ), nettle ( <i>Urtica dioica</i> ) A, dog's mercury ( <i>Mercurialis perennis</i> ), curled dock ( <i>Rumex crispus</i> ), cleavers ( <i>Galium aparine</i> ) A., false oxlip ( <i>Primula x polyantha</i> ).
AB	H-AB-043	h2a - hedgerow (priority habitat)	Hedgerow along a ditch with little water and no flow.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), willow sp. ( <i>Salix</i> sp.), dogwood ( <i>Cornus sanguinea</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), curled dock ( <i>Rumex crispus</i> ), willowherb species ( <i>Epilobium</i> sp.), dog's mercury ( <i>Mercurialis perennis</i> ), cleavers ( <i>Galium aparine</i> ).
AB	H-AB-042	h2a - hedgerow (priority habitat)	Flailed hedgerow with trees. Signs of enrichment. Veteran oak species present.	Blackthorn ( <i>Prunus spinosa</i> ), hazel ( <i>Corylus avellana</i> ), spindle ( <i>Euonymus europaeus</i> ), dog-rose ( <i>Rosa canina</i> ) sp. ( <i>Rosa</i> sp.), veteran oak sp. ( <i>Quercus</i> sp.). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ), cleavers ( <i>Galium aparine</i> ), nettle ( <i>Urtica dioica</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), spurge laurel ( <i>Daphne laureola</i> ).
AB	H-AB-040	h2a - hedgerow (priority habitat)	Shrubby, native species rich hedgerow with trees. Associated with a dry ditch. Hedgerow has been flailed.	Hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), oak sp. ( <i>Quercus</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), hazel ( <i>Corylus avellana</i> ), dog-rose ( <i>Rosa canina</i> ), ash ( <i>Fraxinus excelsior</i> ), dogwood ( <i>Cornus sanguinea</i> ), willow species likely goat willow ( <i>Salix caprea</i> ), sallow willow ( <i>Salix caprea</i> ) R, elm sp. ( <i>Ulmus</i> sp.), sycamore ( <i>Acer pseudoplatanus</i> ), elder ( <i>Sambucus nigra</i> ) R. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ), cleavers ( <i>Galium aparine</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), spurge laurel ( <i>Daphne laureola</i> ) R, wood false-brome ( <i>Brachypodium sylvaticum</i> ) R, garlic mustard ( <i>Alliaria petiolata</i> ), wood spurge ( <i>Euphorbia amygdaloides</i> ) R
AB	H-AB-039	h2a - hedgerow (priority habitat)	Partially replanted hedgerow along a ditch with little water and a slow flow rate.	Hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ), dogwood ( <i>Cornus sanguinea</i> ), field maple ( <i>Acer campestre</i> ), elm sp. ( <i>Ulmus</i> sp.). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), wood false-brome ( <i>Brachypodium sylvaticum</i> )

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-038	h2a - hedgerow (priority habitat)	Partially planted hedgerow. Mature oak species present.	Ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ) D, blackthorn ( <i>Prunus spinosa</i> ), dog-rose ( <i>Rosa canina</i> ), dogwood ( <i>Cornus sanguinea</i> ), mature oak sp. ( <i>Quercus</i> sp.). Ground flora: nettle ( <i>Urtica dioica</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ).
AB	H-AB-037	h2a - hedgerow (priority habitat)	Hedgerow with sparse ground flora.	Ash ( <i>Fraxinus excelsior</i> ) O, oak sp. ( <i>Quercus</i> sp.) R, field maple ( <i>Acer campestre</i> ) F, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) F, ivy ( <i>Hedera helix</i> ) F, rose sp. ( <i>Rosa</i> sp.) F. Ground flora: dock ( <i>Rumex</i> sp.) O, nettle ( <i>Urtica dioica</i> ) O, garlic mustard ( <i>Alliaria petiolata</i> ) F.
AB	H-AB-036	h2a - hedgerow (priority habitat)	Flailed hedgerow along a wire fence. Field grazed closely up to the hedgerow resulting in ground flora similar to the adjacent grassland.	hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ) D with shield, maple sp. ( <i>Acer</i> sp.), rose sp. ( <i>Rosa</i> sp.) O and dogwood F. Ground flora: cowslip ( <i>Primula veris</i> ) A, cow parsley ( <i>Anthriscus sylvestris</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) D, bramble ( <i>Rubus</i> sp.) F. Two willow sp. ( <i>Salix</i> sp.), hazel ( <i>Corylus avellana</i> ) R.
AB	H-AB-034	h2a - hedgerow (priority habitat)	Flailed hedgerow, uniform in height and width.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), field maple ( <i>Acer campestre</i> ), rose sp. ( <i>Rosa</i> sp.), hop ( <i>Humulus lupulus</i> ). Ground flora: cowslip ( <i>Primula veris</i> ) F, lords-and-ladies ( <i>Arum maculatum</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) F, willowherb species ( <i>Epilobium</i> sp.) A, bramble ( <i>Rubus</i> sp.) F, nettle ( <i>Urtica dioica</i> ) F, bristly ox-tongue ( <i>Helminthotheca echioides</i> ) O
AB	H-AB-033	h2a - hedgerow (priority habitat)	Uniform flailed blackthorn hedgerow with some trees.	Blackthorn ( <i>Prunus spinosa</i> ), oak sp. ( <i>Quercus</i> sp.), hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), sycamore ( <i>Acer pseudoplatanus</i> ). Ground flora: cowslip ( <i>Primula veris</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), hogweed ( <i>Heracleum sphondylium</i> ), willowherb species ( <i>Epilobium</i> sp.), dock sp. ( <i>Rumex</i> sp.), dog's mercury ( <i>Mercurialis perennis</i> ), thistle sp. ( <i>Cirsium</i> sp.).
AB	H-AB-032	h2a - hedgerow (priority habitat)	Flailed hedge with large gaps, along dry ditch.	Hawthorn ( <i>Crataegus monogyna</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, rose sp. ( <i>Rosa</i> sp.) O, dogwood ( <i>Cornus sanguinea</i> ) O. Ground flora: cowslip ( <i>Primula veris</i> ) A, nettle ( <i>Urtica dioica</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) F, dog's mercury ( <i>Mercurialis perennis</i> ) F, dock sp. ( <i>Rumex</i> sp.) F, bindweed sp. ( <i>Convolvulus</i> sp.) O
AB	H-AB-031	h2a - hedgerow (priority habitat)	Short, flailed hedgerow along dry ditch.	Blackthorn ( <i>Prunus spinosa</i> ) D, rose sp. ( <i>Rosa</i> sp.) O, hawthorn ( <i>Crataegus monogyna</i> ) F, ash ( <i>Fraxinus excelsior</i> ) R, field maple ( <i>Acer campestre</i> ) R. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) F, hogweed ( <i>Heracleum sphondylium</i> ) F, nettle ( <i>Urtica dioica</i> ) F, lords-and-ladies ( <i>Arum maculatum</i> ) O, cowslip ( <i>Primula veris</i> ) F.
AB	H-AB-030	h2a - hedgerow (priority habitat)	Short, heavily flailed hedgerow.	Field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, rose sp. ( <i>Rosa</i> sp.) F, hawthorn ( <i>Crataegus monogyna</i> ) A. Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ) F, dog's mercury ( <i>Mercurialis perennis</i> ) F, nettle ( <i>Urtica dioica</i> ) O, dock sp. ( <i>Rumex</i> sp.) O, cowslip ( <i>Primula veris</i> ) F, knapweed ( <i>Centaurea nigra</i> ) O.
AB	H-AB-029	h2a - hedgerow (priority habitat)	Native hedgerow with trees.	N/A. Not surveyed
AB	H-AB-028	h2a - hedgerow (priority habitat)	Defunct hedgerow on a hedge bank. Ditch with a small tree line present. Semi-mature and mature present.	Two ash ( <i>Fraxinus excelsior</i> ), oak sp. ( <i>Quercus</i> sp.).
AB	H-AB-027	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
AB	H-AB-026	h2a - hedgerow (priority habitat)	Flailed hedgerow along a ditch and road boundary.	Blackthorn ( <i>Prunus spinosa</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) D, willow sp. ( <i>Salix</i> sp.). Ground flora: nettle ( <i>Urtica dioica</i> ), hogweed ( <i>Heracleum sphondylium</i> ), plantain sp. ( <i>Plantago</i> sp.), dock sp. ( <i>Rumex</i> sp.), moss sp. ( <i>Bryophyta</i> sp.), cowslip ( <i>Primula veris</i> ).

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-025	h2a - hedgerow (priority habitat)	Tall, shrubby hedgerow along and arable field boundary and bordering a garden. Abruptly changes to shorter hedgerow at the end of the garden.	Hawthorn ( <i>Crataegus monogyna</i> ) D, ash ( <i>Fraxinus excelsior</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) F, plum ( <i>Prunus domestica</i> ) O, rose sp. ( <i>Rosa</i> sp.) O, hazel ( <i>Corylus avellana</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, field maple ( <i>Acer campestre</i> ) O. Ground flora: cowslip ( <i>Primula veris</i> ) F, lords-and-ladies ( <i>Arum maculatum</i> ) O, nettle ( <i>Urtica dioica</i> ) F, dog's mercury ( <i>Mercurialis perennis</i> ) F, knapweed ( <i>Centaurea nigra</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) F.
AB	H-AB-023	h2a - hedgerow (priority habitat)	Shrubby hedgerow becoming increasingly sparse towards the eastern end. Surveyed using binoculars due to access constraints.	Blackthorn ( <i>Prunus spinosa</i> ) A, dog's mercury ( <i>Mercurialis perennis</i> ) F, primrose ( <i>Primula vulgaris</i> ) O, hogweed ( <i>Heracleum sphondylium</i> ) F, oak sp. ( <i>Quercus</i> sp.) F, field maple ( <i>Acer campestre</i> ) O, dandelion sp. ( <i>Taraxacum</i> sp.) O, willow sp. ( <i>Salix</i> sp.) O, hazel ( <i>Corylus avellana</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, lords-and-ladies ( <i>Arum maculatum</i> ) O, nettle ( <i>Urtica dioica</i> ) O, ivy ( <i>Hedera helix</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R.
AB	H-AB-022	h2a - hedgerow (priority habitat)	Hedgerow in good condition. Mature ash trees present.	Section 1: blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), oak sp. ( <i>Quercus</i> sp.). Section 2: blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), oak sp. ( <i>Quercus</i> sp.). Section 3: hawthorn ( <i>Crataegus monogyna</i> ), 4x mature ash., bramble ( <i>Rubus</i> sp.), ivy ( <i>Hedera helix</i> ), elder ( <i>Sambucus nigra</i> ), holly ( <i>Ilex aquifolium</i> ). Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), nettle ( <i>Urtica dioica</i> ), dock sp. ( <i>Rumex</i> sp.), dandelion sp. ( <i>Taraxacum</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cowslip ( <i>Primula veris</i> ).
AB	H-AB-021	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow with trees. Associated with a dry ditch. Hedgerow has been flailed.	Ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ), elder ( <i>Sambucus nigra</i> ), oak sp. ( <i>Quercus</i> sp.). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), dock sp. ( <i>Rumex</i> sp.).
AB	H-AB-020	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow with trees. Associated with a dry ditch. Hedgerow has been flailed.	Ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ), elder ( <i>Sambucus nigra</i> ), oak sp. ( <i>Quercus</i> sp.). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), dock ( <i>Rumex</i> sp.).
AB	H-AB-019	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow with trees. Associated with a dry ditch. Hedgerow has been flailed.	Ash ( <i>Fraxinus excelsior</i> ), hawthorn ( <i>Crataegus monogyna</i> ), elder ( <i>Sambucus nigra</i> ), oak sp. ( <i>Quercus</i> sp.). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), nettle ( <i>Urtica dioica</i> ), cleavers ( <i>Galium aparine</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), dock sp. ( <i>Rumex</i> sp.).
AB	H-AB-018	h2 - hedgerow	UKHab category derived from aerial imagery	-
AB	H-AB-017	h2a - hedgerow (priority habitat)	Hedgerow with sparse tree line and veteran oaks.	Veteran oak sp. ( <i>Quercus</i> sp.). Ground flora: nettle ( <i>Urtica dioica</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), cleavers ( <i>Galium aparine</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ).
AB	H-AB-016	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow with trees. Associated with a dry ditch. Hedgerow has been flailed.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ), willow sp. ( <i>Salix</i> sp.), oak sp. ( <i>Quercus</i> sp.). Section 2: blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ). Section 3: oak sp. ( <i>Quercus</i> sp.), hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), primrose ( <i>Primula vulgaris</i> ), nettle ( <i>Urtica dioica</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), cleavers ( <i>Galium aparine</i> ), curled dock ( <i>Rumex crispus</i> ), ivy ( <i>Hedera helix</i> )
AB	H-AB-015	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow along a dry ditch. Hedgerow has been flailed.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ), willow sp. ( <i>Salix</i> sp.), rose sp. ( <i>Rosa</i> sp.). Section 2: hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ), field maple ( <i>Acer campestre</i> ). Section 3: hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ). Ground flora: ivy ( <i>Hedera helix</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), dandelion sp. ( <i>Taraxacum</i> sp.).

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
AB	H-AB-014	h2a - hedgerow (priority habitat)	Native Hedgerow with trees - Associated with bank or ditch. Along garden.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), ash ( <i>Fraxinus excelsior</i> ), Leyland cypress ( <i>Cupressus x leylandii</i> ). Ground flora: ground-ivy ( <i>Glechoma hederacea</i> ), ivy ( <i>Hedera helix</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), daffodil sp. ( <i>Narcissus</i> sp.), dog's mercury ( <i>Mercurialis perennis</i> ), nettle ( <i>Urtica dioica</i> ), bramble ( <i>Rubus</i> sp.).
AB	H-AB-013	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow with trees. Associated with a dry ditch. Hedgerow has been flailed.	hawthorn ( <i>Crataegus monogyna</i> ), ash ( <i>Fraxinus excelsior</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), ground-ivy ( <i>Hedera helix</i> ), bramble ( <i>Rubus</i> sp.), teasel sp. ( <i>Dipsacus</i> sp.), rose sp. ( <i>Rosa</i> sp.).
AB	H-AB-012	h2a - hedgerow (priority habitat)	Native Hedgerow.	N/A. Not surveyed
AB	H-AB-011	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees. Not surveyed.	N/A. Not surveyed
AB	H-AB-010	h2a - hedgerow (priority habitat)	Native, recently planted hedgerow.	Hawthorn ( <i>Crataegus monogyna</i> ) D. Ground flora: dandelion sp. ( <i>Taraxacum</i> sp.).
AB	H-AB-009	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees. Not fully surveyed due to land access constraints.	Hawthorn ( <i>Crataegus monogyna</i> ) D, ivy ( <i>Hedera helix</i> ) D. Ground floral cock's-foot ( <i>Dactylis glomerata</i> ), nettle ( <i>Urtica dioica</i> ).
AB	H-AB-008	h2a - hedgerow (priority habitat)	Flailed native hedgerow bordering footpath.	Hawthorn ( <i>Crataegus monogyna</i> ) D.
AB	H-AB-007	h2a - hedgerow (priority habitat)	Flailed hedgerow bordering footpath and arable field. Heavily managed and defunct.	N/A. Not surveyed
AB	H-AB-004	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees, flailed and coppiced.	Elder ( <i>Sambucus nigra</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) A, coppiced hazel ( <i>Corylus avellana</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) D, immature pedunculate oak ( <i>Quercus robur</i> ) R, field maple ( <i>Acer campestre</i> ) R. Ground flora: moschatel ( <i>Adoxa moschatellina</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), dog's mercury ( <i>Mercurialis perennis</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cleavers ( <i>Galium aparine</i> ), red dead-nettle ( <i>Lamium purpureum</i> ), nettle ( <i>Urtica dioica</i> ), lesser celandine ( <i>Ranunculus ficaria</i> ).
AB	H-AB-002	h2a - hedgerow (priority habitat)	Native species rich hedgerow, along a dry ditch. Planted and flailed.	Hawthorn ( <i>Crataegus monogyna</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) D, elder ( <i>Sambucus nigra</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, hazel ( <i>Corylus avellana</i> ) O, dog-rose ( <i>Rosa canina</i> ). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ), nettle ( <i>Urtica dioica</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), ivy ( <i>Hedera helix</i> ), cleavers ( <i>Galium aparine</i> ), bramble ( <i>Rubus</i> sp.).
C	H-C-01	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
C	H-C-02	h2a - hedgerow (priority habitat)	Planted, native, species rich hedgerow along a road.	Hazel ( <i>Corylus avellana</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) A, field maple ( <i>Acer campestre</i> ) F, holly ( <i>Ilex aquifolium</i> ) R, nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), bramble ( <i>Rubus</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ).
C	H-C-03	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
C	H-C-04	h2a - hedgerow (priority habitat)	Hedgerow at the end of a woodland parcel, along arable field boundary.	Hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), elder ( <i>Sambucus nigra</i> ), oak sp. ( <i>Quercus</i> sp.). Ground flora: nettle ( <i>Urtica dioica</i> ), lesser celandine ( <i>Ranunculus ficaria</i> ), cleavers ( <i>Galium aparine</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ).

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
C	H-C-05	h2a - hedgerow (priority habitat)	Hedgerow along an arable field and road boundary.	Hawthorn ( <i>Crataegus monogyna</i> ), honeysuckle ( <i>Lonicera periclymenum</i> ). Ground flora: cleavers, nettle ( <i>Urtica dioica</i> ), garlic mustard ( <i>Alliaria petiolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), spurge ( <i>Euphorbia</i> ).
C	H-C-06	h2a - hedgerow (priority habitat)	Native hedgerow with trees, flailed.	Hawthorn ( <i>Crataegus monogyna</i> ) D, sapling ash ( <i>Fraxinus excelsior</i> ). Ground flora: garlic mustard ( <i>Alliaria petiolata</i> ), cleavers ( <i>Galium aparine</i> ), nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), white dead-nettle ( <i>Lamium album</i> ), hogweed ( <i>Heracleum sphondylium</i> ), cleavers ( <i>Galium aparine</i> ).
C	H-C-09	h2 - hedgerow	UKHab category derived from aerial imagery	-
D	H-D-01	h2a - hedgerow (priority habitat)	Large unmanaged hedgerow possibly expanding into woodland habitat at edges. Large number of fallen trees. Bordered by arable fields and grassland habitat.	Hawthorn ( <i>Crataegus monogyna</i> ) A, elm sp. ( <i>Ulmus</i> sp.) A, ivy ( <i>Hedera helix</i> ) F. Ground flora: nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) A, creeping thistle ( <i>Cirsium arvense</i> ) F, nettle ( <i>Urtica dioica</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) D, bluebell ( <i>Hyacinthoides</i> sp.) O, cock's-foot ( <i>Dactylis glomerata</i> ) R.
D	H-D-02	h2a - hedgerow (priority habitat)	Hedgerow, not fully surveyed.	Section 1: blackthorn ( <i>Prunus spinosa</i> ), elm sp. ( <i>Ulmus</i> sp.), pedunculate oak ( <i>Quercus robur</i> ), hawthorn ( <i>Crataegus monogyna</i> ), Prunus sp. Ground flora: bluebell ( <i>Hyacinthoides</i> sp.), garlic mustard ( <i>Alliaria petiolata</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), lords-and-ladies ( <i>Arum maculatum</i> ). Section 2: wild cherry ( <i>Prunus avium</i> ), lime ( <i>Tilia x europaea</i> ), elm species ( <i>Ulmus</i> sp.), pedunculate oak ( <i>Quercus robur</i> ), hawthorn ( <i>Crataegus monogyna</i> ), hazel ( <i>Corylus avellana</i> ).
D	H-D-03	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
D	H-D-04	h2a - hedgerow (priority habitat)	Shrubby hedgerow, flailed along a dry ditch.	Field maple ( <i>Acer campestre</i> ), blackthorn ( <i>Prunus spinosa</i> ), wych elm ( <i>Ulmus glabra</i> ), large-leaved lime ( <i>Tilia platyphyllos</i> ), rose sp. ( <i>Rosa</i> sp.), honeysuckle ( <i>Lonicera periclymenum</i> ), holly ( <i>Ilex aquifolium</i> ), blackthorn ( <i>Prunus spinosa</i> ), pedunculate oak ( <i>Quercus robur</i> ) R. Ground flora: cleavers ( <i>Galium aparine</i> ), dandelion sp. ( <i>Taraxacum</i> sp.), bramble ( <i>Rubus</i> sp.), garlic mustard ( <i>Alliaria petiolata</i> ), wood avens ( <i>Geum urbanum</i> ).
D	H-D-05	h2a - hedgerow (priority habitat)	Native, species rich hedgerow, with flailed management. Ditch present at the corner.	Hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), dog-rose ( <i>Rosa canina</i> ), hazel ( <i>Corylus avellana</i> ), sapling ash ( <i>Fraxinus excelsior</i> ), sapling oak sp. ( <i>Quercus</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), holly ( <i>Ilex aquifolium</i> ).
D	H-D-06	h2a - hedgerow (priority habitat)	Hedgerow with trees and hedge bank. A dry ditch.	Trees: sycamore ( <i>Acer pseudoplatanus</i> ), hornbeam ( <i>Carpinus betulus</i> ), pedunculate oak ( <i>Quercus robur</i> ). Section 1: sycamore, hawthorn ( <i>Crataegus monogyna</i> ) D, field maple ( <i>Acer campestre</i> ) R, wild cherry ( <i>Prunus avium</i> ) R, hazel ( <i>Corylus avellana</i> ) R, hornbeam ( <i>Carpinus betulus</i> ) R, eagle fern ( <i>Pteridium aquilinum</i> ) R. Section 2: pedunculate oak ( <i>Quercus robur</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) D, hazel ( <i>Corylus avellana</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, Cherry ( <i>Prunus avium</i> ) R. Section 3: hawthorn ( <i>Crataegus monogyna</i> ) D, blackthorn ( <i>Prunus spinosa</i> ) O, field maple ( <i>Acer campestre</i> ) O, hazel ( <i>Corylus avellana</i> ) R, dog-rose ( <i>Rosa canina</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, common dogwood ( <i>Cornus sanguinea</i> ) R, small-leaved lime ( <i>Tilia cordata</i> ) O.
D	H-D-07	h2a - hedgerow (priority habitat)	Hedgerow along a road with woodland on south-west boundary. Hedgerow connection on the opposite side of the road.	Section 1: hazel ( <i>Corylus avellana</i> ) O, dog-rose ( <i>Rosa canina</i> ) F, elder ( <i>Sambucus nigra</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) D, holly ( <i>Ilex aquifolium</i> ) O, field maple ( <i>Acer campestre</i> ) O, cherry sp. ( <i>Prunus</i> sp.) O. Section 2: hazel ( <i>Corylus avellana</i> ) R, dog-rose ( <i>Rosa canina</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, field maple ( <i>Acer campestre</i> ) R, small-leaved lime ( <i>Tilia cordata</i> ) D, cherry sp. ( <i>Prunus</i> sp.) R, willow sp. ( <i>Salix</i> sp.) D. Section 3: hazel ( <i>Corylus avellana</i> ) F, dog-rose ( <i>Rosa canina</i> ) sp. ( <i>Rosa</i> sp.) O, dogwood ( <i>Cornus sanguinea</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) A, holly ( <i>Ilex aquifolium</i> ) R, cherry sp. ( <i>Prunus</i> sp.) R, wild cherry ( <i>Prunus avium</i> ) R, elm sp. ( <i>Ulmus</i> sp.) R

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
D	H-D-08	h2a - hedgerow (priority habitat)	Hedgerow on a hedge bank. Dead oak present. Unable to survey entire hedgerow length.	Section 1: hazel ( <i>Corylus avellana</i> ) A, dogwood ( <i>Cornus sanguinea</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, field maple ( <i>Acer campestre</i> ) F. Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) R, wood melick ( <i>Melica uniflora</i> ) O, wood false-brome ( <i>Brachypodium sylvaticum</i> ) R, wood avens ( <i>Geum urbanum</i> ) R.
D	H-D-09	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees. Coppiced and runs along a dry ditch.	Oak sp. ( <i>Quercus</i> sp.), hawthorn ( <i>Crataegus monogyna</i> ), wild cherry ( <i>Prunus avium</i> ), holly ( <i>Ilex aquifolium</i> ), hornbeam ( <i>Carpinus betulus</i> ), silver birch ( <i>Betula pendula</i> ), lime species ( <i>Tilia</i> sp.), field maple ( <i>Acer campestre</i> ), ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ), Leyland cypress ( <i>Cupressus × leylandii</i> ), sycamore ( <i>Acer pseudoplatanus</i> ). Ground flora: wild garlic ( <i>Allium ursinum</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), bluebell sp. ( <i>Hyacinthoides</i> sp.), dog's mercury ( <i>Mercurialis perennis</i> ).
D	H-D-10	h2a - hedgerow (priority habitat)	Species rich hedgerow with occasional, hedgerow was not fully surveyed.	Spindle ( <i>Euonymus europaeus</i> ), hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ), cherry plum ( <i>Prunus cerasifera</i> ), willow ( <i>Salix</i> sp.), hazel ( <i>Corylus avellana</i> ), ash ( <i>Fraxinus excelsior</i> ), dogwood ( <i>Cornus sanguinea</i> ), pedunculate oak ( <i>Quercus robur</i> ), Prunus sp. Ground flora: Alexanders ( <i>Smyrnium olusatrum</i> ), cow parsley, garlic mustard ( <i>Alliaria petiolata</i> ), ramsons ( <i>Allium ursinum</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), groundsel sp. ( <i>Senecio</i> ).
D	H-D-11	h2a - hedgerow (priority habitat)	Native hedgerow.	Hawthorn ( <i>Crataegus monogyna</i> ), dogwood ( <i>Cornus sanguinea</i> ), field maple ( <i>Acer campestre</i> ), poplar ( <i>Populus</i> sp.)
D	H-D-13	h2a - hedgerow (priority habitat)	Hedgerow with trees planted on a clear bank. Flailed on western side by arable field with little ground flora. Limited access to the hedgerow, so hedgerow was surveyed within the Order Limits.	Small-leaved lime ( <i>Tilia cordata</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) F, wild cherry ( <i>Prunus avium</i> ) F, field maple ( <i>Acer campestre</i> ), dogwood ( <i>Cornus sanguinea</i> ) R, elder ( <i>Sambucus nigra</i> ) R, crab apple ( <i>Malus sylvestris</i> ) R, semi mature pedunculate oak ( <i>Quercus robur</i> ). Ground flora: nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) R, garlic mustard ( <i>Alliaria petiolata</i> ) O, greater stitchwort ( <i>Stellaria holostea</i> ) R, bluebell sp. ( <i>Hyacinthoides</i> sp.) O, lord-and-ladies ( <i>Arum maculatum</i> ) R.
D	H-D-14	h2a - hedgerow (priority habitat)	Native planted hedgerow with trees. Flailed management.	Willow sp. ( <i>Salix</i> sp.), bramble ( <i>Rubus</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ), birch sp. ( <i>Betula</i> sp.), cleavers ( <i>Galium aparine</i> ), elm sp. ( <i>Ulmus</i> sp.), elder ( <i>Sambucus nigra</i> ), holly ( <i>Ilex aquifolium</i> ), hazel ( <i>Corylus avellana</i> ), oak sp. ( <i>Quercus</i> sp.). Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ), lords-and-ladies ( <i>Arum maculatum</i> ), ground-ivy ( <i>Glechoma hederacea</i> ), thistle sp. ( <i>Cirsium</i> sp.), stitchwort sp. ( <i>Stellaria</i> sp.), ivy ( <i>Hedera helix</i> ), bluebell sp. ( <i>Hyacinthoides</i> sp.), germander speedwell ( <i>Veronica chamaedrys</i> ), dandelion sp. ( <i>Taraxacum</i> sp.), nettle ( <i>Urtica dioica</i> ), dead-nettle sp. ( <i>Lamium</i> sp.), hogweed ( <i>Heracleum sphondylium</i> ), fern sp.
D	H-D-15	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
D	H-D-16	h2a - hedgerow (priority habitat)	Native hedgerow	Hawthorn ( <i>Crataegus monogyna</i> ) A, hazel ( <i>Corylus avellana</i> ) A, field maple ( <i>Acer campestre</i> ) A. Ground flora: nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), cleavers ( <i>Galium aparine</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), thistle sp. ( <i>Cirsium</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ), plantain sp. ( <i>Plantago</i> sp.), red campion ( <i>Silene dioica</i> ), curled dock ( <i>Rumex crispus</i> ), cranes-bill sp. ( <i>Geranium</i> sp.), bramble ( <i>Rubus</i> sp.).
D	H-D-17	h2a - hedgerow (priority habitat)	Hedgerow along field boundary and woodland, merges into woodland and no longer stockproof at one end.	Hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) F, dog-rose ( <i>Rosa canina</i> ) O, mature field maple ( <i>Acer campestre</i> ) R, mature pedunculate oak ( <i>Quercus robur</i> ) R. Ground flora: nettle ( <i>Urtica dioica</i> ) D, hedge parsley sp. ( <i>Torilis</i> sp.), Yorkshire fog ( <i>Holcus lanatus</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ).

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
D	H-D-19	h2a - hedgerow (priority habitat)	Native hedgerow planted and flailed.	Hawthorn ( <i>Crataegus monogyna</i> ) D, bramble ( <i>Rubus</i> sp.) F, field maple ( <i>Acer campestre</i> ) O, cherry laurel ( <i>Prunus laurocerasus</i> ). Ground flora: nettle ( <i>Urtica dioica</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), daisy ( <i>Bellis perennis</i> ), wood speedwell ( <i>Veronica montana</i> ).
D	H-D-20	h2a - hedgerow (priority habitat)	Planted and flailed, native species rich hedgerow.	Wild cherry ( <i>Prunus avium</i> ), hawthorn ( <i>Crataegus monogyna</i> ), holly ( <i>Ilex aquifolium</i> ), elm sp. ( <i>Ulmus</i> sp.), ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ), field maple ( <i>Acer campestre</i> ), elder ( <i>Sambucus nigra</i> ), Ground flora: cock's-foot ( <i>Dactylis glomerata</i> ), nettle ( <i>Urtica dioica</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ), hogweed ( <i>Heracleum sphondylium</i> ), mallow sp. ( <i>Malva</i> sp.), perennial ryegrass ( <i>Lolium perenne</i> ), false oat-grass ( <i>Arrhenatherum elatius</i> ), tufted hair-grass ( <i>Deschampsia cespitosa</i> ).
D	H-D-22	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
D	H-D-23	h2a - hedgerow (priority habitat)	Hedgerow, not fully surveyed due to land access constraints and health and safety.	Pedunculate oak ( <i>Quercus robur</i> ), hornbeam ( <i>Carpinus betulus</i> ), elder ( <i>Sambucus nigra</i> ), field maple ( <i>Acer campestre</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), holly ( <i>Ilex aquifolium</i> ), hawthorn ( <i>Crataegus monogyna</i> ). Ground flora: nettle ( <i>Urtica dioica</i> ), bramble ( <i>Rubus</i> sp.), dog's mercury ( <i>Mercurialis perennis</i> ).
D	H-D-24	h2a - hedgerow (priority habitat)	Small unmanaged hedgerow in grassland with parallel hedgerow to the west and orchard to the north.	Hawthorn ( <i>Crataegus monogyna</i> ) D, hazel ( <i>Corylus avellana</i> ) O, elm sp. ( <i>Ulmus</i> sp.) A, dog-rose ( <i>Rosa canina</i> ) F, ivy ( <i>Hedera helix</i> ) O. Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ) F, creeping thistle ( <i>Cirsium arvense</i> ) F, dandelion sp. ( <i>Taraxacum</i> sp.) O, cow parsley ( <i>Anthriscus sylvestris</i> ) O, hemlock ( <i>Conium maculatum</i> ) O.
D	H-D-025	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees	N/A. Not surveyed
E	H-E-01	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees	N/A. Not surveyed
E	H-E-02	h2a - hedgerow (priority habitat)	Old hedgerow on hedge bank. Separates and orchard and a field. Unmanaged on orchard side and flailed on roadside. Significant elm dieback and regrowth. For practical and access reasons only a representative section of this very long hedgerow has been surveyed	Section 1: hawthorn ( <i>Crataegus monogyna</i> ) R, elm sp. ( <i>Ulmus</i> sp.) A, holly ( <i>Ilex aquifolium</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R. Section 2: hawthorn ( <i>Crataegus monogyna</i> ) F, elm sp. ( <i>Ulmus</i> sp.) D, dog-rose ( <i>Rosa canina</i> ) R. Section 3: holly ( <i>Ilex aquifolium</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, field maple ( <i>Acer campestre</i> ) F, elder ( <i>Sambucus nigra</i> ) O.
E	H-E-03	h2a - hedgerow (priority habitat)	Overgrown, unmanaged hedgerow on a hedge bank with many rabbit warrens present. Approximately 15% dead elm due to Dutch elm disease. Turning into a line of trees.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) A, elm sp. ( <i>Ulmus</i> sp.) R, elder ( <i>Sambucus nigra</i> ) O. Section 2: blackthorn ( <i>Prunus spinosa</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) F, spindle ( <i>Euonymus europaeus</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, field maple ( <i>Acer campestre</i> ) R. Section 3: dog-rose ( <i>Rosa canina</i> ) R, hazel ( <i>Corylus avellana</i> ) R. Section 3: blackthorn ( <i>Prunus spinosa</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) F, elder ( <i>Sambucus nigra</i> ) O.
E	H-E-04	h2a - hedgerow (priority habitat)	UKHab category derived from aerial imagery	-
E	H-E-05	h2a - hedgerow (priority habitat)	Native hedgerow with trees	Rose ( <i>Rosa</i> sp.) R, willow ( <i>Salix</i> sp.) O, elder ( <i>Sambucus nigra</i> ) O, bramble ( <i>Rubus</i> sp.) D. Ground flora: nettle ( <i>Urtica dioica</i> ) D, cleavers ( <i>Galium aparine</i> ) A, willow herb ( <i>Epilobium</i> sp.) F, dead nettle ( <i>Lamium</i> sp.) O, ground ivy ( <i>Glechoma hederacea</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) A
E	H-E-06	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow which has been flailed.	Hawthorn ( <i>Crataegus monogyna</i> ) O, ash ( <i>Fraxinus excelsior</i> ), blackthorn ( <i>Prunus spinosa</i> ) O, holly ( <i>Ilex aquifolium</i> ), ash ( <i>Fraxinus excelsior</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) sapling R, mature pedunculate oak ( <i>Quercus robur</i> ), mallow sp. ( <i>Malva</i> sp.) R. Ground flora: cleavers ( <i>Galium aparine</i> ) A, nettle ( <i>Urtica dioica</i> ) F, hedgerow crane's bill ( <i>Geranium pyrenaicum</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) O, false oat-grass ( <i>Arrhenatherum elatius</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), barren brome ( <i>Bromus sterilis</i> ), black bryony ( <i>Tamus communis</i> )

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
E	H-E-07	h2a - hedgerow (priority habitat)	Hedgerow with some dead elms and uniformly cut.	Elm sp. ( <i>Ulmus</i> sp.) A, field maple ( <i>Acer campestre</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R, old man's beard ( <i>Clematis vitalba</i> ), elder ( <i>Sambucus nigra</i> ), ash ( <i>Fraxinus excelsior</i> ), spindle ( <i>Euonymus europaeus</i> ) O. Ground flora: nettle ( <i>Urtica dioica</i> ) A, cleavers ( <i>Galium aparine</i> ) A, garlic mustard ( <i>Alliaria petiolata</i> ), hedge mustard ( <i>Sisymbrium officinale</i> ), white campion ( <i>Silene latifolia</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), cock's-foot ( <i>Dactylis glomerata</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), false oat-grass ( <i>Arrhenatherum elatius</i> ).
E	H-E-08	h2a - hedgerow (priority habitat)	Native Hedgerow with trees.	N/A. Not surveyed
E	H-E-09	h2a - hedgerow (priority habitat)	Overgrown hedgerow with trees, including some dead English elm.	Holly ( <i>Ilex aquifolium</i> ) R, English elm sp. ( <i>Ulmus</i> sp.) O, field maple ( <i>Acer campestre</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R. Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ), nettle ( <i>Urtica dioica</i> ), black knapweed ( <i>Centaurea nigra</i> ), false oat-grass ( <i>Arrhenatherum elatius</i> ), poppy sp. ( <i>Papaver</i> sp.), cock's-foot ( <i>Dactylis glomerata</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), lesser stitchwort ( <i>Stellaria graminea</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), ribwort plantain ( <i>Plantago lanceolata</i> ), hogweed ( <i>Heracleum sphondylium</i> ), greater stitchwort ( <i>Stellaria holostea</i> ), common mouse-ear ( <i>Cerastium fontanum</i> ), rose sp. ( <i>Rosa</i> sp.).
E	H-E-10	h2a - hedgerow (priority habitat)	Unmanaged hedgerow with some dead elms.	elm sp. ( <i>Ulmus</i> sp.) F, ash ( <i>Fraxinus excelsior</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ), veteran pedunculate oak ( <i>Quercus robur</i> ), elder ( <i>Sambucus nigra</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) R, spindle ( <i>Euonymus europaeus</i> ) R. Ground flora: nettle ( <i>Urtica dioica</i> ) D, cleavers ( <i>Galium aparine</i> ) D, bramble ( <i>Rubus</i> sp.), bracken ( <i>Pteridium aquilinum</i> )
E	H-E-11	h2a - hedgerow (priority habitat)	Hedgerow separating two arable fields.	Field maple ( <i>Acer campestre</i> ), hawthorn ( <i>Crataegus monogyna</i> ), hazel ( <i>Corylus avellana</i> ), wych elm ( <i>Ulmus glabra</i> ), elder ( <i>Sambucus nigra</i> ), old man's beard ( <i>Clematis vitalba</i> ), oak sp. ( <i>Quercus</i> sp.)
E	H-E-12	h2a - hedgerow (priority habitat)	Hedgerow recently flailed. Bordering two arable fields.	Hawthorn ( <i>Crataegus monogyna</i> ) D, field maple ( <i>Acer campestre</i> ) F, ivy ( <i>Hedera helix</i> ) O, oak sp. ( <i>Quercus</i> sp.) R, holly ( <i>Ilex aquifolium</i> ) R. Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ) A, nettle ( <i>Urtica dioica</i> ) D, dead-nettle sp. ( <i>Lamium</i> sp.) O, dandelion sp. ( <i>Taraxacum</i> sp.) O, ground-ivy ( <i>Hedera helix</i> ) O, cleavers ( <i>Galium aparine</i> ) A, lords-and-ladies ( <i>Arum maculatum</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) R.
E	H-E-13	h2a - hedgerow (priority habitat)	Hedgerow with trees (some recently planted) alongside road and grassland.	Oak sp. ( <i>Quercus</i> sp.), field maple ( <i>Acer campestre</i> ), hawthorn ( <i>Crataegus monogyna</i> ), blackthorn ( <i>Prunus spinosa</i> ), crab apple ( <i>Malus sylvestris</i> ), bramble ( <i>Rubus</i> sp.), rose sp. ( <i>Rosa</i> sp.), hazel ( <i>Corylus avellana</i> ), apple sp. ( <i>Malus</i> sp.), ash ( <i>Fraxinus excelsior</i> ), old man's beard ( <i>Clematis vitalba</i> ).
E	H-E-14	h2a - hedgerow (priority habitat)	Shrubby, native species rich hedgerow.	Hawthorn ( <i>Crataegus monogyna</i> ) D, holly ( <i>Ilex aquifolium</i> ) R. Ground flora: cow parsley ( <i>Anthriscus sylvestris</i> ) D, dock sp. ( <i>Rumex</i> sp.) R, nettle ( <i>Urtica dioica</i> ) A, dandelion sp. ( <i>Taraxacum</i> sp.) O, cleavers ( <i>Galium aparine</i> ) A, ground-ivy ( <i>Glechoma hederacea</i> ) O.
E	H-E-15	h2a - hedgerow (priority habitat)	Flailed hedgerow with wet ditch, which flows to the south. Hedgerow adjacent to woodland.	Dogwood ( <i>Cornus sanguinea</i> ) O, oak sp. ( <i>Quercus</i> sp.) R, field maple ( <i>Acer campestre</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) D, nettle ( <i>Urtica dioica</i> ) A, willowherb sp. ( <i>Epilobium</i> sp.) O, cleavers ( <i>Galium aparine</i> ) O, common hogweed ( <i>Heracleum sphondylium</i> ) O.
E	H-E-16	h2a - hedgerow (priority habitat)	Hedgerow along road, recently flailed.	Bramble ( <i>Rubus</i> sp.) D, pedunculate oak ( <i>Quercus robur</i> ), bramble ( <i>Rubus</i> sp.) D, hawthorn ( <i>Crataegus monogyna</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) A, elder ( <i>Sambucus nigra</i> ) R. Ground flora: hemlock ( <i>Conium maculatum</i> ) R, ground-ivy ( <i>Glechoma hederacea</i> ) O, garlic mustard F, cleavers ( <i>Galium aparine</i> ) A, nettle ( <i>Urtica dioica</i> ) D, cow parsley ( <i>Anthriscus sylvestris</i> ) O, dead-nettle sp. ( <i>Lamium</i> sp.) O.

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
E	H-E-017	h2a - hedgerow (priority habitat)	Native Hedgerow	N/A. Not surveyed
F	H-F-01	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees	N/A. Not surveyed
F	H-F-02	h2a - hedgerow (priority habitat)	Native Hedgerow	N/A. Not surveyed
F	H-F-03	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees	N/A. Not surveyed
F	H-F-04	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees	N/A. Not surveyed
F	H-F-05	h2a - hedgerow (priority habitat)	Native Hedgerow	N/A. Not surveyed
F	H-F-06	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees	N/A. Not surveyed
F	H-F-08	h2a - hedgerow (priority habitat)	Native Hedgerow with trees	N/A. Not surveyed
F	H-F-09	h2a - hedgerow (priority habitat)	Hedgerow planted approximately 15 years ago, dividing two arable fields. Tree guards are still visible.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ) D, pedunculate oak ( <i>Quercus robur</i> ) R, ash ( <i>Fraxinus excelsior</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) F, English yew ( <i>Taxus baccata</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, holly ( <i>Ilex aquifolium</i> ) R. Section 2: hawthorn ( <i>Crataegus monogyna</i> ) A, field maple ( <i>Acer campestre</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) A, dog-rose ( <i>Rosa canina</i> ) O
F	H-F-10	h2a - hedgerow (priority habitat)	Hedgerow bordering a field and road with no margin. Flailed on the roadside. Some young emergent trees, mainly holly.	Section 1: elm sp. ( <i>Ulmus</i> sp.) F, pedunculate oak ( <i>Quercus robur</i> ) O, dog-rose R, holly ( <i>Ilex aquifolium</i> ) A, elder ( <i>Sambucus nigra</i> ) R, field maple ( <i>Acer campestre</i> ) O, Leyland cypress ( <i>Cupressus × leylandii</i> ) R. Section 2: elm sp. ( <i>Ulmus</i> sp.) D, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) F, holly ( <i>Ilex aquifolium</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) R. Section 3: elm sp. ( <i>Ulmus</i> sp.) D, hawthorn ( <i>Crataegus monogyna</i> ) R, holly ( <i>Ilex aquifolium</i> ) O
F	H-F-11	h2a - hedgerow (priority habitat)	Not considered important. Native hedgerow with trees.	N/A
F	H-F-12	h2a - hedgerow (priority habitat)	Hedgerow with hedge bank along a ditch, with relatively young trees that were trimmed well but left to grow tall. Track to the west and arable field to the east of the hedgerow.	Section 1: spindle ( <i>Euonymus europaeus</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, rose sp. ( <i>Rosa</i> sp.) R, hawthorn ( <i>Crataegus monogyna</i> ) R. Section 2: holly ( <i>Ilex aquifolium</i> ) R, elm sp. ( <i>Ulmus</i> sp.) likely wych elm ( <i>Ulmus glabra</i> ) A, hazel ( <i>Corylus avellana</i> ) R, field maple ( <i>Acer campestre</i> ). Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ) R.
F	H-F-13	h2b - other hedgerows	Hedgerow in poor condition.	Leyland cypress ( <i>Cupressus × leylandii</i> )
F	H-F-14	h2a - hedgerow (priority habitat)	Hedgerow on public right of way. Replanted in sections with trees. Meadow present on eastern boundary. Turning into a blackthorn scrub.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) A, elm sp. ( <i>Ulmus</i> sp.) R, dog-rose ( <i>Rosa canina</i> ) R. Section 2: blackthorn ( <i>Prunus spinosa</i> ) A, wild cherry ( <i>Prunus avium</i> ) R, cherry sp. ( <i>Prunus</i> sp.) R, hawthorn ( <i>Crataegus monogyna</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) O.
F	H-F-15	h2a - hedgerow (priority habitat)	Hedgerow lacking management and growing horizontally instead of vertically. Wetland area present on eastern section.	Section 1: elder ( <i>Sambucus nigra</i> ) D, goat willow ( <i>Salix caprea</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R.
F	H-F-16	h2a - hedgerow (priority habitat)	Hedgerow could be historical but planted with poplars in tight spaces approximately 1m. Wet ditch present flowing south.	Section 1: mature hybrid black poplar ( <i>Populus x canadensis</i> agg.) D, hawthorn ( <i>Crataegus monogyna</i> ), dog-rose ( <i>Rosa canina</i> ) R, elder ( <i>Sambucus nigra</i> ) R, grey willow ( <i>Salix cinerea</i> ) R, hazel ( <i>Corylus avellana</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) R, holly ( <i>Ilex aquifolium</i> ) R. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) LA.

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
F	H-F-17	h2a - hedgerow (priority habitat)	Hedgerow which is fenced and separating horse paddocks.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) A. Section 2: grey willow ( <i>Salix cinerea</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O.
F	H-F-18	h2a - hedgerow (priority habitat)	Hedgerow separating paddocks, possibly historical as it is growing through central fence on eastern aspect. Likely older in central and western sections. Ditch and hedge bank present from centre to western aspect down from south to north. Bramble growing over dead branches. Dead elm present along the whole hedgerow.	Section 1: wild cherry ( <i>Prunus avium</i> ) R, field maple ( <i>Acer campestre</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, elm sp. ( <i>Ulmus</i> sp.) A, dog-rose ( <i>Rosa canina</i> ) R, elder ( <i>Sambucus nigra</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O.
F	H-F-19	h2a - hedgerow (priority habitat)	Hedgerow separating a field and a road.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ) O, hazel ( <i>Corylus avellana</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, ash ( <i>Fraxinus excelsior</i> ) R, almost dead pedunculate oak ( <i>Quercus robur</i> ) O.
F	H-F-20	h2a - hedgerow (priority habitat)	Hedgerow on hedge bank with ditch. Trees growing over shrub hedgerow species stumping their growth. Mature oaks at regular intervals. Some damage evident from flailing.	Section 1: elm sp. ( <i>Ulmus</i> sp.) F, pedunculate oak ( <i>Quercus robur</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, field maple ( <i>Acer campestre</i> ) R, apple sp. ( <i>Malus</i> sp.) R. Section 2: pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, hazel ( <i>Corylus avellana</i> ) O.
F	H-F-21	h2a - hedgerow (priority habitat)	Hedgerow with hedge bank and bordering a road. Mature oaks present in the southern section.	Section 1: field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, wild cherry ( <i>Prunus avium</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, elm sp. ( <i>Ulmus</i> sp.) R, hazel ( <i>Corylus avellana</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R. Section 2: field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, wild cherry ( <i>Prunus avium</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) F, hazel ( <i>Corylus avellana</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) F.
F	H-F-22	h2a - hedgerow (priority habitat)	Hedgerow with a ditch bordering and agricultural field and a road.	Section 1: field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) A, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, holly ( <i>Ilex aquifolium</i> ) R. Section 2: blackthorn ( <i>Prunus spinosa</i> ) A, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) A, ash ( <i>Fraxinus excelsior</i> ) R, elder ( <i>Sambucus nigra</i> ) R, willow sp. ( <i>Salix</i> sp.) R. Section 3: blackthorn ( <i>Prunus spinosa</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) D, holly ( <i>Ilex aquifolium</i> ) R, elder ( <i>Sambucus nigra</i> ) R, spindle ( <i>Euonymus europaeus</i> ) R. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) O, wood avens ( <i>Geum urbanum</i> ) O
F	H-F-23	h2a - hedgerow (priority habitat)	Hedgerow connected to woodland on southern boundary which consists partly of a garden with non-native species present.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, apple sp. ( <i>Malus</i> sp.) O, hazel ( <i>Corylus avellana</i> ) R. Section 2: blackthorn ( <i>Prunus spinosa</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, field maple ( <i>Acer campestre</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) A, cherry sp. ( <i>Prunus</i> sp.) R, ash ( <i>Fraxinus excelsior</i> ) R, elder ( <i>Sambucus nigra</i> ) R, holly ( <i>Ilex aquifolium</i> ) R. Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) O.
F	H-F-24	h2a - hedgerow (priority habitat)	Hedgerow - likely planted but could have possibly grown naturally. Young woody species on the west of the hedgerow.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) R, dog-rose ( <i>Rosa canina</i> ) R. Section 2: blackthorn ( <i>Prunus spinosa</i> ) R, field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R, spindle ( <i>Euonymus europaeus</i> ) R.
F	H-F-25	h2a - hedgerow (priority habitat)	Ditch with hedge bank 3m in depth from east to west. Canopy grown out and spreading, creating space under the hedgerow in sections but lacking scrub ground cover.	Section 1: apple sp. ( <i>Malus</i> sp.) O, cherry sp. ( <i>Prunus</i> sp.) O, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) O, field maple ( <i>Acer campestre</i> ) F, small-leaved lime ( <i>Tilia cordata</i> ) R. Section 2: cherry sp. ( <i>Prunus</i> sp.) A, wild cherry ( <i>Prunus avium</i> ) R, elm sp. ( <i>Ulmus</i> sp.) R, field maple ( <i>Acer campestre</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) A, ash ( <i>Fraxinus excelsior</i> ) O, elder ( <i>Sambucus nigra</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) O, cherry laurel ( <i>Prunus laurocerasus</i> ) O.

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
F	H-F-26	h2a - hedgerow (priority habitat)	Hedgerow with trees and a ditch present on a hedge bank. Predominately hawthorn hedgerow with 6 oaks, some mature. Lollipop shape as hawthorn is very old and not being laid in recent history.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) F, dog-rose ( <i>Rosa canina</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) R, holly ( <i>Ilex aquifolium</i> ) O. Section 2: hawthorn ( <i>Crataegus monogyna</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, hazel ( <i>Corylus avellana</i> ) R.
F	H-F-27	h2a - hedgerow (priority habitat)	Hedgerow dividing two arable fields. Flailed every year and now developing lollipop form as the stems are very wide and not being laid.	Section 1: elder ( <i>Sambucus nigra</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, hazel ( <i>Corylus avellana</i> ) F, dog-rose ( <i>Rosa canina</i> ) O, holly ( <i>Ilex aquifolium</i> ) O. Section 2: elder ( <i>Sambucus nigra</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) D, hazel ( <i>Corylus avellana</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, field maple ( <i>Acer campestre</i> ) R. Section 3: elder ( <i>Sambucus nigra</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) D, blackthorn ( <i>Prunus spinosa</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, holly ( <i>Ilex aquifolium</i> ) O
F	H-F-28	h2a - hedgerow (priority habitat)	Hedgerow with a ditch.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) R, dog-rose ( <i>Rosa canina</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, holly ( <i>Ilex aquifolium</i> ) D. Section 2: dog-rose ( <i>Rosa canina</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, holly ( <i>Ilex aquifolium</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R. Section 3: blackthorn ( <i>Prunus spinosa</i> ) R, dog-rose ( <i>Rosa canina</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, holly ( <i>Ilex aquifolium</i> ) A, ash ( <i>Fraxinus excelsior</i> ) R. Ground flora: wood false-brome ( <i>Brachypodium sylvaticum</i> ) R
F	H-F-29	h2a - hedgerow (priority habitat)	Hedgerow with trees, including two veteran trees. Borders a dry ditch.	Pedunculate oak ( <i>Quercus robur</i> ) O, bramble ( <i>Rubus sp.</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) O, nettle ( <i>Urtica dioica</i> ) LD, bracken ( <i>Pteridium aquilinum</i> ) O, ivy ( <i>Hedera helix</i> ) F, lime ( <i>Tilia x europaea</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) A, rose sp. ( <i>Rosa sp.</i> ) R, hazel ( <i>Corylus avellana</i> ) F, holly ( <i>Ilex aquifolium</i> ) F, field maple ( <i>Acer campestre</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R, cleavers ( <i>Galium aparine</i> ) F, cherry sp. ( <i>Prunus sp.</i> ) R, elder ( <i>Sambucus nigra</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, cypress sp. ( <i>Cupressus sp.</i> ) R, horse chestnut ( <i>Aesculus hippocastanum</i> ) R, elm sp. ( <i>Ulmus sp.</i> ) O, gorse ( <i>Ulex europaeus</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ), bramble ( <i>Rubus sp.</i> ).
F	H-F-30	h2a - hedgerow (priority habitat)	Hedgerow with trees. Separated from a property by chicken wire.	Elder ( <i>Sambucus nigra</i> ) O, nettle ( <i>Urtica dioica</i> ) A, elm sp. ( <i>Ulmus sp.</i> ) F, willow sp. ( <i>Salix sp.</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) O, ivy ( <i>Hedera helix</i> ) A, bramble ( <i>Rubus sp.</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) O, holly ( <i>Ilex aquifolium</i> ) O, hornbeam ( <i>Carpinus betulus</i> ) R, field maple ( <i>Acer campestre</i> ) R, poplar sp. ( <i>Populus sp.</i> ) R.
F	H-F-32	h2a - hedgerow (priority habitat)	Planted hedgerow with a line of trees within. Running parallel to a line of trees.	Hawthorn ( <i>Crataegus monogyna</i> ) D, bramble ( <i>Rubus sp.</i> ) O.
F	H-F-33	h2a - hedgerow (priority habitat)	Defunct hedgerow with trees along a dry ditch.	Bramble ( <i>Rubus sp.</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) A, ivy ( <i>Hedera helix</i> ) F, cow parsley ( <i>Anthriscus sylvestris</i> ) F, nettle ( <i>Urtica dioica</i> ) O, cleavers ( <i>Galium aparine</i> ) O, hemlock ( <i>Conium maculatum</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, rose sp. ( <i>Rosa sp.</i> ) R, ground-ivy ( <i>Hedera helix</i> ) O.
F	H-F-34	h2a - hedgerow (priority habitat)	Hedgerow with trees.	Cleavers ( <i>Galium aparine</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) A, nettle ( <i>Urtica dioica</i> ) A, cow parsley ( <i>Anthriscus sylvestris</i> ) F, sorrel ( <i>Rumex acetosa</i> ), perennial ryegrass ( <i>Lolium perenne</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) O, rose sp. ( <i>Rosa sp.</i> ) R, bramble ( <i>Rubus sp.</i> ) D
F	H-F-35	h2a - hedgerow (priority habitat)	Shrubby hedgerow with trees on a hedge bank. Cut to 5m below existing power lines. Dividing a commercial orchard and a lane.	Section 1: elm sp. ( <i>Ulmus sp.</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) F, field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, spindle ( <i>Euonymus europaeus</i> ) R. Section 2: elm sp. ( <i>Ulmus sp.</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, ash ( <i>Fraxinus excelsior</i> ) R, dog-rose ( <i>Rosa canina</i> ) R.

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
F	H-F-36	h2a - hedgerow (priority habitat)	Defunct hedgerow with trees.	Rose sp. ( <i>Rosa</i> sp.) R, bracken ( <i>Pteridium aquilinum</i> ) F, bramble ( <i>Rubus</i> sp.) O, field maple ( <i>Acer campestre</i> ) O, hazel ( <i>Corylus avellana</i> ) F, ground-ivy ( <i>Hedera helix</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, dogwood ( <i>Cornus sanguinea</i> ) R, ivy ( <i>Hedera helix</i> ) A, cow parsley ( <i>Anthriscus sylvestris</i> ) O, oak sp. ( <i>Quercus</i> sp.) O, mayweed sp. ( <i>Tripleurospermum</i> sp.) R, speedwell sp. ( <i>Veronica</i> sp.) R, elder ( <i>Sambucus nigra</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, elm sp. ( <i>Ulmus</i> sp.) F, perennial ryegrass ( <i>Lolium perenne</i> ) F.
F	H-F-39	h2a - hedgerow (priority habitat)	Hedgerow with trees and a dry ditch. Mature pedunculate oak present.	Hazel ( <i>Corylus avellana</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) A, ash ( <i>Fraxinus excelsior</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) F, willow sp. ( <i>Salix</i> sp.) R, nettle ( <i>Urtica dioica</i> ) O, bracken ( <i>Pteridium aquilinum</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) F, rose sp. ( <i>Rosa</i> sp.) R, cock's-foot ( <i>Dactylis glomerata</i> ) A, ivy ( <i>Hedera helix</i> ) F, dock ( <i>Rumex</i> sp.) F, ground-ivy ( <i>Hedera helix</i> ) F, garlic mustard ( <i>Alliaria petiolata</i> ) A, field maple ( <i>Acer campestre</i> ) O.
F	H-F-40	h2a - hedgerow (priority habitat)	Shrubby hedgerow with trees.	Bracken ( <i>Pteridium aquilinum</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, ivy ( <i>Hedera helix</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, field maple ( <i>Acer campestre</i> ) R, plum ( <i>Prunus domestica</i> ) R, ash ( <i>Fraxinus excelsior</i> ) R, elm sp. ( <i>Ulmus</i> sp.) O, hazel ( <i>Corylus avellana</i> ) LA, red dead-nettle ( <i>Lamium purpureum</i> ) F.
F	H-F-43	h2a - hedgerow (priority habitat)	Hedgerow not fully surveyed, assumed to be species rich. Veteran oak in the centre.	Veteran pedunculate oak ( <i>Quercus robur</i> ).
F	H-F-44	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow along a dry ditch.	N/A
F	H-F-46	h2a - hedgerow (priority habitat)	Native hedgerow along a dry ditch.	Hawthorn ( <i>Crataegus monogyna</i> ) A, holly ( <i>Ilex aquifolium</i> ) O and cherry sp. ( <i>Prunus</i> sp.) O.
F	H-F-47	h2a - hedgerow (priority habitat)	Shrubby, native hedgerow which has been flailed.	Hawthorn ( <i>Crataegus monogyna</i> ) D.
F	H-F-48	h2a - hedgerow (priority habitat)	Native species rich hedgerow which has been planted and flailed. Along a bank with trees present.	N/A
F	H-F-49	h2a - hedgerow (priority habitat)	Hedgerow surveyed from adjacent land parcel due to lack access constraints. Hedgerow along wooden fence, all evenly aged with a small area of replanting.	Hawthorn ( <i>Crataegus monogyna</i> ) D.
F	H-F-50	h2a - hedgerow (priority habitat)	Hedgerow with trees. Partially runs along a road with a dry ditch.	Plum ( <i>Prunus domestica</i> ), silver birch ( <i>Betula pendula</i> ), blackthorn ( <i>Prunus spinosa</i> ) A, ash ( <i>Fraxinus excelsior</i> ) O, field maple ( <i>Acer campestre</i> ) O, cock's-foot ( <i>Dactylis glomerata</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) O, nettle ( <i>Urtica dioica</i> ) O, bramble ( <i>Rubus</i> sp.) A, holly ( <i>Ilex aquifolium</i> ), ivy ( <i>Hedera helix</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R, elm sp. ( <i>Ulmus</i> sp.) LA, rose sp. ( <i>Rosa</i> sp.) O, pedunculate oak ( <i>Quercus robur</i> ) F, plum ( <i>Prunus domestica</i> ) R, turkey oak ( <i>Quercus cerris</i> ) R, willow sp. ( <i>Salix</i> sp.) O, cow parsley ( <i>Anthriscus sylvestris</i> ) F.
F	H-F-51	h2a - hedgerow (priority habitat)	Hedge with trees transitioning to bramble dominated hedgerow. Mature pedunculate oaks and line of trees present. Along a dry ditch.	Section 1: Non-native Canadian poplar ( <i>Populus x canadensis</i> ), mature pedunculate oak ( <i>Quercus robur</i> ), bramble ( <i>Rubus</i> sp.) D, eyelet, blackthorn ( <i>Prunus spinosa</i> ) F, horse chestnut ( <i>Aesculus hippocastanum</i> ) R, elm sp. ( <i>Ulmus</i> sp.) O, silver birch ( <i>Betula pendula</i> ), 1x black walnut ( <i>Juglans nigra</i> ). Section 2: Northern hedge - 1x sumac species ( <i>Rhus</i> sp.), Canadian poplar ( <i>Populus x canadensis</i> ) O, nettle ( <i>Urtica dioica</i> ) F, pedunculate oak ( <i>Quercus robur</i> ), perennial ryegrass ( <i>Lolium perenne</i> ), cow parsley ( <i>Anthriscus sylvestris</i> ) O, blackthorn ( <i>Prunus spinosa</i> ), rose sp. ( <i>Rosa</i> sp.), sycamore ( <i>Acer pseudoplatanus</i> ), wild cherry ( <i>Prunus avium</i> ), perennial ryegrass ( <i>Lolium perenne</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) R, holly ( <i>Ilex aquifolium</i> ) R, elder ( <i>Sambucus nigra</i> ) O

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
F	H-F-52	h2a - hedgerow (priority habitat)	Hedgerow with trees, along a ditch holding water.	Blackthorn ( <i>Prunus spinosa</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) F, bramble ( <i>Rubus</i> sp.) A, pedunculate oak ( <i>Quercus robur</i> ) R, cow parsley ( <i>Anthriscus sylvestris</i> ) O, rush species ( <i>Juncus</i> sp.) O, hogweed ( <i>Heracleum sphondylium</i> ) F, rose sp. ( <i>Rosa</i> sp.) O, common sorrel ( <i>Rumex acetosa</i> ), willowherb species ( <i>Epilobium</i> sp.) O, ash ( <i>Fraxinus excelsior</i> ) R, nettle ( <i>Urtica dioica</i> )
F	H-F-53	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
F	H-F-54	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
F	H-F-55	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
F	H-F-56	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
F	H-F-57	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
F	H-F-58	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
F	H-F-59	h2 - hedgerow	UKHab category derived from aerial imagery	N/A. Not surveyed
G	H-G-02	h2a - hedgerow (priority habitat)	Hedgerow with a wet ditch and planted willows present on northern side. Trees present within hedgerow.	Section 1: hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, wild cherry ( <i>Prunus avium</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) R, wood false-brome ( <i>Brachypodium sylvaticum</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) R. Section 2: hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) R, dog-rose ( <i>Rosa canina</i> ) O, willow sp. ( <i>Salix</i> sp.) R. Section 3: hazel ( <i>Corylus avellana</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, willow ( <i>Salix</i> sp.) A, dog-rose ( <i>Rosa canina</i> ) A, alder ( <i>Alnus glutinosa</i> ) R.
G	H-G-03	h2a - hedgerow (priority habitat)	Hedgerow on bank with lack of management resulting in the growth of trees. Small dry ditch on southern side which has silted up.	Hazel ( <i>Corylus avellana</i> ) A, field maple ( <i>Acer campestre</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) R, dog's mercury ( <i>Mercurialis perennis</i> ) F, cowslip ( <i>Primula veris</i> ) O, wood false-brome ( <i>Brachypodium sylvaticum</i> ) O, wood avens ( <i>Geum urbanum</i> ) O.
G	H-G-04	h2a - hedgerow (priority habitat)	Hedgerow on a bank with coppiced trees present. Borders possible old orchard.	Hazel ( <i>Corylus avellana</i> ) D, field maple ( <i>Acer campestre</i> ) R, elder ( <i>Sambucus nigra</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) R. Ground flora: nettle ( <i>Urtica dioica</i> ) A, wood false-brome ( <i>Brachypodium sylvaticum</i> ) O, wood avens ( <i>Geum urbanum</i> ) O, lords-and-ladies ( <i>Arum maculatum</i> ) O, garlic mustard ( <i>Alliaria petiolata</i> ) O.
G	H-G-05	h2a - hedgerow (priority habitat)	Hedgerow on a bank between road and grazing pasture. Dense on pasture side. Lack of management resulting in the growth of trees.	Section 1: hazel ( <i>Corylus avellana</i> ) O, mature pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, alder ( <i>Alnus glutinosa</i> ) O, elder ( <i>Sambucus nigra</i> ) R.
G	H-G-06	h2a - hedgerow (priority habitat)	Hedgerow on the southern bank of a stream, partially replanted to the east and non-native species present. Lack of management resulting in the growth of an understorey hedgerow.	Section 1: mature ash ( <i>Fraxinus excelsior</i> ) O, elder ( <i>Sambucus nigra</i> ) O, field maple ( <i>Acer campestre</i> ) O, alder ( <i>Alnus glutinosa</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R. Ground flora: lords-and-ladies ( <i>Arum maculatum</i> ) O, wood false-brome ( <i>Brachypodium sylvaticum</i> ) F, herb-Robert ( <i>Geranium robertianum</i> ) R.
G	H-G-08	h2a - hedgerow (priority habitat)	Hedgerow on a bank with coppiced trees present.	Section 1: Field maple ( <i>Acer campestre</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, hazel ( <i>Corylus avellana</i> ) F C, rose sp. ( <i>Rosa</i> sp.) R, dogwood ( <i>Cornus sanguinea</i> ) R, lords-and-ladies ( <i>Arum maculatum</i> ) R, herb-Robert ( <i>Geranium robertianum</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) O.
G	H-G-09	h2a - hedgerow (priority habitat)	Hedgerow with trees. Bramble smothering ground layer.	Section 1: hazel ( <i>Corylus avellana</i> ) O, goat willow ( <i>Salix caprea</i> ) R, elder ( <i>Sambucus nigra</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) M. Bramble ( <i>Rubus</i> sp.), willowherb ( <i>Epilobium</i> sp.) O, nettle ( <i>Urtica dioica</i> ) O.
G	H-G-11	h2a - hedgerow (priority habitat)	Hedgerow with trees and a ditch. Some maturing horse chestnuts and ancient woodland indicators at the northern end.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) A, hazel ( <i>Corylus avellana</i> ) O, field maple ( <i>Acer campestre</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) R, elm sp. ( <i>Ulmus</i> sp.) R, willow sp. ( <i>Salix</i> sp.) R, horse chestnut ( <i>Aesculus hippocastanum</i> ). Ground flora: wood false-brome ( <i>Brachypodium sylvaticum</i> ) O, wood avens ( <i>Geum urbanum</i> ) R, lords-and-ladies R

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
G	H-G-12	h2a - hedgerow (priority habitat)	Native species rich hedgerow. Not surveyed.	N/A. Not surveyed
G	H-G-13	h2a - hedgerow (priority habitat)	Hedgerow with trees on a bank going up approximately 5m from east to west. Mature trees present on northern side. Abundance reducing towards the southern side.	Section 1: elder ( <i>Sambucus nigra</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, hazel ( <i>Corylus avellana</i> ) O, holly ( <i>Ilex aquifolium</i> ) O, field maple ( <i>Acer campestre</i> ) R, pedunculate oak ( <i>Quercus robur</i> ). Section 2: hazel ( <i>Corylus avellana</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, ash ( <i>Fraxinus excelsior</i> ) R, field maple ( <i>Acer campestre</i> ) O, elm sp. ( <i>Ulmus</i> sp.) R. Ground flora: red campion ( <i>Silene dioica</i> ) R.
G	H-G-14	h2a - hedgerow (priority habitat)	Hedgerow with trees present on a bank from west going up approximately 5m to east.	Section 1: elder ( <i>Sambucus nigra</i> ) O, field maple ( <i>Acer campestre</i> ) D. Section 2: elder ( <i>Sambucus nigra</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) O, hazel ( <i>Corylus avellana</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) R.
G	H-G-16	h2a - hedgerow (priority habitat)	Overgrown hedgerow with trees and a ditch present on a hedge bank. Partially managed hedgerow probably replanted to fill gaps. Some ancient woodland indicators present. Borders an arable margin with well-developed perennial vegetation, topped annually.	Section 1: elm sp. ( <i>Ulmus</i> sp.) O, hawthorn ( <i>Crataegus monogyna</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) O, wild cherry ( <i>Prunus avium</i> ) R, common lime ( <i>Tilia x europaea</i> ) R, rhododendron ( <i>Rhododendron ponticum</i> ) O, small-leaved lime ( <i>Tilia cordata</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, hazel ( <i>Corylus avellana</i> ) R, field maple ( <i>Acer campestre</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R.
G	H-G-17	h2a - hedgerow (priority habitat)	Partially planted hedgerow containing two medium sized oaks (approximately 15m high). Bordering an arable field which is topped annually with blackthorn encroaching from the hedgerow.	Section 1: field maple ( <i>Acer campestre</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) R, hazel ( <i>Corylus avellana</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) F, dog-rose ( <i>Rosa canina</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, spindle ( <i>Euonymus europaeus</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) O. Section 2: field maple ( <i>Acer campestre</i> ) F, elm sp. ( <i>Ulmus</i> sp.) R, blackthorn ( <i>Prunus spinosa</i> ) O, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) F, dogwood ( <i>Cornus sanguinea</i> ) O.
G	H-G-18	h2a - hedgerow (priority habitat)	Native hedgerow - associated with bank or ditch.	N/A. Not surveyed
G	H-G-19	h2a - hedgerow (priority habitat)	Hedgerow with trees and a ditch. Three maturing oaks at the northern end.	Section 1: field maple ( <i>Acer campestre</i> ) D, dog-rose ( <i>Rosa canina</i> ) O, elder ( <i>Sambucus nigra</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) R, elm sp. ( <i>Ulmus</i> sp.) R. Section 2: field maple ( <i>Acer campestre</i> ) F, dogwood ( <i>Cornus sanguinea</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) O, elm sp. ( <i>Ulmus</i> sp.) O, hazel ( <i>Corylus avellana</i> ) O. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) R, wood false-brome ( <i>Brachypodium sylvaticum</i> ) O
G	H-G-20	h2a - hedgerow (priority habitat)	Overgrown hedgerow with a deep ditch and trees present. Bordering a property with a fence.	Section 1: elder ( <i>Sambucus nigra</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) O, hazel ( <i>Corylus avellana</i> ) O, elm sp. ( <i>Ulmus</i> sp.) R, field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) O, dog-rose ( <i>Rosa canina</i> ) O. Section 2: elder ( <i>Sambucus nigra</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) R, elm sp. ( <i>Ulmus</i> sp.) R, field maple ( <i>Acer campestre</i> ) F, ash ( <i>Fraxinus excelsior</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) R
G	H-G-21	h2a - hedgerow (priority habitat)	Shrubby hedgerow with trees. Some maturing trees and small-leaved lime present.	Section 1: beech ( <i>Fagus sylvatica</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) A, ash ( <i>Fraxinus excelsior</i> ) F, field maple ( <i>Acer campestre</i> ) F, dog-rose ( <i>Rosa canina</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, dogwood ( <i>Cornus sanguinea</i> ) R. Section 2: elm sp. ( <i>Ulmus</i> sp.) R, hawthorn ( <i>Crataegus monogyna</i> ) A, ash ( <i>Fraxinus excelsior</i> ) O, field maple ( <i>Acer campestre</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) O, English yew ( <i>Taxus baccata</i> ) R. Section 3: hawthorn ( <i>Crataegus monogyna</i> ) R, field maple ( <i>Acer campestre</i> ) O, dog-rose ( <i>Rosa canina</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O, small-leaved lime ( <i>Tilia cordata</i> ) O, spindle ( <i>Euonymus europaeus</i> ) O, hazel ( <i>Corylus avellana</i> ) O

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
G	H-G-25	h2a - hedgerow (priority habitat)	Native hedgerow with trees.	Blackthorn ( <i>Prunus spinosa</i> ) O, bramble ( <i>Rubus sp.</i> ) O, spindle ( <i>Euonymus europaeus</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, field maple ( <i>Acer campestre</i> ) R, elm sp. ( <i>Ulmus sp.</i> ) F. Ground flora: garlic mustard ( <i>Alliaria petiolata</i> ) F, common nettle ( <i>Urtica dioica</i> ) O, ivy ( <i>Hedera helix</i> ) O, red dead-nettle ( <i>Lamium purpureum</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) O, bluebell sp. ( <i>Hyacinthoides sp.</i> ) O, lords-and-ladies ( <i>Arum maculatum</i> ) O, cow parsley ( <i>Anthriscus sylvestris</i> ) R.
G	H-G-26	h2a - hedgerow (priority habitat)	Hedgerow along old, barbed wire fence, between earth track and narrow wild patch.	Section 1: English elm sp. ( <i>Ulmus sp.</i> ) D, elder ( <i>Sambucus nigra</i> ), holly ( <i>Ilex aquifolium</i> ). Section 2: young pedunculate oak ( <i>Quercus robur</i> ), spindle ( <i>Euonymus europaeus</i> ), blackthorn ( <i>Prunus spinosa</i> ), hazel ( <i>Corylus avellana</i> ), hawthorn ( <i>Crataegus monogyna</i> )
G	H-G-28	h2a - hedgerow (priority habitat)	Native species rich hedgerow which has been flailed.	Section 1: dogwood ( <i>Cornus sanguinea</i> ) R, spindle ( <i>Euonymus europaeus</i> ) R, field maple ( <i>Acer campestre</i> ) R, oak sp. ( <i>Quercus sp.</i> ) R, elm sp. ( <i>Ulmus sp.</i> ) O. Section 2: elm sp. ( <i>Ulmus sp.</i> ), pedunculate oak ( <i>Quercus robur</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, ash ( <i>Fraxinus excelsior</i> ) R. Section 3: Coppiced hazel ( <i>Corylus avellana</i> ) O, elm sp. ( <i>Ulmus sp.</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) R, field maple ( <i>Acer campestre</i> ) R
G	H-G-29	h2a - hedgerow (priority habitat)	Wide, overgrown hedgerow with hedge bank. Remnants of an ancient hedgerow dominated by elm, occasionally 5m high.	Section 1: holly ( <i>Ilex aquifolium</i> ) O, elder ( <i>Sambucus nigra</i> ) R, elm sp. ( <i>Ulmus sp.</i> ) D, blackthorn ( <i>Prunus spinosa</i> ) A, ash ( <i>Fraxinus excelsior</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R. Ground flora: herb-Robert ( <i>Geranium robertianum</i> ) R
G	H-G-30	h2a - hedgerow (priority habitat)	Hedgerow with small section left unmanaged.	Blackthorn ( <i>Prunus spinosa</i> ), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> )
G	H-G-31	h2a - hedgerow (priority habitat)	Hedgerow with outgrown trees 10-15m tall.	Spindle ( <i>Euonymus europaeus</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) D, wych elm ( <i>Ulmus glabra</i> ) O, red campion ( <i>Silene dioica</i> ), pedunculate oak ( <i>Quercus robur</i> ), young wych elm ( <i>Ulmus glabra</i> ), field maple ( <i>Acer campestre</i> ), holly ( <i>Ilex aquifolium</i> ), ash ( <i>Fraxinus excelsior</i> ), dog-rose ( <i>Rosa canina</i> ) R, mallow sp. ( <i>Malva sp.</i> ) R. Intermittent ivy ( <i>Hedera helix</i> ), bramble ( <i>Rubus sp.</i> ).
G	H-G-32	h2a - hedgerow (priority habitat)	Planted hedgerow bordering a fence. Bare ground beneath. Dense vegetation under the pylon.	Hawthorn D, ash ( <i>Fraxinus excelsior</i> ) sapling R, hazel ( <i>Corylus avellana</i> ) sapling R, elder ( <i>Sambucus nigra</i> ) R, mature ash ( <i>Fraxinus excelsior</i> ), field maple ( <i>Acer campestre</i> ), elder ( <i>Sambucus nigra</i> ). Ground flora: crane's-bill sp. ( <i>Geranium sp.</i> ), Yorkshire fog ( <i>Holcus lanatus</i> ), nettle ( <i>Urtica dioica</i> ), field pansy ( <i>Viola arvensis</i> ), scarlet pimpernel ( <i>Anagallis arvensis</i> ), gorse ( <i>Ulex europaeus</i> ), thistle sp. ( <i>Cirsium sp.</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), bugloss ( <i>Anchusa arvensis</i> ), Timothy ( <i>Phleum pratense</i> )
G	H-G-33	h2a - hedgerow (priority habitat)	Native Species Rich Hedgerow with trees. Notable species record within close proximity.	N/A. Not surveyed
G	H-G-34	h2a - hedgerow (priority habitat)	Native Hedgerow - Associated with bank or ditch. Species poor, defunct.	N/A. Not surveyed
G	H-G-35	h2a - hedgerow (priority habitat)	Shrubby hedgerow borders road with gap of 5m. Hedge bank 2m tall with active rabbit warrens present. Non-native 'pink cascade' most likely a garden escapee. Dead elm and significant elm regrowth present. Patches in hedgerow filled in with hawthorn planting.	Section 1: elm sp. ( <i>Ulmus sp.</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) A, dog-rose ( <i>Rosa canina</i> ) R, field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, elder ( <i>Sambucus nigra</i> ) R, wild cherry ( <i>Prunus avium</i> ) R. Section 2: elm sp. ( <i>Ulmus sp.</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) A, field maple ( <i>Acer campestre</i> ) O, ash ( <i>Fraxinus excelsior</i> ) R, elder ( <i>Sambucus nigra</i> ) R, non-native 'pink cascade' ( <i>Tamarix ramosissima</i> ) R.
G	H-G-36	h2a - hedgerow (priority habitat)	Hedgerow with trees.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ) A, elm sp. ( <i>Ulmus sp.</i> ) R, elder ( <i>Sambucus nigra</i> ) R, pedunculate oak ( <i>Quercus robur</i> ) A, blackthorn ( <i>Prunus spinosa</i> ), ivy ( <i>Hedera helix</i> )

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
G	H-G-37	h2a - hedgerow (priority habitat)	Hedgerow with small hedge bank. Dominated by introduced trees, including conifer. Unable to survey part of the hedge due to access constraints.	Walnut sp. ( <i>Juglans</i> sp.), cherry laurel ( <i>Prunus laurocerasus</i> ), lime ( <i>Tilia x europaea</i> ). Section 1: elder ( <i>Sambucus nigra</i> ) O, wild cherry ( <i>Prunus avium</i> ) R, silver birch ( <i>Betula pendula</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R. Section 2: cherry sp. ( <i>Prunus</i> sp.) F, elder ( <i>Sambucus nigra</i> ) O, wild cherry ( <i>Prunus avium</i> ) F.
G	H-G-38	h2a - hedgerow (priority habitat)	Defunct hedgerow with a large dry ditch due to dense canopy cover. Gaps in hedgerow larger than 5m x 3. Flailed on gaps to have moderate condition. Small amounts of dog's mercury present within the ditch.	Section 1: pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) F, field maple ( <i>Acer campestre</i> ) F, dog-rose ( <i>Rosa canina</i> ) R, elm sp. ( <i>Ulmus</i> sp.) O, hazel ( <i>Corylus avellana</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O. Section 2: pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) O, field maple ( <i>Acer campestre</i> ) O, dog-rose ( <i>Rosa canina</i> ) F, elm sp. ( <i>Ulmus</i> sp.) O, blackthorn ( <i>Prunus spinosa</i> ) F. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) R.
G	H-G-39	h2a - hedgerow (priority habitat)	Defunct hedgerow with hedge bank containing rabbit holes present. Trees present within the hedgerow and are clustered at the end.	Section 1: elder ( <i>Sambucus nigra</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) F, dog-rose ( <i>Rosa canina</i> ) O, elm sp. ( <i>Ulmus</i> sp.) O, holly ( <i>Ilex aquifolium</i> ) O, hazel ( <i>Corylus avellana</i> ) R. Section 2: elder ( <i>Sambucus nigra</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) R, hazel ( <i>Corylus avellana</i> ) O, field maple ( <i>Acer campestre</i> ) O. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ).
G	H-G-40	h2a - hedgerow (priority habitat)	Unmanaged hedgerow on a hedge bank with rabbit disturbance and high partridge density. One mature oak, two moderate oaks and some bracken encroachment present. Borders a dirt track.	Wild cherry ( <i>Prunus avium</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) R, elm sp. ( <i>Ulmus</i> sp.) O, dog-rose ( <i>Rosa canina</i> ) F, field maple ( <i>Acer campestre</i> ) A, ash ( <i>Fraxinus excelsior</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R. Section 2: hawthorn ( <i>Crataegus monogyna</i> ) O, elm sp. ( <i>Ulmus</i> sp.) O, dog-rose ( <i>Rosa canina</i> ) F, field maple ( <i>Acer campestre</i> ) A, pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, hazel ( <i>Corylus avellana</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) R.
G	H-G-41	h2a - hedgerow (priority habitat)	Unmanaged hedgerow comprising of scattered trees and shrubs, on steep, dry bank with significant bare ground from partridge erosion. One medium oak and some medium field maples present. Borders a dirt track with almost no herb layer to the side of dirt track.	Section 1: ash ( <i>Fraxinus excelsior</i> ) F, elm sp. ( <i>Ulmus</i> sp.) O, hawthorn ( <i>Crataegus monogyna</i> ) O, field maple ( <i>Acer campestre</i> ) F, pedunculate oak ( <i>Quercus robur</i> ) O, hazel ( <i>Corylus avellana</i> ) R. Ground flora: wood false-brome ( <i>Brachypodium sylvaticum</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) R.
G	H-G-42	h2a - hedgerow (priority habitat)	Overgrown hedgerow with a hedge bank, a very dry ditch. Some small trees and tall marginal vegetation present.	Section 1: field maple ( <i>Acer campestre</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) O, dog-rose ( <i>Rosa canina</i> ) F, willow sp. ( <i>Salix</i> sp.) O, dogwood ( <i>Cornus sanguinea</i> ) F, hazel ( <i>Corylus avellana</i> ) R, elder ( <i>Sambucus nigra</i> ) R. Section 2: field maple ( <i>Acer campestre</i> ) D, hawthorn ( <i>Crataegus monogyna</i> ) R, dog-rose ( <i>Rosa canina</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, ash ( <i>Fraxinus excelsior</i> ) O. Ground flora: wood false-brome ( <i>Brachypodium sylvaticum</i> ) O
G	H-G-43	h2a - hedgerow (priority habitat)	Hedgerow with a ditch. Replanted in sections approximately 15 years old.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) D, dog-rose ( <i>Rosa canina</i> ) F, field maple ( <i>Acer campestre</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) O. Section 2: blackthorn ( <i>Prunus spinosa</i> ) A, dog-rose ( <i>Rosa canina</i> ) R, ash ( <i>Fraxinus excelsior</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O. Section 3: blackthorn ( <i>Prunus spinosa</i> ) A, dog-rose ( <i>Rosa canina</i> ) F, hawthorn ( <i>Crataegus monogyna</i> ) A, dogwood ( <i>Cornus sanguinea</i> ) O, spindle ( <i>Euonymus europaeus</i> ) R
G	H-G-44	h2a - hedgerow (priority habitat)	Long, species rich hedgerow, 2-5 wide with trees and a ditch. Mature oaks present. Bordering an unsettled track and a small aspen plantation at the eastern end of the hedge.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) F, ash ( <i>Fraxinus excelsior</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) O, dog-rose ( <i>Rosa canina</i> ) R, field maple ( <i>Acer campestre</i> ) A, hazel ( <i>Corylus avellana</i> ) O, small-leaved lime O, pedunculate oak ( <i>Quercus robur</i> ) O.
G	H-G-45	h2a - hedgerow (priority habitat)	Coppiced hedgerow with trees and a ditch on a hedge bank. Some coppiced hazel and managed blackthorn and field maple. Mature oak and ash standards. Some ancient woodland indicators present.	Section 1: blackthorn ( <i>Prunus spinosa</i> ) O, ash ( <i>Fraxinus excelsior</i> ) F, hazel ( <i>Corylus avellana</i> ) A, cherry sp. ( <i>Prunus</i> sp.) O, aspen ( <i>Populus tremula</i> ) R, hawthorn ( <i>Crataegus monogyna</i> ) O. Section 2: pedunculate oak ( <i>Quercus robur</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) A, ash ( <i>Fraxinus excelsior</i> ) F, elder ( <i>Sambucus nigra</i> ) R, field maple ( <i>Acer campestre</i> ) O, hazel ( <i>Corylus avellana</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) O.
G	H-G-46	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed

Section	Habitat ID	Habitat Type	Description	Species List with DAFOR
G	H-G-49	h2a - hedgerow (priority habitat)	Overgrown hedgerow and a ditch on a dry hedge bank. Bordering a road. Hedge bank lacking vegetation. Significant elms present, approximately 50 years old. Abundance of holly and some maturing oaks present. Flailed to 4m forming an arch with the hedge opposite	Section 1: spindle ( <i>Euonymus europaeus</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) O, holly ( <i>Ilex aquifolium</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) O, dog-rose ( <i>Rosa canina</i> ) R. Section 2: spindle ( <i>Euonymus europaeus</i> ) R, elm sp. ( <i>Ulmus</i> sp.) A, blackthorn ( <i>Prunus spinosa</i> ) F, holly ( <i>Ilex aquifolium</i> ) A, field maple ( <i>Acer campestre</i> ) O, hawthorn ( <i>Crataegus monogyna</i> ) O, pedunculate oak ( <i>Quercus robur</i> ) O
G	H-G-50	h2a - hedgerow (priority habitat)	Hedgerow with a lot of vertical gaps, young and dead elm trees present.	Section 1: elm sp. ( <i>Ulmus</i> sp.) D, hazel ( <i>Corylus avellana</i> ) R.
G	H-G-052	h2a - hedgerow (priority habitat)	Hedgerow on a hedge bank. Veteran and coppiced trees present.	Section 1: field maple ( <i>Acer campestre</i> ) O, mature oak sp. ( <i>Quercus</i> sp.) F, coppiced hazel ( <i>Corylus avellana</i> ) O, holly ( <i>Ilex aquifolium</i> ) R, dog-rose ( <i>Rosa canina</i> ), blackthorn ( <i>Prunus spinosa</i> ) O. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) O, primrose ( <i>Primula vulgaris</i> ) O, bluebell sp. ( <i>Hyacinthoides</i> sp.) O, hairy brome ( <i>Bromus ramosus</i> ) O
G	H-G-053	h2a - hedgerow (priority habitat)	Hedgerow on a hedge bank with a ditch. Veteran and coppiced trees present.	Section 1: coppiced hazel ( <i>Corylus avellana</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) F, mature sweet chestnut ( <i>Castanea sativa</i> ) R, mature pedunculate oak ( <i>Quercus robur</i> ) F, field maple ( <i>Acer campestre</i> ) F. Ground flora: hairy brome ( <i>Bromus ramosus</i> ) O, primrose ( <i>Primula vulgaris</i> ) O, dog's mercury ( <i>Mercurialis perennis</i> ) O, bluebell sp. ( <i>Hyacinthoides</i> sp.) O.
G	H-G-054	h2a - hedgerow (priority habitat)	Hedgerow on a hedge bank. Borders a dirt track.	Section 1: field maple ( <i>Acer campestre</i> ) O, dogwood ( <i>Cornus sanguinea</i> ) R, blackthorn ( <i>Prunus spinosa</i> ) F, dog-rose ( <i>Rosa canina</i> ) F, willow sp. ( <i>Salix</i> sp.) O. Ground flora: dog's mercury ( <i>Mercurialis perennis</i> ) O
G	H-G-055	h2a - hedgerow (priority habitat)	Shrubby hedgerow which is mostly unmanaged but shows some evidence of flailing. Trees present with an average height of 4m. One mature oak and some semi-mature field maples present. Turning into a line of trees. Bordering a garden. Mostly unmanaged but some evidence of flailing.	Mature oak sp. ( <i>Quercus</i> sp.), some semi-mature field maple ( <i>Acer campestre</i> ).
G	H-G-056	h2a - hedgerow (priority habitat)	Overgrown hedgerow on a steep bank next to a dirt track and ditch. Hedge opposite was counted as part of connection. Greater than 5m wide in sections and dominated by nettle and hogweed. Non-native Canadian poplar present.	Section 1: Canadian poplar ( <i>Populus x canadensis</i> ) R, dog-rose ( <i>Rosa canina</i> ) O, field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) A, elm sp. ( <i>Ulmus</i> sp.) R, pedunculate oak ( <i>Quercus robur</i> ) O, ash ( <i>Fraxinus excelsior</i> ) O, hazel ( <i>Corylus avellana</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R. Section 2: elder ( <i>Sambucus nigra</i> ) R, dog-rose ( <i>Rosa canina</i> ) F, field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) A, hazel ( <i>Corylus avellana</i> ) O. Section 3: dog-rose ( <i>Rosa canina</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) A, hazel ( <i>Corylus avellana</i> ) O
G	H-G-057	h2a - hedgerow (priority habitat)	Hedgerow on a bank. Evidence of historic coppicing but now lacks management resulting in the growth of trees. Flailed on eastern aspect and left to east.	Section 1: ash sapling ( <i>Fraxinus excelsior</i> ), rose sp. ( <i>Rosa</i> sp.) O, wych elm ( <i>Ulmus glabra</i> ) F with dead elm ( <i>Ulmus</i> sp.). Ground flora: wide section of bramble ( <i>Rubus</i> sp.).
G	H-G-058	h2a - hedgerow (priority habitat)	Hedgerow on a bank between field and wood, with a track on wood boundary. Coppiced and pollarded trees present. Dead tree stumps. Ditch present from central section running north but has been silted up over time. Not likely to contain water.	Section 1: hawthorn ( <i>Crataegus monogyna</i> ) R, field maple ( <i>Acer campestre</i> ) O, blackthorn ( <i>Prunus spinosa</i> ) O, rose sp. ( <i>Rosa</i> sp.) R, dog's mercury ( <i>Mercurialis perennis</i> ) O, false brome ( <i>Brachypodium sylvaticum</i> ) R, herb-Robert ( <i>Geranium robertianum</i> ) R. Section 2: hawthorn ( <i>Crataegus monogyna</i> ), field maple ( <i>Acer campestre</i> ) O, coppiced hazel ( <i>Corylus avellana</i> ) R.
G	H-G-059	h2a - hedgerow (priority habitat)	Shrubby, native species rich hedgerow which has been flailed. Trees present. Along a dry ditch.	Section 1: mature oak sp. ( <i>Quercus</i> sp.) F, dog-rose ( <i>Rosa canina</i> ) sp. ( <i>Rosa</i> sp.) F, mature field maple ( <i>Acer campestre</i> ) F, spindle ( <i>Euonymus europaeus</i> ) O, semi-mature ash ( <i>Fraxinus excelsior</i> ) R, elder ( <i>Sambucus nigra</i> ) R, dogwood ( <i>Cornus sanguinea</i> ) R, elm sp. ( <i>Ulmus</i> sp.) R
G	H-G-60	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-61	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-62	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-63	h2 - hedgerow	UKHab category derived from aerial imagery	-

<b>Section</b>	<b>Habitat ID</b>	<b>Habitat Type</b>	<b>Description</b>	<b>Species List with DAFOR</b>
G	H-G-64	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-65	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-66	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-67	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-69	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-70	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-71	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-72	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-73	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-74	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-75	h2 - hedgerow	UKHab category derived from aerial imagery	-
G	H-G-76	h2 - hedgerow	UKHab category derived from aerial imagery	-
H	H-H-01	h2a - hedgerow (priority habitat)	Native hedgerow.	N/A. Not surveyed
H	H-H-02	h2a - hedgerow (priority habitat)	Native hedgerow with trees.	N/A. Not surveyed
H	H-H-03	h2a - hedgerow (priority habitat)	Native hedgerow with trees.	N/A. Not surveyed
H	H-H-04	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
H	H-H-08	h2a - hedgerow (priority habitat)	Hedgerow along an arable field boundary. Two mature oak species.	Two mature oak sp. ( <i>Quercus</i> sp.), wych elm ( <i>Ulmus glabra</i> ), hazel ( <i>Corylus avellana</i> ), rose sp. ( <i>Rosa</i> sp.), field maple ( <i>Acer campestre</i> ), blackthorn ( <i>Prunus spinosa</i> ), dogwood ( <i>Cornus sanguinea</i> ), hawthorn ( <i>Crataegus monogyna</i> )
H	H-H-09	h2a - hedgerow (priority habitat)	Defunct hedgerow (some greater than 5m) with trees. Present on the southern side of a dry ditch.	Hawthorn ( <i>Crataegus monogyna</i> ) O, bramble ( <i>Rubus</i> sp.) O, hazel ( <i>Corylus avellana</i> ) F, field maple ( <i>Acer campestre</i> ) F, mature oak sp. ( <i>Quercus</i> sp.) R, holly ( <i>Ilex aquifolium</i> ) O.
H	H-H-10	h2a - hedgerow (priority habitat)	Hedgerow with trees. Average 10m height, average 2.5m wide.	Section 1: dogwood ( <i>Cornus sanguinea</i> ), aspen ( <i>Populus tremula</i> ), oak sp. ( <i>Quercus</i> sp.), mature hazel ( <i>Corylus avellana</i> ), elm sp. ( <i>Ulmus</i> sp.), blackthorn ( <i>Prunus spinosa</i> ), hawthorn ( <i>Crataegus monogyna</i> ). Section 2: hawthorn ( <i>Crataegus monogyna</i> ), mature hazel ( <i>Corylus avellana</i> ), dogwood ( <i>Cornus sanguinea</i> ), blackthorn ( <i>Prunus spinosa</i> ), elm sp. ( <i>Ulmus</i> sp.), ash ( <i>Fraxinus excelsior</i> ), field maple ( <i>Acer campestre</i> ), holly ( <i>Ilex aquifolium</i> ), bracken ( <i>Pteridium aquilinum</i> ), bramble ( <i>Rubus</i> sp.), rose sp. ( <i>Rosa</i> sp.), willow sp. ( <i>Salix</i> sp.)
H	H-H-11	h2b - other hedgerows	Hedgerow ornamental and non-native.	N/A. Not surveyed
H	H-H-12	h2b - other hedgerows	Hedgerow ornamental and non-native. Not possible to survey up close.	N/A. Not surveyed
H	H-H-13	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees - associated with bank or ditch. Not surveyed.	N/A. Not surveyed
H	H-H-14	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees - associated with bank or ditch. Not surveyed.	N/A. Not surveyed

<b>Section</b>	<b>Habitat ID</b>	<b>Habitat Type</b>	<b>Description</b>	<b>Species List with DAFOR</b>
H	H-H-16	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees - associated with bank or ditch. Not surveyed.	N/A. Not surveyed
H	H-H-17	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees - associated with bank or ditch. Not surveyed.	N/A. Not surveyed
H	H-H-18	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees - associated with bank or ditch. Not surveyed.	N/A. Not surveyed
H	H-H-20	h2a - hedgerow (priority habitat)	Species poor and defunct hedgerow, with some trees. No ditch indicated.	N/A. Not surveyed
H	H-H-21	h2a - hedgerow (priority habitat)	Managed hedgerow with 3 trees along the length. Species diverse with native woody species. Two 6m gaps within hedgerow for field access.	Field maple ( <i>Acer campestre</i> ), hazel ( <i>Corylus avellana</i> ), ash ( <i>Fraxinus excelsior</i> ), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ).
H	H-H-22	h2a - hedgerow (priority habitat)	Intact hedgerow 2m tall x 2m wide with evidence of management. Semi-mature oaks along its length with two 6m gaps for field access.	Field maple ( <i>Acer campestre</i> ), hazel ( <i>Corylus avellana</i> ), ash ( <i>Fraxinus excelsior</i> ), elder ( <i>Sambucus nigra</i> ), hawthorn ( <i>Crataegus monogyna</i> ).
H	H-H-23	h2a - hedgerow (priority habitat)	Intact hedgerow 1.5-2 wide, 2m tall. Semi-mature oak located in the middle.	Semi-mature oak sp. ( <i>Quercus</i> sp.) O, hawthorn ( <i>Crataegus monogyna</i> ) F, blackthorn ( <i>Prunus spinosa</i> ) F, field maple ( <i>Acer campestre</i> ) F, hazel ( <i>Corylus avellana</i> ) F, dog-rose ( <i>Rosa canina</i> ) sp. ( <i>Rosa</i> sp.) O.
H	H-H-24	h2a - hedgerow (priority habitat)	Hedgerow associated with a titch. Hedgerow was 2m tall and 1.5m wide, small margins of grassland adjacent to arable fields.	Field maple ( <i>Acer campestre</i> ) A, blackthorn ( <i>Prunus spinosa</i> ) A, hawthorn ( <i>Crataegus monogyna</i> ) A.
H	H-H-28	h2a - hedgerow (priority habitat)	UKHab category derived from aerial imagery.	-
H	H-H-29	h2a - hedgerow (priority habitat)	Native species rich hedgerow with trees.	N/A. Not surveyed
H	H-H-30	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-31	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-32	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-33	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-34	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-35	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-36	h2 - hedgerow	UKHab category derived from aerial imagery.	-
H	H-H-37	h2 - hedgerow	UKHab category derived from aerial imagery.	-

Page intentionally blank

National Grid House,  
Warwick Technology Park,  
Gallows Hill, Warwick.  
CV34 6DA United Kingdom

Registered in England and Wales  
No. 4031152  
[nationalgrid.com](http://nationalgrid.com)