



# Gatwick Airport Northern Runway Project

Environmental Statement

Appendix 9.6.2: Ecology Survey Report – Part 1

**Book 5**

VERSION: 1.0

DATE: JULY 2023

Application Document Ref: 5.3

PINS Reference Number: TR020005

## Table of Contents

1	Introduction
2	Methodology
3	Results
4	Evaluation
5	Conclusions
6	References
7	Glossary

## Annex

Annex 1 Survey Methodologies

Annex 2 Extended Results

Annex 3 Evaluation

Annex 4 Invertebrate Scoping Survey

Annex 5 Terrestrial Invertebrate Survey

Annex 6 Aquatic Ecology Survey

Annex 7 Figures

## Tables

Table 3.11.1: Bat Emergence Survey Dates, Weather Conditions and Survey Times	16
Table 3.13.1: Bat Static/Automated Survey Summary for Location 1	17
Table 3.13.2: Species Summary for Location 1	17

Table 3.13.3: Bat Static/Automated Survey Summary for Location 2	1
Table 3.13.4: Species Summary for Location 2	1
Table 3.13.5: Bat Static/Automated Survey Summary for Location 3	3
Table 3.13.6: Species Summary for Location 3	27
Table 3.13.7: Bat Static/Automated Survey Summary for Location 4	32
Table 3.13.8: Species Summary for Location 4	32
Table 3.13.9: Bat Static/Automated Survey Summary for Location 5	33
Table 3.13.10: Species Summary for Location 5	
Table 3.13.11: Bat Static/Automated Survey Summary for Location 6	
Table 3.13.12: Species Summary for Location 6	
Table 3.13.13: Bat Static/Automated Survey Summary for Location 7	
Table 3.13.14: Species Summary for Location 7	
Table 3.13.15: Bat Static/Automated Survey Summary for Location 8	
Table 3.13.16: Species Summary for Location 8	
Table 3.13.17: Bat Static/Automated Survey Summary for Location 9	16
Table 3.13.18: Species Summary for Location 9	17
Table 3.13.19: Bat Static/Automated survey summary for Location 10	17
Table 3.13.20: Species Summary for Location 10	

Table 3.13.21: Bat Static/Automated Survey Summary for Location 11	18
Table 3.13.22: Species Summary for Location 11	18
Table 3.13.19: Bat Static/Automated survey summary for Location 12	18
Table 3.13.20: Species Summary for Location 12	19
Table 3.13.19: Bat Static/Automated survey summary for Location 13	19
Table 3.13.20: Species Summary for Location 13	19
Table 3.13.19: Bat Static/Automated survey summary for Location 14	20
Table 3.13.20: Species Summary for Location 14	21
Table 3.13.19: Bat Static/Automated survey summary for Location 15	21
Table 3.13.20: Species Summary for Location 15	21
Table 3.13.19: Bat Static/Automated survey summary for Location 16	21
Table 3.13.20: Species Summary for Location 16	22
Table 3.13.23: Total Bat Passes	22
Table 3.13.24 Breakdown of bat passes from River Mole	22
Table 3.13.25: Breakdown of bat passes from Riverside Garden Park	23
Table 4.1.1: Breeding Bird Assemblage Diversity Criteria	23
	23
	23

## 1 Introduction

- 1.1.1 The document forms **ES Appendix 9.6.2: Ecology Survey Report** (Doc Ref. 5.3) of the Environmental Statement (ES) prepared on behalf of Gatwick Airport Limited (GAL) for the proposal to make best use of Gatwick Airport's existing runways and infrastructure (referred to within this report as 'the Project').
- 1.1.2 This report provides details of ecological surveys undertaken on land within and around Gatwick Airport, Horley, West Sussex to inform the design of the Project, as described in **ES Chapter 5: Project Description** (Doc Ref. 5.1).
- 1.1.3 The areas surveyed included the land within the Project site boundary and adjacent areas of potential ecological interest, where access allowed. Where an area is outwith the Project site boundary, this is signposted in the text and shown on the relevant figures. All figures are presented in Annex 7 of this report.
- 1.1.4 The ecological surveys for protected or notable habitats or species comprised:
- Phase 1 habitat survey;
  - National Vegetation Classification (NVC) survey;
  - hedgerow survey;
  - breeding bird survey;
  - wintering bird survey;
  - reptile survey;
  - great crested newt *Triturus cristatus* survey;
  - dormouse *Muscardinus avellanarius* survey;
  - otter *Lutra lutra* survey;
  - water vole *Arvicola amphibius* survey;
  - badger *Meles meles* survey;
  - bat roost assessment;
  - bat emergence/re-entry surveys;
  - bat activity transect surveys;
  - bat crossing point surveys;
  - bat static/automated surveys;
  - terrestrial invertebrate survey;
  - aquatic invertebrate survey; and
  - fish survey.
- 1.1.5 Methodologies and results for the latter three surveys are presented in Annexes 5 and 6 of this report.
- 1.1.6 Additional surveys have also been undertaken for bats, including thermal imaging of bat activity on the runway and bat trapping/bat

tracking (**ES Appendix 9.6.3: Bat Trapping and Radio Tracking Surveys** (Doc Ref. 5.3)).

- 1.1.7 The methodologies and results of these surveys are described and presented within this report.

## 2 Methodology

### 2.1 Phase 1 Habitat Survey

- 2.1.1 The methodology and habitat descriptions used were based on the standard Joint Nature Conservation Committee (JNCC) Phase 1 Habitat Survey methodology 'Handbook for Phase 1 Habitat Survey' (JNCC, 2010).
- 2.1.2 The Phase 1 Habitat Survey was carried out on 18–22 March and 10 & 11 July 2019. The Phase 1 survey covered the area within the Project site boundary and adjacent habitats considered to be of potential ecological interest (Riverside Garden Park, for example).
- 2.1.3 Habitats identified during the survey were described using the categories set out in the Phase 1 handbook (JNCC, 2010).

### 2.2 NVC Survey

- 2.2.1 A NVC survey was carried out following the methodology and guidelines detailed in the JNCC's NVC User's Handbook (Rodwell *et al.*, 2006).
- 2.2.2 Fieldwork was carried out in April, July and August 2019 by a qualified ecologist and botanist. The survey was undertaken during the optimal times for both grassland and woodland botanical surveys.

- 2.2.3 The survey methodology is described in Annex 1.

### 2.3 Hedgerow Survey

- 2.3.1 A survey of all hedgerows within the Project site boundary was carried out in accordance with the methodology and guidelines set out in the Hedgerow Survey Handbook (Department for Environment, Food and Rural Affairs (Defra), 2007) to identify Important hedgerows, as defined in the Hedgerow Regulations 1997.
- 2.3.2 The assessment was carried out on the 5–8 August 2019.

- 2.3.3 Any hedgerows qualifying for protection under the Hedgerow Regulations 1997 were identified.

- 2.3.4 The survey methodology is described in Annex 1.

### 2.4 Breeding Bird Survey

- 2.4.1 The breeding bird survey undertaken was based on a standard territory mapping methodology as described in Gilbert *et al.* (1998) and Bibby *et al.* (2000).
- 2.4.2 This method is based on the principle that the majority of species are territorial during the breeding season. This takes into account birds occupying discrete territories and displaying various behaviours (eg conspicuous song, visual display and periodic disputes with neighbouring individuals) allowing their location and abundance to be estimated.
- 2.4.3 Surveys for breeding birds were undertaken in spring/summer 2019 with a total of seven survey visits taking place.
- 2.4.4 The survey area, as shown in Figure 2.4.1, was walked at a slow pace in order to locate and identify all individual birds. Visits were undertaken early in the morning, finishing before midday.
- 2.4.5 The survey methodology is described in Annex 1.

### 2.5 Wintering Bird Survey

- 2.5.1 The wintering bird surveys were based on a transect survey methodology as detailed in Bibby *et al.* (2000) and Gilbert *et al.* (1998).
- 2.5.2 The transect route was selected to include all field boundaries and visit all areas of the Project to within 200 metres, where possible. Visits were undertaken early in the morning.
- 2.5.3 All bird species were recorded and mapped across the whole Project site area, where accessible.
- 2.5.4 The survey methodology is described in Annex 1.

### 2.6 Reptile Survey

- 2.6.1 The reptile survey followed the recommended methodology described in the Herpetofauna Worker's Manual (JNCC, 2003) and Froglife's Surveying for Reptiles (Froglife, 2016).
- 2.6.2 It was undertaken by experienced ecologists and was conducted within areas of the Project identified as containing the most favourable habitat for reptiles.

<p>2.6.3 Reptiles are best surveyed from April following hibernation until June and then again in September and October.</p> <p>2.6.4 The survey methodology is described in Annex 1.</p> <p><b>2.7 Great Crested Newt Survey</b></p> <p>2.7.1 Each pond within the Project site boundary was assessed for its potential to support great crested newts, where accessible.</p> <p>2.7.2 Surveys were undertaken following the advice given in Froglife's 'Great Crested Newt Conservation Handbook' (2001), English Nature's 'Great Crested Newt Mitigation Guidelines' (English Nature, 2001) and the 'Herpetofauna Workers Manual' (Gent and Gibson, 2003).</p> <p>2.7.3 The survey methodology is described in in Annex 1.</p> <p><b>2.8 Dormouse Survey</b></p> <p>2.8.1 A dormouse nest tube survey was undertaken based on methodology and best practice guidelines set out in the Dormouse Conservation Handbook, second edition (Bright, Morris and Mitchell-Jones, 2006). Survey visits were undertaken regularly in suitable weather conditions between May and October 2019. Further visits were also undertaken on 27-28<sup>th</sup> May, 28-29<sup>th</sup> June, 18-19<sup>th</sup> July, 23-24<sup>th</sup> August, 20-21<sup>st</sup> September, 20<sup>th</sup>, 24<sup>th</sup> October, 28<sup>th</sup>, 29<sup>th</sup> and 30<sup>th</sup> November 2022. The survey methodology is described in Annex 1.</p> <p><b>2.9 Aquatic Mammal Survey</b></p> <p><b>Otter Survey</b></p> <p>2.9.1 The otter survey was undertaken on the 13<sup>th</sup> and 14<sup>th</sup> May 2019 by suitably experienced ecologists. Further surveys were also undertaken on the 3-6<sup>th</sup>, 11<sup>th</sup>, 13<sup>th</sup>, 19<sup>th</sup>, 20<sup>th</sup>, and 21<sup>st</sup> October 2022. The survey was based on the methodology described in the Design Manual for Roads and Bridges (DMRB), Volume 10 Section 4, Part 4 (Highways Agency <i>et al.</i>, 1999). Whilst the DMRB guidance has since been withdrawn and replaced by Volume 10, Section 4, Part 1 (LA 118) (Highways England <i>et al.</i>, 2019), no specific methodology in relation to otters has been revised. As such, the methodology contained within the former Volume 10, Section 4, Part 4 (Highways Agency <i>et al.</i>, 1999) remains relevant. The methodology was developed for linear schemes which are likely to affect otter habitats or populations but was adopted for this site.</p>	<p>2.9.2 The suitable areas along the River Mole and Gatwick Stream were walked and examined in detail for evidence of the presence of otters in the form of characteristic field signs.</p> <p>2.9.3 The survey methodology is described in Annex 1.</p> <p><b>Water Vole Survey</b></p> <p>2.9.4 The water vole survey was carried out on the 13<sup>th</sup> and 14<sup>th</sup> May 2019 by suitably experienced ecologists. Further surveys were also undertaken on the 3-6<sup>th</sup>, 11<sup>th</sup>, 13<sup>th</sup>, 19<sup>th</sup>, 20<sup>th</sup>, and 21<sup>st</sup> October 2022.</p> <p>2.9.5 The surveys were carried out in accordance with guidelines of best practice set out in the Water Vole Conservation Handbook – Third Edition (Strachan <i>et al.</i>, 2011).</p> <p>2.9.6 The suitable areas along the River Mole were walked and examined in detail for evidence of the presence of water vole in the form of characteristic field signs.</p> <p>2.9.7 Wherever possible, the banks were inspected on both sides, from the water's edge to the top of the bank.</p> <p>2.9.8 The survey methodology is described in Annex 1.</p> <p><b>2.10 Badger Survey</b></p> <p>2.10.1 The site was systematically searched for evidence of badgers during walkover surveys. This involved looking for setts, latrines, hairs, footprints, runs, and any other signs of badger activity. Any evidence recorded was mapped.</p> <p>2.10.2 Further details of the badger survey methodologies and results are provided in confidential <b>ES Appendix 9.6.4: Confidential Badger Survey</b> (Doc Ref. 5.3).</p> <p><b>2.11 Preliminary Bat Roost Assessment</b></p> <p><b>Buildings</b></p> <p>2.11.1 An assessment of the suitability of the buildings within the landside and airside areas of the Project site boundary for bat roosting potential was undertaken at the same time as the Phase 1 Habitat Survey.</p> <p>2.11.2 The survey included a thorough ground level inspection of the exterior of all accessible buildings and the features of the buildings listed below were noted:</p> <ul style="list-style-type: none"> <li>▪ type;</li> </ul>	<ul style="list-style-type: none"> <li>▪ wall construction, in particular the type of material used;</li> <li>▪ form of the roof, in particular the presence of gable ends, hipped roofs etc and the nature and condition of the roof; and</li> <li>▪ the general condition of the building.</li> </ul> <p>2.11.3 The methodology is detailed in full within Annex 1.</p> <p><b>Trees</b></p> <p>2.11.4 A ground-level Preliminary Bat Roost Assessment (PBRA) of trees along the A23 within the site boundary was undertaken in November and December 2022.</p> <p>2.11.5 Following guidance from the Bat Conservation Trust Bat Survey: Good Practice Guidelines (BCT, 2016) trees were assessed as having the potential to support bat roosts if they had potential bat roost features.</p> <p>2.11.6 The suitability of the trees for roosting bats was also assessed by examining the surrounding habitat.</p> <p>2.11.7 When suitable features were identified, they were inspected for signs indicating use or possible use by bats.</p> <p>2.11.8 Details on the methodology for bat roost assessment of trees is to follow.</p> <p><b>2.12 Bat Emergence/Re-entry Surveys</b></p> <p>2.12.1 In order to comply with best practice guidelines (Collins, 2016) emergence surveys were carried out on any buildings considered to have bat roosting potential. Surveys were undertaken between May and October 2019. The aims of these surveys were to determine the use of the buildings (if any) by roosting bats, the egress locations of any bats emerging from the buildings and the species assemblage within the Project site boundary.</p> <p>2.12.2 The methodology is described in Annex 1.</p> <p><b>2.13 Bat Activity Transect Surveys</b></p> <p>2.13.1 A total of five transect routes were devised to cover a broad range of the habitat types present on site but focusing on those likely to be of greatest value to bats, including woodland, woodland edges, river corridors and open grassland. Descriptions of each transect are in Annex 1.</p>
--	---	---

- 2.13.2 Each transect was surveyed twice per month between April and October 2019.
- 2.13.3 The methodology is described in Annex 1.
- 2.13.4 In addition, three further transects were completed in August to October 2020 and April to July 2021 within areas not surveyed in 2020. The transects are described in Annex 1.
- 2.13.5 A further transect was carried out in May to October 2022. This transect was to the north of the site taking in grassland, scattered tress and riverside habitats. The transect is described in Annex 1.
- 2.14 Bat Static/Automated Surveys**
- 2.14.1 A total of 11 Elekon Batlogger A units were deployed across the Project site between April and October 2019 for a minimum of five nights. The units were positioned at various locations, in order to sample a broad range of the habitat types present on site but focusing on those likely to be of greatest value to bats.
- 2.14.2 In addition, 5 new locations were monitored between May and October 2022. Locations are described in Section 3.13 below.
- 2.14.3 The methodology is described in Annex 1.
- 2.15 Bat Crossing Point Surveys**
- 2.15.1 Bat Crossing Point surveys were undertaken at two locations, the River Mole corridor and Riverside Park, in August 2020, September 2020, May 2021 and June 2021.
- 2.15.2 The methodology is provided in Annex 1.
- 2.16 Invertebrate Scoping Survey**
- 2.16.1 The invertebrate scoping survey was carried out by Marcel Ashby and Tristan Bantock, for Colin Plant Associates.
- 2.16.2 The survey assessed the potential for the Project site to support Species of Principal Importance in England, as defined within Section 41 of the Natural Environment and Rural Communities (NERC) Act 2006, although species included in other conservation categories were also considered.
- 2.16.3 The survey report, including the methodology, is at Annex 4.
- 2.17 Terrestrial Invertebrate Surveys**
- 2.17.1 Walk-over surveys for terrestrial invertebrates were completed by Ecus Ltd. on six occasions during 2020 – 27th May, 19th June,

- 22th June, 30th June, 10th September and 14th September. These focused on areas along the River Mole and the Gatwick Stream. On each occasion, the areas were walked by an experienced entomologist who sampled along each transect using sweep netting, a beating tray and a stout trowel.
- 2.17.2 The survey concentrated on the following major groups (orders): Coleoptera (beetles), Diptera (flies), Hemiptera (bugs, froghoppers, etc), Hymenoptera (bees, wasps and ants) and Lepidoptera (butterflies and moths). Some examples of other groups were noted if found.
- 2.17.3 Samples were collected for later laboratory identification.
- 2.17.4 The survey report, including the methodology, is at Annex 5.
- 2.18 Aquatic Macroinvertebrate Surveys**
- 2.18.1 Following an initial scoping walk-over, 100 m sections of both the River Mole and Gatwick Stream were identified for detailed survey as representative of the site. Three survey visits were undertaken during 2020 by Ecus Ltd.; 4th June, 29th July and 29th September. Samples were collected at each of the sites using the Whalley Hawkes Paisley Trigg (WHPT) method comprising a standard three-minute kick sample using a long-handled pond net with 1 mm mesh size, which was supplemented by a one-minute hand search.
- 2.18.2 The survey report, including the methodology, is at Annex 6.
- 2.19 Fish Survey**
- 2.19.1 Fish surveys were undertaken by Ecus Ltd. using the catch depletion method in order to assess species composition, age structure and to estimate population size. Surveys were undertaken by an accredited electric fishing team comprising three members of staff. Surveys and analysis conformed to the relevant guidance outlined in BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity (British Standards Institute, 2003).
- 2.19.2 Surveys were undertaken in summer (4th June) and autumn (29th September) 2020 along the same 100 m stretches used for the aquatic invertebrate surveys.
- 2.19.3 The survey report, including the methodology, is at Annex 6.

- 2.20 Veteran Trees**
- 2.20.1 All surveyed trees were assessed in general accordance with the requirements set out in BS 5837:2012 “Trees in Relation to Design, Demolition and Construction – Recommendations”, by a fully qualified and experienced Arboriculturist.
- 2.20.2 The veteran tree surveys were carried out on 16<sup>th</sup> and 24<sup>th</sup> May and 14<sup>th</sup>, 16<sup>th</sup> and 30<sup>th</sup> June 2022. All information was digitally captured on site, using a tablet running Axciscape 4.02 software. This is a program specifically designed for arboricultural surveying, which allows trees to be located directly onto a digital copy of a site’s topographical survey.
- 2.20.3 The methodology is included in Annex 1.
- 3 Results**
- 3.1 Phase 1 Habitat Surveys**
- 3.1.1 The Project site is located on the Sussex/Surrey border, west of the M23 and east of Charlwood. The majority of the site is a working airport comprising large areas of hardstanding associated with the runways and taxiways; buildings including terminals, hangars and other buildings associated with airport activities; and well-maintained amenity grassland surrounding the runway. All of these areas were surrounded by security fencing.
- 3.1.2 Around the main airport the site comprised broadleaved and mixed woodland; neutral semi-improved, poor semi-improved, marshy and improved grassland; scattered and dense scrub; tall ruderal vegetation; running water; areas of standing water; dry ditches; species poor and species rich hedgerows; individual trees; dry ditches; fences; residential and commercial buildings; and areas of hardstanding.
- 3.1.3 The site was divided into eight areas (A1-A8) that were based on land use types and land ownership boundaries.
- 3.1.4 Descriptions of the habitats identified within each of the eight areas during the survey are given below. The locations of each habitat within each survey area are shown on the Phase 1 habitat plan (Figures 3.1.1 and 3.1.2). The locations of ponds are shown on Figure 3.1.3a and b.
- 3.1.5 A list of target notes is provided in Table 4 in Annex 2 and these are referred to within the text. The locations of the target notes are shown on Figures 3.1.1 and 3.1.2.

## A1 – Fields to the North and South of the M23 for Construction Laydown plus Link to Junction 9

### A1.2 Broadleaved Plantation Woodland

3.1.16 Along the embankments of Airport Way, highway associated planting of trees was present from the B2036 in the east stretching to the railway boundary in the west. Trees in this area included semi-mature poplar *Populus sp.* around the northern edge of the roundabout, hawthorn *Crataegus monogyna*, pedunculate oak *Quercus robur*, ash *Fraxinus excelsior* and field maple *Acer campestre*.

3.1.17 Scattered through the trees, bramble *Rubus fruticosus* scrub was dense in patches.

3.1.18 At the far eastern end of the M23 spur, part of the junction had been planted with poplar and silver Birch *Betula pendula*, east of the scrub.

### A2.1 Dense Scrub

3.1.19 Along the bottom of the southern M23 spur bank, dense bramble scrub over a wooden fence formed the boundary between highway land and the public footpath through the fields south of the road. Within the scrub there were occasional oak and ash saplings growing through the brambles.

3.1.10 Bramble scrub dominated the north western end of the spur road bank.

3.1.11 Directly south, the scrub became less dense and the vegetation turned more ruderal with rosebay willowherb *Chamaenerion angustifolium*, swards of Yorkshire fog *Holcus lanatus*, thistle *Cirsium sp.* and broad-leaved dock *Rumex obtusifolius*.

### A2.2 Scattered Scrub

3.1.12 Five areas of scattered scrub were present around the M23 and the spur road.

3.1.13 In the east, along the M23, scattered scrub was present in the piece of land between the motorway and the northbound slip road. Bramble scrub dominated here with occasional hawthorn and blackthorn *Prunus spinosa*.

3.1.14 On the western half of the junction where the spur meets the motorway, bramble and blackthorn scrub dominated with poor semi-improved grass in places.

3.1.15 In the south east of the M23 spur area, there was a small patch of bramble scrub (TN14) with Yorkshire fog, perennial rye-grass *Lolium perenne*, cleavers *Galium aparine*, primrose *Primula vulgaris*, ivy *Hedera helix* and dock also present.

3.1.16 Along the field boundary next to the railway, north of the M23 spur, there was scattered scrub dominated by bramble, field rose *Rosa arvensis* and elder *Sambucus nigra*.

3.1.17 Along the northern boundary of Airport Way, bramble scrub was interspersed between young and semi-mature trees associated with highways planting.

### A3.1 Scattered Broadleaved Trees

3.1.18 Within the roundabout that connects the M23 spur to Airport Way, and north of the Long Stay South Car Park entrance, several poplar trees had been planted as ornamental features.

3.1.19 Along the eastern edge of Car Park 'B' a planted treeline between the edge of the car park and the western edge of the public footpath within this area comprised ash, downy birch *Betula pubescens*, silver birch and immature sycamore *Acer pseudoplatanus*. Bramble and ivy were also present.

### B2.2 Neutral Semi-improved Grassland

3.1.20 Directly south of the M23 spur road and east of the B2036, one field, with a public footpath going through, was dominated by cock's foot *Dactylus glomerata* and Yorkshire fog. However, there were also a number of herbaceous species throughout the grass including white clover *Trifolium repens*, Cut-leaved cranesbill *Geranium dissectum*, stitchwort *Stellaria sp.*, perennial rye-grass, shining cranesbill *Geranium lucidum*, vetch *Vicia sp.*, buttercup *Ranunculus sp.*, birds-foot trefoil *Lotus corniculatus*, creeping cinquefoil *Potentilla reptans*, dock and pyramidal orchids *Anacamptis pyramidalis*. Bramble also ran through the grass.

### B4 Improved Grassland

3.1.21 The improved grassland pasture fields north of the roundabout, east of the B2036 and the pasture fields south of the M23 spur were dominated by Yorkshire fog and annual meadow-grass *Poa annua* with abundant dandelion *Taraxacum officinale* and occasional spear thistle *Cirsium vulgare*.

### F1 Swamp

3.1.22 The area immediately surrounding Pond E11 was dominated by reedmace *Typha latifolia* creating a swamp habitat.

### G1 Standing Water

3.1.23 An artificial attenuation pond (Pond E11) had been created at the eastern end of the spur road. The margins were dominated by reedmace *Typha sp.* and common reed *Phragmites australis*.

### J1.2 Amenity Grassland

3.1.24 Amenity grassland was present within some parts of the central reservation and associated with the roundabout linking the M23 spur and Airport Way.

### J2.3.2 Species-poor Hedge with Trees

3.1.25 To the north of the M23 spur road, along the northern boundary of a public footpath, an old hedge with mature trees was present. The hedgerow was an oak and sycamore dominated treeline with a hawthorn hedge running underneath. Field rose was occasionally present.

3.1.26 A hedge with trees was present along the northern Project site boundary. This was located to the north of the roundabout linking the M23 spur to Airport Way and comprised oak, horse chestnut *Aesculus hippocastanum* and copper beech *Fagus sylvatica*.

### J2.4 Fence

3.1.27 Around much of the area wooden fencing was present, mainly along the edge of public footpaths and road boundaries. Along the edge of the railway line and in staff Car Park B more secure metal fencing was used.

### J2.6 Dry Ditch

3.1.28 At the time of survey seven field boundary ditches were dry. These ditches were mainly located in the land south of the M23 spur. Three of these ran north to south within the improved grassland field. One dry ditch ran east to west along the south embankment of the spur road, one east to west under a treeline south of the neutral semi-improved grassland and one north to south along the east edge of the B2036 Balcombe Road.

3.1.29 One dry ditch was identified north of the spur road running in a north to south direction along the western edge of the B2036 Balcombe Road.

3.1.30 No aquatic vegetation was present within the ditches. The two ditches along the B2036 had common nettle *Urtica dioica* and hogweed *Heracleum sphondylium* growing out of them. The ditches under treelines were choked with fallen leaf litter.

<p><b>J4 Bare Ground</b></p> <p>3.1.31 Part of the northern highway’s embankment had been cleared at the eastern end of the M23 spur to form a site compound for the M23 Smart Motorway upgrade works.</p>	<p>3.1.39 Throughout the Long Stay South car parks, woodland had been planted around remnants of old field boundaries with mature trees incorporated amongst the newer planting.</p>	<p>3.1.50 Areas of shorter managed grass were a regular occurrence within the northern section of ring road south, and a further area of grassland in the west of the site, between the railway and Pond F, was also identified as having a regular management regime.</p>
<p><b>J5 Other (Hardstanding)</b></p> <p>3.1.32 The M23, Airport Way, B2036 and staff Car Park B were all tarmacked surfaces with heavy use.</p> <p>3.1.33 The main London to Brighton trainline ran north to south between the northern pasture field and staff Car Park B.</p>	<p><b>A3.1 Scattered Broadleaved Trees</b></p>	<p>3.1.51 Pennyroyal <i>Mentha pulegium</i> was found growing within this area of grassland (TN1). Information on this species protected status can be found in Annex 1.</p>
<p><b>A2 – Eastern Car Parking and Associated Surface Water Features</b></p> <p><b>A1.1.1 Semi-natural Broadleaved Woodland</b></p>	<p>3.1.40 Towards the eastern side of Pentagon Field, two isolated, mature oak trees were present.</p>	<p><b>J1.4 Introduced Shrub</b></p>
<p>3.1.34 To the south of the Long Stay South Car Park, there was a large area of semi-natural broadleaved woodland that formed the northern portion of Horleyland Wood and Lower Pickett’s Wood. These two areas have a similar range of species, with the canopy dominated by oak.</p>	<p>3.1.41 There were planted semi-mature and mature trees around the roads, car parks and within the roundabout linking the Long Stay South Car Park to the M23 Spur road. Trees were planted in small groups over non-native shrubs and amenity grassland.</p>	<p>3.1.52 Within the parking areas and along some of ring road south, non-native shrubs and hedgerows had been planted in borders. All the shrubs were relatively low in height and well maintained. Due to the non-native planting, species within these areas were not recorded.</p>
<p>3.1.35 West of Pentagon Field was a large triangular area of woodland, within the Long Stay South Car Park. This area of woodland was predominantly on a raised earth bank with ditches around the northern and southern bases. An access track heading east to west split the woodland from a line of trees further north. Blackthorn, yew <i>Taxus baccata</i> and bramble scrub covered the eastern and southern banks.</p>	<p>3.1.42 Further young and mature trees were present throughout the Parking area east of the railway. Some of the tree lines were associated with old field boundaries with mature oak and beech. Some of the younger treelines were present over ornamental shrubs and amenity grassland.</p>	<p><b>J2.4 Fence</b></p>
<p>3.1.36 Heading from this woodland south along the western edge of Pentagon Field, the woodland continued until it joined with Lower Pickett’s Wood.</p>	<p><b>B6 Poor Semi-improved Grassland</b></p>	<p>3.1.53 Around Long Stay South Car Park large metal security fences lined the boundary of GAL owned land and other land ownership boundaries, including highway land and the railway.</p>
<p><b>A1.1.2 Plantation Broadleaved Woodland</b></p>	<p>3.1.43 To the east of the Long Stay South Car Park and west of the B2036, Pentagon Field was a large, open grazed field. The majority of the field was dominated by cock’s foot and perennial rye-grass. Areas around the eastern boundary and in places through the field were wetter and had rushes <i>Juncus sp.</i> and sedges <i>Carex sp.</i> colonising.</p>	<p><b>J2.8 Earth Banks</b></p>
<p>3.1.37 Along the boundaries of the M23 spur strips, woodland had been planted on the bank of the carriageway. These areas spanned from the London to Brighton railway in the west to the B2036 in the East.</p>	<p><b>G1 Standing Water</b></p>	<p>3.1.54 The majority of the woodland sections were on raised earth banks. The banks were approximately 1.5 metres high and varied in width and length.</p>
<p>3.1.38 Forming the southern boundary and a sizable portion of the western boundary of Pentagon Field, oak and hazel <i>Corylus avellana</i> had been planted in rows. The western boundary planting being older than the southern boundary plantation.</p>	<p>3.1.44 Wet ditches surrounded the western, northern and eastern boundaries of Pentagon Field. These ditches were on the road and footpath side of the field boundary fences.</p>	<p>3.1.55 A larger earth bank (TN2) was identified within a section of woodland in the east of Long Stay South Car Park. The bank here was approximately 3 metres high and 25 metres x 55 metres in extent.</p>
	<p>3.1.45 Pond G was located within the eastern car parking zone. Ditches that held water were also present around the majority of the car parks.</p>	<p><b>J3.6 Buildings</b></p>
	<p>3.1.46 The majority of the water bodies were associated with flood management and were man made.</p>	<p>3.1.56 A range of building types were identified throughout the wider eastern car parking area. These buildings were associated with commercial businesses, hotels, airport car parking and private office blocks. The majority of these were large multi-storey buildings.</p>
	<p>3.1.47 The northern-most of these water bodies was a man-made holding lake with a barrier across it. The banks were vegetated with willow <i>Salix sp.</i>, reedmace and common reed.</p>	<p><b>J5 Other (Hardstanding)</b></p>
	<p>3.1.48 A description of the waterbodies can be found in Annex 1.</p>	<p>3.1.57 The majority of the car parks were large open tarmacked areas with walkways and raised planting. The southern and western most car parks had steel multi-storey parking within the parking areas.</p>
	<p><b>J1.2 Amenity Grassland</b></p>	<p>3.1.58 A raised walkway ran along the western edge of Horleyland Wood connecting the wood to the car parks further north.</p>
	<p>3.1.49 A large area of well kept, regularly mown grassland was identified within the roundabout connecting the M23 spur to Airport Way.</p>	

### A3 – Land East of the Railway Line

#### A1.1.1 Semi-natural Broadleaved Woodland

- 3.1.59 Within the biodiversity area there were two distinct areas of semi-natural broadleaved woodland, which included areas of ancient and semi-natural woodland. These two areas are referred to as Horleyland Wood (TN3) and Upper Pickett's Wood (TN4).
- 3.1.60 Both Horleyland Wood and Upper Pickett's Wood were predominantly oak with beech, birch and ash also throughout. The ground flora differs between the woodlands however with Horleyland Wood being dominated by bracken *Pteridium aquilinum*. Upper Pickett's Wood had a more diverse woodland ground flora with species such as wood avens *Geum urbanum*, enchanter's nightshade *Circaea lutetiana* and ground ivy *Glechoma hederacea*.
- 3.1.61 Located to the south of Upper Pickett's Wood, two Notable species; Solomon's seal *Polygonatum odoratum* and narrow-lipped helleborine *Epipactis leptochila* were present (TN5), both of which were found within 20 metres of one another and are designated as Nationally Scarce. Bluebell *Hyacinthoides non-scripta* (a species listed under Schedule 8 of the Wildlife and Countryside Act 1981) was also found throughout the woodland.
- 3.1.62 South of the biodiversity car park an area of woodland surrounded a field on the western, eastern and southern boundaries. This area was dominated by oak and ash with occasional hawthorn, blackthorn, elder, beech and hazel present.
- 3.1.63 Little ground flora was present in this area of woodland, but occasional fern species were scattered throughout.

#### A1.1.2 Broadleaved Woodland Plantation

- 3.1.64 Several small areas of plantation woodland were recorded.
- 3.1.65 The area to the east of Gatwick Stream consisted of willow and alder *Alnus glutinosa* (TN6a).
- 3.1.66 The area to the south of the Crawley Sewage Treatment works consisted of coppiced hazel (TN6b).
- 3.1.67 The area located between these two areas consisted of beech (TN6c).

#### A2.1 Dense Scrub

- 3.1.68 Areas of scrub dominated by young stands of willow and alder with bramble were situated throughout the attenuation fields.

These stands had been planted between the retained earth banks and were bounded by a wooden fence.

#### A3.1 Scattered Broadleaved Trees

- 3.1.69 Mature oak trees were scattered throughout the attenuation fields upon earth mounds. Their locations are shown on the phase one habitat plan (Figures 3.1.1 and 3.1.2).

#### B2.2 Neutral Semi-improved Grassland

- 3.1.70 The grassland to the south of the biodiversity area and around the perimeter of the attenuation fields was neutral semi-improved grassland.
- 3.1.71 The grassland around the edge of the attenuation fields was herb rich and appears to originate from a seed mix. Here, wild carrot *Daucus carota*, black knapweed *Centaurea nigra*, oxeye daisy *Leucanthemum vulgare* and yellow rattle *Rhianthus minor* were the dominant herbs throughout, with timothy grass *Phleum pratense* and false oat-grass *Arrhenatherum elatior* as the dominant grass species. Crested dog's-tail *Cynosurus cristatus* and sweet vernal-grass *Anthoxanthum odoratum* were also frequent throughout.
- 3.1.72 The northern-most fields contained large patches of ruderal species with spear thistle *Cirsium vulgare* and dock species being the dominant herbaceous species. False oat-grass was the dominant grass species in this area of the fields.
- 3.1.73 The field to the south of the car park differed in species composition with common bent *Agrostis capillaris* being the dominant species with grass vetchling *Lathyrus nissola* and birds-foot trefoil *Lotus corniculatus* the most dominant herbaceous species.

#### B5 Marsh/Marshy Grassland

- 3.1.74 The attenuation fields south of Crawley Sewage Treatment Works had a different species composition to that of the raised banks that ran around the perimeter of the area.
- 3.1.75 Dominant species within the attenuation area were both hard rush *Juncus inflexus* and soft rush *Juncus effusus* with floating sweet-grass *Glyceria fluitans* and common water-plantain *Alisma plantago-aquatica* frequent within the wetter areas.
- 3.1.76 The drier areas within the attenuation fields had a varying species composition with grass species such as sweet vernal-grass, timothy grass, red fescue *Festuca rubra* and crested dog's-tail abundant throughout.

- 3.1.77 Forb species such as knapweed, meadow buttercup *Ranunculus acris* and cuckoo flower *Cardamine pratensis* were also frequent throughout the attenuation fields.

- 3.1.78 Throughout Upper Pickett's Wood were open areas of wet grassland. Young oak was scattered throughout with bramble growing around the edges of the openings. Both soft and hard rush were frequent throughout the grassland with species such as glaucous sedge *Carex flacca*, hairy sedge *Carex hirta*, marsh thistle *Cirsium palustre* and meadowsweet *Filipendula ulmaria* all indicative of marshy habitats. Other species found here were cock's-foot, red fescue and common knapweed among others.

#### C3.1 Tall Ruderal

- 3.1.79 Multiple stands of ruderal species were identified across the fields in the biodiversity area.
- 3.1.80 A long stretch of tall ruderal vegetation ran along the eastern boundary of the railway line. The species here were predominantly a mix of spear thistle and broad-leaved dock.
- 3.1.81 Within the areas of semi-improved grassland south of the sewage works were many areas of ruderal vegetation. These habitats were all similar in species composition with spear thistle, creeping thistle, ragwort *Senecio vulgaris*, burdock *Arctium minor* and various dock species present.

#### G1 Standing Water

- 3.1.82 A number of ponds were present throughout the biodiversity area. The locations of the ponds are shown on Figure 3.1.3a and b.
- 3.1.83 Pond 8N8 was located north of the Old Lagoon and south of the New Lagoon. See Annex 2 for the pond description.
- 3.1.84 The Old Lagoon was located within the biodiversity woodland and formed part of the sewage works. See Annex 2 for the pond description.
- 3.1.85 Ponds AA20 and AA21 were located within the area of mixed woodland to the south of Upper Pickett's Wood. See Annex 2 for the pond descriptions.
- 3.1.86 Ponds 1WH and NU1 were located to the south of the biodiversity area in a small patch of woodland located along the road to the Crawley Sewage Treatment Works. See Annex 2 for the pond descriptions.
- 3.1.87 Pond 30P was located within woodland south of Upper Pickett's Wood. See Annex 2 for the pond description.



### G2 Running Water

- 3.1.88 The Gatwick Stream ran between the attenuation fields and the neutral grassland to the east. The stream was around 5 m across and was fast flowing. The banks were steep and covered in vegetation. Himalayan balsam *Impatiens glandulifera* (a non-native invasive species listed in Schedule 9 of the Wildlife and Countryside Act 1981) was found growing along the banks of the stream.

### J1.2 Amenity Grassland

- 3.1.89 The grassland around the sewage works lagoon was heavily managed and cut short. The species here were predominantly grasses with species such as perennial rye-grass being dominant.

### J2.4 Fence

- 3.1.90 The raised banks throughout the attenuation field had a wooden fence around the outside.
- 3.1.91 The areas of scrub in the attenuation fields had a wooden fence around the outside.
- 3.1.92 A wooden fence ran along the northern edge of the marshy grassland within the attenuation fields east of the rail line.
- 3.1.93 A metal security fence ran along both sides of the Gatwick Stream between the Crawley Sewage Treatment Works and the attenuation fields east of the railway line. Security fencing also surrounded the Crawley Sewage Treatment Works and the Old Lagoon and New Lagoon.

### J2.8 Earth Banks

- 3.1.94 Throughout the attenuation fields were retained earth banks. Atop each bank stood a mature oak. The ground flora here differed from that of the surrounding habitat as woodland species such as dog's mercury *Mercurialis perennis* were present. Other species included bracken, fox glove *Digitalis purpurea*, bramble and false oat-grass.
- 3.1.95 Numerous linear earth banks were also situated throughout the fields to the south of the Crawley Sewage Treatment Works.

### J4 Bare Ground

- 3.1.96 An area of bare ground in the biodiversity area was used as a car park by the biodiversity team.

- 3.1.97 A bare ground path also ran through Upper Picketts Wood, to the east of the Old Lagoon and continued through the woodland.

### J5 Other (Hardstanding)

- 3.1.98 The access road to the Crawley Sewage Treatment works split the biodiversity area into two. The road was tarmacked with treelines down both sides.

### A4 – Airside

#### A1.1.1 Semi-natural Broadleaved Woodland

- 3.1.99 Crawler's Wood (TN13) was located along the southern boundary of the airside land parcel. The woodland had a high diversity of broadleaved tree species, the most prevalent being sycamore, field maple, birch and ash. The ground flora was relatively species poor with a dense mat of ivy covering the ground.

#### A1.3.2 Mixed Plantation Woodland

- 3.1.100 To the north of the airside land parcel ran a raised earth bank. This had a ground cover of amenity grassland with mixed planted woodland growing along the bank. The tree species consisted of mainly beech and oak with conifers growing throughout.

#### A2.1 Dense Continuous Scrub

- 3.1.101 An area of dense scrub was located around Pond FFJ. This was dominated by bramble and young shrubs such as hawthorn and willow.

#### A2.2 Scattered Scrub

- 3.1.102 The raised earth bank to the far east of the runway was covered in scattered scrub. Species such as young willow, gorse *Ulex europaeus*, and oak were scattered throughout with bramble and ruderal species such as sow-thistle *Sonchus sp.* and broad-leaved dock also present.

#### B6 Poor Semi-improved Grassland

- 3.1.103 An area of poor semi-improved grassland ran between Crawler's Wood and the amenity grassland associated with the airfield. This section of grassland was much longer with a greater species diversity than that of the amenity grassland areas.
- 3.1.104 The species here consisted of Yorkshire fog, false oat-grass, common bent, and cock's-foot with flowering species such as white clover, red clover *Trifolium pratense*, thistle, knapweed, hogweed and bird's-foot trefoil.

### G1 Standing Water

- 3.1.105 Pond FFJ was located to the north of the runway near the fire training area. It was surrounded by dense bramble scrub and marginal vegetation such as pond sedge *Carex riparia* and reedmace.

### G2 Running Water

- 3.1.106 Crawler's Brook, a 3 m wide stream, ran along the southern boundary of the main runway. The banks were regularly cleared of vegetation and the brook dredged. Some marginal vegetation was present in places with Himalayan balsam and rushes growing along the bank.

### J1.2 Amenity Grassland

- 3.1.107 The grassland surveyed around the runways was identified as being amenity grassland due to the high levels of management. The grassland was regularly mown to around 10-14 cm with selective herbicide applied.
- 3.1.108 The grassland was made up of Yorkshire fog, false oat-grass and common bent with flowering species such as clover, thistle, hogweed and bird's-foot trefoil.
- 3.1.109 To the north of airside was a raised bank of well managed amenity grassland with planted trees.

### J2.4 Fence

- 3.1.110 Around the airfield and buildings associated with the airfield, were metal security gates topped with razor wire.

### J2.7 Earth Bank

- 3.1.111 Multiple earth banks were identified in the north west corner of the airfield.
- 3.1.112 The largest of the banks was dominated by scrub with gorse, bramble, hogweed and young oak.

- 3.1.113 Many smaller banks were also noted. These however were not scrubby in habit with the vegetation being consistent with the amenity grassland.

### J3.6 Buildings

- 3.1.114 A large number of buildings were across the airside parcel. J5 Other (Hardstanding)
- 3.1.115 All the runways, taxiways and roads airside were tarmacked and in good condition.

## A5 – Non-Airside South and Land East of the Aviation Museum

### A1.1.1 Semi-natural Broadleaved Woodland

- 3.1.116 The strip of woodland that ran between Gatwick Aviation Museum Field and the River Mole was dominated by oak, birch and sycamore with hawthorn and blackthorn understorey. The ground flora was sparse with ivy and bramble being common throughout.
- 3.1.117 The woodland to the west of the Fire Training Ground was young woodland with the dominant species being a mixture of birch and sycamore. The understorey was dominated by dense patches of bramble. There was much deadwood present throughout the woodland.
- 3.1.118 Crawter’s Wood ran along the southern edge of the site. The species composition here was consistent with that of the woodland along the western side of the River Mole, with sycamore and oak among other species.

### A3.1 Scattered Broadleaved Trees

- 3.1.119 Along the northern perimeter of Car Park X, young willow and alder trees lined a wet ditch.
- 3.1.120 Lining the emergency access to Car Park X a mature treeline was present either side of the access track. The line was dominated by ash and oak.
- 3.1.121 Individual oak trees were scattered throughout Car Park X.
- 3.1.122 Within the land north of the Fire Training Ground there were a number of scattered mature oak trees along an old field boundary north of the marshy grassland area.

### A2.1 Dense/Continuous Scrub

- 3.1.123 A patch of dense scrub was located around Pond 29A, north east of the Fire Training Ground and west of the River Mole. These areas had long grass, mainly false oat-grass and cock’s-foot with bramble and young hawthorn growing throughout.

### A2.1 Scattered Scrub

- 3.1.124 There were areas of scrub throughout the land east of the Aviation Museum all predominantly bramble. The areas of scrub were situated along the margins of woodland and are shown on the Phase 1 Habitat plan (Figures 3.1.1 and 3.1.2a – 3.1.2l).

## B6 Poor Semi-improved Grassland

- 3.1.125 The fenced off area to the west and north east of the Fire Training Ground was a mix of habitat types, a large section of which was semi-improved grassland. The dominant species here were creeping thistle *Cirsium arvense* and false-oat grass. Hard rush was also frequent throughout the wetter areas.

## B4 Improved Grassland

- 3.1.126 The fields across the land east of the Aviation Museum were improved grassland. The species composition consisted of mainly Yorkshire fog and perennial rye-grass with forbs such as daisy *Bellis perennis* frequent throughout.

## B5 Marshy Grassland

- 3.1.127 An area of marshy grassland was found within the south of the land east of the Aviation Museum. This was seasonally wet with wet depressions throughout. The dominant species was false oat-grass with stands of rushes around the wetter areas. Ruderal species such as common nettle and thistle were also frequent throughout the grassland.

## C3.1 Tall Ruderal

- 3.1.128 A large patch of tall ruderal vegetation dominated by spear thistle and broad-leaved dock was found within the fenced off section west of the Fire Training Ground.

## G1 Standing Water

- 3.1.129 Pond 29A was located between the Gatwick Aviation Museum Field and the River Mole. The pond was man made with steep sides. Minimal vegetation was growing within the pond with only a small amount of soft rush.
- 3.1.130 Pond AVF was located to the south of the Aviation Museum Field. Marginal vegetation surrounded the pond, the dominant species was reedmace.
- 3.1.131 A wet drainage ditch runs though the secure area to the west of the Fire Training Ground. The species here are typical of wet ground with soft rush and reedmace present.
- 3.1.132 Pond MHA was located in the south west of Car Park X. The pond was seasonal and held water during wetter times of the year. Woodland and scrub surrounded the pond.

## J1.2 Amenity Grassland

- 3.1.133 Patches of heavily managed short amenity grassland was identified along the southern boundary as grass verges along roads and around car parks. The species were consistent throughout with perennial rye-grass, cock’s-foot, buttercup, dandelion and dock.

## J2.1.1 Native Species-rich Hedge

- 3.1.134 One hedgerow located along the western boundary of the Project site within the land east of the Gatwick Aviation Museum Field was found to be species rich. This contained hawthorn, blackthorn, dog rose *Rosa canina*, ash, dogwood *Cornus sanguinea* and oak. A hedgerow assessment was carried out and it was found not to be an important hedgerow.

## J2.3.2 Species-poor Hedge with Trees

- 3.1.135 Within the land east of the Gatwick Aviation Museum Field, the majority of hedgerows were found to be species poor. They were predominantly made up of a mix of hawthorn and blackthorn. These hedgerows had mature oak and ash trees scattered throughout, some starting to take over and forming taller hedgerows, especially along Man’s Brook and the hedgerows running south from there.

## J2.4 Fence

- 3.1.136 Poorly kept wooden fences bordered the fields within the land east of the Gatwick Aviation Museum Field. Hedges and trees had encroached and caused the fences to be in a state of disrepair.
- 3.1.137 Around the southern edge of Crawter’s Wood, metal security fencing was present.

## J5 Other (Hardstanding)

- 3.1.138 Large areas of hardstanding were recorded throughout the car parks and roads across the site.

## A6 – The North West Zone, containing the River Mole Corridor and Brockley Wood Biodiversity Area

### A1.1.1 Semi-natural Broadleaved Woodland

- 3.1.139 Brockley wood (TN7) was located along the River Mole corridor, between Gatwick airside and the River Mole. The woodland was dominated by oak with birch also being present. The ground flora was a mix of species typical of woodland habitats such as ivy, ground ivy and wood avens.

3.1.140 Along the northern bank of the River Mole corridor runs a long continuous stretch of semi-natural broadleaved woodland. The species composition changed in dominance throughout with the more dominant species being sycamore, oak, ash, birch and willow. The understorey species consisted mainly of hawthorn, dog rose, honeysuckle *Lonicera periclymenum* and bramble. The ground flora varied throughout the woodland with some areas of bare ground. The dominant species throughout the woodland were mainly ground ivy, ivy, common nettle, lords and ladies *Arum maculatum* and wood avens among others.

3.1.141 The woodland along the south of the River Mole was very similar in species composition to that of the woodland to the north. The main difference was the south bank was much more steeply sloping and black poplar *Populus nigra* was present.

#### A1.1.2 Broadleaved Plantation Woodland

3.1.142 The southern embankment of the River Mole flood plain was planted with native broadleaved tree species following the realignment of the River Mole to its current course. The planting extends from Brockley Wood in the south to London Road in the north.

3.1.143 The trees have grown creating a dense woodland dominated by oak, silver birch, willow, poplar, hawthorn and blackthorn.

3.1.144 The ground flora within this woodland comprised wild garlic *Allium ursinum*, hogweed, broad-leaved dock, bramble, lesser celandine *Ficaria verna*, daffodil *Narcissus sp.*, cuckoo flower, perennial rye-grass, compact rush *Juncus conglomeratus*, small-leaved nettle *Urtica urens*, reedmace, creeping cinquefoil *Potentilla reptans*, dove's-foot crane's-bill *Geranium molle*, cherry laurel *Prunus laurocerasus*, common vetch and sedges *Carex sp.*

#### A2.1 Dense Continuous Scrub

3.1.145 An area of dense scrub was situated adjacent to Brockley Wood (TN8). The area contained stands of bramble with hawthorn present throughout. Areas of raised banks and ditches ran through this area with rushes growing in the wetter parts. False oat-grass was common throughout as well as thistle.

3.1.146 Areas of dense scrub were present in the field to the north of Longbridge Roundabout, comprising blackthorn, hawthorn and bramble, with occasional elder.

#### A2.2 Scattered Scrub

3.1.147 Scattered scrub was growing on the banks of a large earth bank south of Brockley Wood. The species consisted of mainly bramble with young saplings such as willow, oak and hawthorn. False oat-grass, wild carrot and common knapweed were frequent in the less scrubby areas.

3.1.148 Within the marshy area of Pond C24, scattered willow and alder scrub was growing.

3.1.149 Patchy areas of scattered bramble scrub were present along the eastern bank of the field to the north of Longbridge Roundabout.

#### A3.1 Scattered Broadleaved Trees

3.1.150 Scattered mature trees were present along the southern, eastern and northern boundaries of the field to the north of Longbridge Roundabout, with species comprising oak, ash, field maple and elder.

#### A3.3 Mixed Scattered Trees

3.1.151 Within Longbridge Roundabout a mix of semi-mature broadleaved and coniferous trees had been planted. Tree species included oak, silver birch and Leyland cypress.

3.1.152 Around the north west corner of the roundabout, south east of the Holiday Inn, coniferous trees lined the eastern side of the amenity grassland, west of the pavement. A single Leyland cypress, a sycamore and a cherry were within the line of conifers.

#### B2.2 Semi-improved Neutral Grassland

3.1.153 The semi-improved grassland along the River Mole runs along the south bank. The lower lying areas of grassland contained a higher number of wetland species associated with regular flooding.

3.1.154 The grassland had a diverse mix of species, the dominant grass species was tufted hair-grass *Deschampsia cespitosa* which dominated large swards. In areas where this was less dominant, other grasses such as false oat-grass, timothy grass and meadow foxtail *Alopecurus pratensis* were frequent. In these areas the most common forb species were wild carrot, ox-eye daisy, greater bird's-foot trefoil *Lotus pedunculatus* and red bartsia *Odontites vernus*.

3.1.155 Lesser quaking grass *Briza minor* and ragged robin *Lychnis flos-cuculi* were both found within this area (TN9). They are

designated as Nationally Scarce and Near Threatened respectively.

#### B5 Marshy Grassland

3.1.156 Areas of marshy grassland were present towards the south of the River Mole.

3.1.157 A large area of marshy grassland was located to the south of Brockley Wood (TN10a) (Figure 3.1.2j). The grassland here was seasonally wet and was relatively species poor, dominated with hard rush.

3.1.158 A smaller section of marshy grassland (TN10b) (Figure 3.1.2j) was located south east of Brockley Wood. The dominant species here was common reed. Purple loose-strife *Lythrum salicaria* and thistle were also present throughout.

3.1.159 The southern margins of the River Mole were relatively diverse (TN10c) (Figure 3.1.2j). This area contained a range of species including purple loose-strife, common reed, marsh woundwort *Stachys palustris* and hard rush. Himalayan balsam was also abundant in this section of the River Mole.

3.1.160 There was a small area of marshy grassland situated around Pond C24. This was dominated by common reed. Reedmace was also present growing in the pond.

#### B6 Poor Semi-improved Grassland

3.1.161 A small section of poor semi-improved grassland was located around Pond M. The grassland here was less diverse than that along the River Mole and appeared to be managed more heavily.

3.1.162 Longbridge Roundabout comprised managed grassland and mature trees. Species comprised cock's-foot, ribwort plantain *Plantago lanceolata*, common speedwell *Veronica persica*, dandelion and dock.

3.1.163 To the north of Longbridge Roundabout was a semi-improved field. The field was divided into two separate areas; the western half was managed as a paddock for horses, dominated by annual meadow-grass, with occasional dandelion, Yorkshire fog, bristly ox-tongue *Helminthotheca echioides*, field speedwell and red clover.

#### C3.1 Tall Ruderal

3.1.164 The eastern half of the paddock was less frequently managed and contained a greater variety of ruderal and scrub species such as spear thistle, creeping thistle, oxeye daisy, broadleaved dock,

ragwort, buttercup, ribwort plantain, shepherd's purse *Capsella bursa-pastoris*, red dead nettle, hogweed, common nettle and cleavers. A large stand of Himalayan balsam was present on the southern boundary of the field and adjacent to the River Mole, along the eastern boundary.

**F2.2.1 Marginal Vegetation**

3.1.165 Marginal vegetation ran along the edge of the River Mole. This was dominated by common reed and reedmace. Himalayan balsam and hemlock water-dropwort *Oenanthe crocata* were also occasional along the River Mole.

**G1 Standing Water**

3.1.166 Ponds were present along the River Mole Corridor. See Annex 2 for the pond references and descriptions. The locations of the ponds are shown on Figure 3.1.3a and b.

3.1.167 Pond A was situated south of Brockley Wood and east of the Fire Training Ground. Pond A was used as an attenuation pond to hold run-off from the airfield.

3.1.168 Pond M was located along the southern boundary of the River Mole. It was situated to the west of the north stay car park and used as a water reservoir.

3.1.169 Pond C24 was located along the northern edge of the River Mole Corridor. It was situated within semi-improved grassland.

3.1.170 Pond D was situated along the northern boundary of the River Mole Corridor, near the Travelodge hotel. Both ponds were used as reservoirs.

3.1.171 Pond AAA4 was located along the northern edge of the River Mole Corridor. It was situated within the northern area of semi-natural broadleaved woodland.

**G2 Running Water**

3.1.172 The River Mole was fast flowing and up to 3 metres wide. It flowed north to south and within the river corridor had shallow banks with floodplains and marshy grassland areas to the east, and woodland, with public footpaths to the west.

**J1.2 Amenity Grassland**

3.1.173 Amenity grassland was identified in several areas, it was mainly situated around the reservoirs along the southern side of the River Mole Corridor. The grass in these areas was heavily managed.

**J2.4 Fence**

3.1.174 Around the southern edge of Brockley Wood, deer fencing had been erected.

3.1.175 Along the top of the southern section of the River Mole floodplain, a wooden fence was present.

**J2.8 Earth Bank**

3.1.176 South of Brockley Wood and east of the River Mole, a large 8 metre tall earth bank was located (TN11). The bank was steep sided with scattered scrub growing over semi-improved grassland.

3.1.177 An earth track led to the top of the bank, which was flat with pooling water in places.

**J4 Bare Ground**

3.1.178 Areas of bare ground were identified as earth tracks running along the southern section of the River Mole.

**J5 Other (Hardstanding)**

3.1.179 Areas of hardstanding were located along the River Mole Corridor. This included the Long Stay North car parking, parking off Charlwood Road, Perimeter Road North and the bridge spanning the River Mole.

**A7 – Non-airside North**

**A1.1.1 Semi-natural Broadleaved Woodland**

3.1.180 Dog Kennel Wood (TN12) was a small area of woodland in the north east of the non-airside north section. Canopy species within this area of woodland were predominantly oak, horse chestnut and sweet chestnut *Castanea sativa*. Downy birch, goat willow *Salix caprea*, cherry *Prunus sp.*, ash and yew were also occasionally present within the canopy. The understorey comprised bramble, holly *Ilex aquifolium*, elder, rose, gorse and hazel with nettle. Bluebell, dog's mercury, violet *Viola sp.*, daffodil and arum were present in the ground flora.

**A2.2 Scattered Scrub**

3.1.181 Within the man-made ditches around the western-most Long Stay North Car Parks scattered willow and silver birch shrubs dominated the banks. Rosebay willowherb was occasionally present.

3.1.182 Along the western-most ditch was a patch of bramble.

**A2.3 Dense Scrub**

3.1.183 Bramble had choked the southern ditches between the hedgerows and treelines.

3.1.184 In the south east corner of Long Stay North, a triangle shaped area of dense bramble, elder and hawthorn scrub had grown up around a couple of lines of trees.

**A3.1 Scattered Broadleaved Trees**

3.1.185 Within the north of Long Stay North, there were three lines of mature trees including oak, lime and horse chestnut.

3.1.186 The southern-most of these mature lines formed a rectangular area and included beech and ash.

3.1.187 Along the northern side of the southern ditches, within Long Stay North, trees had been planted in rows. Most of the trees were ornamental or non-native and young to semi-mature in age.

3.1.188 Along the western edge of Long Stay North, six mature oaks lined the fence.

**B6 Poor Semi-improved Grassland**

3.1.189 The grassland around the bottom of Pond M and west of the security fence contained areas of taller, less managed grassland with false oat-grass, cock's-foot and perennial rye-grass with occasional birds-foot trefoil, ragwort, annual meadow-grass and dock.

3.1.190 The banks of the ditches in the west of Long Stay North were managed grassland with occasional bramble growing through.

**F2.1 Marginal Vegetation**

3.1.191 Within the bottom of Dog Kennel Pond a variety of marginal species dominated the lower banks and pond bed including willow, willowherb, pendulous sedge *Carex pendula*, water mint *Mentha aquatica*, field horsetail *Equisetum arvense*, reedmace, teasel, gypsywort *Lycopus europaeus*, soft rush, purple loosestrife, false fox-sedge *Carex otrubae*, redshank *Persicaria maculosa*, round-fruited rush *Juncus compressus*, common reed, marsh horsetail *Equisetum palustre*, common spike-rush *Eleocharis palustris* and wood club-rush *Scirpus sylvaticus*

**G1 Standing Water**

3.1.192 Pond M was present north east of Brockley Wood in the west of the non-airside north area. The banks of the pond were well kept grassland.

- 3.1.193 Dog Kennel Pond was a small manmade attenuation pond with steep banks showing high levels of maintenance. A diverse mix of aquatic and marginal vegetation was found within the pond.
- 3.1.194 Within the western-most Long Stay North Car Park there was a man-made ditch. The ditch was present in the northern third of this area of car park and along the western and southern edge of the car park. The banks were 1 metre high made from crushed stone and tarmac. Less than 25 cm depth of water was present. Reedmace dominated the wetter areas.
- J1.2 Amenity Grassland**
- 3.1.195 Around the banks of Dog Kennel Pond in the west of the non-airside area was well managed grassland with cock's-foot, perennial rye-grass, annual meadow-grass, birds-foot trefoil, cut-leaved cranesbill and occasional bristly ox-tongue.
- J1.4 Introduced Shrub (Ornamental Planting)**
- 3.1.196 There were areas of non-native ornamental planting within the Long Stay North Car Park.
- J2.1.2 Species-poor Hedgerow**
- 3.1.197 Along the southern boundary and some of the ditches, blackthorn, silver birch, hazel and willow hedgerows had been left to go patchy.
- J2.3.6 Dry Ditch**
- 3.1.198 Dry ditches were present throughout Long Stay North Car Park, these ditches were associated with flood alleviation and were predominantly in the west and south of the Car Park.
- 3.1.199 All had similar characteristics of being 1 to 2 m deep with scattered scrub along the edges. Some had hedgerows and treelines along them as well.
- J3.6 Buildings**
- 3.1.200 A range of building types were around the northern area of the terminal.
- 3.1.201 All buildings were associated with airport activity such as offices, terminals and industrial sections of the airport. One building was associated with the Police dog kennels.
- J5 Other (Hardstanding)**
- 3.1.202 The north east of the non-airside north area of Gatwick was the Long Stay North car parks and walkways.
- 3.1.203 A number of roads and access roads were located to the north of the airfield perimeter security fencing that linked Brockley Wood in the east to the Airport Way/London Road slipway roundabout in the west. There were also several service roads and hardstandings linked to the industries around this area.
- 3.1.204 South of the new Boeing hangar was a material store for Tarmac.
- A8 – Riverside Garden Park**
- 3.1.205 The majority of Riverside Garden Park is not included in the Project site boundary but was surveyed during an early phase of the Project. As such, details are provided below for context.
- A1.1.1 Semi-natural Broadleaved Woodland**
- 3.1.206 Riverside Garden Park is largely semi-natural broadleaved woodland (see Figure 3.1.2I). The woodland was dominated by oak and sycamore with ash, hazel, goat willow, cherry and alder also occurring. Turkey oak *Quercus cerris* was recorded around the lake and along the river towards the southern edge of the park.
- 3.1.207 The ground flora was largely dominated by bramble with much fallen leaf litter. Some *Buddleia* was present. Herb species included lesser celandine, herb robert *Geranium robertianum*, common nettle, dandelion, Yorkshire fog, ivy, cleavers, holly, wild garlic *Allium ursinum*, geranium species, lords and ladies, hart's tongue *Asplenium scolopendrium*, pendulous sedge, and lady fern *Dryopteris Felix-femina*.
- A1.1.2 Broadleaved Plantation Woodland**
- 3.1.208 Where the highway embankment rose to approximately 4 metres high at the southern end of Riverside Garden Park, a mixture of young and semi-mature oak, sycamore, elder, blackthorn, hazel and field maple had been planted. Dog rose was occasional.
- A2.1 Dense/Continuous Scrub**
- 3.1.209 Patches of dense and continuous scrub was present within the Riverside Garden Park survey area. Along the edge of the eastern London Road to Airport Way slip road banking, the western-most 300 metres of the bank was continuous bramble and gorse scrub with elder and hawthorn occurring frequently. Ribwort plantain and young hazel were also recorded.
- 3.1.210 At the north western end of Riverside Garden Park, in the areas where tree cover was limited, areas of bramble had colonised. The most prevalent area of scrub was a large section of the western bank of the Gatwick Stream that had become dominated by bramble and hawthorn. Himalayan balsam was also scattered through the scrub along the bank of the Gatwick Stream.
- 3.1.211 Bramble scrub formed a transitional habitat from woodland to grassland around the margins of some of the open glades, the largest of these areas being around the top of the northern-most glade.
- 3.1.212 A break in the tree cover had allowed brambles to take over and become dense within an old entrance to Riverside Garden Park from London Road.
- 3.1.213 An area of overgrown bramble and rose dominated part of the southern glade within Riverside Garden Park (TN15) (see Figure 3.1.2I).
- 3.1.214 Two further areas of dense scrub were identified within an old poorly kept paddock, east of the Gatwick Stream and south of the Riverside Road residential parking area. One area was located along the north eastern boundary of the paddock. The other area of scrub was along the top of the eastern bank of the Gatwick Stream.
- B6 Poor Semi-improved Grassland**
- 3.1.215 Several large open areas within the woodland were managed and mown regularly. These areas were dominated by perennial rye-grass and annual meadow-grass. Other occasional grass species in these areas included rough meadow-grass *Poa trivialis* and sweet vernal-grass. Localised patches of cock's-foot, wall barley *Hordium murialis* and meadow foxtail were also present.
- 3.1.216 Herb species which also occurred included white clover, creeping buttercup, creeping thistle, greater plantain, curled dock, spear thistle, dandelion, musk mallow *Malva moschata*, cow parsley, agrimony *Agrimonia eupatoria*, cut-leaved cranesbill, white dead nettle *Lamium album*, common sorrel *Rumex acetosa*, square-stalked St. John's wort *Hypericum tetrapterum*, early forget-me-not *Myosotis ramossima* and birds-foot trefoil.
- 3.1.217 A track/footpath passed through the north of the site from the car park towards London Road. It formed a ride as it passed through the woodland and generally had similar characteristics to the open grassland areas with a similar species composition. Some additional localised species were observed here including hedge woundwort *Stachys sylvatica*, dog rose, ribwort plantain, common selfheal *Prunella vulgaris*, meadowsweet, wood avens and common knapweed.

3.1.218 Several footpaths through areas of open grass had been worn down to bare ground.

### C3.1 Tall Ruderal

3.1.219 Common nettle, cleavers, curled dock, hogweed and willowherb were present in an area along the western bank of the Gatwick Stream in the north of the site. This area was between the northern-most glade and areas of continuous scrub further north towards the confluence of the River Mole and Gatwick Stream.

3.1.220 Ruderal vegetation was present on the earth banks surrounding the carpark. These were localised to the north west corner and south west corner of the bank. Species in these areas included common nettle, dock, hogweed, bindweed, white dead-nettle, white clover, dandelion and buttercup. Burdock was localised to the north west corner of the bank only.

3.1.221 Within the horse paddock east of Riverside Garden Park an area of ruderal vegetation with dock and common nettle was present along the eastern boundary of the paddock.

### G1 Standing Water

3.1.222 A large fishing lake with several wooded islands was located in the centre of the park. The banks of the river were shallow and bare, with occasional aquatic vegetation close to the margins.

### G2 Running Water

3.1.223 The Gatwick Stream ran the length of the eastern side of the park. The stream was 3-5 m wide and steeply banked along the majority of its length. The stream was culverted to the south of the park as it went under the railway line, terminals and airport car parks.

3.1.224 The Gatwick Stream formed a distributary of the River Mole that in the northern most part of the Riverside Garden Park splits from the Mole as the River Mole continues west towards the runways and down the River Mole corridor.

3.1.225 Aquatic vegetation associated with the Gatwick Stream included yellow-flag Iris *Iris pseudacorus*, lesser water-parsnip *Berula erecta* and Himalayan balsam.

### J2.8 Earth Banks

3.1.226 Around the car park, earth banks were present on all sides, and on either side of the car park entrance after the bridge. The banks were dominated by grasses and ruderal vegetation.

### J2.3.1 Native Species-rich Hedge with Trees

3.1.227 Along the eastern side of the London Road footpath was a planted native hedge dominated by hazel, field maple and hawthorn. The planted hedgerow continued from the southern end of the park to approximately halfway along the park boundary. Elder and ash were also occasional.

3.1.228 Towards the northern end of the hedgerow mature silver birch were spaced at regular intervals within the hedge.

3.1.229 The hedge was underlined by a mixture of woven hazel and wooden fencing.

### J5 Other (Hardstanding)

3.1.230 There was a tarmac footpath/cycle way around the north-eastern side of Riverside Garden Park. This joined the underpass towards the short stay south carparks and bus station. The cycle way was lined by street lamps at approximately 50-100 m apart.

3.1.231 West of the park, London Road was aligned north to south with an associated footpath on the eastern carriageway. At the southern end of the park, where the slip road rose on an artificial bank, the tarmacked footpath continues south going through an underpass towards Gatwick train station.

### J4 Bare Ground

3.1.232 Riverside Garden Park car park was a small rectangular car park, the ground of which was compacted earth and rubble with earth banks surrounding it. A worn-down footpath had been created in the western earth bank.

## 3.2 NVC Surveys

### Site Description

3.2.1 The northern-most stretch of the grassy habitat along the southern bank of the River Mole, as shown in the Phase 1 plan (Figures 3.1.1 and 3.1.2a – 3.1.2l), was identified as having a botanically interesting mix of grassland habitats and so a NVC survey was carried out.

3.2.2 The site consisted of a range of grassland habitats depending on the soil conditions. In wetter areas such as along the edge of the River Mole and the more low-lying areas, large continuous stands of marginal vegetation were dominant. In drier areas the vegetation changed with much more diverse grassland being present. At the upper most reaches of the river corridor the grassland gave way to scrub and woodland.

### NVC Categories

#### S4 – *Phragmites australis* Swamp and Reed-beds

3.2.3 The marginal vegetation along the banks of the River Mole was consistent with the NVC category S4 *Phragmites australis* swamp and reed-beds. S4 is described as having an overwhelming dominance of *P. australis*. This was characteristic of the vegetation along the banks of the River Mole, as *P. australis* was the dominant species and in most cases the only species present.

3.2.4 S4 is described as being a species poor habitat with large continuous stands of *P. australis*, often clonal. *Typha latifolia* was also present in a small stand along the River Mole. This was consistent with the S4 NVC community.

3.2.5 S4 shows a strong affinity with that of the habitat on site, as there were large stands of *P. australis* present.

#### MG9b - *Holcus lanatus* – *Deschampsia cespitosa* grassland. *Arrhenatherum elatior* Sub-community.

3.2.6 Stands of MG9b *Deschampsia cespitosa* grassland. *Arrhenatherum elatius* sub-community were identified along the Mole corridor. These stands were found to be present on the dryer raised areas of ground.

3.2.7 MG9b is characteristic of permanently moist and periodically inundated soils in British lowlands. It is commonly found on sloping ground in pastures and meadows along water bodies. This matched the habitat surveyed along the stretch of the Mole corridor.

3.2.8 MG9b is described as being dominated by tussocky grasses such as *D. cespitosa*, *Holcus lanatus*, *Dactylis glomerata* and *A. elatius*. In shorter areas of vegetation, the species composition varies with *Alopecurus pratensis* and *Agrostis stolonifera* being present. Many forb species were also present such as *Centaurea nigra*, *Ranunculus acris*, *Lathyrus pratensis* and *Plantago lanceolata*.

3.2.9 The species described were all found on site showing the grassland on site had a good affinity with MG9b.

3.2.10 The species composition within the quadrats observed is detailed within Annex 2.

<p><b>M27c - <i>Filipendula ulmaria</i> - <i>Angelica sylvestris</i> mire. <i>Juncus effusus</i> – <i>Holcus lanatus</i> Sub-community.</b></p>	<p>3.2.18 British Plant Communities Vol. 4 (Rodwell, 1995) describes many areas of swamp and tall herb fen as difficult to classify due to the species poor nature of these habitats. It suggests in cases where they do not fit any particular NVC category instead the area should be grouped and labelled as a society of the dominant species. This approach has been used for these areas of vegetation and the habitat has been classified as <i>Calamagrostis epigejos</i> society.</p>	<p>3.5.2 Twenty-four species recorded during the surveys meet at least one of a range of criteria relating to conservation importance. These species, and the relevant criteria of conservation importance, are shown in Table 19, Annex 2 and their distribution is shown on Figure 3.5.1.</p>
<p>3.2.11 A small patch of M27c - <i>Filipendula ulmaria</i> - <i>Angelica sylvestris</i> mire. <i>Juncus effusus</i> – <i>Holcus lanatus</i> sub-community was located in a low-lying area that appeared to be a drainage ditch. This NVC community was very localised to the wet ditch so was therefore a result of its construction.</p>	<p>3.2.19 The species composition within the quadrats observed is detailed in Annex 2. The classification areas are shown on Figure 3.2.1.</p>	<p>3.5.3 The conservation status used for the analysis of the wintering bird survey is based on the criteria when the survey was undertaken in and the data analysed in 2019, updated to the 2021 list (BoCC 2021).</p>
<p>3.2.12 M27c is described as having <i>F. ulmaria</i> as a dominant to abundant species. This was not the case with the habitat recorded on site as although it was present (outside of the quadrats) it was not the dominant species. Other species described in M27c were, however, present and matched that described in 'British Plant Communities Vol. 2'. <i>Juncus effusus</i>, <i>Holcus lanatus</i>, <i>Mentha aquatica</i> and <i>Oenanthe crocata</i> were all present in the stand.</p>	<p><b>3.3 Hedgerow Surveys</b></p> <p>3.3.1 None of the hedgerows surveyed were found to comprise important hedgerows.</p>	<p>3.5.4 One species (red kite), afforded special statutory protection under Annex 1 of the EU Birds Directive (Directive 2009/147/EC), was recorded flying over the Project area during the winter bird surveys.</p>
<p>3.2.13 M27c is described as occurring in moist, rich soils protected from grazing, being found across lowland Britain. It is typical of slow-moving streams, dykes and roadside ditches. This fitted the habitat on site as it was found along a low-lying wet ditch.</p>	<p><b>3.4 Breeding Bird Surveys</b></p> <p>3.4.1 A total of 72 species were recorded during the survey of breeding birds in 2019. Of these species, 48 were confirmed to be breeding and three possibly breeding (peregrine, little ringed plover and firecrest).</p>	<p>3.5.5 Nine species of principal importance listed under Section 41 of the NERC Act (2006), and also listed as UK Biodiversity Action Plan (BAP) Priority Species, were recorded during wintering bird surveys comprising bullfinch, dunnock, herring gull, house sparrow, lapwing, marsh tit, skylark, song thrush and starling.</p>
<p>3.2.14 Although the dominant species does not match the description of M27c, the habitat description, and a large proportion of the less dominant species have a good match. M27c therefore shows an affinity with the habitat on site.</p>	<p>3.4.2 Table 16 of Annex 2 provides a summary of the breeding and conservation status of the 72 species recorded during the course of the survey, with the numbers of territories identified (or estimated in the case of probable and possible records). The conservation status used for the analysis of the breeding bird survey is based on the criteria when the survey was undertaken and the data analysed in 2019 (BoCC 2015).</p>	<p>3.5.6 Eleven species recorded during the wintering bird surveys are included on the BoCC Red List and 12 species are included on the BoCC Amber List (BoCC 2021).</p>
<p>3.2.15 The species composition within the quadrats observed is detailed in Annex 2.</p>		<p>3.5.7 Further discussion of the species of conservation concern identified within the Project site boundary is provided below.</p>
<p><b><i>Calamagrostis epigejos</i> Society</b></p>		
<p>3.2.16 Across much of the surveyed area of the River Mole corridor were continuous stands of <i>Calamagrostis epigejos</i>. These stands were very species poor and in most cases were pure stands of this one species. When analysing these stands of vegetation using Modular Analysis of Vegetation Information System (MAVIS) no clear NVC community was identified, and those that were suggested all had low co-efficiency values, with the communities not matching that of the habitat on site.</p>	<p>3.4.3 The locations of territories of NERC Act Species of Principal Importance and Birds of Conservation Concern (BoCC) Red or Amber listed species recorded breeding within the survey area are shown in Figures 3.4.1a and 3.4.1b. The location of Annex 1 and/or Schedule 1 species recorded as possibly breeding within the survey area are shown in Figure 3.4.1c.</p> <p>3.4.4 Detailed results are provided in Annex 3.</p>	<p>3.5.8 Bullfinch is a common resident breeding and wintering bird in the UK with an estimated population of 220,000 birds (Musgrove <i>et al.</i>, 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.</p>
<p>3.2.17 The dominant species <i>C. epegejos</i> only appears within the floristic tables for S24, a community of tall herb and fen vegetation. This category is a better fit than that of the three suggested and better described the habitat found on site. However, the species composition varies significantly with S24 described as species rich.</p>	<p><b>3.5 Wintering Bird Surveys</b></p> <p>3.5.1 A total of 61 species were recorded during the wintering bird surveys undertaken between October 2018 and March 2019. A summary of the species recorded, together with the peak and mean counts of species, is provided in Table 18, Annex 2.</p>	<p>3.5.9 Black-headed gull is a frequent breeding bird in the UK with an estimated breeding population of 140,000 birds (Musgrove <i>et al.</i>, 2013). Over winter, the UK population of black-headed gulls significantly increases up to an estimated 2.3 million birds. As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.</p> <p>3.5.10 Common Gull is a relatively frequent resident breeding bird in the UK, with an estimated population of 49,000 birds (Musgrove <i>et al.</i>, 2013). Common gull is however a common winter visitor with an estimated winter population size of 710,000 birds (Musgrove</p>

	<i>et al.</i> , 2013). Only one observation of common gull was recorded during the surveys and therefore considered unremarkable and broadly representative of the species in the wider landscape.		considered unremarkable and broadly representative of the species in the wider landscape.		the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.11	Dunnock is a common resident breeding and wintering bird in the UK with an estimated population size of 2,500,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.18	Kestrel is a widespread resident breeding and wintering bird in the UK with an estimated population size of 46,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.24	Meadow pipit is a common resident breeding and wintering bird in the UK with an estimated population size of 2,000,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.12	Fieldfare is a rare breeding bird in the UK but a common winter visitor with an estimated winter population size of 720,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.19	Lapwing is one of the most widespread non-breeding wintering waders with an estimated over wintering population of around 650,000 birds. In general, lapwings tend to be concentrated in central and southern Britain during the winter (Lack, 1986). A high proportion of the birds that winter in Britain are of Scandinavian, Danish, Dutch and North German origin (Imboden, 1974). Lapwings respond rapidly to cold weather, and the numbers and distribution of non-breeding birds are strongly influenced by weather patterns in the UK as well as in continental Europe (Kirby and Lack, 1993). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in pastoral farmland in the South East of the UK. However, the site was considered likely to have some minor importance for wintering lapwing due to the likely suitable foraging habitat it supports.	3.5.25	Red kite is a restricted resident breeding and wintering bird in the UK with an estimated population size of 1,600 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.13	Green sandpiper is a rare breeding bird in the UK with only the occasional pair recorded breeding each year. Population estimates of wintering green sandpiper suggest that fewer than 1,000 birds spend the winter in the UK, although rather more are seen on passage (Musgrove <i>et al.</i> , 2013). Only one observation of green sandpiper was recorded during the surveys and, therefore, considered unremarkable and broadly representative of the species in the wider landscape.	3.5.20	Lesser black-backed gull is a widespread resident breeding and wintering bird in the UK with an estimated population size of between 110,000 and 130,000 birds. As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.26	Redwing is a rare breeding bird in the UK but a common winter visitor with an estimated winter population size of 690,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.14	Greylag goose is a relatively frequent breeding bird in the UK, with an estimated breeding population of 46,000 birds. Greylag goose is however a common winter visitor with an estimated winter population size of 230,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.21	Mallard is a common and widespread resident breeding bird in the UK, with an estimated population of between 61,000 and 146,000 birds (Musgrove <i>et al.</i> , 2013). Mallard is also a common winter visitor with an estimated winter population size of 710,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.27	Skylark is a common resident breeding and wintering bird in the UK with an estimated population size of 1,500,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.15	Grey wagtail is a common resident bird in the UK with an estimated population of 38,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.22	Marsh tit is a common resident breeding and wintering bird in the UK with an estimated population of 41,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.28	Snipe is a relatively frequent resident breeding bird in the UK, with an estimated population of 80,000 birds (Musgrove <i>et al.</i> , 2013). Snipe is however a common winter visitor with an estimated winter population size of 1.1 million birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.16	Herring gull is a widespread breeding bird in the UK and a common winter visitor with an estimated winter population size of 740,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.	3.5.23	Mistle thrush is a widespread resident breeding and wintering bird in the UK with an estimated population size of 170,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during	3.5.29	Song thrush is a common resident breeding and wintering bird in the UK with an estimated population size of 1,200,000 birds (Musgrove <i>et al.</i> , 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.
3.5.17	House sparrow is a widespread but declining resident bird in the UK with an estimated population of around 5.3 million birds. As such the numbers recorded during the winter bird surveys are			3.5.30	Starling is a common resident breeding bird in the UK with an estimated population size of 1,800,000 birds (Musgrove <i>et al.</i> , 2013) this population swells in the winter with an additional influx of continental birds although no official estimate of the wintering population is available. As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.



3.5.31 Woodcock is a restricted resident breeding bird and widespread wintering bird in the UK with an estimated wintering population size of 1,400,000 birds (Musgrove *et al.*, 2013). As such the numbers recorded during the winter bird surveys are considered unremarkable and broadly representative of the species in the wider landscape.

## 3.6 Reptile Surveys

### Habitat Assessment

3.6.1 The Phase 1 Habitat Survey identified a range of habitats within the Project site boundary that provided good hibernating, basking, and foraging habitat for reptiles.

3.6.2 The areas that were deemed to have the best reptile habitat were the River Mole corridor and adjoining habitats, land south of the M23 spur road, the area of habitat west of the Fire Training Ground and the open areas around the Land East of the Railway Line woodland and biodiversity wetland, and the east of the site.

3.6.3 These were the areas that were chosen to be surveyed and where reptile refugia were placed. The locations of the reptile refugia are shown on Figures 3.6.1a – 3.6.1e.

### Survey Results

3.6.4 The results of the survey are provided in Annex 2 and shown on Figures 3.6.1a – 3.6.1e.

3.6.5 A peak count of ten grass snakes *Natrix natrix* was identified during a single site visit. Following the population class size assessment, this was considered to be a 'Good' sized population.

3.6.6 The majority of the grass snakes were recorded along the River Mole corridor with a few being recorded in the fields south east of the Land east of the Railway Line woodland.

3.6.7 No other reptile species were recorded.

## 3.7 Great Crested Newt Surveys

### Water Body Assessment

3.7.1 A total of 36 ponds were identified either within the Project site boundary or with connectivity to it during the Phase 1 Habitat Survey, a desk-based study of ordnance survey maps and aerial photography, and through identification during other protected species surveys. These are described in Table 21 of Annex 2.

3.7.2 Further waterbodies were identified within the Project site boundary, these included widespread networks of ditches through the car parks and airfield, roadside ditches and rivers. These waterbodies were not assessed for great crested newts in 2019.

3.7.3 Of the 36 ponds, 29 ponds were identified as requiring assessment for their potential to support great crested newt.

3.7.4 An additional nine ponds were identified within 250 metres of land identified for development and 22 ponds within 500 metres of land identified for development. These were outside the original Project site boundary and have not been assessed. Many of these are separated from development areas within the Project site boundary by busy A roads and rivers which are considered a barrier to GCN dispersal.

3.7.5 The locations of the ponds and waterbodies, the reference codes used to identify them, and the results of the survey are shown on Figures 3.7.1a - 3.7.1j.

### Habitat Suitability Index Results

3.7.6 Great Crested Newt Habitat Suitability Index (HSI) surveys were undertaken on 26 water bodies in 2019. The scores are shown in Annex 2.

3.7.7 Nine ponds surveyed had a 'poor' HSI score, three had a 'below average' score, seven had an 'average' score, five ponds had a 'good' score and two had an 'excellent' habitat suitability score.

3.7.8 Three ponds; Old Lagoon, New Lagoon and Pond M, were all ruled out during the field visit due to them being concrete lined, large man-made ponds that had no aquatic vegetation and heavily managed surrounding terrestrial habitat. Levels of water in these ponds varied significantly and being linked to the Crawley sewage treatment works New Lagoon and Old Lagoon, the quality of water was considered poor. No great crested newt surveys were therefore carried out on these ponds.

3.7.9 Six further ponds were not surveyed for HSI as there were access restrictions to the land that they were within.

3.7.10 One pond, AAA4, was a newly created wildlife pond and so had not developed sufficiently for an accurate HSI assessment.

3.7.11 Great Crested Newt Habitat Suitability Index (HSI) surveys were undertaken in 2021 on the further waterbodies identified within the Project site boundary, including the widespread networks of ditches through the car parks and airfield, roadside ditches and

rivers. Twelve ponds had a 'good' score or 'excellent' habitat suitability score. The scores are shown in Annex 2.

3.7.12 Five HSI surveys were carried out in 2022. Pond 1, pond 6 and additional pond right next to the M23 had 'excellent' scores and ponds 4 and 5 had a score of 'poor'. The full pond descriptions and results are shown in Annex 2.

### Presence/Absence Results

3.7.13 From the HSI scores ponds were chosen to be surveyed further because they had a score of 'average' or better. Pond AAA4 was included within the presence/absence surveys as it was created as a wildlife pond and was in close proximity to other ponds with 'excellent' HSI scores.

### eDNA Results

3.7.14 During 2019, eDNA surveys were conducted on Ponds 30Z, 8N8, A, AA21, FFJ and AVF. Only Pond 8N8 provided a positive result.

3.7.15 Due to the negative eDNA results, surveys on the other ponds were not continued.

3.7.16 Four ponds were found to either contain the eDNA of great crested newts or great crested newts were found during survey visits; Ponds 8N8, W46, K5F and TTD. Population size class surveys were undertaken on these ponds to determine the size of the great crested newt populations present. A summary of the results is provided in Table 3.7.1 below and the full results for all the ponds are provided in Annex 2.

3.7.17 Further eDNA surveys were undertaken in 2021 on the twelve ponds which had a 'good' score or 'excellent' habitat suitability score. eDNA surveys were carried out on ponds A1 Ditch 1; A2 Ditch 7; A2 Ditch 8; A2 Ditch 14; A5 Ditch 1; A5 Ditch 2; A5 Ditch 3; B2036 Ditch 3; Ditch, North Boundary; Car Park X Ditch; Southern Boundary Car Park X; North Long Stay A6/A7 Ditch 2; and North Long Stay A6/A7 Pond 1. In all cases the eDNA results were negative.

3.7.18 In 2022 eDNA surveys of a further 3 ponds (Ponds 4, 5 and 6) gave negative results for the presence of GCN eDNA. Although pond 1 had an 'excellent' HSI score the water was too shallow to collect for eDNA analysis.

3.7.19 Table 3.7.1: Great Crested Newt Population Size Class

Pond No.	Maximum Great Crested Newt Count	Great Crested Newt Population Size Class
8N8	0 (but positive eDNA), GCN eggs present	Small
W46	13	Medium
K5F	8, GCN eggs present	Small
TTD	10, GCN eggs present	Small

- 3.7.20 The maximum great crested newt count on one night using one survey method for each pond was zero for Pond 8N8, 13 for Pond W46, eight for Pond K5F and ten for Pond TTD.
- 3.7.21 Using the Great Crested Newt Population Size Class assessment (Froglife, 2001) this equates to a medium great crested newt population size for Pond W46 and small great crested newt population size for Ponds 8N8, K5F and TTD.
- 3.7.22 Two common toads were recorded on 3<sup>rd</sup> June 2019 within pond W46. One common toad was also recorded on 2<sup>nd</sup> October 2019 during a reptile survey, under an artificial refugia within the field east of the River Mole and south of Brockley Wood.

### 3.8 Dormouse Surveys

- 3.8.1 No evidence of dormouse was found within any of the surveyed areas. The survey areas are shown on Figure 3.8.1a and b.

### 3.9 Aquatic Mammal Surveys

#### Otter Surveys

- 3.9.1 No evidence of otter was found along the River Mole, within the surveyed area, as shown on Figure 3.9.1.

#### Water Vole Surveys

- 3.9.2 No evidence of water vole was found along the River Mole, within the surveyed area, as shown on Figure 3.9.1.

### 3.10 Preliminary Bat Roost Assessment

#### Buildings

- 3.10.1 Two buildings within the Project site boundary were identified as having features suitable to support roosting bats: one, a disused Control Tower (Building JW9) located in the north west of the Project site (landside), adjacent to Control Tower Road and east of the River Mole; and the second, a disused ancillary building (Building D9H) located along the southern boundary of the airside

perimeter fencing, adjacent to Crawler's Brook and Staff Car Park Z. The buildings surveyed are shown on Figure 3.10.1.

#### Trees

- 3.10.2 Thirty two trees along the A23 within the Project site boundary were identified as having features suitable to support roosting bats: eight trees with high bat roost potential, nineteen trees with moderate bat roost potential and five trees with low bat roost potential

### 3.11 Bat Emergence/Re-entry Surveys

- 3.11.1 As recommended by the BCT guidance, three dusk emergence surveys were undertaken on each of the two buildings identified within the Project site boundary as having bat roosting potential.
- 3.11.2 The surveys were undertaken to determine whether a bat roost was present and the species and number of bats using it.
- 3.11.3 A summary of the survey dates, weather conditions and sunset times is provided in Table 3.11.1 below.

**Table 3.11.1: Bat Emergence Survey Dates, Weather Conditions and Survey Times**

Building ref.	Date	Weather	Sunset time	Survey start	Survey end
D9H	15/07/19	22°C, light cloud, no rain	21:15	21:00	22:45
D9H	20/08/19	16°C, light winds, dry, fair	20:20	20:05	21:50
D9H	26/09/19	17°C, windy, clear skies	18:55	18:40	20:25
JW9	15/07/19	22°C, light cloud, no rain	21:15	21:00	22:45
JW9	07/08/19	18°C, cloudy, dry	20:41	20:26	22:11
JW9	02/10/19	13°C, dry, clear, light winds	18:37	18:22	20:07

#### Building JW9 (Landside)

- 3.11.4 No bats were seen emerging from the building but bats were detected foraging nearby at low levels during the emergence surveys on 15th July and 20th August 2019.
- 3.11.5 Bat activity was recorded at low levels during the emergence surveys on 15th July and 20th August 2019.

- 3.11.6 On the emergence survey of 26 September 2019, bat activity was recorded at moderate levels during the survey; although no bats were seen. It was presumed that bats were foraging near to the grassland area to the west of the building. Common pipistrelle, soprano pipistrelle, noctule, Leisler's bat and *Myotis* species were recorded.

- 3.11.7 Further details of the results of these surveys are provided in Annex 2.

#### Building D9H (Airside)

##### Bat Emergence Survey 15th July 2019

- 3.11.8 The bat emergence survey on 15th July commenced at 21:00, 15 minutes before sunset and finished at 22:45.
- 3.11.9 No bats were seen emerging from the building during any of the surveys. However bats were detected at low levels, foraging and commuting nearby. Common pipistrelle and noctule were recorded.
- 3.11.10 Further details of the results of these surveys are provided in Annex 2.

### 3.12 Bat Activity Transect Surveys

- 3.12.1 A summary of the survey dates, weather conditions and sunset times is provided in Table 3.9.1 in Annex 2.
- 3.12.2 For the surveys completed in 2019 (Transects 1-5), the results for each transect route have been grouped into pre-maternity (April and May, Figures 3.12.1a – 3.12.1f), maternity (June and July, Figures 3.12.2a – 3.12.2f) and post-maternity (August-October, 3.12.3a – 3.12.3f) seasons.
- 3.12.3 For surveys completed in 2020/21 and 2022, data are presented for the full seasons for each transect; Transect 6 Figure 3.12.4a-k; Transect 7 Figure 3.12.5a-k; Transect 8 Figures 3.12.6a-k; and Transect 9 Figures 3.12.7a-j.
- 3.12.4 The locations of the transects are shown on the above mentioned figures and are briefly described below:

- Transect 1: Horleyland Wood, Upper Pickets Wood and Lower Pickets Wood;
- Transect 2: Gatwick BAP Area, Land East of the Railway Line (LERL);
- Transect 3: Riverside Garden Park and Perimeter Road East;

- Transect 4: Perimeter Road South;
  - Transect 5: Museum Field and other land west of the River Mole.
  - Transect 6: Brockley Wood
  - Transect 7: Riverside Garden Park
  - Transect 8: Industrial Area, Airport Way roundabout
  - Transect 9: Dairy Farm and Church Meadows
- 3.12.5 Overall, moderate levels of bat activity were recorded across all transects during the pre-maternity, maternity and post-maternity seasons, except for Transect 4 which consistently recorded very low levels of activity.
- 3.12.6 The highest number of bat passes recorded in the pre-maternity season was along Transect 3, with 286 passes. In the maternity and post-maternity seasons, the highest number of passes was recorded along Transect 1, with 400 and 508 passes respectively. The fewest number of passes across all seasons was recorded along Transect 4 (24, 23 and 52 passes respectively).
- 3.12.7 Across Transects 1 and 5, the overall levels of bat activity were considerably higher in the maternity season, compared to the pre-maternity season, whereas activity levels across Transect 3 were considerably lower. The activity levels along Transects 2 and 4 remained constant across both seasons.
- 3.12.8 In the maternity season, significantly higher levels of bat activity were recorded along Transect 5, adjacent to the River Mole corridor and woodland strip, which are well connected with Brockley Wood.
- 3.12.9 Generally, high levels of bat activity were recorded within the woodland areas associated with Transects 1, 2 and 3, including Horleyland Wood and Upper Pickett's Wood, adjacent to and north of the sewage treatment works and woodland associated with Riverside Garden Park, in the north east of the Project site.
- 3.12.10 Higher levels of commuting activity were also recorded along linear features, notably the railway line adjacent to Transect 2, mature hedgerow and tree lines, and the river corridors, including the River Mole, Man's Brook, Crawter's Brook and Gatwick Stream.
- 3.12.11 Foraging activity was generally concentrated along mature hedgerows, through open canopy areas within woodland, woodland edges and adjacent/close to waterbodies, including the lake within Riverside Garden Park and the Crawley Sewage Treatment Works.

- 3.12.12 Lower levels of bat activity were observed in areas of open pasture, such as those associated with Transect 5 and habitat that comprised large, exposed areas of hardstanding with little canopy cover, such as those found along Transect 4.
- 3.12.13 Common pipistrelle *Pipistrellus pipistrellus* was the most frequently recorded species across all transect routes, with peak counts of 777, 1,005 and 1,232 passes recorded during the pre-maternity, maternity and post-maternity seasons respectively. Noctule *Nyctalus noctula* was also recorded in moderate numbers, with a peak count of 19 bat passes recorded along Transect 2 in the pre-maternity season. Lower numbers of soprano pipistrelle *Pipistrellus pygmaeus*, *Myotis sp.* and other big bat species (including serotine *Eptesicus serotinus* and Leisler's *Nyctalus leisleri* bats) were detected throughout the transect surveys. Single Nathusius' pipistrelle passes were recorded along Transect 2 in the maternity season and along Transect 3 in the post-maternity season.
- 3.12.14 Pipistrelle bats and *Myotis sp.* were generally associated with woodland areas and woodland edges, whereas noctule, serotine and Leisler's bat passes were more frequently recorded in open areas of grassland and pasture.

### 3.13 Bat Static/Automated Surveys

- 3.13.1 Within the Project site boundary, 16 static bat detectors were set out. The locations of these detectors are shown on Figure 3.13.1.

#### Location 1 – Land West of the Fire Training Ground

- 3.13.2 A summary of the survey dates, number of nights deployed, and bat passes for Location 1 is provided in Table 3.13.1 below. A summary of the number and species of bats recorded at Location 1 is provided in Table 3.13.2. Full details of passes per night are provided in Annex 2.

**Table 3.13.1: Bat Static/Automated Survey Summary for Location 1**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
24 April 2019 – 30 April 2019	7	59	8
10 May 2019 - 15 May 2019	6	566	94

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
11 June 2019 – 15 June 2019	5	189	38
12 July 2019 – 16 July 2019	5	745	149
13 Aug 2019 – 18 Aug 2019	6	282	47
25 Sept 2019 – 29 Sept 2019	5	357	71
14 Oct 2019 – 18 Oct 2019	5	138	28

**Table 3.13.2: Species Summary for Location 1**

Survey Month	Bb	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	0	51	0	1	0	3	1	3	0	0	59
May	0	532	19	4	2	0	2	7	0	0	566
June	0	177	4	0	0	0	0	8	0	0	189
July	0	555	8	1	6	14	3	151	5	2	745
August	0	222	3	3	0	17	3	31	0	3	282
Sept	1	34	0	0	3	7	0	312	0	0	357
Oct	1	103	3	0	8	7	0	16	0	0	138
<b>Species total</b>	<b>2</b>	<b>1,674</b>	<b>37</b>	<b>9</b>	<b>19</b>	<b>48</b>	<b>9</b>	<b>528</b>	<b>5</b>	<b>5</b>	<b>2,336</b>

Bb – barbastelle, Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - *Myotis* bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.3 A total of 2,336 bat passes were recorded at Location 1 between April and October 2019.
- 3.13.4 The highest number of passes for all species was recorded in July (745 passes) during the maternity season. The fewest number of passes was recorded in April (59 passes) during the pre-maternity season.
- 3.13.5 On average, higher numbers of bat passes were recorded during the maternity season (467 passes) than in the pre- and post-maternity seasons (313 and 259 passes respectively) at Location 1.
- 3.13.6 Common pipistrelle was the most frequently recorded species at this location across all seasons accounting for 72% of the overall bat passes. Peak counts of 532 and 555 common pipistrelle passes were recorded in May and July respectively. The overall

number of passes at this location was comparatively fewer in April and June. Fewer numbers of soprano pipistrelle were recorded at Location 1 and these accounted for between 1 and 3% of the species passes at this location.

- 3.13.7 Nathusius' pipistrelle *Pipistrellus nathusiusii* was recorded in very low numbers throughout the year, with a peak count of eight passes, recorded in October.
- 3.13.8 Noctule accounted for 11% of the species composition at Location 1; a peak count of 312 noctule passes was recorded in September.
- 3.13.9 Moderate levels of *Myotis* sp. Bats were recorded throughout the season, with the highest counts recorded in July (14 passes) and August (17 passes).
- 3.13.10 Lower level of activity for long-eared *Plecotus* sp., serotine and Leisler's bats were recorded at Location 1, with overall counts of nine, five and five passes respectively. Collectively, these species accounted for less than 1% of the overall species composition.
- 3.13.11 Barbastelle *Barbastella barbastellus* was recorded twice during the post-maternity season, with a single pass in September and October.

### Location 2 – Land South West of the River Mole

- 3.13.12 A summary of the survey dates, number of nights deployed, and bat passes for Location 2 is provided in Table 3.13.3 below. A summary of the number and species of bats recorded at Location 2 is provided in Table 3.13.4. Full details of passes per night are provided in Annex 2.

**Table 3.13.3: Bat Static/Automated Survey Summary for Location 2**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
24 April 2019 – 30 April 2019	7	110	15
10 May 2019 - 15 May 2019	6	1,101	184
12 June 2019 – 16 June 2019	5	730	146

12 July 2019 – 16 July 2019	5	1,269	254
13 Aug 2019 – 18 Aug 2019	6	330	55
25 Sept 2019 – 27 Sept 2019	3	291	97
14 Oct 2019 – 18 Oct 2019	5	35	7

**Table 3.13.4: Species Summary for Location 2**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	96	5	0	0	8	0	1	0	0	110
May	948	48	0	0	7	1	97	0	0	1,101
June	66	5	0	3	13	1	642	0	0	730
July	1,183	20	0	1	18	2	41	0	4	1,269
August	149	15	0	39	53	16	69	13	12	330
Sept	42	1	0	1	7	2	238	0	0	291
October	24	5	0	0	5	0	0	1	0	35
<b>Species total</b>	<b>2,508</b>	<b>99</b>	<b>0</b>	<b>44</b>	<b>111</b>	<b>22</b>	<b>1,088</b>	<b>14</b>	<b>16</b>	<b>3,866</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.13 A total of 3,866 bat passes was recorded at Location 2 between April and October 2019.
- 3.13.14 The highest number of passes for all species was recorded in July (1,269 passes) during the maternity season. The fewest number of passes was recorded in October (35 passes) during the post-maternity season.
- 3.13.15 On average, higher numbers of bat passes were recorded during the maternity season (1,000 passes) than in the pre- and post-maternity seasons (606 and 219 passes, respectively) at Location 2.
- 3.13.16 Overall, common pipistrelle was the most frequently recorded species at this location with a total of 2,508 passes and a peak count of 1,183 passes recorded in July. The total number of passes of common pipistrelle showed a marked difference between the pre-maternity/maternity and post-maternity seasons.
- 3.13.17 In April, May and July, common pipistrelles accounted for between 86% and 93% of the total species composition at Location 2. However, in June, only 9% (66 passes) were from

common pipistrelles. In the post-maternity season, the number of common pipistrelle passes averaged 33%.

- 3.13.18 Fewer numbers of soprano pipistrelle were recorded at this location which accounted for between 1% and 14% of the overall species composition at this location. A peak count of 48 passes was recorded in May.
- 3.13.19 Noctule accounted for 28% (1,088 passes) of the overall bat assemblage at Location 2 with peak counts of 642 and 238 passes in June and September respectively.
- 3.13.20 Rarer species including serotine and Leisler's bats were also recorded at this location. The peak count for both species was in August with 12 and 13 passes respectively.
- 3.13.21 Low numbers of long-eared bat species were recorded across all seasons, with a peak count of 16 passes recorded in August.
- 3.13.22 Nathusius' pipistrelle was not recorded at Location 2.

### Location 3 – Brockley Wood

- 3.13.23 A summary of the survey dates, number of nights deployed, and bat passes for Location 3 is provided in Table 3.13.5 below. A summary of the number and species of bats recorded at Location 3 is provided in Table 3.13.6. Full details of passes per night are provided in Annex 2.

**Table 3.13.5: Bat Static/Automated Survey Summary for Location 3**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
25 April 2019 – 1 May 2019	7	2,410	344
10 May 2019 – 14 May 2019	5	19,553	3,911
12 June 2019 – 16 June 2019	5	2,358	472
12 July 2019 – 16 July 2019	5	9,914	1,983
13 Aug 2019 – 18 Aug 2019	6	4,330	722
25 Sept 2019 – 29 Sept 2019	5	1,393	279
14 Oct 2019 – 18 Oct 2019	5	1,787	357

**Table 3.13.6: Species Summary for Location 3**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	2061	228	0	1	117	0	3	0	0	2410
May	15612	529	0	3	3102	234	68	5	0	19553
June	1302	268	0	1	639	4	109	0	0	2323
July	7688	455	1	5	1728	3	34	0	0	9914
August	2339	904	0	535	541	3	6	0	2	4330
Sept	333	83	0	670	145	0	161	0	1	1393
October	455	268	0	53	1005	5	1	0	0	1787
<b>Species total</b>	<b>29790</b>	<b>2735</b>	<b>1</b>	<b>1268</b>	<b>7277</b>	<b>249</b>	<b>382</b>	<b>5</b>	<b>3</b>	<b>41710</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.24 A total of 41,710 bat passes was recorded at Location 3 between April and October 2019, which is the highest number of passes recorded across all static detector locations.
- 3.13.25 The highest number of passes for all species was recorded in May during the pre-maternity season, with a total of 19,553 passes. The fewest number of passes was recorded in September during the post-maternity season, with 1,393 passes.
- 3.13.26 On average, higher numbers of bat passes were recorded during the pre-maternity season (10,982 passes) than in the maternity and post-maternity seasons (6,119 and 2,503 passes respectively) at Location 3.
- 3.13.27 Very high levels of common pipistrelle activity and high levels of soprano pipistrelle and *Myotis* sp. activity were recorded across all seasons, with the highest proportion of common pipistrelle passes recorded in May (80%), for soprano pipistrelles in August (21%) and for *Myotis* sp. in October (56%).
- 3.13.28 A single pass from Nathusius' pipistrelle was recorded in July.
- 3.13.29 Moderate levels of noctule activity were recorded at Location 3, with a peak count of 161 passes in September. Leisler's bat and serotine were recorded in lower numbers with a total of five and three passes respectively. Collectively, these species accounted for less than 1% of the overall composition at Location 3.
- 3.13.30 Low levels of *Plecotus* sp. passes were recorded across all months, with the exception of May, when a total of 234 passes were recorded.

**Location 4 – North of the Long Stay North Car Park**

- 3.13.31 A summary of the survey dates, number of nights deployed, and bat passes for Location 4 is provided in Table 3.13.7 below. A summary of the number and species of bats recorded at Location 4 is provided in
- 3.13.32 Table 3.13.8. Full details of passes per night are provided in Annex 2.

**Table 3.13.7: Bat Static/Automated Survey Summary for Location 4**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
25 April 2019 – 30 April 2019	6	3,093	516
10 May 2019 – 15 May 2019	6	3,781	630
12 June 2019 – 15 June 2019	4	141	35
12 July 2019 – 16 July 2019	5	470	94
13 Aug 2019 – 18 Aug 2019	6	520	87
25 Sept 2019 – 27 Sept 2019	3	123	41
14 Oct 2019 – 18 Oct 2019	5	53	11

**Table 3.13.8: Species Summary for Location 4**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	2795	147	5	1	96	39	6	1	3	3093
May	2405	1310	3	29	31	0	1	2	0	3781
June	99	2	1	0	9	4	26	0	0	141
July	299	23	2	5	78	18	36	0	9	470
August	385	13	0	6	67	3	32	10	4	520
Sept	38	4	0	0	12	6	63	0	0	123
October	16	2	0	0	15	0	20	0	0	53

Species total	6037	1501	11	41	308	70	184	13	16	8181
Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat										

- 3.13.33 A total of 8,181 bat passes was recorded at Location 4 between April and October 2019.
- 3.13.34 The highest number of passes was recorded in May (3,871 passes) during the pre-maternity season. Only 141 passes were recorded in June and 53 passes were recorded in October.
- 3.13.35 On average, higher numbers of bat passes were recorded during the pre-maternity season (3,437 passes) than in the maternity and post-maternity seasons (306 and 232 passes respectively) at Location 4.
- 3.13.36 Common pipistrelle was the most frequently recorded species at this location with a total of 6,037 passes and a peak count of 2,795 passes recorded in May. Across all months, common pipistrelle accounted for between 63 and 90% of the total species composition at this location.
- 3.13.37 High numbers of both common and soprano pipistrelle were recorded in May with 2,405 and 1,310 passes respectively. Moderate numbers of *Myotis* and *Plecotus* sp. bats were recorded across all months, with peak counts of 96 and 39 passes recorded in April.
- 3.13.38 On average moderate numbers of noctule were recorded during the maternity and post-maternity seasons (31 and 38 passes, respectively) compared to the pre-maternity season (4 passes).
- 3.13.39 Low levels of activity were also recorded for Leisler's and serotine bats with a total of 13 and 16 passes respectively.
- 3.13.40 Low levels of activity were also recorded for Nathusius' pipistrelle with a total count of 11 passes and a peak count of five passes in April.

**Location 5 – Riverside Garden Park**

- 3.13.41 Location 5 in Riverside Garden Park is outwith the Project site boundary. A summary of the survey dates, number of nights deployed, and bat passes for Location 5 is provided in Table 3.13.9 below. A summary of the number and species of bats recorded at Location 5 is provided in Table 3.13.10. Full details of passes per night are provided in Annex 2.

**Table 3.13.9: Bat Static/Automated Survey Summary for Location 5**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
0 May 2019 – 15 May 2019	6	3694	616
12 July 2019 – 16 July 2019	5	3321	664
13 Aug 2019 – 18 Aug 2019	6	564	94
25 Sept 2019 – 29 Sept 2019	5	305	61
14 Oct 2019 – 18 Oct 2019	5	68	14

**Table 3.13.10: Species Summary for Location 5**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	0	0	0	0	0	0	0	0	0	0
May	3461	35	0	0	6	7	1	0	0	3694
June	0	0	0	0	0	0	0	0	0	0
July	3060	144	0	2	8	0	16	0	91	3321
August	462	31	0	4	44	3	17	1	2	564
Sept	168	28	0	79	11	4	15	0	0	305
October	47	6	0	6	5	4	0	0	0	68
<b>Species total</b>	<b>7198</b>	<b>244</b>	<b>0</b>	<b>91</b>	<b>74</b>	<b>18</b>	<b>49</b>	<b>1</b>	<b>93</b>	<b>7952</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.42 A total of 7,952 bat passes was recorded at Location 5 during May and between July and October 2019. No data was recorded during April or June due to equipment failure and malfunction.
- 3.13.43 The highest number of passes was recorded in May (3,694 passes) during the pre-maternity season. The lowest number of passes was recorded in October (68 passes) during the post-maternity season.
- 3.13.44 On average, higher numbers of bat passes were recorded during the pre-maternity season (3,694 passes) than in the maternity and post-maternity seasons (3,321 and 312 passes respectively) at Location 5.
- 3.13.45 Common pipistrelles accounted for the highest number of species passes at this location, with between 55% and 94% of the species composition across all seasons at Location 5.
- 3.13.46 Low numbers of soprano pipistrelles were recorded at Location 5, with a total of 244 passes (3% of total passes).
- 3.13.47 Moderate to low numbers of *Myotis* and *Plecotus* sp., serotine and noctule were also recorded at this location with a total of 74, 18, 93 and 49 respectively. A peak count of 91 serotine bat passes was recorded in July.
- 3.13.48 Nathusius' pipistrelle was not recorded at Location 5.

**Location 6 – Land West of the Railway**

- 3.13.49 A summary of the survey dates, number of nights deployed, and bat passes for Location 6 is provided in Table 3.13.11 below. A

summary of the number and species of bats recorded at Location 6 is provided in Table 3.13.12. Full details of passes per night are provided in Annex 2.

**Table 3.13.11: Bat Static/Automated Survey Summary for Location 6**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
25 April 2019 – 27 April 2019	3	269	89
10 May 2019 – 12 May 2019	3	5,093	1,698
12 June 2019 – 16 June 2019	5	7,876	1,575
12 July 2019 – 16 July 2019	5	4,691	938
13 Aug 2019 – 18 Aug 2019	6	7,897	1,316
24 Sept 2019 – 28 Sept 2019	5	2,920	584
14 Oct 2019 – 19 Oct 2019	6	379	63

**Table 3.13.12: Species Summary for Location 6**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	266	2	1	0	0	0	0	0	0	269
May	4839	223	2	0	26	1	2	0	0	5093
June	7754	13	51	1	8	2	46	0	1	7876
July	4583	18	9	0	15	4	60	0	2	4691
August	7772	5	0	0	19	2	96	2	1	7897
Sept	2872	21	0	2	6	0	19	0	0	2920
October	346	29	0	0	2	0	2	0	0	379
<b>Species total</b>	<b>28432</b>	<b>31</b>	<b>63</b>	<b>3</b>	<b>76</b>	<b>9</b>	<b>225</b>	<b>2</b>	<b>4</b>	<b>28845</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.50 A total of 28,845 bat passes was recorded at Location 6 between April and October 2019.

- 3.13.51 The highest number of passes was recorded in June (7,876 passes) during the maternity season. The lowest number of passes was recorded in April (269 passes) during the pre-maternity season.
- 3.13.52 On average, higher numbers of bat passes were recorded during the maternity season (6,284 passes) than in the pre-maternity and post-maternity seasons (2,681 and 3,732 passes, respectively) at Location 6.
- 3.13.53 Common pipistrelle was the most frequently recorded species at this location with a total of 28,432 passes across all months and a peak count of 7,772 passes recorded in August. Across all months, common pipistrelle accounted for between 95% and 99% of the total species composition at this location.
- 3.13.54 Generally low numbers of soprano pipistrelle were recorded across all months, except for May, when 223 passes were recorded. Noctule was also recorded in moderate numbers with peak counts of 60 and 96 passes in July and August, respectively.
- 3.13.55 Moderate numbers of Nathusius' pipistrelle and *Myotis* sp. were recorded at Location 6 with a total of 63 and 76 passes respectively. In June, a peak count of 51 Nathusius' pipistrelle passes was recorded.
- 3.13.56 Low activity levels were recorded for *Plecotus* sp., Leisler's and serotine bats with nine, two and four passes respectively.

**Location 7 – Horleyland Wood**

- 3.13.57 A summary of the survey dates, number of nights deployed, and bat passes for Location 7 is provided in Table 3.13.13 below. A summary of the number and species of bats recorded at Location 7 is provided in Table 3.13.14. Full details of passes per night are provided in Annex 2.

**Table 3.13.13: Bat Static/Automated Survey Summary for Location 7**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
25 April 2019 – 30 April 2019	6	8,459	134
10 May 2019 – 13 May 2019	4	12,878	3,220
12 June 2019 – 16 June 2019	5	8,221	1,644
12 July 2019 – 15 July 2019	4	5,250	1,313

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
13 Aug 2019 – 18 Aug 2019	6	2,421	404
25 Sept 2019 – 27 Sept 2019	3	250	83
15 Oct 2019 – 20 Oct 2019	6	488	81

**Table 3.13.14: Species Summary for Location 7**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	8021	337	0	0	98	3	0	0	0	8459
May	12570	290	1	0	11	4	2	0	0	12878
June	7883	250	0	0	61	7	20	0	0	8221
July	5104	38	0	5	12	19	8	0	64	5250
August	2154	27	0	72	116	16	0	0	25	2421
Sept	148	2	0	84	1	5	10	0	0	250
October	436	42	0	2	7	0	1	0	0	488
<b>Species total</b>	<b>36316</b>	<b>986</b>	<b>1</b>	<b>163</b>	<b>306</b>	<b>54</b>	<b>41</b>	<b>0</b>	<b>89</b>	<b>37967</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.58 A total of 37,967 bat passes was recorded at Location 7 between April and October 2019.
- 3.13.59 The highest number of passes was recorded in May (12,878 passes) during the pre-maternity season. The lowest number of passes was recorded in September (250 passes), during the post-maternity season.
- 3.13.60 On average, higher numbers of bat passes were recorded during the pre-maternity season (10,669 passes) than in the maternity and post-maternity seasons (6,736 and 1,053 passes respectively) at Location 7.
- 3.13.61 Common pipistrelle was the most frequently recorded species at this location with a total of 36,316 passes and a peak count of 12,570 passes recorded in May. Across all seasons, common pipistrelle accounted for between 59% and 98% of the species composition at this location.
- 3.13.62 High activity levels of soprano pipistrelle were recorded in the pre-maternity season, which average 314 passes, compared to

the maternity and post-maternity seasons, which averaged 144 and 24 passes respectively.

- 3.13.63 A single Nathusius' pipistrelle pass was recorded in May.
- 3.13.64 Activity levels of *Myotis* sp. bats remained relatively low throughout the season with a peak count of 116 passes in August and 98 passes in April. Only one *Myotis* sp. pass was recorded in September.

- 3.13.65 Moderate levels of activity were recorded for all other species at this location including for *Plecotus* sp. (54 passes), serotine (89 passes) and noctule (41 passes). Leisler's bat was not recorded during surveys at this location.

### Location 8 – Land East of the Railway Line Wetland

- 3.13.66 A summary of the survey dates, number of nights deployed, and bat passes for Location 8 is provided in Table 3.13.15 below. A summary of the number and species of bats recorded at Location 8 is provided in

- 3.13.67 Table 3.13.16. Full details of passes per night are provided in Annex 2.

**Table 3.13.15: Bat Static/Automated Survey Summary for Location 8**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
24 April 2019 – 1 May 2019	8	1,758	219
12 May 2019 – 15 May 2019	4	2,121	530
11 July 2019 – 16 July 2019	6	203	34
14 Aug 2019 – 18 Aug 2019	5	14	3
25 Sept 2019 – 29 Sept 2019	5	1,775	355
14 Oct 2019 – 19 Oct 2019	6	889	148

**Table 3.13.16: Species Summary for Location 8**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	1,728	2	1	7	0	0	20	0	0	1,758
May	2,118	3	0	0	0	0	0	0	0	2,121
June	0	0	0	0	0	0	0	0	0	0

July	37	0	0	0	0	0	164	0	0	203
August	14	0	0	0	0	0	0	0	0	14
Sept	679	19	9	43	1	0	1,015	8	1	1,775
October	793	24	22	3	8	0	38	1	0	889
<b>Species total</b>	<b>5,369</b>	<b>48</b>	<b>32</b>	<b>53</b>	<b>9</b>	<b>0</b>	<b>1,237</b>	<b>9</b>	<b>1</b>	<b>6,760</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.68 A total of 6,760 bat passes was recorded at Location 8 between April and May and between July and October 2019. No data was recorded during June due to equipment failure and malfunction.

- 3.13.69 The highest number of passes was recorded in May (2,121 passes) during the pre-maternity season. The lowest number of passes was recorded in August (14 passes) during the post-maternity season.

- 3.13.70 On average, higher numbers of bat passes were recorded during the pre-maternity season (1,940 passes) than in the maternity and post-maternity seasons (203 and 893 passes respectively). In general, the number of bat passes at this location was comparatively lower than at the others.

- 3.13.71 Common pipistrelle was the most frequently recorded species at this location with a total of 5,369 passes and a peak count of 2,118 passes recorded in May.

- 3.13.72 Similar numbers of soprano pipistrelle and Nathusius' pipistrelle were recorded at Location 8, with similar numbers of bats recorded in September (19 and nine passes, respectively) and October (24 and 22 passes, respectively).

- 3.13.73 In July and September, noctule was more frequently recorded than any other species, accounting for 81% and 57% of the species composition at this location.

- 3.13.74 There were no recorded passes from *Plecotus* sp. bats and very few passes from serotine and Leisler's bats (one and nine passes, respectively).

### Location 9 – Perimeter Road South

- 3.13.75 A summary of the survey dates, number of nights deployed, and bat passes for Location 9 is provided in Table 3.13.17 below. A summary of the number and species of bats recorded at Location 9 is provided in Table 3.13.18. Full details of passes per night are provided in Annex 2.



**Table 3.13.17: Bat Static/Automated Survey Summary for Location 9**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
25 April 2019 – 1 May 2019	7	22	3
10 May 2019 – 15 May 2019	6	2,089	348
11 June 2019 – 16 June 2019	6	2,828	471
12 July 2019 – 16 July 2019	5	259	52
13 <sup>th</sup> Aug 2019 – 18 <sup>th</sup> Aug 2019	6	108	18
25 <sup>th</sup> Sept 2019 – 29 <sup>th</sup> Sept 2019	5	132	26
15 <sup>th</sup> Oct 2019 – 16 <sup>th</sup> Oct 2019	2	3	2

**Table 3.13.18: Species Summary for Location 9**

Survey Month	Bb	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	0	22	0	0	0	0	0	0	0	0	22
May	0	2,086	1	0	0	2	0	0	0	0	2,089
June	1	2,794	19	9	0	2	0	3	0	0	2,828
July	0	238	2	0	0	0	0	19	0	0	259
August	0	104	1	0	0	3	0	0	0	0	108
Sept	0	126	0	0	6	0	0	0	0	0	132
October	0	2	0	0	0	1	0	0	0	0	3
<b>Species total</b>	<b>1</b>	<b>5,372</b>	<b>23</b>	<b>9</b>	<b>6</b>	<b>8</b>	<b>0</b>	<b>22</b>	<b>0</b>	<b>0</b>	<b>5,441</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.76 A total of 5,441 bat passes was recorded at Location 9 between April and October 2019.
- 3.13.77 The highest number of passes was recorded in June (2,828 passes) during the maternity season. The lowest number of passes was recorded in October (three passes), during the post-maternity season.
- 3.13.78 On average, higher numbers of bat passes were recorded during the maternity season (1,544 passes) than in the pre-maternity and post-maternity seasons (1056 and 81 passes respectively).
- 3.13.79 Common pipistrelle was the most frequently recorded species at this location with a total of 5,372 passes across all months and a peak count of 2,794 passes recorded in June.

3.13.80 The species diversity across all other months was generally quite low with low numbers of soprano pipistrelle (23 passes), Nathusius' pipistrelle (nine passes), *Myotis* sp. (eight passes) and noctule (22 passes) recorded.

- 3.13.81 A single barbastelle pass was recorded at this location in June.
- 3.13.82 Neither Leisler's bat nor serotine were recorded at Location 9.

**Location 10 – Land West of Car Park X**

3.13.83 A summary of the survey dates, number of nights deployed, and bat passes for Location 10 is provided in Table 3.13.19 below. A summary of the number and species of bats recorded at Location 10 is provided in Table 3.13.20. Full details of passes per night are provided in Annex 2.

**Table 3.13.19: Bat Static/Automated survey summary for Location 10**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
10 May 2019 – 15 May 2019	6	2,646	441
12 July 2019 – 16 July 2019	5	2,823	564
13 Aug 2019 – 15 Aug 2019	3	1,407	469
25 Sept 2019 – 29 Sept 2019	5	698	140
14 Oct 2019 – 18 Oct 2019	5	99	20

**Table 3.13.20: Species Summary for Location 10**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	0	0	0	0	0	0	0	0	0	0
May	2,293	345	0	4	3	0	1	0	0	2,646
June	0	0	0	0	0	0	0	0	0	0
July	2,656	136	0	1	6	1	23	0	0	2,823
August	1,227	125	0	12	19	7	12	2	3	1,407
Sept	491	74	1	2	9	2	117	2	0	698
October	78	4	0	0	5	0	11	1	0	99
<b>Species total</b>	<b>6,745</b>	<b>684</b>	<b>1</b>	<b>19</b>	<b>42</b>	<b>10</b>	<b>164</b>	<b>5</b>	<b>3</b>	<b>7,673</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

3.13.84 A total of 7,673 bat passes was recorded at Location 10 during May and between July and October 2019. No data was recorded during April or June due to equipment failure and malfunction.

- 3.13.85 The highest number of passes was recorded in July (2,823 passes) during the maternity season. The lowest number of passes was recorded in October (99 passes) during the post-maternity season.

3.13.86 Common pipistrelle was the most frequently recorded species at this location with 6,745 passes in total and accounting for between 70% and 94% of the species composition across each month. Low to moderate numbers of soprano pipistrelle were recorded throughout the survey season with a peak count of 345 passes recorded in May.

3.13.87 A single Nathusius' pipistrelle pass was recorded in September.

3.13.88 Low numbers of *Myotis* sp. and noctule were recorded at Location 10, with peak counts of 19 *Myotis* sp. passes in August and 117 noctule passes in September.

**Location 11 – Crawter's Wood**

3.13.89 A summary of the survey dates, number of nights deployed, and bat passes for Location 11 is provided in Table 3.13.21 below. A summary of the number and species of bats recorded at Location 11 is provided in Table 3.13.22. Full details of passes per night are provided in Annex 2.

**Table 3.13.21: Bat Static/Automated Survey Summary for Location 11**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
24 April 2019 – 1 May 2019	8	2,037	255
11 May 2019 – 15 May 2019	5	60	12
13 June 2019 – 16 June 2019	4	945	236
12 July 2019 – 16 July 2019	5	4,538	908
13 Aug 2019 – 17 Aug 2019	5	1,290	258
25 Sept 2019 – 27 Sept 2019	3	3,745	1,248
14 Oct 2019 – 19 Oct 2019	6	1546	258

**Table 3.13.22: Species Summary for Location 11**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
April	2,011	7	2	5	4	2	6	0	0	2,037
May	60	0	0	0	0	0	0	0	0	60
June	928	0	0	0	0	0	17	0	0	945
July	4,361	3	7	11	121	2	33	0	0	4,538
August	1,210	1	0	37	13	1	27	0	1	1,290
Sept	2,895	4	0	246	58	3	539	0	0	3,745
October	1,456	9	0	2	59	0	20	0	0	1,546
<b>Species total</b>	<b>12,921</b>	<b>24</b>	<b>9</b>	<b>301</b>	<b>255</b>	<b>8</b>	<b>642</b>	<b>0</b>	<b>1</b>	<b>14,161</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

3.13.90 A total of 14,161 bat passes was recorded at Location 11 during between April and October 2019.

3.13.91 The highest number of passes was recorded in July (4,538 passes) during the maternity season. The lowest number of passes was recorded in May (60 passes), during the pre-maternity season, all of which were from common pipistrelle.

3.13.92 Overall, common pipistrelle was the most frequently recorded species at this location and accounted for 91% of the species assemblage (12,921 passes).

3.13.93 Moderate to high levels of activity from *Myotis* sp. and noctule were recorded across all months, with a peak count of 121 *Myotis* sp. passes in July and 539 noctule passes in September. Low numbers of all other bat species were recorded including soprano pipistrelle (24 passes), Nathusius' pipistrelle (nine passes), *Plecotus* sp. (eight passes) and serotine (one pass).

**Location 12 – River Mole (Bund)**

3.13.94 A summary of the survey dates, number of nights deployed, and bat passes for Location 11 is provided in Table 3.13.19 below. A summary of the number and species of bats recorded at Location 10 is provided in Table 3.13.20. Full details of passes per night are provided in Annex 2.

**Table 3.13.23: Bat Static/Automated survey summary for Location 12**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
28 May 2022 – 02 June 2022	6	118	19
27 July 2022 – 31 July 2022	5	336	67
18 Aug 2022 – 23 Aug 2022	6	16,336	2723

**Table 3.13.24: Species Summary for Location 12**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
May	88	3	0	0	2	0	25	0	0	118
July	227	7	0	0	17	0	73	3	9	336
August	13,014	3,242	0	0	45	1	3	0	31	16,336
<b>Species total</b>	<b>13,329</b>	<b>3,252</b>	<b>0</b>	<b>0</b>	<b>64</b>	<b>1</b>	<b>101</b>	<b>3</b>	<b>40</b>	<b>16,790</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

3.13.95 A total of 16,790 bat passes was recorded at Location 12 during May and between July and August 2022.

3.13.96 The highest number of passes was recorded in August (16,336 passes) during the post-maternity season. The lowest number of passes was recorded in May (118 passes), during the pre-maternity season.

3.13.97 Common pipistrelle was the most frequently recorded species at this location with 13,329 passes in total and accounting for between 68% and 81.5% of the species composition across each month at this location.

3.13.98 Moderate numbers of soprano pipistrelle, noctule, *Myotis* sp. and serotine were recorded throughout the survey season with a peak count of 3,242 soprano pipistrelle passes recorded in August, a peak count of 73 noctule passes recorded in July, a peak count of 45 *Myotis* passes recorded in August and a peak count of 31 serotine passes recorded in August.

3.13.99 Low numbers of Leisler's bats and brown long-eared bats were also recorded with a peak count of 3 Leisler's bat passes recorded in July and 1 brown long-eared bat pass counted in August.

**Location 13 – Riverside Gardens**

3.13.100 A summary of the survey dates, number of nights deployed, and bat passes for Location 13 is provided in Table 3.13.19 below. A summary of the number and species of bats recorded at Location 13 is provided in Table 3.13.20. Full details of passes per night are provided in Annex 2.

**Table 3.13.25: Bat Static/Automated survey summary for Location 13**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
27 May 2022 – 02 June 2022	6	251	41
22 June 2022 – 27 June 2022	6	5,897	982
23 Aug 2022 – 27 Aug 2019	5	3,944	788
08 Sept 2022 – 12 Sept 2022	5	1,807	361
18 Oct 2019 – 24 Oct 2019	6	194	32

**Table 3.13.26: Species Summary for Location 13**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
May	241	2	0	0	0	0	5	3	0	251
June	5,744	104	0	0	10	0	37	1	1	5,897
August	3,819	57	0	8	6	0	40	3	11	3,944
Sept	1,668	85	1	0	3	2	43	4	1	1,807
October	173	14	0	4	0	0	3	0	0	194
<b>Species total</b>	<b>11,645</b>	<b>262</b>	<b>1</b>	<b>12</b>	<b>19</b>	<b>2</b>	<b>128</b>	<b>11</b>	<b>13</b>	<b>12,093</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

3.13.101 A total of 12,093 bat passes was recorded at Location 13 during May, June, August, September and October 2022.

3.13.102 The highest number of passes was recorded in June (5,897 passes) during the maternity season. The lowest number of passes was recorded in October (194 passes), during the post-maternity season.

- 3.13.103 Common pipistrelle was the most frequently recorded species at this location with 11,645 passes in total and accounting for between 90% and 97% of the species composition across each month at this location.
- 3.13.104 Low numbers of *Myotis sp.* and serotine were recorded throughout the survey season, with a peak count of 10 *Myotis sp.* passes in June and 13 serotine passes in August.
- 3.13.105 Moderate numbers of noctule were recorded throughout the survey season, with a peak count of 43 passes in September.

**Location 14 – Services (Surrey CC)**

- 3.13.106 A summary of the survey dates, number of nights deployed, and bat passes for Location 14 is provided in Table 3.13.19 below. A summary of the number and species of bats recorded at Location 14 is provided in Table 3.13.20. Full details of passes per night are provided in Annex 2.

**Table 3.13.27: Bat Static/Automated survey summary for Location 14**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
21 July 2022 – 30 July 2022	10	207	20
23 Aug 2022 – 27 Aug 2022	5	10,177	2.035
08 Sept 2022 – 14 Sept 2022	6	3,455	575
18 Oct 2022 – 23 Oct 2022	6	789	131

**Table 3.13.28: Species Summary for Location 14**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
July	161	4	0	0	0	0	42	0	0	207
August	9,806	275	1	39	4	0	52	0	0	10,177
Sept	2,945	494	1	0	6	0	6	1	2	3,455
October	767	12	0	0	10	0	0	0	0	789
<b>Species total</b>	<b>13,679</b>	<b>785</b>	<b>2</b>	<b>39</b>	<b>20</b>	<b>0</b>	<b>100</b>	<b>1</b>	<b>2</b>	<b>14,628</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.107 A total of 14,628 bat passes were recorded at Location 14 between July and October 2022.

- 3.13.108 The highest number of passes was recorded in August (10,177 passes) during the post maternity season. The lowest number of passes was recorded in July (207 passes), during the maternity season.
- 3.13.109 Common pipistrelle was the most frequently recorded species at this location with 13,679 passes in total and accounting for between 78% and 97% of the species composition across each month at this location.
- 3.13.110 Moderate numbers of noctule were recorded throughout the survey season, with a peak count of 52 passes in August.
- 3.13.111 Low numbers of *Myotis sp.*, serotine and Leisler's bat were recorded throughout the survey season, with a peak count of 10 *Myotis sp.* passes in October, a peak count of 2 serotine passes in September and a single Leisler's bat pass in September.

**Location 15 – Dairy Farm Location 1 (Farm)**

- 3.13.112 A summary of the survey dates, number of nights deployed, and bat passes for Location 15 is provided in Table 3.13.19 below. A summary of the number and species of bats recorded at Location 15 is provided in Table 3.13.20. Full details of passes per night are provided in Annex 2.

**Table 3.13.29: Bat Static/Automated survey summary for Location 15**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
18 May 2022 – 24 May 2022	7	6782	968
10 June 2022 – 19 June 2022	8	17045	2130
07 July 2022 – 10 July 2022	4	7907	1976
07 Aug 2022 – 11 Aug 2022	5	13598	2720
21 Sept 2022 – 25 Sept 2022	5	9,243	1,848
20 Oct 2022 – 27 Oct 2022	5	5,192	1,038

**Table 3.13.30: Species Summary for Location 15**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
May	5,022	1,703	0	0	3	1	0	0	53	6,782

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
June	10,433	6,556	12	0	21	1	1	9	12	17,045
July	6,543	1,321	0	0	20	1	7	3	11	7,906
August	6,535	6,991	0	0	33	1	19	4	15	13,598
Sept	4,255	4,855	0	0	83	34	14	0	2	9,243
October	3,016	2,150	0	0	25	0	1	0	0	5,192
<b>Species total</b>	<b>35,804</b>	<b>23,576</b>	<b>12</b>	<b>0</b>	<b>185</b>	<b>38</b>	<b>42</b>	<b>16</b>	<b>93</b>	<b>59,766</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.113 A total of 59,766 bat passes were recorded at Location 15 between May and October 2022.
- 3.13.114 The highest number of passes was recorded in June (17,045 passes) during the maternity season. The lowest number of passes was recorded in October (5,192 passes), during the post-maternity season.
- 3.13.115 Common pipistrelle was the most frequently recorded species at this location with 35,804 passes in total and accounting for between 46% and 83% of the species composition across each month.
- 3.13.116 Moderate numbers of *Myotis sp.* and serotine were recorded throughout the survey season, with a peak count of 83 *Myotis sp.* passes in September and 53 serotine passes in May.
- 3.13.117 Low numbers of noctule and brown long-eared bats were recorded throughout the survey season, with a peak count of 19 noctule passes in August and 34 brown long-eared bat passes in September.

**Location 16 – Dairy Farm Location 2 (Gate)**

- 3.13.118 A summary of the survey dates, number of nights deployed, and bat passes for Location 16 is provided in Table 3.13.19 below. A summary of the number and species of bats recorded at Location 16 is provided in Table 3.13.20. Full details of passes per night are provided in Annex 2.

**Table 3.13.31: Bat Static/Automated survey summary for Location 16**

Survey dates	Number of nights detector deployed	Total number of bat passes	Average bat passes / night
19 May 2022 – 22 May 2022	3	22	7
10 June 2022 – 15 June 2022	6	3,549	591
10 July 2022 – 15 July 2022	6	4,651	775
18 Aug 2022 – 23 Aug 2022	6	1,824	304
18 Oct 2022 – 23 Oct 2022	6	5,721	953

**Table 3.13.32: Species Summary for Location 16**

Survey Month	Pp	Ppy	Pn	Psp	Msp	PI	Nn	NI	Es	Total
May	22	0	0	0	0	0	0	0	0	22
June	3,297	243	0	0	5	0	2	1	1	3,549
July	3,192	1,425	0	0	2	0	12	5	15	4,651
August	1,441	375	0	4	2	0	1	0	1	1,824
October	4,321	1,386	0	0	13	0	0	0	1	5,721
<b>Species total</b>	<b>12,273</b>	<b>3,429</b>	<b>0</b>	<b>4</b>	<b>22</b>	<b>0</b>	<b>15</b>	<b>6</b>	<b>18</b>	<b>15,767</b>

Pp -common pipistrelle, Ppy - soprano pipistrelle, Pn - Nathusius' pipistrelle, Psp - pipistrelle bats, Msp - Myotis bats, PI - long-eared bats, Nn - noctule, NI - Leisler's bat, Es - serotine bat

- 3.13.119 A total of 15,767 bat passes was recorded at Location 16 between May and August, and October 2022.
- 3.13.120 The highest number of passes was recorded in October (5,721 passes) during the post maternity season. The lowest number of passes was recorded in May (99 passes), during the pre-maternity season.
- 3.13.121 Common pipistrelle was the most frequently recorded species at this location with 12,273 passes in total and accounting for between 68.5% and 100% of the species composition across each month at this location.
- 3.13.122 Low numbers of *Myotis sp.*, serotine, noctule and Leisler's were recorded throughout the survey season, with a peak count of 13 *Myotis sp.* passes in October, a peak count of 15 serotine passes in July, a peak count of 12 noctule passes in July, and 5 Leisler's bat passes in July.

### Bat Crossing Point Survey

3.13.123 A total of 2459 bat passes was recorded across both locations, 2437 of which were observed using the target features (ie passing within 5m distance of the feature). A breakdown of the total number of passes observed at each crossing point is provided in Table 3.13.23 below.

**Table 3.13.33: Total Bat Passes**

Crossing point	Number of survey visits	Number of passes observed	Number of passes observed using the feature
River Mole	3	1298	1278
Riverside Garden Park	3	1161	1159

### River Mole Crossing Point

3.13.124 A total of 1278 bat passes was observed using the feature, of which 220 (17%) were considered to be flying within the river corridor, 659 (52%) were considered to be flying directly above the river corridor and 399 (31%) were flying at a height above 5m from ground level. These data are presented for each species / species group in Table 3.13.24 below.

**Table 3.13.34 Breakdown of bat passes from River Mole**

Species	Number of passes	Number of passes using the feature	Number of passes within river corridor	Number of passes directly above river corridor	Number of passes at a height higher than 5m from ground level
Myotis species	26	26	1	23	2
Noctule	53	53	0	7	40

Species	Number of passes	Number of passes using the feature	Number of passes within river corridor	Number of passes directly above river corridor	Number of passes at a height higher than 5m from ground level
Brown long-eared bat	3	3	0	3	0
Common pipistrelle	1017	1003	213	573	217
Soprano pipistrelle	24	24	5	19	0
Pipistrelle species	5	0	0	0	0
Unknown	170	170	1	34	135

3.13.125 Bats of at least five species were observed using the feature including *Myotis sp.*, noctule, brown long-eared bat, common pipistrelle and soprano pipistrelle. No additional species were confirmed as present by sound analysis.

### Riverside Garden Park Crossing Point

3.13.126 A total of 1159 bat passes was observed using the feature, of which 216 (19%) were observed passing at an "unsafe height" and 943 (81%) were observed passing at a safe height. These data are presented per species / species group in Table 3.13.25 below.

**Table 3.13.35: Breakdown of bat passes from Riverside Garden Park**

Species	Number of passes	Number of passes using the feature	Number of passes at height below or equal to 5m height	% of passes at height below or equal to 5m height	Number of passes at height above 5m	% of passes at height above 5m
<b>Myotis species</b>	18	18	5	27.78	13	72.22
<b>Brown long-eared bat</b>	3	3	1	33.33	2	66.67
<b>Common pipistrelle</b>	655	653	142	21.75	511	78.25
<b>Soprano pipistrelle</b>	431	430	51	11.86	379	88.14
<b>Pipistrelle species</b>	10	10	7	70	3	30
<b>Unknown</b>	45	45	10	22.22	35	77.78

3.13.127 Bats of at least four species / species groups were observed using the feature including *Myotis sp.*, brown long-eared bat, common pipistrelle and soprano pipistrelle. An additional species, noctule, was confirmed by sound analysis as present, although not observed.

#### Invertebrate Scoping Survey

3.13.128 The results of the invertebrate scoping survey are provided in Annex 4.

#### Terrestrial Invertebrate Survey

3.13.129 The results of the terrestrial invertebrate survey are provided in Annex 5.

#### Aquatic Invertebrate Survey

3.13.130 The results of the aquatic invertebrate survey are provided in Annex 6.

#### Fish Survey

3.13.131 The results of the fish survey are provided in Annex 6.

#### Veteran Tree Surveys

3.13.132 No veteran trees were found within the survey area. The results are in **ES Appendix 8.8.1: Outline Landscape and Ecology Management Plan** (Doc Ref. 5.3) along with the other tree survey data.

## 4 Evaluation

### 4.1 Breeding Bird Surveys

4.1.1 Seventeen of the 51 species recorded during the survey qualify as being of 'conservation interest' by meeting one or more of the criteria listed in Annex 2. The following accounts relate to those species confirmed as breeding, or considered to be possibly breeding, within the survey area in 2019 that are included on one or more of the lists of species either afforded special statutory protection or denoting a species of high conservation importance.

4.1.2 The conservation status used for the analysis of the breeding bird survey is based on the criteria when the survey was undertaken and the data analysed in 2019 (BoCC 2015).

#### Specially Protected Species

4.1.3 Although no Annex 1 or Schedule 1 species were confirmed to be breeding within the survey area, three species (little ringed plover, peregrine and firecrest) were recorded within the Project site boundary and could possibly have bred.

4.1.4 Little ringed plover - one adult was recorded on visit five flying over the main lagoon east of Crawley Sewage Treatment Works in an area not accessible during the survey, so birds may have been present on previous surveys and not detected.

4.1.5 Peregrine - one male was recorded on visit three on top of Pier 3, just north of the South Terminal building. As only one observation was recorded, and due to restrictions of access around airport buildings and high noise levels restricting possibilities of detecting adults, it was not possible to confirm signs of breeding during the surveys but was suspected from discussions with GAL staff.

4.1.6 Firecrest - single singing males were recorded at the eastern fringe of Horleyland Wood on visit two and in Upper Pickett's Wood on visit three. These observations could relate to territorial

males that failed to find a mate or passage migrants as there were no further records beyond late April.

#### Species of Principal Importance

4.1.7 Nine species, confirmed as breeding within the survey area (skylark, dunnoek, song thrush, marsh tit, starling, house sparrow, linnnet, bullfinch and reed bunting) are listed in Section 41 of the NERC Act 2006 as being of principal importance for the conservation of biodiversity in England.

#### Species of Conservation Concern

4.1.8 Eight species confirmed breeding within the survey area are included on the BoCC Red list. The species and reasons for Red list status are given below.

- Marsh tit – moderate breeding population decline over 25 years (-43%) and severe breeding population decline over the longer term (-72%).
- Skylark – moderate breeding population decline over 25 years (-32%) and severe breeding population decline over the longer term (-62%).
- Starling – severe breeding population decline over 25 years (-70%) and the longer term (-83%).
- Song thrush – severe breeding population decline over the longer term (-59%).
- Mistle thrush – moderate breeding population decline over 25 years (-45%) and severe breeding population decline over the longer term (-62%).
- House sparrow – moderate breeding population decline over 25 years (-32%) and severe breeding population decline over the longer term (-66%).
- Grey wagtail – moderate breeding population decline over 25 years (-33%) and severe breeding population decline over the longer term (-57%).
- Linnnet – severe breeding population decline over the longer term (-60%).

4.1.9 Six species recorded during the survey are included on the BoCC Amber List. The species and reasons for Amber list status are given below:

- Mallard – moderate non-breeding population decline over 25 years (-38%).
- Stock dove – UK breeding population is of international importance.
- Kestrel – moderate breeding population decline over 25 years (-33%) and the longer term (-46%).

- Dunnock – moderate breeding population decline over the longer term (-31%).
- Bullfinch – moderate breeding population decline over the longer term (-39%).
- Reed bunting – moderate breeding population decline over the longer term (-38%).

### Geographic Importance

- 4.1.10 The following geographical frames of reference and selection criteria (based on the Guidelines for Ecological Impact Assessment in the United Kingdom (CIEEM, 2016)) are used to ascribe nature conservation value or potential value to the bird populations within the survey area.
- International importance - a species which is cited as part of the designated interest of a SPA and occurs in internationally or nationally important numbers.
  - National importance - a species which is cited as part of the designated interest of a SSSI and occurs in nationally important numbers.
  - Regional importance – NERC Species of Principal Importance, BoCC Red List species or UK BAP Priority species that regularly occur in regionally important numbers.
  - County importance - NERC Species of Principal Importance, BoCC Red List species, UK or Hampshire BAP Priority Species that regularly occur in numbers that are important on a county basis.
  - Local importance - NERC Species of Principal Importance, BoCC Red or Amber List species, UK or Hampshire BAP Priority Species which occur regularly in locally sustainable populations.
  - Site - all common and widespread species.
- 4.1.11 For the purposes of this evaluation the number of breeding territories recorded during the survey is compared to the species' national, regional (South East England) and county (Surrey and Sussex) population estimates (where available).
- 4.1.12 National breeding population estimates are based on Holling *et al.* (2018), Musgrove *et al.* (2013) and Wilson *et al.* (2018). For those species where data are available, regional breeding population estimates are based on Conway *et al.* (2008), Holling *et al.* (2018) and Wilson *et al.* (2018). For those species where data are available, county breeding population estimates are based on Holling *et al.* (2018), in addition, a descriptive county status has been derived from the Surrey and Sussex bird lists (Surrey Bird Club, 2019; Sussex Ornithological Society, 2016).

- 4.1.13 Where no regional or county population estimates are available, professional judgment and comparisons with population estimates at higher geographical levels have been used to inform this assessment.
- 4.1.14 Table 1 of Annex 4 summarises the abundance of species of conservation interest recorded during the survey, the national and/or regional population estimate and county status for these species and the geographical importance of the populations within the survey area as derived from the criteria outlined above.
- 4.1.15 The level of geographical importance of the breeding populations of species of conservation interest is local for all species except little ringed plover, peregrine, marsh tit and firecrest. Peregrine was possibly present in numbers of regional importance; little ringed plover was possibly present in numbers of county importance; firecrest was possibly present in numbers of county importance; and marsh tit was confirmed as present in numbers of county importance.
- 4.1.16 A single adult little ringed plover was recorded on visit five near Crawley sewage treatment works (in an area of restricted access). The breeding population of little ringed plover is stable in the UK although, in recent decades, the species has expanded its range further into Wales, northern England and south and east Scotland.
- 4.1.17 A single observation of peregrine falcon was recorded just north of the South Terminal building during visit three. The UK population of peregrine has increased in recent years, particularly lowland populations as found in Surrey and Sussex. Reasons for increases in populations of peregrines in the lowlands include increasing use of human structures as breeding sites (eg pylons), abundant availability of prey and a lack of conflict with humans.
- 4.1.18 The confirmed marsh tit territory was recorded within Upper Pickett's Wood on the eastern side of the Project area. Marsh tit populations in the UK (including Surrey and Sussex) have undergone severe declines. Contributory factors in these declines include habitat loss, increased woodland isolation, loss of woodland understorey and reductions in dead wood availability (Vanhinsbergh *et al.*, 2001).
- 4.1.19 Two observations of singing firecrests were recorded during the survey; one on the eastern side of Horleyland Wood on visit two and the other in Upper Pickett's Wood on visit three. Firecrest populations in the UK (including Surrey and Sussex) have increased rapidly in recent years.

- 4.1.20 With the exception of the four species discussed above, the bird community recorded during the survey was considered typical for the habitats present within the survey area. Whilst the majority of species recorded are common and widespread in Surrey and Sussex, the habitats within the survey area do provide breeding habitat for an assemblage of species of conservation importance.

### Diversity of the Breeding Bird Assemblage

- 4.1.21 The number of species recorded in an area is a simple measure of diversity that can indicate the site's importance. Table 4.1.1 shows the criteria outlined in Fuller (1980) for breeding bird assemblages to indicate the importance of sites at various geographic levels.

**Table 4.1.1: Breeding Bird Assemblage Diversity Criteria**

	National Importance	Regional importance	County importance	Local importance
Number of species	85+	70-84	50-69	25-49

- 4.1.22 Based on Fuller's criteria, the breeding bird assemblage of 48-51 species recorded within the survey area in 2019 was at the lower limit of county importance and upper limit of local importance. However, it should be noted that Fuller's analysis was developed in the 1970's since when species diversity has declined significantly. As a result, Fuller's thresholds are considered too high for today's breeding bird populations. Taking this into consideration, the diversity of the breeding assemblage should be considered as of county importance.
- 4.1.23 Overall, the breeding bird assemblage within the Project site boundary was considered to be of county importance due to the diversity of species present and the presence of three species breeding, or possibly breeding, in numbers of county importance and one species possibly breeding in numbers of regional importance.
- 4.1.24 Conclusion
- 4.1.25 The survey of breeding birds recorded a breeding assemblage of 51 species in 2019. The survey undertaken from March-July 2019 was undertaken during the peak breeding period.
- 4.1.26 Of the 51 species recorded as breeding or possibly breeding within the survey area, 17 species meet at least one of a range of

	criteria relating to special statutory protection or conservation importance.	4.1.37	The overall wintering bird population within the site was considered as being of no more than local importance.		
4.1.27	No breeding population of any species within the survey area approaches the 1% level of the national population. Therefore, no species considered to be breeding or possibly breeding are present in nationally important numbers.	4.2	<b>Reptile Surveys</b>		
		4.2.1	A good size population of grass snake was identified in grassland habitats along the River Mole in the west of the Project site.		
4.1.28	One species (peregrine), possibly breeding within the survey area meets the 1% level of the regional population and was considered to be possibly breeding in regionally important numbers.	4.2.2	Individual grass snakes were also identified around wetland habitats in the east of the site suggesting a separate low size population.	4.4.2	Although <i>Myotis</i> species are notoriously difficult to distinguish from sound analysis alone, a number of calls were characteristic of those of Brandt's/whiskered bat, Daubenton's bat and natterer's bat. Therefore, these species have been included in the account below as they are likely to be present but from bat sound analysis alone their presence cannot be confirmed.
4.1.29	Two species (little ringed plover and firecrest), possibly breeding within the survey area meet the 1% level of the county (Surrey and/or Sussex) populations and are considered to be possibly breeding in numbers of county importance.	4.2.3	The two areas where grass snake was recorded were disconnected from each other. The habitats between them were associated with the airport and comprised low value habitats for grass snake. Therefore, the survey results indicate that two separate populations are present.	4.4.3	Desk study records confirmed the presence of two additional species within the search area, Bechstein's <i>Myotis bechsteinii</i> and Alcahioe <i>Myotis alcahioe</i> , which are considered very rare and rare species.
4.1.30	One species (marsh tit) was confirmed breeding within the survey area and met the 1% level of the county (Surrey and/or Sussex) population and was considered to be breeding in numbers of county importance.	4.3	<b>Great Crested Newt Surveys</b>	4.4.4	Species classified as very rare, rare, scarce and uncommon are as such because of restricted distribution and/or low to moderate populations.
		4.3.1	Thirty-six ponds were identified within the Project site boundary.	4.4.5	Bats not identified to species level comprised pipistrelle bats and <i>Myotis</i> species.
4.1.31	Records of black redstart were provided in the desk study data from 2013, with the last breeding records from 2012. Black redstarts were previously recorded around the Old Control Tower (landside). The habitat around this area was found to continue to provide some suitable breeding and foraging opportunities for black redstart; however, none were recorded during surveys undertaken between winter and spring 2019.	4.3.2	Four ponds were identified as having great crested newts present. Pond W46 was identified as having a medium sized great crested newt population, Ponds K5F and TTD were identified as having small populations of great crested newt.	4.4.6	The highest levels of activity were recorded at Location 3 (Brockley Wood) (41,710 passes), Location 7 (Horleyland Wood) (37,967 passes), Location 6 (Perimeter Road East) (28,845 passes) and Location 11 (Crawter's Wood) (14,161 passes), indicating the importance of these woodland habitats to the bat assemblage within the Project site boundary and wider area.
4.1.32	The diversity of species present within the survey area was at a level indicative of county importance for breeding birds.	4.3.3	Although no great crested newt adults were identified within Pond 8N8, great crested newt eggs were identified within the marginal vegetation and an eDNA survey produced a positive result for great crested newt.	4.4.7	A total of three barbastelle passes were recorded during the static surveys, comprising two passes at Location 1 (Land west of the Fire Training Ground) in September and one pass at Location 9 (Perimeter Road South) in July. This indicates that barbastelle use the woodland and woodland edge immediately to the south and west of the airport. Barbastelle's use a range of habitats and forage over a wide area.
4.1.33	Wintering Bird Surveys	4.3.4	Both ponds K5F and TTD had great crested newt eggs recorded within their marginal vegetation meaning that these along with Pond 8N8 are viable populations.	4.4.8	Nathusius' pipistrelles were recorded during both transect and static surveys. During static surveys peak counts of 63 and 32 passes were recorded at Location 6 (Perimeter Road East) and Location 8 (Land East of the Railway Line Wetland) respectively. The majority of passes (83) were recorded in June, July and August. This coincides with likely higher levels of activity associated with the maternity season. In general, higher numbers
4.1.34	A total of 61 species were recorded within the survey area during the wintering bird surveys undertaken between October 2018 and March 2019.	4.3.5	The distribution of the ponds indicates that two great crested newt metapopulations are present.		
4.1.35	No wintering species were recorded in any numbers which were considered to be of national or international significance (ie >1% of the wintering population) and in all cases the numbers of birds recorded were considerably below this threshold.	4.3.6	Common toad was also recorded within the survey area. The toads were located in Pond W46 and within the field south of Brockley Wood.		
4.1.36	The area within the Project site boundary was considered to be of site-level importance for wintering lapwing based on the peak counts for this species and its current conservation status. Lapwings were predominantly recorded around the Crawley Sewage Treatment Works in the east of the Project site.	4.4	<b>Bat Surveys</b>		
			<b>Bat Assemblage</b>		
		4.4.1	Field surveys undertaken between 2019 and 2022 confirmed the presence of at least 12 species of bat within the Project site boundary and surrounding area. These include:		
					<ul style="list-style-type: none"> <li>▪ one very rare species – barbastelle;</li> <li>▪ three rare species – Nathusius' pipistrelle, Brandt's bat and whiskered bat;</li> <li>▪ one scarce species – Leisler's bat;</li> <li>▪ two uncommon species – noctule and serotine; and</li> <li>▪ five common species – common pipistrelle, soprano pipistrelle, Daubenton's bat, Natterer's bat and <i>Plecotus</i> sp.</li> </ul>

- of Nathusius' pipistrelle were recorded to the north and east of the Project site boundary, suggesting that they use the woodland associated with the Crawley Sewage Treatment Works, Riverside Garden Park and railway corridors for foraging and commuting.
- 4.4.9 Common and widespread species such as common pipistrelle and soprano pipistrelle were abundant throughout the survey area with moderate to high levels of activity recorded during both transect and static surveys. Common pipistrelle was the most frequently recorded species during static surveys, accounting for over 65% of the species composition across all locations. At Locations 6 and 9, common pipistrelle accounted for over 98% of all bat passes recorded between April-October.
- 4.4.10 Pipistrelle bats are generally flexible in their habitat requirements for foraging and commuting and are able to utilise a range of habitats in both urban and rural landscapes.
- 4.4.11 *Plecotus sp.* bats are generally woodland species, although can be found using other habitats such as parkland. Peak counts of *Plecotus sp.* bats were recorded at Location 3 (Brockley Wood) (249 passes), Location 4 (North of the Long Stay North car park) adjacent to the River Mole corridor (70 passes) and Location 7 (54 passes). *Plecotus sp.* were recorded in generally low numbers throughout the Project site and were not recorded at either Location 8 or Location 9 during static surveys.
- 4.4.12 Higher numbers of *Myotis sp.* passes were generally associated with the woodland areas such as Brockley Wood, Horleyland Wood, Upper Pickett's Wood and Riverside Garden Park. A total of 7,277 *Myotis sp.* passes was recorded at Location 3 (Brockley Wood) with peak counts of 3,102 passes and 1,728 passes recorded in May and July respectively. These periods coincide with likely higher levels of activity associated with the pre-maternity and maternity seasons. A large number of *Myotis sp.* bat passes was also recorded along the woodland edges associated with the River Mole Corridor in the west of the survey area. Although not confirmed through sound analysis, a proportion of these calls (particularly within the woodland) are likely to be from Bechstein's bats, which are known to be present in the area and typically roost and forage within deciduous woodland.
- 4.4.13 Daubenton's bats, which are commonly associated with habitats found within the area including broadleaved woodland and standing water, were detected during transect surveys, notably along Transect 2 and Transect 5.
- 4.4.14 Some of the *Myotis sp.* calls were characteristic of Whiskered and/or Brandt's bats, which were recorded along Transects 1 and 5. Both species are characteristic of woodland habitat, although to a lesser extent for Whiskered bats. Brandt's bats tend to forage at low and medium heights in the woodland canopy and are more likely to forage over open water, whereas Whiskered bats favour woodland edges, close to vegetation, hedgerows and open habitats, including flowing water.
- 4.4.15 Several calls of *Myotis sp.* bats were characteristic of natterer's bat and recorded along Transects 1, 3 and 5. As much of the species' prey is taken from foliage and they normally fly at low altitudes (less than 5 metres), the woodland around Transects 1 and 3 and the woodland edge along the eastern and south western boundary of Transect 5 provide suitable foraging habitats for Natterer's bats.
- 4.4.16 Moderate to high levels of bat activity of scarce and uncommon species, Leisler's, noctule and serotine, were recorded predominantly in areas of open riparian habitat, in comparison to those recorded along linear features (such as the river and railway corridors). These species often fly over open habitat, making them easier to detect.
- 4.4.17 Peak counts of noctules were recorded at Location 2 (Land south west of the River Mole) with 1,088 passes and Location 8 with 1,237 passes; the detectors in these locations were situated in more open areas of habitat. Noctules are a fast, high-flying species when foraging and commuting. They are typically associated with broadleaved woodland and open pasture and it was unlikely that the fragmentation of habitats would impact upon this species.
- Foraging Habitat**
- 4.4.18 Areas of significant bat foraging activity were recorded within the woodland areas across the survey area and water bodies (Old Lagoon and New Lagoon) associated with Crawley Sewage Treatment Works in the east of the Project site. The patchy wooded landscape and associated riparian habitats are likely to provide optimum foraging habitat for a variety of species including *Myotis sp.* bats, pipistrelles and long-eared bats.
- 4.4.19 High levels of foraging activity were recorded along Transect 5, adjacent to the Aviation Museum; the boundary habitats here comprised mature trees and hedgerows, woodland edge and the River Mole along the eastern boundary of the transect route.
- 4.4.20 The landscape in the area generally comprised large areas of woodland and interconnecting hedgerows and other linear features which provide links to high value habitat across the wider area.
- 4.4.21 The presence of less common and rare species suggests that the overall quality of the habitats present is able to support populations of large numbers of bats and a high diversity of species, which contributes to the importance of foraging habitat in this area for bats.
- Commuting Habitat**
- 4.4.22 The woodland compartments connected by watercourses, mature hedgerows and tree lines provided suitable habitat to support the bat assemblage in this area.
- 4.4.23 Significantly lower levels of commuting activity were recorded along Transect 4, with only a few common pipistrelle passes recorded. This was likely to be due to the lack of suitable habitat and the presence of strong artificial light and noise emanating from the airport and surrounding ancillary buildings.
- 4.4.24 Overall, the continuity of connective habitat was likely to provide an extensive network of habitat features suitable for a wide range of commuting bats, providing links to the wider landscape in this area.
- Crossing Point Surveys**
- River Mole**
- 4.4.25 A total of 1278 bat passes from at least five species were observed using the feature over three survey visits, with the highest total number of passes from common pipistrelle (1017) and the lowest total number from brown long-eared bat (3).
- 4.4.26 Nineteen roosting locations for [REDACTED] bats were identified in 2020 and 2021 using bat radio tracking survey within [REDACTED].
- 4.4.27 The River Mole was identified as a core foraging area for this species from radio-tracking surveys undertaken in 2019, 2020 and 2021. It was identified as a core foraging area in 2019 for a male and a peripheral foraging area for two males, out of the seven Bechstein's bats which were radio-tracked in 2019. It was identified as a core foraging area for three out of the fourteen Bechstein's bats which were radio-tracked in 2020 and 2021. These included a lactating female, a post-lactating female and an adult male.



4.4.28 Flightlines along the River Mole were identified for three males and one of the non-breeding females in 2019, out of the seven bats which were radio-tracked. No flightlines were recorded from bat roosts to foraging areas in 2020 and 2021 as the majority of bats were recorded close to their roosting locations

4.4.29 Twenty-four passes of *Myotis sp.* bats were recorded within the river corridor or directly above it. This, in conjunction with results of the bat radio tracking survey, indicates that *Myotis sp.* bats, likely including Bechstein's bats, are using the River Mole corridor to move across the landscape and for foraging.

**Riverside**

4.4.30 A total of 1159 passes from at least five species were observed using the feature over three survey visits, with the highest total number from common pipistrelle (654) and the lowest from brown long-eared bat (2).

4.4.31 Of the passes of bats observed using the feature, 19% were observed passing at an "unsafe height" and 81% were observed passing at a safe height. The definition of safe and unsafe height is based on the assumption that the proposed road improvements in the areas of the surveys will be at the current height of the ground.

4.4.32 The Riverside Park was identified as a core foraging area for Bechstein's bats using bat radio tracking survey in 2019. This area was identified for one of the seven Bechstein's bats, an adult male radio-tracked in 2019. None of the fourteen Bechstein's bats radio-tracked in 2020 and 2021 was recorded foraging within Riverside Park.

4.4.33 A total of 18 passes of *Myotis* species were recorded within Riverside Park. This, in conjunction with the results of bat radio tracking survey, indicates that *Myotis sp.* bats, likely to include Bechstein's bats, are using Riverside Park for foraging and commuting.

**Assessment of importance**

**River Mole**

4.4.34 This location is confirmed as an important commuting route and foraging area for bats. The River Mole is considered to be an important commuting route at regional level for common pipistrelle, at county level for noctule and soprano pipistrelle, and at local level for brown long-eared bat. The River Mole is considered to be an important foraging area at county level for

noctule and common pipistrelle, and at local level for brown long-eared bat and soprano pipistrelle.

4.4.35 Although it was not possible to distinguish between *Myotis* species on the basis of call parameters, twenty-four passes of *Myotis sp.* bats were recorded and the radio-tracking surveys confirmed that three Bechstein's bats out of a sample size of seven bats used the Rive Mole corridor as a core or peripheral foraging area in 2019, and that three Bechstein's bats out of a sample size of fourteen bats used the River Mole corridor as a core foraging area in 2020 and 2021. This indicates that the River Mole is likely to be an important foraging area for Bechstein's bats at regional level.

**Riverside Park**

4.4.36 This location is confirmed as an important commuting route and foraging area for bats. Riverside Park is considered to be an important commuting route at regional level for common pipistrelle and soprano pipistrelle and at local level for brown long-eared bat. This feature is considered to be an important foraging area at county level for common pipistrelle and soprano pipistrelle, and at local level for brown long-eared bat.

**4.5 Invertebrate Scoping Survey**

4.5.1 Several areas within the Project site presented features of potential value to invertebrates which were considered to have a moderate invertebrate interest that would likely be raised above the expected regional background level.

**4.6 Terrestrial Invertebrate Survey**

4.6.1 Surveys of the North West Zone and Land East of the Railway Line identified a diverse assemblage of terrestrial invertebrates in these areas, including a range of scarce and unusual species.

**4.7 Aquatic Invertebrate Survey**

4.7.1 Several species designated under Section 41 of the NERC Act (2006) were identified by the desk study.

4.7.2 In 2019, the invertebrate habitat appraisal identified that Pond M and the ditches adjacent to Pentagon Field had features of moderate invertebrate interest above the expected regional background level.

4.7.3 Further detailed assessment of the River Mole and Gatwick Stream found both watercourses supported macroinvertebrate communities indicative of moderately polluted conditions,

exacerbated by relatively low flow conditions and high levels of sedimentation. Dense macrophyte growth on the River Mole is contributing to acute reductions in dissolved oxygen which are impacting on the macroinvertebrate assemblage.

4.7.4 There is one record from 2013 of shining ram's-horn snail, an IUCN Red List species and UK species of principal importance under the 2006 NERC Act. Although not recorded during the survey, there remains a possibility that the species may occur at the site of the 2013 record at the downstream end of the desk study area.

4.7.5 The Gatwick Stream appears to be impacted by both organic pollution and silt deposition, possibly from a storm water discharge outlet from a nearby industrial area.

4.7.6 The invasive New Zealand mud snail was identified at the River Mole and Gatwick Stream sites, and signal crayfish were observed at both the Gatwick Stream sites during each visit

**4.8 Fish Survey**

4.8.1 The desk study identified that brown trout had previously been recorded within the Project site boundary, although it was not recorded in surveys in 2020. Brown trout is listed under Section 41 of the NERC Act (2006).

4.8.2 Both the River Mole and Gatwick Stream had consistently high fish populations. This is likely to be a consequence of stable temperature and DO conditions caused by shading and potentially high abundances of pollution tolerant macroinvertebrates such as Oligochaete worms as a food source.

**4.9 Limitations**

4.9.1 A number of limitations were experienced during the surveys. The most frequently occurring limitations across a range of surveys included:

- a lack of access to certain areas required to complete surveys;
- unsuitable weather conditions; and
- high levels of noise and lighting.

4.9.2 Further details of specific limitations affecting each survey are given below.

### Breeding Birds

- 4.9.3 There were some limitations to the 2019 Gatwick breeding bird survey which included:
- restrictions on land access from landowners and restricted areas airside. This included no access south of the runway during survey visit two due to visibility restrictions in place at the time;
  - excessive noise levels from aircraft and associated activities, particularly during airside surveys, which may have reduced / impeded distance to which vocalisations were detected; and
  - moderate to poor weather conditions during the first day of survey visits two and four.

- 4.9.4 In light of the above limitations to the survey methodology, consideration has been made during the analysis of survey data and in the writing of this report with regard to:
- the possibility of species presence not being detected during the survey; and
  - the likelihood of a reduced number of territories being detected.

### Wintering Birds

- 4.9.5 During the sixth and seventh survey visits within part of area A3 Land East of the Railway Line, a group of travellers had gained access to the fields south of Upper Picketts Wood, so these areas were not surveyed in the last two surveys.

### Great Crested Newt Surveys

- 4.9.6 Several ponds could not be accessed due to restrictions on the surrounding land. Without further surveys of these ponds, the possibility that they may support great crested newt cannot be ruled out.
- 4.9.7 Ponds C24 and 29A were not included within the original surveys. However, HSI conducted outside of the optimal survey season (September 2019) identified that Pond C24 had a 'good' habitat score and Pond 29A had an 'average' habitat score for great crested newt.

### Bat Transects

- 4.9.8 The routes for Transects 1, 3 and 5 were modified between April and May, and 3 and 5 were changed again from June onwards.

4.9.9 Along Transect 3, a minor deviation was made to the route through the Riverside Garden Park to include a broader range of habitats along the northern and southern boundaries of the park. For Transect 5, the route deviated to incorporate an area of land to the north of the original transect route, which included a large area of pasture land and wooded hedgerows adjacent to Man's Brook. Transect 1 was changed due to the original transect being too complex to reliably duplicate over subsequent transects. The deviations from the original route are not thought to be a limitation to the results, as the new routes incorporated a larger and more diverse area, which could potentially be utilised by a greater variety of species.

4.9.10 The first post-maternity survey along Transect 2 was cancelled due to access constraints. Although the survey was not rescheduled, this is not considered to have significantly affected the overall assessment.

4.9.11 A number of static detectors failed to record data for a minimum of five nights due to equipment failure and malfunction. Where this occurred, the species assessment is conservative to account for gaps in the information.

4.9.12 Some species, such as long-eared bats, are likely to be underrepresented in the analysis. This is due to their call characteristics which are comparatively quiet compared to that of other species. In order for the detectors to record long-eared bat calls, bats must fly within 3 metres of the microphone.

4.9.13 In addition, the calls of *Myotis sp.* and long-eared bat species are difficult to distinguish and therefore calls were only analysed down to species level where they were clearly characteristic of that species and within suitable habitat. Although Bechstein's bats were not confirmed through sound analysis of activity data, it is likely that a proportion of unidentified *Myotis* bat calls are from Bechstein's bats which are known to be present within the woodlands surveyed.

4.9.14 Additional survey techniques, including bat trapping and radio-tracking surveys ensured that the presence of this species was accounted for and included within the assessment of the overall bat assemblage at Gatwick.

4.9.15 Sound analysis was not possible for a small number of transect surveys during the pre-maternity and maternity seasons due to equipment failure and malfunction, therefore some species' accounts and interpretation are based on field observations only. This was not thought to be a significant limitation to the results

and the species assessment was conservative to account for these gaps in information.

## 5 Conclusions

- 5.1.1 The ecology surveys undertaken on the Project site boundary found that the majority of the centre of the site, associated with the airport and infrastructure, comprised buildings, areas of hardstanding, amenity grassland and introduced shrubs and trees. They provided some areas of suitable habitat for breeding birds but were otherwise of overall low ecological value.
- 5.1.2 The habitats within the Project site boundary that surrounded the airport supported a number of higher value habitats, including semi-natural broadleaved woodland, scrub and trees, semi-improved neutral grassland, marshy grassland, ponds, rivers and hedgerows.
- 5.1.3 These areas were considered higher value habitats which supported a more diverse flora and fauna, especially within the associated Land East of the Railway Line wetland and woodland in the south east of the Project site boundary and the areas around the River Mole. They supported a variety of breeding birds, including species of conservation concern and were well used by foraging and commuting bats, including some rare bat species.
- 5.1.4 Populations of great crested newt and grass snake were found in these habitats within the Project site boundary.

## 6 References

- Bat Conservation Trust (2011) Statement on the impact and design of artificial light on bats. Bat Conservation Trust, London.
- Bat Conservation Trust (2014) Artificial lighting and wildlife Interim Guidance: Recommendations to help minimise the impact of artificial lighting on bats. Bat Conservation Trust, London.
- Berthinussen A. and Altringham, J. (2015). Development of a cost-effective method for monitoring the effectiveness of mitigation for bats crossing linear transport infrastructures. Defra Research Project WC1060]
- Bibby, C.J., Burgess, N.D., Hill, D.A. & Mustoe, S.H. (2000) *Bird Census Techniques, second edition*. Academic Press, London.

Bicker, R. (2018) Gatwick Biodiversity Action Plan: Five Year Review 2012-2017.

CIEEM (2016) *Guidelines for Ecological Impact Assessment in the UK and Ireland, second edition*. Chartered Institute of Ecology and Environmental Management.

Collins J. (ed.) (2016) *Bat surveys for Professional Ecologists: Good practice guidelines (3rd Edition)*. Bat Conservation Trust, London.

Conway, G.J., Burton, N.H.K., Handschuh, M. and Austin, G.E. (2008) *UK population estimates from the 2007 Breeding Little Ringed Plover and Ringed Plover Surveys, BTO Research Report No. 510*. British Trust for Ornithology, Thetford, Norfolk.

Department for Environment, Food and Rural Affairs (Defra) (2007) *Hedgerow Survey Handbook: A standard procedure for local surveys in the UK*. DEFRA, London.

Eaton, M., Aebischer, N., Brown, A., Hearn, R., Lock, L., Musgrove A., Noble D., Stroud, D. and Gregory, R. (2015) Birds of Conservation Concern 4. The population status of birds in the United Kingdom, Channel Islands and Isle of Man. *British Birds* **108**, 708-746

English Nature (2001) Great Crested Newt mitigation guidelines. English Nature, Peterborough.

Fawcett-Williams, K. (2019) *Thermal Imaging: Bat Survey Guidelines*. Bat Conservation Trust, London

Fuller, R. J. (1980) A Method for Assessing the Ornithological Interest of Sites for Conservation. *Biological Conservation* **17**, 229-239.

Gilbert, G., Gibbons, D.W. and Evans, J. (1998) *Bird Monitoring Methods: A manual of techniques for key species*. RSPB/BTO/JNCC/WWT/ITE/The Seabird Group. RSPB/BTO, Sandy, Beds.

Holling, M. and the Rare Breeding Birds Panel (2018) Rare breeding birds in the United Kingdom in 2016. *British Birds* **111**, 644-694.

Imboden, C. (1974) Resettlement and Breeding Season of the Northern lapwing *Vanellus vanellus* in Europe. *Orn. Beob.* **71**, 5-134.

Joint Nature Conservation Committee (JNCC) (2010) *Handbook for Phase 1 Habitat survey: a technique for environmental audit (revised reprint)*. Joint Nature Conservation Committee, Peterborough.

Kirby, S. and Lack, P. C. (1993) Spatial dynamics of wintering Lapwings and Golden Plovers in Britain and Ireland, 1981/82 to 1983/84. *Bird Study* **40**, 38-50. Lack, P.C. (1986) *The atlas of wintering birds in Britain and Ireland*. T. & A.D. Poyser, Calton.

Met Office (2021). Cool, wet May brings a Spring of marked contrasts. Available at: <https://www.metoffice.gov.uk/about-us/press-office/news/weather-and-climate/2021/cool-wet-may-concludes-spring-of-marked-contrasts> [ Accessed 07.06.2021]

Musgrove, A., Aebischer, N., Eaton, M., Hearn, R., Newson, S., Noble, D., Parsons, M., Risely, K, and Stroud, D. (2013) Population estimates of birds in Great Britain and the United Kingdom. *British Birds* **106**, 64-100.

Rodwell, S. (1995) *British Plant Communities: Volume 4 Aquatic Communities, Swamps and Tall-Herb Fens*. Cambridge University Press, Cambridge.

Surrey Bird Club (2019) The Surrey List <https://surreybirdclub.org.uk/records/surrey-list/> [accessed September 2019].

Sussex Ornithological Society (2016) The Sussex List <https://www.sos.org.uk/the-sussex-list> [accessed September 2019].

Vanhinsbergh, D., Fuller, R.J & Noble, D.G. (2001) *A Review of Possible Causes of Recent Changes in Populations of Woodland Birds in Britain, BTO Research Report 245*. British Trust for Ornithology, Thetford, Norfolk.

Wilson, M. W., Balmer, D. E., Jones, K., King, V. A., Raw, D., Rollie, C. J., Rooney, E., Ruddock, M., Smith, G. D., Stevenson, A., Stirling-Aird, P. K., Wernham, C. V., Weston, J. M., & Noble, D. G. (2018) The breeding population of Peregrine Falcon (*Falco peregrinus*) in the United Kingdom, Isle of Man and Channel Islands in 2014. *Bird Study* **65**: 1-19.

## 7 Glossary

### 7.1 Glossary of terms

Term	Description
<b>ARGUK</b>	Amphibian and Reptile Group UK
<b>BAP</b>	UK Biodiversity Action Plan Priority Species
<b>BCT</b>	Bat Conservation Trust
<b>BoCC</b>	Birds of Conservation Concern
<b>BTO</b>	British Trust for Ornithology
<b>CIEEM</b>	Chartered Institute of Ecology and Environmental Management
<b>DMRB</b>	Design Manual for Roads and Bridges
<b>eDNA</b>	Environmental DNA
<b>EIA</b>	Environmental Impact Assessment
<b>ES</b>	Environmental Statement
<b>GAL</b>	Gatwick Airport Limited
<b>HSI</b>	Habitat Suitability Index
<b>JNCC</b>	Joint Nature Conservation Committee
<b>LNR</b>	Local Nature Reserve
<b>MAVIS</b>	Modular Analysis of Vegetation Information System
<b>NERC</b>	Natural Environment and Rural Communities Act
<b>NNRs</b>	National Nature Reserve
<b>NVC</b>	National Vegetation Classification
<b>PBRA</b>	Preliminary Bat Roost Assessment
<b>PEIR</b>	Preliminary Environmental Information Report
<b>SAC</b>	Special Area of Conservation
<b>SNCI</b>	Site of Nature Conservation Importance
<b>SPA</b>	Special Protection Area
<b>SSSI</b>	Site of Special Scientific Interest
<b>WHPR</b>	Whalley Hawkes Paisley Trigg

## Annex 1

### Survey Methodologies

## A1.1 Survey Methodologies

### National Vegetation Classification Surveys

- A1.1.1 A National Vegetation Classification (NVC) survey was carried out following the methodology and guidelines set out in the Joint Nature Conservation Committee's (JNCC) NVC User's Handbook (Rodwell *et al.*, 2006).
- A1.1.2 Fieldwork was carried out on the 8–12 April, 8–12 July and 6–8 August 2019 by a suitably qualified ecologist and botanist. The survey was undertaken during the optimal time for both grassland and woodland botanical surveys.
- A1.1.3 A general walkover of the site was carried out to identify homogenous stands of vegetation within the survey boundary.
- A1.1.4 Quadrat data was collected (1 metre x 1 metre quadrats) from within their represented stands of homogenous vegetation. Plant species within the quadrats were recorded following the nomenclature in Stace (2010). Percentage cover and DOMIN values were also recorded for each species. A breakdown of DOMIN values and their estimated percentage cover are outlined in Table 2.1.1.
- A1.1.5 The habitat community was identified for each homogenous stand of vegetation using the computer programme TABLEFIT. The TABLEFIT programme computes 'Goodness of Fit' between quadrat data (from sampled vegetation) and the published NVC tables (which define the NVC communities and sub-communities). This gives an initial indication of which NVC types the data are most likely to have been drawn from – the highest coefficient does not necessarily indicate the correct NVC diagnosis.
- A1.1.6 It was then necessary to identify the NVC type through careful consideration of the NVC descriptions in British Plant Communities (Rodwell, 1991, 1992, 1995, 2000; Rodwell *et al.*, 2000). There is no guarantee that the highest coefficient corresponds to the 'correct' NVC diagnosis.
- A1.1.7 DOMIN scale and percentage cover estimates:

**Table A1.1.1: DOMIN Scale and Percentage Cover Estimates**

Cover	DOMIN value
<4 % (few individuals)	1
<4 % (several individuals)	2
<4% (many individuals)	3
4 – 10 %	4
11 – 25 %	5
26 – 33 %	6
34 – 50 %	7
51 – 75 %	8
76 – 90 %	9
90 – 100 %	10

### Hedgerow Surveys

- A1.1.8 The hedgerow survey followed the methodology and guidance set out in the Hedgerow Survey Handbook: A standard procedure for local surveys in the UK (Department for Environmental, Food and Rural Affairs (Defra), 1997) and involved surveying 30 metre lengths of hedgerow.
- A1.1.9 A hedgerow is defined as any boundary line of trees or shrubs over 20 metres long and less than 5 metres wide at the base, provided that at one time the trees or shrubs were more or less continuous. All hedgerows consisting of at least one woody UK native species are UK BAP priority habitats.
- A1.1.10 Hedgerow surveys were undertaken on 5–8 August 2019 and all hedgerows were surveyed to assess whether they qualified as protected hedgerows or not. The method is based on definable lengths of hedgerow between two end points, that were identified as:
- any point of connection between two, or more, hedgerows or to other features eg fences, walls, ditches, roads;
  - the point at which a hedgerow stops and there is a gap of more than 20 metres to the next hedgerow (eg where the hedgerow ends in the middle of a field); and
  - the point at which the hedgerow links to a woodland or other semi-natural habitat such as a pond.
- A1.1.11 Three additional end points were included in the assessment, where there was significant variation in the characteristics of the hedgerow. These were:

- the point at which the hedgerow changes character from one hedgerow type to another for 20 metres or more;
- where there is a distinct change in hedgerow height for lengths of 20 metres or more; and
- the ends of lengths (20 metres or more) of recent planting, coppicing or laying.

A1.1.12 Each section between two end points was considered a separate hedgerow and was surveyed as such. 30 metre lengths of each hedgerow were identified. Where the hedge was 30 metres or less in length then the whole hedge was surveyed. If the hedge was between 30-100 metres then the central 30 metres of hedgerow was surveyed. If the hedge was between 100-200 metres long then the hedgerow was divided into two and the central 30 metres of the two sections was surveyed. Finally, if the hedge was over 200 metres in length it was divided into three sections and the central 30 metres of each of the thirds was surveyed.

A1.1.13 To be considered protected, the hedgerow had to exhibit one of the following:

- it had an average of seven or more woody species in the surveyed section(s);
- it had an average of six woody species in the surveyed section(s) and three or more features from:
  - a wall or bank along half or more of the length;
  - o a ditch along half or more of the length;
  - an average of one standard tree or more per 50 metres of hedgerow;
  - o gaps which do not add up to more than 10% of the hedge;
  - three woodland understorey species;
  - o a parallel hedge within 15 metres; or
  - connections scoring four points. Connections to a hedge scores one point. Connections to a pond or wood score two points;
- it had six woody species and one of the following rare trees – black poplar, large leaved lime, small leaved lime, wild service tree;
- it had an average of five woody species on average in the surveyed section(s) and has four or more features listed above (bullet point two); and
- it had four woody species on average in the surveyed section(s); is adjacent to a footpath, bridleway, byway open to all traffic (but not necessarily a normal adopted vehicular

highway unless it also is one of these) and has two or more features listed above (bullet point two).

### Breeding Bird Surveys

- A1.1.14 The breeding bird survey was based on a standard territory mapping methodology as outlined in Gilbert *et al.* (1998) and Bibby *et al.* (2000).
- A1.1.15 This method is based on the principle that the majority of species are territorial during the breeding season. This results in birds occupying discrete territories and displaying various behaviours (eg conspicuous song, visual display and periodic disputes with neighbouring individuals) allowing their location and abundance to be estimated.
- A1.1.16 The survey area (Project site boundary), as shown in Figure 2.4.1, was walked at a slow pace in order to locate and identify all individual birds. Visits were undertaken early in the morning, finishing before midday. All of the site was covered where land access was granted or where it was safe to do so given the constraints of the operational airport. There was no access airside on visit one of the survey so an extra visit to site to make up for this was carried out on the 27<sup>th</sup> of June. On the second visit, there was no access to the south-side of the main runway due to a necessary enforcement of a visibility restriction preventing movement of security vehicles. No extra visits were conducted to cover this. Suitable optical equipment was used to observe bird behaviour and all accessible parts of the survey area were approached to within 50-100 metres. Survey routes were mapped and the direction walked alternated on each visit, to ensure that all areas were covered at various times of morning across the duration of the survey. All species encountered within the survey area were recorded and mapped.
- A1.1.17 Surveys for breeding birds were undertaken in spring/summer 2019 with a total of seven survey visits taking place. The survey visits were as follows:
  - Visit 1: 27 and 28 March 2019;
  - Visit 2: 9 and 10 April 2019;
  - Visit 3: 23 and 24 April 2019;
  - Visit 4: 7 and 8 May 2019;
  - Visit 5: 21 and 22 May 2019;
  - Visit 6: 5 and 6 June 2019; and
  - Visit 7: 27 June 2019 ('airside' only).
- A1.1.18 On each visit, registrations were recorded directly into ESRI Arcpad GIS software loaded on handheld PDA devices, with a

1:10,000 scale Ordnance Survey base map of the Project area and adjacent land. A fresh map was used for each survey. Registrations of birds were recorded using standard British Trust for Ornithology (BTO) two letter species codes (BTO, 2009). Specific codes were also used to denote singing, calling, movement between areas, flight, carrying food, nest building, aggressive encounters and other behaviour.

- A1.1.19 The expected outcome of a territory mapping survey is that mapped registrations fall into clusters approximately coinciding with territories. A cluster is generally a spatially distinct group of registrations that represent the activity of not more than one territorial male or pair. Ideally, clusters include registrations of territorial behaviour across all visits and are clearly demarcated from adjacent clusters by simultaneous recording of neighbouring birds. Where a species exhibits high territory density, the mapping of simultaneously singing birds becomes essential. Territory boundaries are assumed to be between such birds.
- A1.1.20 Territory mapping methods produce analysis maps of non-overlapping ellipses encircling clusters of records thought to relate to separate territorial males or breeding pairs. These ellipses may not show the entire extent of a pair's actual breeding territory, which may be significantly larger; however, they are likely to show those areas in which the pair is most active.
- A1.1.21 On completion of the surveys analysis maps were produced for each species consisting of all registrations recorded during the survey. From these species maps, the number of territories was calculated by identifying the number of clusters present.
- A1.1.22 Standard registration mapping techniques were also used to record non-breeding species.
- A1.1.23 The following definitions have been used to identify the breeding status of the species recorded.
  - Confirmed breeding: includes species for which territories were positively identified as a result of the number of registrations, the location of an active nest, and the presence of recently fledged young or downy young.
  - Probable breeding: includes a pair observed in suitable nesting habitat in the breeding season, or agitated behaviour / anxiety calls from adults suggesting probable presence of a nest or young nearby. Behaviour was observed on insufficient occasions to confirm the presence of a territory.
  - Possible breeding: includes species observed in the breeding season in suitable nesting habitats or a singing

male present (or breeding calls heard) in the breeding season in suitable breeding habitat.

- Non-breeding: fly-over species observed but suspected to be on migration, or species observed but suspected to be summering non-breeder.

### Assessment Criteria

- A1.1.24 The assessment of the breeding bird community within the Project site boundary includes a focus on species that are afforded special statutory protection or those included on one, or more, of the lists of species of conservation interest, these include:
  - Species listed on Annex 1 of the EC Birds Directive (Directive 2009/147/EC).
  - Species listed on Schedule 1 of the Wildlife and Countryside Act 1981 (as amended).
  - Species included on the Section 41 list of Species of Principal Importance of the Natural Environment and Rural Communities (NERC) Act 2006.
  - Species included in the Birds of Conservation Concern (BoCC) Red and Amber Lists (Eaton *et al.*, 2015).
  - Species occurring in nationally, regionally or locally important numbers.
- A1.1.25 Annex 1 species are those for which the UK Government is required to take special measures, including the designation of Special Protection Areas (SPAs), to ensure the survival and reproduction of these species throughout their area of distribution.
- A1.1.26 Schedule 1 species are those which, along with their nests, eggs and dependant young, are afforded additional protection during the breeding season.
- A1.1.27 The NERC list of Species of Principal Importance is used to guide decision-makers such as public bodies, including local and regional authorities, in implementing their duty under section 40 of the NERC Act 2006; under Section 40 every public authority (eg a local authority or local planning authority) must, in exercising its functions, have regard, so far as is consistent with the proper exercise of those functions, to the purpose of conserving biodiversity. In addition, with regard to those species on the list of Species of Principal Importance prepared under Section 41, the Secretary of State must:
 

*“(a) take such steps as appear to the Secretary of State to be reasonably practicable to further the conservation*

of the living organisms and types of habitat included in any list published under this section”, or

“(b) promote the taking by others of such steps.”

- A1.1.28 Species listed on the BoCC Red List are those that have declined in numbers by 50% over the last 25 years, those that have shown a historical population decline between years 1800 and 1995 and species that are of global conservation concern. The 67 species on the Red List are of the most urgent conservation concern.
- A1.1.29 Species listed on the BoCC Amber List, of which there are currently 96, include those that have shown a moderate decline in numbers (25%-49%) over the last 25 years and those with total populations of less than 300 breeding pairs. Also included are those species which represent a significant proportion (greater than 20%) of the European breeding or wintering population, those for which at least 50% of the British population is limited to ten sites or less, and those of unfavourable conservation status in Europe.
- A1.1.30 The remaining species are placed on the Green List, indicating that they are of low conservation priority. These species still receive full protection through the provisions of the Wildlife and Countryside Act 1981, as amended.

### Wintering Bird Surveys

- A1.1.31 The wintering bird surveys were based on a transect survey methodology as detailed in Bibby *et al.* (2000) and Gilbert *et al.* (1998).
- A1.1.32 The survey area (Project site boundary) is shown on Figure 2.4.1.
- A1.1.33 The transect route was selected to include all field boundaries and to visit all areas of the Project site to within 200 metres, where possible. Visits were undertaken early in the morning.
- A1.1.34 On each visit the route was walked at a slow pace with start and finish times noted. All birds seen and heard were recorded directly onto an ArcGIS base map using ESRI software on hand-held PDA devices, with a 1:10,000 scale Ordnance Survey base map of the study area (and adjacent land). A fresh map was used for each survey. Registrations of birds were recorded using standard BTO two letter species codes.
- A1.1.35 All bird species were recorded and mapped across the whole Project site, where accessible.

A1.1.36 Surveys for wintering birds were undertaken between October 2018 and March 2019. A total of five survey visits were undertaken, each over two consecutive days. The survey visits were as follows:

- Visit 1: 30 and 31 October 2018;
- Visit 2: 22 and 23 November 2018;
- Visit 3: 19 and 20 December 2018;
- Visit 4: 23 and 24 January 2019; and
- Visit 5: 20 and 21 March 2019.

A1.1.37 An assessment of the ornithological importance of the survey area during the winter period was made by evaluating the species recorded against the following criteria:

- Annex 1 of the EU Birds Directive;
- UK BAP priority bird species;
- NERC Species of Principal Importance; and
- BoCC Red and Amber Lists (Eaton *et al.*, 2015).

A1.1.38 Reference is not made to species afforded special protection under Schedule 1 of the Wildlife and Countryside Act (1981) as the protection measures only apply to these species during the breeding season.

### Reptile Surveys

A1.1.39 The reptile survey followed the recommended methodology described in the Herpetofauna Worker’s Manual (Joint Nature Conservation Committee (JNCC), 2003) and Froglife’s Surveying for Reptiles (Froglife, 2016). It was undertaken by experienced ecologists and was conducted in areas of the site identified as containing the most favourable habitat for reptiles.

A1.1.40 Reptiles are best surveyed from April following hibernation until June and then again in September. At this time of year, the sun is often shining but air temperatures are low, so reptiles spend a long time basking and are therefore more easily observed.

A1.1.41 The reptile survey was conducted using artificial refugia made from corrugated tin or roofing felt measuring 50 cm x 50 cm and 50 cm x 100 cm. These provide shelter and basking opportunities for reptiles which can be recorded on or under the refugia in suitable weather conditions.

A1.1.42 On the 26-28 March, 29 May and 7 August reptile refugia were placed in areas identified as providing the greatest suitability for reptiles and which had optimal basking opportunities. The locations of the refugia are shown on Figures 3.6.1a – 3.6.1e.

A1.1.43 The refugia were left undisturbed for ten days prior to the first survey being undertaken in order to allow them to bed down and to give them time for reptiles to find them. In order to conform to best practice guidelines, the refugia were inspected on seven separate survey visits and a visual search was undertaken when the refugia were being laid.

A1.1.44 On each of the visits every refugia was inspected for reptiles basking on top and was then lifted to identify any reptiles beneath. The number, species, age class and where possible, sex of each reptile observed were recorded.

A1.1.45 Visit times were selected to coincide with suitable weather conditions and times of day when refugia would be acting as heat traps which would attract reptiles to use them whilst basking. Periods of strong wind or heavy rain were avoided, and surveys were typically undertaken during periods of sunshine and when air temperatures were between 10°C and 18°C.

A1.1.46 Froglife (1999) provides a basic index of relative abundance of reptiles based on peak survey counts (Table 1 of the guidance). The figures in Table 1 of the guidance refer to the maximum number of adults seen by direct observation and/or on or under refuges by one person in one day.

### Great Crested Newt Surveys

#### Habitat Suitability Index (HSI)

A1.1.47 HSI assessments of all ponds within 250 metres of the Project boundary were undertaken where access was allowed.

A1.1.48 An HSI is a numerical index, between 0 and 1 where 0 indicates unsuitable habitat and 1 represents optimal habitat.

- <0.5: poor;
- 0.5 – 0.59: below average;
- 0.6 – 0.69: average;
- 0.7 – 0.79: good; and
- >0.8: excellent.

A1.1.49 The HSI methodology for great crested newts has been developed to assess the suitability of ponds for use as breeding sites. The assessments were made in accordance with the methodology set out in Advice Note 5 published by the Amphibian and Reptile Group UK (ARGUK) (ARGUK, 2010).

A1.1.50 The HSI incorporates ten suitability indices, all of which are factors thought to affect the likelihood of great crested newt

presence. The ten indices are location, pond area, pond drying, water quality, shade, waterfowl, fish, other ponds within 1 km, terrestrial habitat and macrophytes.

### eDNA

- A1.1.51 The surveys were conducted by great crested newt license holders. The surveys followed the eDNA surveying and laboratory analysis techniques as described by Biggs *et al.* (2014).
- A1.1.52 Water samples were collected using sampling kits supplied by NatureMetrics Ltd and Surescreen.
- A1.1.53 Surveyors collected 30ml water samples from the waterbodies using a sterile ladle. Surveyors collected the sample from the bank edge and did not enter the water.
- A1.1.54 Where access allowed, the samples were collected from points evenly spaced along the waterbody. Samples were spread out as much as possible to ensure a representative sample was collected and to ensure the effectiveness of the survey was not compromised.
- A1.1.55 The surveyors used the ladle to gently agitate the water to mix the water column, whilst taking care not to disturb and collect any sediment. The samples collected were emptied into a sterile plastic bag and homogenised by gently shaking the bag to ensure eDNA was evenly mixed through the sample.
- A1.1.56 A pipette was used to collect 15 ml subsamples of the pond water from the bag into sterile tubes already containing 35 ml of ethanol to preserve the eDNA sample.
- A1.1.57 The samples were then removed from site and sent to NatureMetrics Ltd or Surescreen for analysis. The water samples were analysed using the quantitative Polymerase Chain Reaction (qPCR) eDNA test.
- A1.1.58 Biggs *et al.* (2014) has demonstrated the effectiveness of eDNA in the detection of great crested newt. In detailed field studies eDNA detected great crested newt 99.3% of the time in ponds where they were known to occur.

### Bottle Trapping

- A1.1.59 Bottle traps constructed from 2 L plastic drinks bottles supported on bamboo canes were located at approximately 2 metre intervals around the edge of each pond. On each survey visit traps were set out before dusk and were emptied and removed the next morning before 10 am. Traps were always placed ¾

submerged so that they contained at least ¼ air; they also had air holes in the exposed ends. The species, number and sex of newts captured in the traps were recorded and the newts were carefully released back into the pond from which they were caught.

### Torch Survey

- A1.1.60 The shoreline of each water body was scanned after dusk using a high powered torch of 1,000,000 candlepower. The perimeter and centre of the pond were slowly scanned with the torch and the number, and where possible sex, of any amphibians seen was recorded.

### Egg Searching

- A1.1.61 Egg searches were undertaken by searching for folded leaves on marginal and aquatic vegetation around the perimeter of a pond and carefully opening them up to reveal newt eggs.
- A1.1.62 The eggs of great crested newts can be distinguished from those of other species by their size and colour.

### Dormouse Surveys

- A1.1.63 A dormouse nest tube survey was undertaken based on the methodology and best practice guidelines set out in the dormouse conservation handbook, second edition (Bright, Morris and Mitchell-Jones, 2006).
- A1.1.64 A total of 684 dormouse tubes were set out on 1–8 April and 22 May 2019. They were tied to suitable vegetation around the site following standard survey guidelines (English Nature 2006) to provide nesting opportunities for any dormice present (Figure 3.8.1a and b). During the 2022 surveys 179 tubes were set out in total, 162 of which were placed along the A23 corridor.
- A1.1.65 Survey visits were undertaken regularly in suitable weather conditions between May and June, with additional monthly visits until November 2019. Following the table of probability of finding dormice in the Dormouse Conservation Handbook (Table 5 – English Nature, 2006) and assuming the tubes are in situ until September, this gives a score in excess of 20 which is considered an acceptable level of survey.

- A1.1.66 Further surveys were carried out in 2022, on the 27-28th May, 28-29th June, 18-19th July, 23-24th August, 20-21st September, 20th, 24th October, 28th, 29th and 30th November 2022. No signs of dormouse were recorded during these surveys.

## Aquatic Mammal Surveys

### Otter Survey

- A1.1.67 The otter survey was undertaken with regard to the methodology described in the Design Manual for Roads and Bridges, Volume 10 Section 4, Part 4 (Highways Agency *et al.*, 2001). The methodology was developed for linear schemes which are likely to affect otter habitats or populations but was adopted for this site.
- A1.1.68 The suitable areas along the River Mole were walked and examined in detail for evidence of the presence of otter in the form of characteristic field signs.
- A1.1.69 The following field signs were searched for:
- spraints;
  - prints;
  - holts and couches;
  - slides and runs; and
  - feeding remains.

### Water Vole Surveys

- A1.1.70 Although they do not hibernate, water voles are not very active above ground during the winter, so surveys are best carried out between April and October when field signs are most abundant.
- A1.1.71 The survey was carried out in accordance with guidelines of best practice set out in the Water Vole Conservation Handbook – Third Edition (Strachan *et al.*, 2011).
- A1.1.72 The suitable areas along the River Mole were walked and examined in detail for evidence of the presence of water vole in the form of characteristic field signs.
- A1.1.73 Wherever possible, the banks were inspected on both sides, from the water's edge to the top of the bank.
- A1.1.74 American mink is a non-native, invasive species listed under Section 9 of the Wildlife and Countryside Act 1981 (as amended). They predate on water voles and are considered to be to one of the main reasons for the dramatic decline in the size of the water vole population in the UK. Therefore, incidental signs of this species observed during the survey were also noted.
- A1.1.75 The following water vole field signs were sought during the survey work.



## Field Signs

### Latrines

- A1.1.76 Droppings are the most distinctive field sign. These are about 8-12 mm long and 4-5 mm wide, cylindrical and symmetrical with blunt ends. Colouration varies from green, brown, black and even purple, depending on what food has been eaten and its water content. They have the texture of putty when fresh but when dry may show green concentric rings of fine plant material if broken open. Rat droppings are always larger than water vole droppings and have an unpleasant odour. Most droppings are deposited at discrete latrine sites near the nest, at range boundaries and where they enter and leave the water. Latrines are established and maintained from February to November. Scent from the lateral flank glands is deposited on the latrine when the vole drum marks with its hind feet, so that many latrines often show a flattened mass of old droppings, topped with fresh ones.

### Feeding Stations

- A1.1.77 Food items are often brought to favoured feeding stations along voles' pathways or at haul-out platforms along the water edge. These show feeding remains as a neat pile of chewed lengths of vegetation. The sections are typically up to 10 cm long showing the marks of the two large incisors and are quite good field signs of the presence of water vole. These chopped sections of vegetation are often taken into the burrow entrances by the voles and laid up as stores along the tunnels or in chambers.

### Burrows

- A1.1.78 Water vole burrow entrances are typically wider than high with a diameter of between 4-8 cm. At the water's edge the entrances may occasionally appear larger due to erosion, but the tunnel soon contracts down to the size of two fingers. Externally the burrow system appears as a series of holes along the water's edge, some at or just above the water level on steep banks, some opening below the water line and others occurring within the vegetation up to five metres from the water's edge (for access to food and for ventilation). At the water's edge spoil excavated from the burrows tends to be washed away while those burrows opening high on the bank are probably dug from underground as no spoil can normally be found around them.

### Lawns

- A1.1.79 Around burrow entrances on land, grazed 'lawns' can be found. These frequently occur when the female is nursing young and time away from the nest is kept to a minimum. The female grazes the vegetation short to the ground within easy reach of the hole; often by not fully leaving the hole and being wary to dart back should danger threaten.

### Nests

- A1.1.80 Both males and females take bedding underground to line nest chambers in the burrow system. Nurseries consist of a large ball of finely shredded grasses or reeds and the chamber entrance may be plugged by the female with loose soil or grass. Where vegetation cover is dense and the water table is high, nests roughly the same size and shape as a rugby ball can sometimes be found above ground, often woven into the bases of rushes, sedges or reed.

### Footprints

- A1.1.81 Although footprints may be readily found along the soft margin of a watercourse (of many species besides water voles) they are not the easiest field sign to use. Large adult water vole tracks will appear very similar to those of juvenile rats. As with all rodents, the imprints show four toes in a star arrangement from the forefoot and five toes of the hind foot with the outer ones splayed, but often the tracks of the hind feet partially overlap those of the fore. The hind foot typically measures between 26-34 mm and is noticeably smaller than that of the Common Rat at 40-45 mm (heel to claw measurements). The Brown Rat is also heavier and so leaves a deeper impression.

### Runways in Vegetation

- A1.1.82 These are most often found within 2 metres of the water's edge and take the form of low tunnels pushed through the vegetation. Pathway width may be 5-9 cm and they often branch many times, leading to the water's edge, burrow entrances or favoured feeding areas. Rat runs on the other hand are usually very obvious as clear or bare pathways linking burrows and often running along the bank away from the water's edge.

### Assessing Population Size

- A1.1.83 Water voles live in colonies, but distribute themselves along a watercourse through a series of contiguous territories. Breeding

female water voles are territorial but may share territory with their female offspring. Males have home ranges which overlap with the territories of a number of females and other males. A female's territory length typically varies between 30 metres to 150 metres and a male's home range from 60 metres to 300 metres.

- A1.1.84 The number of water vole latrines present gives an indication of the strength of the water vole colony. Approximately six latrines are maintained per breeding female. However, larger and more robust populations show a large number of closely packed latrines. Typically, fewer maintained latrines are present when water vole populations are small and fragmented.

## Preliminary Bat Roost Assessment

### Buildings

- A1.1.85 An assessment of the suitability of the buildings for bat roosting potential, within the landside and airside areas of the Project site, was undertaken at the same time as the Phase 1 Habitat Survey on 18th–22nd March and 10th & 11th July 2019
- A1.1.86 The survey included a thorough, ground level inspection of the exterior of all accessible buildings and the features of the building listed below were noted:
- type;
  - age;
  - wall construction, in particular the type of material used;
  - form of the roof, in particular the presence of gable ends, hipped roofs etc and the nature and condition of the roof; and
  - the general condition of the building.
- A1.1.87 The above information would inform the potential for roost features to be present and identify potential bat access points and roost places and field signs of bats being present.
- A1.1.88 When suitable features were identified, they were inspected for signs indicating use or possible use by bats including tiny scratches, staining and flies around the entry points, bat droppings and feeding remains in, around and below entrances, distinctive smell of bats and the smoothing of surfaces around cavities.
- A1.1.89 Guidance from the Bat Conservation Trust (BCT) (2016) on the features of buildings which correlate with their use by bats was considered.

A1.1.90 Preliminary bat roost assessments of buildings can be carried out at any time of year; however, summer surveys are more likely to reveal signs of bat activity.

### Trees

A1.1.91 A ground-level Preliminary Bat Roost Assessment (PBRA) of trees along the A23 within the site boundary was undertaken in November and December 2022.

A1.1.92 Trees were assessed as having the potential to support bat roosts if they had features such as holes, cavities, split/broken limbs, trunk hollows, knot holes, flaking bark and woodpecker holes.

A1.1.93 The suitability of the trees for roosting bats was also assessed by examining the surrounding habitat. Important habitat features surrounding a tree which may influence bat roost potential include whether the tree is in a semi-rural or parkland location, its proximity to significant linear habitat features such as a watercourse, mature hedgerow, wooded lane or an area of woodland.

A1.1.94 When suitable features were identified, they were inspected for signs indicating use or possible use by bats including tiny scratches, staining and flies around the entry points, bat droppings and feeding remains in, around and below entrances, distinctive smell of bats and the smoothing of surfaces around cavities.

A1.1.95 Guidance from the Bat Conservation Trust Bat Survey: Good Practice Guidelines (BCT, 2016) on the features of trees which correlate with their use by bats was considered.

A1.1.96

### Bat Emergence/Re-entry Surveys

A1.1.97 In order to comply with best practice guidelines (Collins, 2016) emergence surveys were carried out on any buildings considered to have bat roosting potential. Surveys were undertaken between May–October 2019. The aim of these surveys was to determine the use of the buildings (if any) by roosting bats, the species assemblage within the Project site and the egress locations of any bats emerging from the buildings.

A1.1.98 Observations were made outside the buildings from where it was considered bats might emerge. The dusk survey commenced 15

minutes before sunset, and lasted for approximately 90 minutes, to record any bats that may emerge from the buildings.

A1.1.99 During each survey visit, the building was continuously surveyed by up to three experienced ecologists and visual observations were made of where any bats emerged/re-entered and in what direction they were flying to or from. Behavioural observations were also recorded for any bats encountered on site or within the vicinity, including direction of flight and activity observed eg foraging or commuting.

A1.1.100 Elekon Batlogger and Anabat bat detectors were used to record bat echolocation calls of any emerging bats and identify species, where possible.

A1.1.101 Bat activity can be strongly dependent on weather conditions; therefore, the surveys were only carried out in favourable conditions when bat activity was deemed to be likely (sunset temperature 10°C or above, no rain or strong wind).

### Bat Activity Transect Surveys

A1.1.102 A total of five transect routes were devised to cover a broad range of the habitat types present on site but focusing on those likely to be of greatest value to bats, including woodland, woodland edges, river corridors and open grassland. A brief overview of each transect route is as follows.

#### Transect 1

A1.1.103 Transect 1 was devised to include potentially high value habitat in the south eastern part of the Project site (within the airport's biodiversity area), including the old and new lagoons associated with the Crawley Sewage Treatment Works (hereafter referred to as 'the sewage works') and immediately surrounding woodland areas, including Upper Pickett's Wood to the south east and Horleyland Wood to the north west.

A1.1.104 The lagoons and open water habitats were considered to provide good opportunities for a wide variety of bats to forage, as was the woodland. Additionally, the woodland habitat would provide foraging and roosting opportunities for a wide variety of species associated with this habitat type, including pipistrelle and *Myotis* species.

A1.1.105 The transect route also incorporated the corridors between woodland compartments, including those to the north of the Crawley Sewage Treatment Works and adjacent to the Long Stay

South Car Park. These areas would likely be used for foraging and commuting bats.

#### Transect 2

A1.1.106 Transect 2 focussed on areas of potentially high value habitat, immediately to the south of Transect 1 and the Crawley Sewage Treatment Works.

A1.1.107 The transect route covered large areas of open grassland to the east of the railway corridor, the woodland edges and connecting habitat which linked woodland corridors and linear features, such as the Gatwick Stream and railway line, to the Crawley Sewage Treatment Works and Horleyland Wood.

A1.1.108 These linear features were considered to potentially provide good commuting corridors between woodland compartments, further linking to the wider area. The large areas of open grassland would likely provide foraging opportunities for larger species of bats, such as noctule, Leisler's and serotine bats which typically exploit these types of habitats.

#### Transect 3

A1.1.109 Transect 3 covered areas of potentially high value habitat in the north eastern part of the Project site.

A1.1.110 The transect included a narrow strip of woodland between the railway line and A23 London Road, heading north towards the main trunk road leading to the M23. From there, the transect covered Riverside Garden Park, which included areas of both dense and open canopy woodland, grassland and a large pond in the south of the park. Gatwick Stream ran adjacent to the transect route along the north eastern edge of the Project site boundary.

A1.1.111 The habitats present along the transect route were considered likely to provide a range of opportunities for both woodland and larger bat species to forage. The woodland edges, railway corridor and stream would likely provide linear features for bats to commute along to access areas of high value habitat to the north and south of the Project site boundary, including the woodland compartments adjacent to Crawley Sewage Treatment Works.

#### Transect 4

A1.1.112 Transect 4 covered an area in the southern part of the site, adjacent to buildings and car parks (X and Z) associated with the airport.

A1.1.113 The transect also included small strips of woodland located between the A23 London Road and Perimeter Road South, and between the staff car park and Charlwood Road. Part of the transect incorporated the length of Crawter's Brook which sits between Perimeter Road South and the southern boundary of the airport.

A1.1.114 The transect route aimed to incorporate habitat which could potentially be used as commuting corridors between different woodland compartments, linking to further suitable habitat within the wider landscape.

#### Transect 5

A1.1.115 Transect 5 covered a broadly open area of habitat, Land East of the Aviation Museum, in the north west of the Project site.

A1.1.116 The transect incorporated areas of potentially high value habitat such as mature hedgerows and tree lines, Man's Brook to the north of the Project site and the River Mole and woodland corridor along the east of the transect route. The River Mole and woodland corridor further linked to large areas of woodland to the north and south of the Project site, including Brockley Wood.

A1.1.117 The woodland corridor to the east of the transect route would likely provide strong commuting links, for all bat species, to areas of suitable foraging and roosting habitat within the wider area, particularly to the open and diverse mosaic of habitat to the north west of the airport. The large areas of open grassland within the Project site boundary would likely provide foraging opportunities for larger bat species, such as noctule, Leisler's and serotine bats which typically exploit these types of habitats.

#### Transect 6

A1.1.118 Transect 6 focussed on potentially high value habitat south of Brockley Wood, which is surrounded by the River Mole to the south and west.

A1.1.119 The transect route covered the more open areas of grassland that surround Brockley Wood and runs next to the River Mole. The transect followed the southern woodland edges until Man's Brook where the route then followed the meandering section of the River Mole.

A1.1.120 The River Mole and Brockley Wood potentially provide good commuting corridors which could extend to the whole northwest of the airport, further linking to the wider area. Further the open water habitat is likely to be utilised by bats for foraging. The

woodland habitat also provides opportunities for foraging bats, as well as providing roosting opportunities for a wide variety of species associated with this habitat, including pipistrelle and *Myotis* species.

#### Transect 7

A1.1.121 Transect 7 was located around the potentially high value habitat of Riverside Garden Park and along the woodland that surrounds the A23.

A1.1.122 The route followed the more open grassland that is surrounded by woodland within the park, the woodland edges next to the A23, ending next to the Gatwick Stream and man-made lake.

A1.1.123 The open water within Riverside Garden Park provides potential foraging habitat for bats. The woodland habitat is likely to provide roosting potential, alongside the more open grassland that supplies further foraging habitat. Further the woodland corridor bordering the A23 provides good commuting opportunities, which also connects to the wider area.

#### Transect 8

A1.1.124 Transect 8 focussed on the woodland patches and hedges that surround the industrial area next to the Airport Way roundabout.

A1.1.125 The route followed the tree line on the north side of the M23, the woodland patches around the urbanised areas which includes passing a small lake and ending within a more open grassland habitat with corridors of trees.

A1.1.126 The grassland habitat provides open foraging space. The woodland patches are likely to provide roosting potential. The woodland patches also provide a good commuting corridor to allow bats to access the wider habitat. The small lake is also likely to provide foraging opportunities.

#### Transect 9

A1.1.127 The Dairy Farm transect was centred around areas of potentially high value habitat of Church Meadows, and the land immediately to the west where the River Mole is also present.

A1.1.128 The transect route covered the areas of open grassland to the east and west of the River Mole. The route followed the treelines along each field boundary and the most northern boundaries, where the trees meet the A23.

A1.1.129 The treelines and hedges within Church Meadows and the surrounding areas potentially provide areas suitable for commuting bats, which would allow them to further access the wider area. The meadowland and the River Mole would likely provide valuable foraging habitat for various species of bat. The trees could also provide roosting opportunities.

#### Methods

A1.1.130 Each transect was surveyed twice per month between April-October. Each visit commenced at least fifteen minutes before sunset and continued until at least two hours after sunset.

A1.1.131 On each visit, two ecologists walked along the transect at a steady speed starting at opposite points of the transect to ensure the full transect route was covered and to reduce bias associated with levels of bat activity at particular times of day/night. Several spot-sampling locations were included, distributed evenly along the transect route. The surveyors stopped at each of these points for between three and five minutes and recorded all bat activity seen and/or heard.

A1.1.132 Visual observations for bats were undertaken by scanning the skyline, and bat detectors were used to listen to and record echolocation calls. Elekon Batlogger and Anabat bat detectors were used, and recordings were made. For any bats encountered, notes were made on location, species or species group, behavioural observations (eg direction of flight, habitat) and activity heard (eg feeding buzzes or social calls).

A1.1.133 Bat activity can be strongly dependent on weather conditions; therefore, the surveys were only carried out in favourable conditions when bat activity was deemed to be likely (sunset temperature 10°C or above, no rain or strong wind).

#### Bat Static/Automated Surveys

A1.1.134 Elekon Batlogger A units were deployed across the site between April-October for a minimum of five nights. The units were positioned at various locations in order to sample a broad range of the habitat types present on site but focusing on those likely to be of greatest value to bats.

A1.1.135 The automated bat detectors were programmed to commence recording approximately 15 minutes before sunset and terminating 30 minutes after sunrise. This period covered the peak time bats would be commuting to and from their roosts.

### Bat Data Analysis

- A1.1.136 The bat passes were recorded, and all bats were identified to species level, where possible, on site. Echolocation calls were subsequently analysed using computer software (BatExplorer and Kaleidoscope) for confirmation of species. Where possible, additional notes on size, flight height, type of flight (such as commuting, foraging, fast or slow) and direction of flight were also recorded.
- A1.1.137 All sound files were subject to manual analysis by an experienced bat ecologist. Where possible, identification was carried out to species level. Bats of the species group *Myotis* and long-eared bat species are difficult to distinguish and therefore calls were only analysed down to species level where they were characteristic of that species and present within suitable habitat.
- A1.1.138 The number of bat passes recorded is not representative of the number of bats present within any given area, as a single bat may have made many passes. Therefore, descriptions of bat species assemblage represent the minimum number present rather than a definite list of all species present.
- A1.1.139 Where several bat species were present within a call segment, then all the species were tagged in the results spreadsheet. For example, a common pipistrelle, soprano pipistrelle and *Myotis* bat all calling simultaneously would result in three individual bat registrations for calculating bat pass counts.

### Bat Crossing Point Surveys

- A1.1.140 The methods generally follow the standard best practice (Berthinussen and Altringham, 2015). Any specific deviations due to objectives of the surveys are detailed where necessary.
- A1.1.141 Crossing Point surveys were undertaken at two locations, River Mole corridor and Riverside Park, in August 2020, September 2020, May 2021 and June 2021. Table A1.1.2 summarises survey dates and locations (easting and northing) for both crossing point locations.

**Table A1.1.2: Crossing Point Locations**

Crossing Point location	Easting	Northing	Dates of surveys
River Mole A	525699	140500	18/08/2020, 01/09/2020, 05/05/2021 (not valid due to unsuitable weather), 18/05/2021 and 01/06/2021
River Mole B	526002	140589	18/08/2020, 01/09/2020, 05/05/2021 (not valid due to unsuitable weather), 18/05/2021, 01/06/2021
Riverside Park A	527619	142393	19/08/2020, 22/09/2020, 06/05/2021, 02/06/2021
Riverside Park B	527629	142392	19/08/2020, 22/09/2020, 06/05/2021, 02/06/2021

- A1.1.142
- A1.1.143 The locations were selected using the results of trapping and radio-tracking surveys undertaken in 2019, which recorded Bechstein's bats flying along the River Mole and foraging within Riverside Park as well as due to potential impacts to the areas in relation to a new flood mitigation strategy and North Terminal Junction improvements.
- A1.1.144 The River Mole runs alongside a bare ground vehicle track and presents steep vegetated embankment at this crossing point location. Wildlife netting was present on both embankments at both locations A and B. It had also been installed across the river corridor at location A.
- A1.1.145 Riverside Park is an area of public open space comprising broadleaved woodland with grassland glades and paths. It is bounded by Gatwick Stream which runs south-east to north-west. The crossing point is located in the north-west of Riverside Park along a public footpath, with woodland to the north, east and south, and Gatwick Stream to the west with its vegetated steep embankments covered in tall ruderals.
- A1.1.146 The Defra research report (Berthinussen and Altringham, 2015) recommends undertaking a preliminary dusk and dawn survey at each location to determine if a feature can be considered a flight

path when certain conditions are met. These conditions include that when surveying the feature at dusk and dawn on the same night more than either 10 bats of common species or five bats of rarer species must be recorded using the habitat feature per survey to consider it to be a flight path. After the first dusk visit these conditions were met and therefore it was considered that the lack of data at dawn is not a constraint to the survey results.

- A1.1.147 Visits to each crossing point comprised observing bats at dusk, with surveys commencing 15 minutes before sunset and continuing for 120 minutes after sunset as advised by the Defra research report (Berthinussen and Altringham, 2015) when woodland-adapted bat species are present within the area.
- A1.1.148 Two surveyors monitored each crossing point, one at either side of the habitat feature used by commuting bats.
- A1.1.149 Each surveyor was provided with a Elekon Batlogger M full spectrum detector and with a thermal camera (FLIR T1020).
- A1.1.150 The ultrasonic bat detectors were set to automatically record ultrasound between 13 and 155 kHz and signals were digitised at a rate of 312 kHz with 16-bit sampling depth. This allowed for the recording of any bat passes which were in close proximity to the surveyor.
- A1.1.151 The thermal equipment was used with a combination of 45 degree lenses, thermal sensitivity <20mK at 30°C and an infrared resolution of 1024 x 768 pixels allowing for a maximum detection distance for a bat in flight to be 104 metres (Fawcett-Williams, 2019).
- A1.1.152 The data in radiometric format captured during the surveys was stored on an SD card inside the camera. Radiometric data allows analysis (Flir Tools) of thermal patterns in the images through thermal tuning and the use of different colour palettes. The object of interest can then be enhanced through thermal tuning with non-target objects falling outside the scale (Infrared Training Centre, 2017). The colour palettes allow different colours to be assigned to mark specific temperature levels. For this study, high contrast palettes were utilised to enhance small temperature

<sup>1</sup> A single bat pass is defined as one or more clearly recognisable echolocation calls from a single species, separated from the next one by a gap of at least a second (Berthinussen and Altringham, 2015).

differences and improve the detectability of small moving objects (bats) against a varied background.

- A1.1.153 Each surveyor/analyst recorded direct observation of bats, their species (where this could be accurately determined) and their flight behaviour, ground-level distance from the feature and height above the ground when observed. The closest distance the bat came to the feature was recorded, and for flight height during crossing, the lowest height was recorded. Incidental records of bat activity near the surveyor locations were also collected. Each passing bat was recorded as a separate observation, regardless of whether the same bat had clearly passed the surveyor more than once.
- A1.1.154 Each pass was assigned to species either by the surveyor in the field, or by matching recordings of the passing bat whether on the proforma, or the time the bat was observed during analysis. Recordings were analysed post-survey to determine the bat species they represent using Elekon BatExplorer. The output from sound analysis was subsequently checked against the surveyor's identification, and changes made were necessary, favouring the identification made manually.
- A1.1.155 For instances where recordings revealed more than one bat species present (eg a *Myotis* sp. and a *Pipistrellus* sp.) passes were included for each, but flight behaviour data was left as unknown for those which were not observed by surveyors/analysts.
- A1.1.156 The output of sound and radiometric analysis was subject to a quality assurance process; a minimum of 10% of the sound files identified to each species / genus and a minimum of 30 minutes of radio-metric data were verified by a principal grade analyst using BatExplorer or Flir Tools software.
- A1.1.157 Data for each crossing point were categorised and presented the following information for each survey and location:
  - Total number of passing bats observed;
  - Total number of passing bats observed per species;
  - Number of bats using (passing within 5m distance of the feature) and not using (passing further than 5m distance) the habitat feature;

- Number of bats flying at a 'safe' height at Riverside Park and therefore not at risk of collision, defined as passing at a height higher than 5 metres from the ground and based on the assumption that the proposed road improvements will be at the current height of the ground and any impacts from traffic would be within 5 metres height;
- Number of bats using the River Mole (passing within 5m distance of the feature) at a height below or equal to ground level (within the river corridor), at a height between ground level and 5 metres (directly above the river corridor) and at a height above 5 metres<sup>2</sup>; and
- Species of bats heard but not seen.

A1.1.158 Safe and unsafe height are defined with reference to the maximum height for a heavy goods vehicle on UK roads which is 4.95m. It has been rounded to 5m for the purposes of analysis. Bats passing above this threshold were considered to not be at risk of collision.

#### Assessment of importance

A1.1.159 The value of the River Mole and Riverside Park as commuting routes and foraging areas for bat species have been calculated using Wray et al. (2010) where the rarity of the species involved, the approximate numbers of bats using them (based on crossing point survey data), the proximity of known roosts, and the nature and complexity of linear features in the landscape are all taken into account. The importance of the areas are assessed at a geographical level eg local, regional, national.

A1.1.160 It was not possible to carry out this assessment for species groups as rarity of single species within the groups varies greatly and sound analysis does not allow separation of species on the basis of call parameters. However, an assessment was made for Bechstein's bats using crossing point survey data for *Myotis* species in conjunction with radio-tracking survey data from 2019, 2020 and 2021.

#### Veteran Trees

A1.1.161 The survey assesses individual trees and groups of trees for quality and benefits within the context of proposed development. The quality of each tree or group of trees has been recorded by

allocating it to one of four categories in general accordance with the requirements set out in BS 5837:2012:

- A1.1.162 A: Trees/Vegetation of high quality and value
- A1.1.163 B: Vegetation of moderate quality and value
- A1.1.164 C: Trees/Vegetation of low quality and value
- A1.1.165 U: Those in such a condition that they cannot realistically be retained as living trees in the context of the current land use for longer than 10 years

A1.1.166 When assessing any tree's potential category, its possible status as a veteran tree is also considered. This is done following guidance from the Ancient Tree Forum and Natural England:

A1.1.167 'Ancient and Other Veteran Trees: further guidance on management' Published by The Ancient Tree Forum and edited by David Lonsdale in 2013.

A1.1.168 'Veteran Trees: A guide to good management (IN13)' published by Natural England on 1 February 2000.

A1.1.169 Although there is not a clearly defined, measurable definition within this guidance of what a veteran tree is, they are generally considered to be both large in girth and of a condition that indicates they are in decline.

A1.1.170 Government guidance suggests that a buffer zone can be used to protect veteran trees:

A1.1.171 'A buffer zone around an ancient or veteran tree should be at least 15 times larger than the diameter of the tree. The buffer zone should be 5m from the edge of the tree's canopy if that area is larger than 15 times the tree's diameter'.

#### Methodology

A1.1.172 When surveying potential veteran trees, the two key features used to identify them are a large girth & a declining condition that raises their ecological value. These two attributes must both be present before it is classed as veteran.

<sup>2</sup> This analysis is related to the proposed diversion of the River Mole and increase use of aircraft which would have potential impacts on commuting and foraging bats.

A1.1.173 Girth- When considering a tree for veteran status the girth of the tree is the most obvious and measurable indicator of a tree being advanced in age.

A1.1.174 Where possible, the girth of a tree is also compared to the “Ancient Tree Size Criteria Table” (Lonsdale, ATF 2013)

A1.1.175 If the girth of a tree is considered large enough that the tree may be notable, then the condition of the tree will be assessed to confirm whether or not the tree will be classed as a veteran.

A1.1.176 Condition- The main determining factor in whether or not a tree can be classed as a veteran or not, is its physical condition. In ‘Ancient and Other Veteran Trees: further guidance on management’ David Lonsdale states that:

A1.1.177 ‘In order to qualify as a veteran, the tree should show crown retrenchment and signs of decay in the trunk, branches or roots, such as exposed dead wood or fungal fruit bodies.’ (Lonsdale, ATF 2013)

A1.1.178 This makes the two most significant indicators of veteran status:

A1.1.179 Retrenchment: The progressive deterioration of the outer crown of a tree through dieback and limb loss that reduces the overall height/ spread of the tree and increases stability.

A1.1.180 Decay: When wood rots and decomposes; usually caused by a parasitic fungus or bacteria.

A1.1.181 Both of these defects create good habitat, for fauna such as invertebrates or birds, raising its ecological value and indicating veteran status.

A1.1.182 A tree may still be considered for veteran status even if it does not possess one or both of these features, if it still has a number of defects that would also be considered veteran features, such as, but not limited to:

- A large quantity of deadwood in the crown
- Hollows/ Cracks without signs of decay
- Bark Loss
- Saprophytic Fungi
- Habitat spaces (such as woodpecker holes)
- Storm Damage

### Data validity and limitations

A1.1.183 It is important to note that even where data are held, a lack of records for a defined geographical area does not necessarily mean that there is a lack of ecological interest; the area may be simply under-recorded. Bats are highly mobile animals and can move roost sites both within and between years.

## Annex 2

### Extended Results

## A2.1 Extended Results

### Phase 1 Habitat Surveys

#### Annex 2.1.1: Pond Descriptions

Gatwick Pond ID No.	Ecology ID number	Description
FCZ	P1	No access was granted to this pond.
9VG	P2	No access was granted to this pond.
Pond F	P3	A Large man-made attenuation pond, a barrier crosses the middle running north to south and marginal vegetation was present around all the sides.
SM7	P4	Small pond behind services with poor water quality and little aquatic vegetation.
981	P5	Large pond within a woodland. No aquatic vegetation and little woodland ground flora. Mature trees surround pond and a film of algae across pond.
Pond G	P6	Shallow pond with a silt bed that was mostly dry.
30Z	P7	Large pond within Horleyland Wood. Pond has shallow banks and marginal vegetation.
8N8	P8	woodland pond flooded over into surrounding woodland and cutting off footpath to rest of woodland
W46	P9	Small man-made wildlife pond, lots of aquatic vegetation along the southern bank. Banks relatively steep sided with a fallen tree across middle.
Old Lagoon	P10	Y shaped lagoon, manmade, amenity grass banks and steep sided.
E11	P11	Long linear settlement pond, wider at eastern end, linked to the M23 spur road. Reeds and bulrush dominated.
Pond E	P12	No access granted to this pond
A0A	P13	Pond within Police training area, swamped with willow and surrounded by woodland.
MHA	P14	Circular pond in the middle of the southern staff car park. The pond was surrounded with vegetation.
JCT	P15	Outside Project boundary, no access granted.
Pond A	P16	A pond located to the north of the runway near the fire training area. It was surrounded by dense bramble scrub and marginal vegetation such as pond sedge and bulrush
New Lagoon	P17	A circular sewage pond known as 'New lagoon' was a man-made, steep sided amenity grassland settlement lagoon.
Pond M	P18	Settlement pond east of the biodiversity wall. A manmade structure with steep concrete walls. Semi-improved grassland surrounded the pond. Only the eastern half of the pond held water.
WP9	P19	No access granted to this pond
AA20	P20	Awaiting further detail
AA21	P21	Awaiting further detail
K5F	P22	A long pond with 0.5m high banks around the northern side. The southern bank was covered with scrub and inaccessible. Around all sides there was a large amount of aquatic vegetation.
TTD	P23	A small circular manmade pond surrounded by willow and pine trees. Aquatic vegetation was present around the eastern, northern and southern sides. A concrete outflow was identified in the south east corner of the pond.
C24	P24	Large pond around 30 metres x 20 metres. Lots of marginal vegetation mainly bulrush, completely dry willow and ash growing around edge
Pond D	P25	A rectangular attenuation pond. It was concrete sided with an outflow into the Mole corridor.
Pond D	P26	A triangular attenuation pond made from concrete and steep sided. The pond was surrounded by managed grassland.
293	P27	Large open fishing lake in middle of public park. Two islands within middle of the lake densely covered with trees.
FFJ	P28	A small attenuation pond for the runway. Marginal vegetation was present here with rushes being dominant.
29A	P29	A long thin man-made channel. 5m high sides with a fence line around the top of it. Water was swamped by algae and had little aquatic vegetation, the banks were vegetated with tufted grass.
30P	P30	Murky shallow pond with clear animal tracks leading to it.
AVF	P31	A large pond within the Land East of the Aviation Museum covered with reeds and willowherb and algae topped. Nettle and willowherb ruderal surrounds the pond.



Gatwick Pond ID No.	Ecology ID number	Description
<b>Dog Kennel Pond</b>	P32	A small manmade attenuation pond with steep banks showing high levels of maintenance. A diverse mix of aquatic and marginal vegetation was found within the pond.
<b>Dog Kennel Pond</b>	P33	A small manmade attenuation pond with steep banks showing high levels of maintenance. A diverse mix of aquatic and marginal vegetation was found within the pond.
<b>AAA4</b>	P34	Newly created pond along Mole corridor.
<b>1WH</b>	P35	Pond in centre of eastern part of woodland had a small amount of water in with a heavy covering of duckweed.
<b>NU1</b>	P36	Pond in centre of eastern part of woodland had a small amount of water in with a heavy covering of duckweed. Linked to Pond 1WH
<b>A1 Ditch 1</b>		Ditch goes along the boundary of a small area of woodland. No water or aquatic vegetation present. Visible from adjacent parallel footpath to the south.
<b>A1 Ditch 2</b>		Field boundary ditch with trees and scrub on either side. No aquatic vegetation. Dry in places. Completely shaded. Visible from adjacent parallel footpath to the south.
<b>A1 Ditch 3</b>		Ditch on field boundary at the base of a wooded bank. Shaded by adjacent trees. No aquatic vegetation. Some stagnant water. Ditch continues west, parallel to the A23 but is mostly dry.
<b>Un-named additional ditch 1</b>		Additional ditch recorded under bridge.
<b>A1 Ditch 5</b>		Ditch parallel to the B2036. Wet with some aquatic vegetation. Visible along the side of the road.
<b>A1 Ditch 4</b>		Wet ditch at the base of an earth bank for the A23. No aquatic vegetation. Quite shaded by trees either side. Visible from the B2036.
<b>A1 services ditch</b>		Dry ditch parallel to the A23/Airport Way, at the bottom of the earth bank for the A23. Surrounded by trees. Access from adjacent footpath.
<b>A1 Services lake</b>		Large lake behind the services area. No aquatic vegetation. Wildfowl present, kingfisher seen twice. Banks are lined with trees. Access from adjacent footpath.
<b>A1 additional ditch by M23</b>		Wet ditch with reeds adjacent to the M23 slip road. Trees and scrub either side.
<b>A2 Ditch 5</b>		Wet ditch parallel to a railway embankment. Occasional adjacent trees and scrub. No aquatic vegetation. No direct access, viewed through fence.
<b>A2 SUDS area</b>		Large SUDS area in the middle of the car park. Surrounded by trees, scrub and grassland.
<b>A2 Ditch 6</b>		Dry ditch along the edge of some woodland. No aquatic vegetation. Also continues along the southern edge of the car park.
<b>A2 Ditch 7</b>		Wet ditch with some emergent vegetation. Surrounded by car parking with some adjacent trees.
<b>A2 Ditch 8</b>		Ditch surrounded by car parking. Partly wet at eastern end with some aquatic vegetation. Adjacent trees along south side of ditch.
<b>A2 Ditch 9</b>		Dry ditch along the edge of an access track with mature trees on its' northern bank. No aquatic vegetation.
<b>A2 Ditch 10</b>		Partly wet ditch along the edge of an access track. No aquatic vegetation. Mature trees present on the southern bank of the ditch.
<b>A2 additional ditch south of D10</b>		Additional ditch parallel to D10, to the south, along the edge of a car park. Partly wet but no aquatic vegetation.
<b>A2 Ditch 11</b>		There are two ditches here, either side of the road. Both are dry with no aquatic vegetation.
<b>A2 Ditch 12</b>		Wet ditch through a car park with adjacent mature trees and dense bracken. No aquatic vegetation.
<b>A2 Ditch 13 - West</b>		Dry ditch along a tree line. No aquatic vegetation.
<b>A2 Ditch 13 - East</b>		Wet ditch along a tree line with scrub either side. No aquatic vegetation.
<b>A2 Ditch 14</b>		Wet ditch surrounded by scrub and self-seeded tree saplings. No aquatic vegetation present.
<b>A2 Pond 1</b>		Large woodland pond with some emergent vegetation.
<b>A2 Pond 2</b>		Small woodland pond with over hanging trees. No aquatic vegetation present.
<b>A3 Pond 1</b>		New environmental pond. Lined. No aquatic vegetation visible apart from duckweed. Pond situated in an area of woodland, adjacent to a large wet meadow. Habitat connectivity to P2.
<b>A3 Pond 2</b>		Woodland pond surrounded by tussocky grassland and trees. No aquatic vegetation. GCN recorded in 2021 during bottle trapping.
<b>A3 Green Ditch 1</b>		Wet ditch within a large field of tussocky grassland / wet meadow. Some young willow trees growing either side of the ditch. No aquatic vegetation but surrounding habitat very good, with habitat connectivity to P2 with confirmed GCN presence.
<b>A3 Purple Ditch 1</b>		Ditch along the north side of a grass access track. Wet but no aquatic vegetation. A hedge is on the north side of the ditch.

Gatwick Pond ID No.	Ecology ID number	Description
<b>A3 Purple Ditch 2</b>		Ditch along the south side of a grass access track. Wet but no aquatic vegetation. A hedge is on the south side of the ditch.
<b>A3 Purple Ditch 3</b>		Ditch along the east side of a grass access track. Wet but no aquatic vegetation. A hedge is on the east side of the ditch.
<b>A3 Purple Ditch 4</b>		Ditch along the west side of a grass access track. Wet but no aquatic vegetation. Woodland is adjacent to the west side of the ditch.
<b>A5 Ditch 1</b>		Wet ditch along the edge of a grass field, with adjacent scrub and trees. No aquatic vegetation. Ditch is only adjacent to the southern field boundary, there is no ditch along the eastern field boundary as marked on the map.
<b>A5 Ditch 2</b>		Wet ditch with grass banks along the edges of a field. No aquatic vegetation.
<b>A5 Ditch 3</b>		Dry ditch with grass banks along the edge of a field. Adjacent trees and hedge. Visible from Aviation museum field to the west. Direct access via clay shooting school.
<b>A5 Pond 1</b>		No pond at the location shown on the map.
<b>A5 additional Pond 2</b>		Large pond with dense reeds surrounded by grassland.
<b>A5 additional Pond 3</b>		Small pond surrounded by trees and tussocky grassland. No aquatic vegetation.
<b>B2036 Ditch 1</b>		Unknown location, not labelled on map.
<b>B2036 Ditch 2</b>		Unknown location, not labelled on map.
<b>B2036 Ditch 3</b>		????
<b>B2036 Ditch 4</b>		????
<b>Crawter's wood- west</b>		No access available. This pond is airside so security access and a security escort need to be pre-arranged for access.
<b>Crawters wood- east</b>		No access available. This pond is airside so security access and a security escort need to be pre-arranged for access.
<b>Detention Centre Pond</b>		No access available.
<b>Ditch, North boundary, Car park X</b>		Partly wet ditch with bulrushes. Grass and ruderal vegetation on the banks. Surrounded by hard-standing.
<b>Ditch, Southern boundary, Car park X</b>		Wet ditch along southern edge of car park. Trees and scrub adjacent to the south. No aquatic vegetation.
<b>Ditch, Western boundary, Car park X</b>		Wet ditch along the edge of the car park with some aquatic vegetation and adjacent scrub.
<b>Pond, Car Park X</b>		Additional pond. In the centre of the car park. Surrounded by trees and grassland. No aquatic vegetation.
<b>North long stay A6/A7- Ditch 1</b>		Ditch runs through the centre of a car park. No connecting habitat. The ditch holds water but no aquatic vegetation is present. The banks are lined with self-seeded tree saplings and some scrub.
<b>North long stay A6/A7- Ditch 2</b>		Ditch runs along the southern and western boundaries of the car park. Wet, but no visible aquatic vegetation. Bordered by scrub and self-seeded tree saplings. More open along western boundary.
<b>North long stay A6/A7- Ditch 3</b>		Ditch runs parallel to southern edge of car park. Some water but no visible aquatic vegetation. Heavily shaded by adjacent trees and scrub.
<b>North long stay A6/A7- Ditch 4</b>		Ditch between two car parks. Some water but no visible aquatic vegetation. Heavily shaded by adjacent trees and scrub.
<b>North long stay A6/A7- Pond 1</b>		Woodland pond surrounded by trees and grassland. Some emergent vegetation present.
<b>Woodland behind Premier Inn-Ditch 1</b>		A8 South – Shallow wet ditch through an area of woodland. No aquatic vegetation present.

Gatwick Pond ID No.	Ecology ID number	Description
Woodland behind Premier Inn- Riverside Garden		A8 North – Ditch runs parallel to the A23 London Road, through an area of woodland. Dry in places. No aquatic vegetation. Access from within Riverside Gardens
A8 Island ditch		Island of land surrounded by road. There are a few metres of amenity grassland before a ditch, which encircles the western half of the island. The majority of the island is wooded. The ditch itself is deep but mostly dry, with no aquatic vegetation.
Pond 1		Large pond in the corner of an arable field with lots of emergent vegetation and adjacent trees.
Pond 4		Large pond in the corner of a grass field surrounded by trees. No aquatic vegetation is present.
Pond 5		Small pond in the corner of a grass field, surrounded by trees. No aquatic vegetation is present.
Pond 6		Large garden pond with extensive emergent vegetation. Surrounded by amenity grassland. Mature trees line the garden boundary.
Additional pond right next to the M23		Large drainage pond adjacent to the earth bank of the M23. Extensive emergent vegetation present. Trees and scrub are present along the northern edge of the pond.

**Annex 2.1.2: Protected or Notable Species identified during Botanical Survey**

Species Name	Common Name	Protected or Notable Status	Location
Briza minor	Lesser quaking grass	Nationally scarce	River Mole
Epipactis leptochila	Narrow-lipped helleborine	Nationally scarce	LERL Biodiversity area (woodland)
Hyacinthoides non-scripta	Bluebell	WCA Schedule 8	LERL Biodiversity area (woodland, attenuation field)
Lychnis flos-cuculi	Ragged robin	Near Threatened	River Mole
Mentha pulegium	Pennyroyal	UK BAP, Nationally Scarce, Endangered, Schedule 8, NERC S.41, Critically Endangered	Grassland along rail line
Polygonatum odoratum	Solomon's seal	Nationally scarce	LERL Biodiversity area (Woodland)

**Annex 2.1.3: Invasive Plant Species identified within the Gatwick Project Boundary**

Species Name	Common Name	Protected or Notable Status	Location
Impatiens glandulifera	Himalayan balsam	WCA Schedule 9	River Mole, Gatwick Stream, Airside Stream

**Annex 2.1.4: Target Notes**

Target Note Ref.	Description
TN1	Location of pennypoyal
TN2	Large vegetated earth bank within Eastern Carparking
TN3	Horleyland Wood
TN4	Upper Pickett's Wood
TN5	Solomons seal, narrow-lipped helleborine and bluebell locations
TN6a	Plantation woodland 1

Target Note Ref.	Description
TN6b	Plantation Woodland 2
TN6c	Plantation Woodland 3
TN7	Brockley Wood
TN8	Large Area of Scrub near Brockley Wood
TN9	Lesser quaking grass and ragged robin location
TN10a	Western marshy grassland
TN10b	Eastern marshy grassland
TN10c	Marshy grassland along the River Mole
TN11	Large, 8 metres tall earth bank south west of Brockley Wood
TN12	Dog Kennel Wood
TN13	Crawter's Wood
TN14	Area of isolated dense scrub
TN15	Area of dense, overgrown bramble and rose encroaching onto open grassy glade.

NVC Surveys

Annex 2.1.5: Quadrat 1

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Lotus pedunculatus</b>	Greater bird's foot trefoil	20
<b>Juncus conglomeratus</b>	Compact rush	20
<b>Dactylis glomerata</b>	Cock's foot	20
<b>Holcus lanatus</b>	Yorkshire fog	10
<b>Centaurea nigra</b>	Common knapweed	10
<b>Carex otrubae</b>	False fox-sedge	10
<b>Alopecurus pratensis</b>	Meadow foxtail	10
<b>Vicia cracca</b>	Bird vetch	>1
<b>Poa trivialis</b>	Rough meadow-grass	>1
<b>Lathyrus nissolia</b>	Grass vetchling	>1
<b>Arrhenatherum elatius</b>	False oat-grass	>1
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

Annex 2.1.6: Quadrat 2

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Centaurea nigra</b>	Common knapweed	40
<b>Lotus pedunculatus</b>	Greater bird's-foot trefoil	15

Homogenous Stand 1		
<b>Potentilla reptans</b>	Creeping cinquefoil	15
<b>Juncus conglomeratus</b>	Compressed rush	10
<b>Agrostis stolonifera</b>	Creeping bent	10
<b>Arrhenatherum elatius</b>	False oat-grass	5
<b>Holcus lanatus</b>	Yorkshire fog	5
<b>Phleum pratensis</b>	Timothy grass	5
<b>Deschampsia cespitosa</b>	Tufted Hair grass	5
<b>Trifolium pratense</b>	Red clover	>1
<b>Oenanthe crocata</b>	Hemlock water dropwort	>1
<b>Ranunculus acris</b>	Meadow buttercup	>1
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

**Annex 2.1.7: Quadrat 3**

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Centaurea nigra</b>	Common knapweed	30
<b>Anthoxanthum odoratum</b>	Sweet vernal grass	15
<b>Plantago lanceolata</b>	Ribwort plantain	15
<b>Agrostis capillaris</b>	Common bent	10
<b>Galium verum</b>	Lady's bedstraw	10
<b>Achillea millefolium</b>	Yarrow	10
<b>Lotus pedunculatus</b>	Greater bird's-foot trefoil	10
<b>Holcus lanatus</b>	Yorkshire fog	5
<b>Briza minor</b>	Lesser quaking grass	5
<b>Deschampsia cespitosa</b>	Tufted hair grass	5
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

**Annex 2.1.8: Quadrat 4**

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Plantago lanceolata</b>	Ribwort plantain	20
<b>Centaurea nigra</b>	Common knapweed	20
<b>Stachys palustris</b>	Marsh woundwort	20
<b>Briza minor</b>	Lesser quaking grass	20
<b>Galium verum</b>	Lady's bedstraw	10
<b>Odontites vernus</b>	Red bartsia	10
<b>Festuca rubra</b>	Red fescue	>1
<b>Agrostis stolonifera</b>	Creeping bent	>1
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

**Annex 2.1.9: Quadrat 5**

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Agrostis stolonifera</b>	Creeping bent	20
<b>Briza minor</b>	Lesser quaking grass	20
<b>Centaurea nigra</b>	Common knapweed	15
<b>Arrhenatherum elatius</b>	False oat-grass	15
<b>Stachys palustris</b>	Marsh woundwort	10
<b>Galium verum</b>	Lady's bedstraw	10
<b>Anthoxanthum odoratum</b>	Sweet vernal grass	10
<b>Plantago lanceolata</b>	Ribwort plantain	10
<b>Hypericum perforatum</b>	Perforate St john's-wort	5
<b>Agrimonia eupatoria</b>	Agrimony	5
<b>Calamagrostis epigejos</b>	Wood small reed	5
<b>Agrostis stolonifera</b>	Creeping bent	20
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

\*Other Species: *Primula sp.*, Primrose, *Cynosurus cristatus*, Crested Dog's Tail, *Sanguisorba officinalis*, Great Burnet.

**Annex 2.1.10: Quadrat 6**

Homogenous Stand 2		
Species Name	Common Name	% Cover
Calamagrostis epigejos	Wood small reed	80
Phleum pratense	Timothy grass	10
Juncus conglomeratus	Compact rush	5
Lotus pedunculatus	Greater bird's-foot trefoil	5
Calamagrostis epigejos Society		

**Annex 2.1.11: Quadrat 7**

Homogenous Stand 2		
Species Name	Common Name	% Cover
Calamagrostis epigejos	Wood small reed	80
Deschampsia cespitosa	Tufted hair grass	15
Alopecurus pratensis	Meadow foxtail	10
Calamagrostis epigejos Society		

**Annex 2.1.12: Quadrat 8**

Homogenous Stand 2		
Species Name	Common Name	% Cover
Lotus pedunculatus	Greater bird's-foot trefoil	40
Centaurea nigra	Common knapweed	15
Briza minor	Lesser quaking grass	10
Calamagrostis epigejos	Wood small reed	10
Arrhenatherum elatius	False oat-grass	10
Holcus lanatus	Yorkshire fog	10
Anthoxanthum odoratum	Sweet vernal grass	5
Ranunculus acris	Meadow buttercup	5
Calamagrostis epigejos Society		

**Annex 2.1.13: Quadrat 9**

Homogenous Stand 3		
Species Name	Common Name	% Cover
Lotus pedunculatus	Greater bird's-foot trefoil	20

Homogenous Stand 3		
<b>Briza minor</b>	Lesser quaking grass	10
<b>Leucanthemum vulgare</b>	Oxeye daisy	10
<b>Anthoxanthum odoratum</b>	Sweet vernal grass	10
<b>Agrostis stolonifera</b>	Creeping bent	10
<b>Centaurea nigra</b>	Common knapweed	10
<b>Juncus effusus</b>	Soft rush	10
<b>Calamagrostis epigejos</b>	Wood small reed	10
<b>Daucus carota</b>	Wild carrot	5
<b>Ranunculus acris</b>	Meadow buttercup	5
<b>Holcus lanatus</b>	Yorkshire fog	>1
NVC Category: M27c Filipendula ulmaria-Angelica sylvestris mire, Juncus effusus – Holcus lanatus sub-community.		

**Annex 2.1.14: Quadrat 10**

Homogenous Stand 3		
Species Name	Common Name	% Cover
<b>Juncus effusus</b>	Soft rush	40
<b>Oenanthe crocata</b>	Hemlock water-dropwort	20
<b>Calamagrostis epigejos</b>	Wood small reed	20
<b>Epilobium hirsuta</b>	Greater willowherb	10
<b>Potentilla anserina</b>	Silverweed	10
<b>Scrophularia auriculata</b>	Water figwort	5
NVC Category: M27c Filipendula ulmaria-Angelica sylvestris mire, Juncus effusus – Holcus lanatus sub-community.		

**Annex 2.1.15: Quadrat 11**

Homogenous Stand 3		
Species Name	Common Name	% Cover
<b>Juncus effusus</b>	Soft rush	70
<b>Mentha aquatica</b>	Water mint	20
<b>Oenanthe crocata</b>	Hemlock water dropwort	10
<b>Lythrum salicaria</b>	Purple loosestrife	5
<b>Lychnis flos-cuculi</b>	Ragged robin	>1
NVC Category: M27c Filipendula ulmaria-Angelica sylvestris mire, Juncus effusus – Holcus lanatus sub-community.		



Breeding Bird Surveys

Annex 2.1.16: Breeding Status, Abundance and Conservation Status of Birds Recorded within the Gatwick Airport Survey Area in 2019

Species	Breeding status	No. of territories	Annex 1 EU Birds Directive	Schedule 1 WCA	NERC Species of Principal Importance	BoCC 5 Red and Amber species
Great Crested Grebe	Non-breeding	-	-	-	-	-
Cormorant	Non-breeding	-	-	-	-	-
Grey Heron	Non-breeding	-	-	-	-	-
Greylag Goose	Confirmed	3	-	-	-	- 1
Canada Goose	Confirmed	3	-	-	-	-
Mallard	Confirmed	9	-	-	-	Amber
Red Kite	Non-breeding	-	■	■	-	-
Common Buzzard	Confirmed	2	-	-	-	-
Kestrel	Confirmed	4	-	-	-	Amber
Peregrine	Possible	1	■	■	-	-
Sparrowhawk	Non-breeding	-	-	-	-	Amber**
Red-legged Partridge	Non-breeding	-	-	-	-	-
Pheasant	Confirmed	3	-	-	-	-
Moorhen	Confirmed	5	-	-	-	Amber**
Coot	Confirmed	3	-	-	-	-
Little ringed plover	Possible	1	-	■	-	-
Snipe	Non-breeding	-	-	-	-	Amber
Black-headed gull	Non-breeding	-	-	-	-	Amber
Herring gull	Non-breeding	-	-	-	■	Amber
Lesser black-backed gull	Non-breeding	-	-	-	-	Amber
Feral rock dove	Confirmed	6	-	-	-	-
Stock dove	Confirmed	3	-	-	-	Amber
Woodpigeon	Confirmed	37	-	-	-	Amber**
Collared dove	Confirmed	2	-	-	-	-
Swift	Non-breeding	-	-	-	-	Red**
Ring-necked parakeet	Non-breeding	-	-	-	-	-
Green woodpecker	Confirmed	3	-	-	-	-
Great spotted woodpecker	Confirmed	11	-	-	-	-
Skylark	Confirmed	12	-	-	■	Red
Swallow	Non-breeding	-	-	-	-	-
House martin	Non-breeding	-	-	-	-	Red**
Pied wagtail	Confirmed	5	-	-	-	-
Grey wagtail	Confirmed	1	-	-	-	Red
Wren	Confirmed	74	-	-	-	Amber**
Duncock	Confirmed	18	-	-	■	Amber

Species	Breeding status	No. of territories	Annex 1 EU Birds Directive	Schedule 1 WCA	NERC Species of Principal Importance	BoCC 5 Red and Amber species
Robin	Confirmed	100	-	-	-	-
Nightingale	Non-breeding	-	-	-	-	Red
Wheatear	Non-breeding	-	-	-	-	Amber**
Song thrush	Confirmed	19	-	-	■	Amber**
Redwing	Non-breeding	-	-	■	-	Amber**
Mistle thrush	Confirmed	2	-	-	-	Red
Blackbird	Confirmed	58	-	-	-	-
Garden warbler	Confirmed	2	-	-	-	-
Blackcap	Confirmed	43	-	-	-	-
Lesser whitethroat	Confirmed	2	-	-	-	-
Whitethroat	Confirmed	9	-	-	-	-
Reed warbler	Confirmed	1	-	-	-	-
Willow warbler	Non-Breeding	-	-	-	-	Amber
Chiffchaff	Confirmed	12	-	-	-	-
Goldcrest	Confirmed	9	-	-	-	-
Firecrest	Possible	1	-	■	-	-
Great tit	Confirmed	72	-	-	-	-
Coal tit	Confirmed	8	-	-	-	-
Blue tit	Confirmed	89	-	-	-	-
Marsh tit	Confirmed	1	-	-	■	Red
Long-tailed tit	Confirmed	15	-	-	-	-
Nuthatch	Confirmed	7	-	-	-	-
Treecreeper	Confirmed	7	-	-	-	-
Magpie	Confirmed	23	-	-	-	-
Jay	Confirmed	4	-	-	-	-
Jackdaw	Confirmed	11	-	-	-	-
Rook	Non-breeding	-	-	-	-	Amber**
Carrion crow	Confirmed	15	-	-	-	-
Starling	Confirmed	2	-	-	■	Red
House sparrow	Confirmed	4	-	-	■	Red
Chaffinch	Confirmed	8	-	-	-	-
Linnet	Confirmed	1	-	-	■	Red
Goldfinch	Confirmed	10	-	-	-	-
Greenfinch	Non-breeding	-	-	-	-	Red**
Siskin	Non-breeding	-	-	-	-	-
Bullfinch	Confirmed	1	-	-	■	Amber
Reed bunting	Confirmed	2	-	-	■	Amber

Note: 1. The native population of Greylag Goose in the UK is amber listed, however, the birds recorded during the survey are part of the introduced feral population and, as such, do not meet the criteria relating to species of conservation importance. Any BoCC statuses marked with \*\* have had an updated status within BoCC 5 since this report was first written.

Annex 2.1.17: Alphabetical List of Bird Species Recorded During the Survey in 2019

English name	Scientific name
Blackbird	<i>Turdus merula</i>
Blackcap	<i>Sylvia atricapilla</i>
Black-headed gull	<i>Chroicocephalus ridibundus</i>
Blue Tit	<i>Cyanistes caeruleus</i>
Bullfinch	<i>Pyrrhula pyrrhula</i>
Buzzard	<i>Buteo buteo</i>
Canada goose	<i>Branta canadensis</i>
Carrion crow	<i>Corvus corone</i>
Chaffinch	<i>Fringilla coelebs</i>
Chiffchaff	<i>Phylloscopus collybita</i>
Coal tit	<i>Periparus ater</i>
Collared dove	<i>Streptopelia decaocto</i>
Coot	<i>Fulica atra</i>
Cormorant	<i>Phalacrocorax carbo</i>
Dunnock	<i>Prunella modularis</i>
Feral Dove	<i>Columba livia</i>
Firecrest	<i>Regulus ignicapilla</i>
Garden warbler	<i>Sylvia borin</i>
Goldcrest	<i>Regulus regulus</i>
Goldfinch	<i>Carduelis carduelis</i>
Great crested grebe	<i>Podiceps cristatus</i>
Great spotted woodpecker	<i>Dendrocopos major</i>
Great tit	<i>Parus major</i>
Green woodpecker	<i>Picus viridis</i>
Greenfinch	<i>Chloris chloris</i>
Grey heron	<i>Ardea cinerea</i>
Grey wagtail	<i>Motacilla cinerea</i>
Greylag goose	<i>Anser anser</i>
Herring gull	<i>Larus argentatus</i>
House martin	<i>Delichon urbicum</i>
House sparrow	<i>Passer domesticus</i>
Jackdaw	<i>Coloeus monedula</i>
Jay	<i>Garrulus glandarius</i>
Kestrel	<i>Falco tinnunculus</i>
Lesser black-backed gull	<i>Larus fuscus</i>
Lesser whitethroat	<i>Sylvia curruca</i>
Linnet	<i>Linaria cannabina</i>
Little ringed plover	<i>Charadrius dubius</i>

English name	Scientific name
Long-tailed tit	<i>Aegithalos caudatus</i>
Magpie	<i>Pica pica</i>
Mallard	<i>Anas platyrhynchos</i>
Marsh tit	<i>Poecile palustris</i>
Mistle thrush	<i>Turdus viscivorus</i>
Moorhen	<i>Gallinula chloropus</i>
Nightingale	<i>Luscinia megarhynchos</i>
Nuthatch	<i>Sitta europaea</i>
Peregrine	<i>Falco peregrinus</i>
Pheasant	<i>Phasianus colchicus</i>
Pied wagtail	<i>Motacilla alba</i>
Red kite	<i>Milvus milvus</i>
Red-legged partridge	<i>Alectoris rufa</i>
Redwing	<i>Turdus iliacus</i>
Reed bunting	<i>Emberiza schoeniclus</i>
Reed warbler	<i>Acrocephalus scirpaceus</i>
Ring-necked parakeet	<i>Psittacula krameri</i>
Robin	<i>Erithacus rubecula</i>
Rook	<i>Corvus frugilegus</i>
Siskin	<i>Spinus spinus</i>
Skylark	<i>Alauda arvensis</i>
Snipe	<i>Gallinago gallinago</i>
Song thrush	<i>Turdus philomelos</i>
Sparrowhawk	<i>Accipiter nisus</i>
Starling	<i>Sturnus vulgaris</i>
Stock dove	<i>Columba oenas</i>

Wintering Bird Surveys

Annex 2.1.18: Summary Count Data of Birds Recorded During Survey - October 2018 and March 2019

Species	Peak Count	Mean Count	Species	Peak Count	Mean Count
Blackbird	54	44.4	Kestrel	4	3
Bullfinch	7	2.8	Red kite	1	0.2
Black-headed gull	110	34.6	Lapwing	240	48
Blue tit	140	98.2	Lesser black-backed gull	2	0.4
Buzzard	3	2	Long-tailed tit	58	24.4
Carrion crow	42	31.4	Mistle thrush	3	1.2
Chiffchaff	15	3	Mallard	17	14.2
Collared dove	2	0.6	Magpie	36	22.6

Species	Peak Count	Mean Count	Species	Peak Count	Mean Count
Canada goose	28	7.2	Moorhen	8	3.6
Chaffinch	6	2.6	Mandarin duck	2	0.4
Common gull	1	0.2	Meadow pipit	31	7.8
Coal tit	10	5	Marsh tit	3	0.6
Dunnock	15	10.4	Nuthatch	11	6.4
Feral rock dove	9	2.2	Pheasant	3	0.8
Egyptian goose	2	0.4	Pied wagtail	10	5
Firecrest	1	0.2	Robin	81	57.8
Fieldfare	19	7.8	Reed bunting	1	0.2
Green woodpecker	3	1.8	Redwing	75	20.4
Goldcrest	33	16.4	Ring-necked parakeet	2	0.6
Green sandpiper	1	0.4	Rook	27	8
Greylag goose	5	1	Skylark	13	3.2
Grey wagtail	3	1.4	Starling	55	26.8
Goldfinch	12	7.4	Sparrowhawk	2	0.4
Greenfinch	2	0.4	Siskin	23	5.2
Great spotted woodpecker	11	7.6	Snipe	7	1.4
Great tit	83	64	Song thrush	17	14.2
Grey heron	3	1.4	Treecreeper	7	4.8
Herring gull	10	3	Woodcock	1	0.2
House sparrow	3	1.2	Woodpigeon	102	62.2
Jay	15	7.2	Wren	38	21.8
Jackdaw	175	75.6			

Annex 2.1.19: Conservation Status of Birds Recorded within the Project Area - October 2018 and March 2019

Species	Annex 1 EU Birds Directive	UK BAP Priority Species	NERC Species of Principal Importance	Birds of Conservation Concern
Bullfinch		•	•	Amber
Black-headed gull				Amber
Common gull				Amber
Dunnock		•	•	Amber
Fieldfare				Red
Green sandpiper				Amber
Greylag goose				Amber
Grey wagtail				Red
Herring gull		•	•	Amber

Species	Annex 1 EU Birds Directive	UK BAP Priority Species	NERC Species of Principal Importance	Birds of Conservation Concern
House sparrow		•	•	Red
Kestrel				Amber
Lapwing		•	•	Red
Lesser black-backed gull				Amber
Mallard				Amber
Marsh tit		•	•	Red
Mistle thrush				Red
Meadow pipit				Amber
Red kite	•			N/A
Redwing				Amber**
Skylark		•	•	Red
Snipe				Amber
Song thrush		•	•	Amber**
Starling		•	•	Red
Woodcock				Red

Reptile Surveys

Annex 2.1.20: Reptile Survey Results

Survey	Survey Area	Date	Weather	Species recorded
1	A3, A5	17/04/19	15C, Wind F2, Cloud 3/8	None
	A6 River Mole Corridor	18/04/19	14C, Wind F3, Cloud 2/8	None
	A1	12/06/19	14C, Wind F1, Cloud 7/8	None
	A6 Field south of Brockley Wood	03/09/19	19C, Wind F1, Cloud 7/8	None
2	A3	01/05/19	10-11C, Wind F1, Cloud 1/8-3/8	None
	A5			None
	A6 River Mole Corridor			3 female grass snake, 1 sub-adult male grass snake, 3 juvenile grass snake and 3 grass snake
	A1	18/06/19	16C, Wind F2, Cloud 2/8	None
	A6 Field south of Brockley Wood	05/09/19	15C, Wind F2, Cloud 3/8	None
3	A3	13/05/19	15C, Wind F1, Cloud 2/8	None
	A5			None
	A6 River Mole Corridor			4 grass snake, 1 juvenile grass snake, 3 adult grass snake and 2 sub-adult grass snake
	A1	26/06/19	18C, Wind F1, Cloud 7/8	1 grass snake, 1 juvenile grass snake
	A6 Field south of Brockley Wood	10/09/2019	16C, Wind F1, Cloud 3/8	None
4	A5	03/06/19	19C, Wind F3, Cloud 3/8	2 juvenile grass snake
	A3, A6 River Mole Corridor	13/06/19	13C, wind F3, Cloud 6/8	None
	A1	08/08/2019	18C, Wind F1, Cloud 1/8	None

Survey	Survey Area	Date	Weather	Species recorded
	A6 Field south of Brockley Wood	16/09/2019	15C, Wind F1, Cloud 5/8	None
<b>5</b>	A3	26/06/19	18C, wind F1, Cloud 7/8	1 grass snake, 1 juvenile grass snake
	A5			None
	A6 River Mole Corridor			None
	A1	16/09/2019	15C, Wind F1, Cloud 5/8	None
	A6 Field south of Brockley Wood	19/09/2019	15C, Wind F1, Cloud 1/8	None
<b>6</b>	A5	06/08/19	20C, wind F4, Cloud 4/8	None
	A3	08/08/19	18C, Wind F3, Cloud 6/8	None
	A6 River Mole Corridor			2 grass snake slough
	A6 Field south of Brockley Wood	19/09/2019	15C, Wind F1, Cloud 1/8	None
<b>7</b>	A1	26/09/2019	18C, Wind F3, Cloud 3/8	None
	A3, A5, A6 River Mole Corridor, A6 Field south of Brockley Wood	02/10/2019	14C, Wind F2, Cloud 1/8	None

Great Crested Newt Surveys

Annex 2.1.21: HSI Scores for All Ponds within Project Boundary

Pond No.	Description	HSI score
FCZ	No access was granted to this pond.	N/S
9VG	No access was granted to this pond.	N/S
Pond F	A large man-made attenuation pond, a barrier crosses the middle running north to south and marginal vegetation was present around all the sides.	Poor
SM7	Small pond behind services with poor water quality and little aquatic vegetation.	Poor
981	Large pond within a woodland. No aquatic vegetation and little woodland ground flora. Mature trees surround pond and a film of algae across pond.	Below average
Pond G	Shallow pond with a silt bed that was mostly dry	Below average
30Z	Large pond within Horleyland Wood. Pond has shallow banks and marginal vegetation.	Average
8N8	Woodland pond flooded over into surrounding woodland and cutting off footpath to rest of woodland	Good
W46	Small man-made wildlife pond, lots of aquatic vegetation along the southern bank. Banks relatively steep sided with a fallen tree across middle.	Average
Old Lagoon	Y shaped lagoon, man made, amenity grass banks and steep sided.	N/S
E11	Long linear settlement pond, wider at eastern end, linked to the M23 spur road. Reeds and bulrush dominated.	Average
Pond E	No access granted to this pond,	N/S
A0A	Pond within Police training area, swamped with willow and surrounded by woodland.	Below average
MHA	Circular pond in the middle of the southern staff car park. The pond was surrounded with vegetation.	Poor

Pond No.	Description	HSI score
JCT	Outside Project boundary, no access granted.	N/S
Pond A	A pond located to the north of the runway near the fire training area. It was surrounded by dense bramble scrub and marginal vegetation such as pond sedge and bulrush	Good
New Lagoon	A circular sewage pond known as 'New lagoon' was a man-made, steep sided amenity grassland settlement lagoon.	N/S
Pond M	Settlement pond east of the biodiversity wall. A man-made structure with steep concrete walls. Semi-improved grassland surrounded the pond. Only the eastern half of the pond held water.	N/S
WP9	No access granted to this pond.	N/S
AA20	Awaiting details.	Poor
AA21	Awaiting details.	Poor
K5F	A long pond with 0.5m high banks around the northern side. The southern bank was covered with scrub and inaccessible. Around all sides there was a large amount of aquatic vegetation.	Excellent
TTD	A small circular man-made pond surrounded by willow and pine trees. Aquatic vegetation was present around the eastern, northern and southern sides. A concrete outflow was identified in the south east corner of the pond.	Excellent
C24	Large pond around 30 x 20 m. Lots of marginal vegetation mainly bulrush completely dry willow and ash growing around edge.	Good*
Pond D	A rectangular attenuation pond. It was concrete sided with an outflow into the Mole corridor.	Poor
Pond D	A triangular attenuation pond made from concrete and steep sided. The pond was surrounded by managed grassland.	Poor

Pond No.	Description	HSI score
293	Large open fishing lake in middle of public park. Two islands within middle of the lake densely covered with trees.	Poor
FFJ	A small attenuation pond for the runway. Marginal vegetation was present here with rushes being dominant.	Good
29A	A long thin man-made channel. 5 metre high sides with a fence line around the top of it. Water was swamped by algae and had little aquatic vegetation The banks were vegetated with tufted grass.	Average*
30P	Murky shallow pond, with clear animal tracks leading to it.	Poor
AVF	A large pond within the Land East of the Gatwick Aviation Museum Field covered with reeds and willowherb and algae topped. Nettle and willowherb ruderal surrounded the pond.	Good
Dog Kennel Pond	A small manmade attenuation pond with steep banks showing high levels of maintenance. A diverse mix of aquatic and marginal vegetation was found within the pond.	Average
AAA4	Newly created pond along Mole corridor.	N/S
1WH	Pond in centre of eastern part of woodland had a small amount of water in with a heavy covering of duckweed.	Average
NU1	Pond in centre of eastern part of woodland had a small amount of water in with a heavy covering of duckweed. Linked to Pond 1WH	Average
A1 Ditch 1	Ditch goes along the boundary of a small area of woodland. No water or aquatic vegetation present. Visible from adjacent parallel footpath to the south.	Good
A1 Ditch 2	Field boundary ditch with trees and scrub on either side. No aquatic vegetation. Dry in places. Completely	Average



Pond No.	Description	HSI score
	shaded. Visible from adjacent parallel footpath to the south.	
<b>A1 Ditch 3</b>	Ditch on field boundary at the base of a wooded bank. Shaded by adjacent trees. No aquatic vegetation. Some stagnant water. Ditch continues west, parallel to the A23 but is mostly dry.	Average
<b>Un-named additional ditch 1</b>	Additional ditch recorded under bridge.	????
<b>A1 Ditch 5</b>	Ditch parallel to the B2036. Wet with some aquatic vegetation. Visible along the side of the road.	????
<b>A1 Ditch 4</b>	Wet ditch at the base of an earth bank for the A23. No aquatic vegetation. Quite shaded by trees either side. Visible from the B2036.	Below average
<b>A1 services ditch</b>	Dry ditch parallel to the A23/Airport Way, at the bottom of the earth bank for the A23. Surrounded by trees. Access from adjacent footpath.	????
<b>A1 Services lake</b>	Large lake behind the services area. No aquatic vegetation. Wildfowl present, kingfisher seen twice. Banks are lined with trees. Access from adjacent footpath.	????
<b>A1 additional ditch by M23</b>	Wet ditch with reeds adjacent to the M23 slip road. Trees and scrub either side. Access – can see from adjacent footpath but surrounded by dense scrub.	????
<b>A2 Ditch 5</b>	Wet ditch parallel to a railway embankment. Occasional adjacent trees and scrub. No aquatic vegetation. No direct access, viewed through fence.	Average
<b>A2 SUDS area</b>	Large SUDS area in the middle of the car park. Surrounded by trees, scrub and grassland.	????
<b>A2 Ditch 6</b>	Dry ditch along the edge of some woodland. No aquatic vegetation. Also	Average

Pond No.	Description	HSI score
	continues along the southern edge of the car park.	
<b>A2 Ditch 7</b>	Wet ditch with some emergent vegetation. Surrounded by car parking with some adjacent trees.	Good
<b>A2 Ditch 8</b>	Ditch surrounded by car parking. Partly wet at eastern end with some aquatic vegetation. Adjacent trees along south side of ditch.	Good
<b>A2 Ditch 9</b>	Dry ditch along the edge of an access track with mature trees on its northern bank. No aquatic vegetation.	Average
<b>A2 Ditch 10</b>	Partly wet ditch along the edge of an access track. No aquatic vegetation. Mature trees present on the southern bank of the ditch.	Average
<b>A2 additional ditch south of D10</b>	Additional ditch parallel to D10, to the south, along the edge of a car park. Partly wet but no aquatic vegetation.	????
<b>A2 Ditch 11</b>	There are two ditches here, either side of the road. Both are dry with no aquatic vegetation.	Average
<b>A2 Ditch 12</b>	Wet ditch through a car park with adjacent mature trees and dense bracken. No aquatic vegetation.	Below average
<b>A2 Ditch 13 - West</b>	Dry ditch along a tree line. No aquatic vegetation.	Average
<b>A2 Ditch 13 - East</b>	Wet ditch along a tree line with scrub either side. No aquatic vegetation.	????
<b>A2 Ditch 14</b>	Wet ditch surrounded by scrub and self-seeded tree saplings. No aquatic vegetation present.	Good
<b>A2 Pond 1</b>	Large woodland pond with some emergent vegetation.	Average
<b>A2 Pond 2</b>	Small woodland pond with over-hanging trees. No aquatic vegetation present.	????
<b>A3 Pond 1</b>	New environmental pond. Lined. No aquatic vegetation visible apart from duckweed. Pond situated in an area of	Below average

Pond No.	Description	HSI score
	woodland, adjacent to a large wet meadow. Habitat connectivity to P2.	
<b>A3 Pond 2</b>	Woodland pond surrounded by tussocky grassland and trees. No aquatic vegetation. GCN recorded in 2021 during bottle trapping.	Average
<b>A3 Green Ditch 1</b>	Wet ditch within a large field of tussocky grassland / wet meadow. Some young willow trees growing either side of the ditch. No aquatic vegetation but surrounding habitat very good, with habitat connectivity to P2 with confirmed GCN presence.	Below average
<b>A3 Purple Ditch 1</b>	Ditch along the north side of a grass access track. Wet but no aquatic vegetation. A hedge is on the north side of the ditch.	????
<b>A3 Purple Ditch 2</b>	Ditch along the south side of a grass access track. Wet but no aquatic vegetation. A hedge is on the south side of the ditch.	????
<b>A3 Purple Ditch 3</b>	Ditch along the east side of a grass access track. Wet but no aquatic vegetation. A hedge is on the east side of the ditch.	????
<b>A3 Purple Ditch 4</b>	Ditch along the west side of a grass access track. Wet but no aquatic vegetation. Woodland is adjacent to the west side of the ditch.	????
<b>A5 Ditch 1</b>	Wet ditch along the edge of a grass field, with adjacent scrub and trees. No aquatic vegetation. Ditch is only adjacent to the southern field boundary, there is no ditch along the eastern field boundary as marked on the map.	Excellent
<b>A5 Ditch 2</b>	Wet ditch with grass banks along the edges of a field. No aquatic vegetation.	Good
<b>A5 Ditch 3</b>	Dry ditch with grass banks along the edge of a field. Adjacent trees and hedge. Visible from Aviation museum	Excellent

Pond No.	Description	HSI score
	field to the west. Direct access via clay shooting school.	
<b>A5 Pond 1</b>	No pond at the location shown on the map.	Below average
<b>A5 additional Pond 2</b>	Large pond with dense reeds surrounded by grassland.	????
<b>A5 additional Pond 3</b>	Small pond surrounded by trees and tussocky grassland. No aquatic vegetation.	????
<b>B2036 Ditch 1</b>	Unknown location, not labelled on map.	Below average
<b>B2036 Ditch 2</b>	Unknown location, not labelled on map.	Poor
<b>B2036 Ditch 3</b>	????	Good
<b>B2036 Ditch 4</b>	????	Below average
<b>Crawter's Wood- west</b>	No access available. This pond is airside so security access and a security escort need to be pre-arranged for access.	Poor
<b>Crawter's Wood- east</b>	No access available. This pond is airside so security access and a security escort need to be pre-arranged for access.	Poor
<b>Detention Centre Pond</b>	No access available.	Below average
<b>Ditch, North boundary, Car park X</b>	Partly wet ditch with bulrushes. Grass and ruderal vegetation on the banks. Surrounded by hard-standing.	Good
<b>Ditch, Southern boundary, Car park X</b>	Wet ditch along southern edge of car park. Trees and scrub adjacent to the south. No aquatic vegetation.	Good
<b>Ditch, Western boundary, Car park X</b>	Wet ditch along the edge of the car park with some aquatic vegetation and adjacent scrub.	????
<b>Pond, Car Park X</b>	Additional pond. In the centre of the car park. Surrounded by trees and grassland. No aquatic vegetation.	????

Pond No.	Description	HSI score
<b>North long stay A6/A7- Ditch 1</b>	Ditch runs through the centre of a car park. No connecting habitat. The ditch holds water but no aquatic vegetation is present. The banks are lined with self-seeded tree saplings and some scrub.	Below average
<b>North long stay A6/A7- Ditch 2</b>	Ditch runs along the southern and western boundaries of the car park. Wet, but no visible aquatic vegetation. Bordered by scrub and self-seeded tree saplings. More open along western boundary.	Excellent
<b>North long stay A6/A7- Ditch 3</b>	Ditch runs parallel to southern edge of car park. Some water but no visible aquatic vegetation. Heavily shaded by adjacent trees and scrub.	Average
<b>North long stay A6/A7- Ditch 4</b>	Ditch between two car parks. Some water but no visible aquatic vegetation. Heavily shaded by adjacent trees and scrub.	Average
<b>North long stay A6/A7- Pond 1</b>	Woodland pond surrounded by trees and grassland. Some emergent vegetation present.	Good
<b>Woodland behind Premier Inn- Ditch 1</b>	A8 South – Shallow wet ditch through an area of woodland. No aquatic vegetation present.	Below average
<b>Woodland behind Premier Inn- Riverside Garden</b>	A8 North – Ditch runs parallel to the A23 London Road, through an area of woodland. Dry in places. No aquatic vegetation. Access from within Riverside Gardens	Below average
<b>A8 Island ditch</b>	Island of land surrounded by road. There are a few metres of amenity grassland before a ditch, which encircles the western half of the island. The majority of the island is wooded. The ditch itself is deep but mostly dry, with no aquatic vegetation.	????
<b>Pond 1</b>	Large pond in the corner of an arable field with lots of emergent vegetation and adjacent trees.	Excellent

Pond No.	Description	HSI score
<b>Pond 4</b>	Large pond in the corner of a grass field surrounded by trees. No aquatic vegetation is present.	Poor
<b>Pond 5</b>	Small pond in the corner of a grass field, surrounded by trees. No aquatic vegetation is present.	Poor
<b>Pond 6</b>	Large garden pond with extensive emergent vegetation. Surrounded by amenity grassland. Mature trees line the garden boundary.	Excellent
<b>Additional pond right next to the M23</b>	Large drainage pond adjacent to the earth bank of the M23. Extensive emergent vegetation present. Trees and scrub are present along the northern edge of the pond.	Excellent

**Annex 2.1.22: Protected or Notable Species identified during Botanical Survey**

Species Name	Common Name	Protected or Notable Status	Location
<b>Briza minor</b>	Lesser quaking grass	Nationally scarce	River Mole
<b>Epipactis leptochila</b>	Narrow-lipped helleborine	Nationally scarce	LERL Biodiversity area (woodland)
<b>Hyacinthoides non-scripta</b>	Bluebell	WCA Schedule 8	LERL Biodiversity area (woodland, attenuation field)
<b>Lychnis flos-cuculi</b>	Ragged robin	Near Threatened	River Mole
<b>Mentha pulegium</b>	Pennyroyal	UK BAP, Nationally Scarce, Endangered, Schedule 8, NERC S.41, Critically Endangered	Grassland along rail line
<b>Polygonatum odoratum</b>	Solomon's seal	Nationally scarce	LERL Biodiversity area (Woodland)

**Annex 2.1.23: Invasive Plant Species identified within the Gatwick Project Boundary**

Species Name	Common Name	Protected or Notable Status	Location
<b>Impatiens glandulifera</b>	Himalayan balsam	WCA Schedule 9	River Mole, Gatwick Stream, Airside Stream

**Annex 2.1.24: Target Notes**

Target Note Ref.	Description
<b>TN1</b>	Location of pennypoyal
<b>TN2</b>	Large vegetated earth bank within Eastern Carparking
<b>TN3</b>	Horleyland Wood
<b>TN4</b>	Upper Pickett's Wood
<b>TN5</b>	Solomons seal, narrow-lipped helleborine and bluebell locations
<b>TN6a</b>	Plantation woodland 1
<b>TN6b</b>	Plantation Woodland 2
<b>TN6c</b>	Plantation Woodland 3
<b>TN7</b>	Brockley Wood
<b>TN8</b>	Large Area of Scrub near Brockley Wood
<b>TN9</b>	Lesser quaking grass and ragged robin location
<b>TN10a</b>	Western marshy grassland
<b>TN10b</b>	Eastern marshy grassland
<b>TN10c</b>	Marshy grassland along the River Mole
<b>TN11</b>	Large, 8 metres tall earth bank south west of Brockley Wood
<b>TN12</b>	Dog Kennel Wood
<b>TN13</b>	Crawter's Wood
<b>TN14</b>	Area of isolated dense scrub
<b>TN15</b>	Area of dense, overgrown bramble and rose encroaching onto open grassy glade.

NVC Surveys

Annex 2.1.25: Quadrat 1

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Lotus pedunculatus</b>	Greater bird's foot trefoil	20
<b>Juncus conglomeratus</b>	Compact rush	20
<b>Dactylis glomerata</b>	Cock's foot	20
<b>Holcus lanatus</b>	Yorkshire fog	10
<b>Centaurea nigra</b>	Common knapweed	10
<b>Carex otrubae</b>	False fox-sedge	10
<b>Alopecurus pratensis</b>	Meadow foxtail	10
<b>Vicia cracca</b>	Bird vetch	>1
<b>Poa trivialis</b>	Rough meadow-grass	>1
<b>Lathyrus nissolia</b>	Grass vetchling	>1
<b>Arrhenatherum elatius</b>	False oat-grass	>1
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

Annex 2.1.26: Quadrat 2

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Centaurea nigra</b>	Common knapweed	40
<b>Lotus pedunculatus</b>	Greater bird's-foot trefoil	15
<b>Potentilla reptans</b>	Creeping cinquefoil	15
<b>Juncus conglomeratus</b>	Compressed rush	10
<b>Agrostis stolonifera</b>	Creeping bent	10
<b>Arrhenatherum elatius</b>	False oat-grass	5
<b>Holcus lanatus</b>	Yorkshire fog	5
<b>Phleum pratensis</b>	Timothy grass	5
<b>Deschampsia cespitosa</b>	Tufted Hair grass	5
<b>Trifolium pratense</b>	Red clover	>1
<b>Oenanthe crocata</b>	Hemlock water dropwort	>1
<b>Ranunculus acris</b>	Meadow buttercup	>1
NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.		
Arrhenatherum elatius sub-community.		

**Annex 2.1.27: Quadrat 3**

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Centaurea nigra</b>	Common knapweed	30
<b>Anthoxanthum odoratum</b>	Sweet vernal grass	15
<b>Plantago lanceolata</b>	Ribwort plantain	15
<b>Agrostis capillaris</b>	Common bent	10
<b>Galium verum</b>	Lady's bedstraw	10
<b>Achillea millefolium</b>	Yarrow	10
<b>Lotus pedunculatus</b>	Greater bird's-foot trefoil	10
<b>Holcus lanatus</b>	Yorkshire fog	5
<b>Briza minor</b>	Lesser quaking grass	5
<b>Deschampsia cespitosa</b>	Tufted hair grass	5

NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.  
Arrhenatherum elatius sub-community.

**Annex 2.1.28: Quadrat 4**

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Plantago lanceolata</b>	Ribwort plantain	20
<b>Centaurea nigra</b>	Common knapweed	20
<b>Stachys palustris</b>	Marsh woundwort	20
<b>Briza minor</b>	Lesser quaking grass	20
<b>Galium verum</b>	Lady's bedstraw	10
<b>Odontites vernus</b>	Red bartsia	10
<b>Festuca rubra</b>	Red fescue	>1
<b>Agrostis stolonifera</b>	Creeping bent	>1

NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.  
Arrhenatherum elatius sub-community.

Annex 2.1.29: Quadrat 5

Homogenous Stand 1		
Species Name	Common Name	% Cover
<b>Agrostis stolonifera</b>	Creeping bent	20
<b>Briza minor</b>	Lesser quaking grass	20
<b>Centaurea nigra</b>	Common knapweed	15
<b>Arrhenatherum elatius</b>	False oat-grass	15
<b>Stachys palustris</b>	Marsh woundwort	10
<b>Galium verum</b>	Lady's bedstraw	10
<b>Anthoxanthum odoratum</b>	Sweet vernal grass	10
<b>Plantago lanceolata</b>	Ribwort plantain	10
<b>Hypericum perforatum</b>	Perforate St john's-wort	5
<b>Agrimonia eupatoria</b>	Agrimony	5
<b>Calamagrostis epigejos</b>	Wood small reed	5
<b>Agrostis stolonifera</b>	Creeping bent	20

NVC Category: MG9b Holcus lanatus – Deschampsia cespitosa grassland.  
 Arrhenatherum elatius sub-community.

\*Other Species: *Primula sp.*, Primrose, *Cynosurus cristatus*, Crested Dog's Tail, *Sanguisorba officinalis*, Great Burnet.

**Annex 2.1.30: Quadrat 6**

Homogenous Stand 2		
Species Name	Common Name	% Cover
Calamagrostis epigejos	Wood small reed	80
Phleum pratense	Timothy grass	10
Juncus conglomeratus	Compact rush	5
Lotus pedunculatus	Greater bird's-foot trefoil	5
Calamagrostis epigejos Society		

**Annex 2.1.31: Quadrat 7**

Homogenous Stand 2		
Species Name	Common Name	% Cover
Calamagrostis epigejos	Wood small reed	80
Deschampsia cespitosa	Tufted hair grass	15
Alopecurus pratensis	Meadow foxtail	10
Calamagrostis epigejos Society		

**Annex 2.1.32: Quadrat 8**

Homogenous Stand 2		
Species Name	Common Name	% Cover
Lotus pedunculatus	Greater bird's-foot trefoil	40
Centaurea nigra	Common knapweed	15
Briza minor	Lesser quaking grass	10
Calamagrostis epigejos	Wood small reed	10
Arrhenatherum elatius	False oat-grass	10
Holcus lanatus	Yorkshire fog	10
Anthoxanthum odoratum	Sweet vernal grass	5
Ranunculus acris	Meadow buttercup	5
Calamagrostis epigejos Society		

**Annex 2.1.33: Quadrat 9**

Homogenous Stand 3		
Species Name	Common Name	% Cover
Lotus pedunculatus	Greater bird's-foot trefoil	20
Briza minor	Lesser quaking grass	10

Homogenous Stand 3		
<b>Leucanthemum vulgare</b>	Oxeye daisy	10
<b>Anthoxanthum odoratum</b>	Sweet vernal grass	10
<b>Agrostis stolonifera</b>	Creeping bent	10
<b>Centaurea nigra</b>	Common knapweed	10
<b>Juncus effusus</b>	Soft rush	10
<b>Calamagrostis epigejos</b>	Wood small reed	10
<b>Daucus carota</b>	Wild carrot	5
<b>Ranunculus acris</b>	Meadow buttercup	5
<b>Holcus lanatus</b>	Yorkshire fog	>1
NVC Category: M27c Filipendula ulmaria-Angelica sylvestris mire, Juncus effusus – Holcus lanatus sub-community.		

**Annex 2.1.34: Quadrat 10**

Homogenous Stand 3		
Species Name	Common Name	% Cover
<b>Juncus effusus</b>	Soft rush	40
<b>Oenanthe crocata</b>	Hemlock water-dropwort	20
<b>Calamagrostis epigejos</b>	Wood small reed	20
<b>Epilobium hirsuta</b>	Greater willowherb	10
<b>Potentilla anserina</b>	Silverweed	10
<b>Scrophularia auriculata</b>	Water figwort	5
NVC Category: M27c Filipendula ulmaria-Angelica sylvestris mire, Juncus effusus – Holcus lanatus sub-community.		

**Annex 2.1.35: Quadrat 11**

Homogenous Stand 3		
Species Name	Common Name	% Cover
<b>Juncus effusus</b>	Soft rush	70
<b>Mentha aquatica</b>	Water mint	20
<b>Oenanthe crocata</b>	Hemlock water dropwort	10
<b>Lythrum salicaria</b>	Purple loosestrife	5
<b>Lychnis flos-cuculi</b>	Ragged robin	>1
NVC Category: M27c Filipendula ulmaria-Angelica sylvestris mire, Juncus effusus – Holcus lanatus sub-community.		



Breeding Bird Surveys

Annex 2.1.36: Breeding Status, Abundance and Conservation Status of Birds Recorded within the Gatwick Airport Survey Area in 2019

Species	Breeding status	No. of territories	Annex 1 EU Birds Directive	Schedule 1 WCA	NERC Species of Principal Importance	BoCC 5 Red and Amber species
Great Crested Grebe	Non-breeding	-	-	-	-	-
Cormorant	Non-breeding	-	-	-	-	-
Grey Heron	Non-breeding	-	-	-	-	-
Greylag Goose	Confirmed	3	-	-	-	- 1
Canada Goose	Confirmed	3	-	-	-	-
Mallard	Confirmed	9	-	-	-	Amber
Red Kite	Non-breeding	-	■	■	-	-
Common Buzzard	Confirmed	2	-	-	-	-
Kestrel	Confirmed	4	-	-	-	Amber
Peregrine	Possible	1	■	■	-	-
Sparrowhawk	Non-breeding	-	-	-	-	Amber**
Red-legged Partridge	Non-breeding	-	-	-	-	-
Pheasant	Confirmed	3	-	-	-	-
Moorhen	Confirmed	5	-	-	-	Amber**
Coot	Confirmed	3	-	-	-	-
Little ringed plover	Possible	1	-	■	-	-
Snipe	Non-breeding	-	-	-	-	Amber
Black-headed gull	Non-breeding	-	-	-	-	Amber
Herring gull	Non-breeding	-	-	-	■	Amber
Lesser black-backed gull	Non-breeding	-	-	-	-	Amber
Feral rock dove	Confirmed	6	-	-	-	-
Stock dove	Confirmed	3	-	-	-	Amber
Woodpigeon	Confirmed	37	-	-	-	Amber**
Collared dove	Confirmed	2	-	-	-	-
Swift	Non-breeding	-	-	-	-	Red**
Ring-necked parakeet	Non-breeding	-	-	-	-	-
Green woodpecker	Confirmed	3	-	-	-	-
Great spotted woodpecker	Confirmed	11	-	-	-	-
Skylark	Confirmed	12	-	-	■	Red
Swallow	Non-breeding	-	-	-	-	-
House martin	Non-breeding	-	-	-	-	Red**
Pied wagtail	Confirmed	5	-	-	-	-
Grey wagtail	Confirmed	1	-	-	-	Red
Wren	Confirmed	74	-	-	-	Amber**
Duncock	Confirmed	18	-	-	■	Amber
Robin	Confirmed	100	-	-	-	-
Nightingale	Non-breeding	-	-	-	-	Red

Species	Breeding status	No. of territories	Annex 1 EU Birds Directive	Schedule 1 WCA	NERC Species of Principal Importance	BoCC 5 Red and Amber species
Wheatear	Non-breeding	-	-	-	-	Amber**
Song thrush	Confirmed	19	-	-	■	Amber**
Redwing	Non-breeding	-	-	■	-	Amber**
Mistle thrush	Confirmed	2	-	-	-	Red
Blackbird	Confirmed	58	-	-	-	-
Garden warbler	Confirmed	2	-	-	-	-
Blackcap	Confirmed	43	-	-	-	-
Lesser whitethroat	Confirmed	2	-	-	-	-
Whitethroat	Confirmed	9	-	-	-	-
Reed warbler	Confirmed	1	-	-	-	-
Willow warbler	Non-Breeding	-	-	-	-	Amber
Chiffchaff	Confirmed	12	-	-	-	-
Goldcrest	Confirmed	9	-	-	-	-
Firecrest	Possible	1	-	■	-	-
Great tit	Confirmed	72	-	-	-	-
Coal tit	Confirmed	8	-	-	-	-
Blue tit	Confirmed	89	-	-	-	-
Marsh tit	Confirmed	1	-	-	■	Red
Long-tailed tit	Confirmed	15	-	-	-	-
Nuthatch	Confirmed	7	-	-	-	-
Treecreeper	Confirmed	7	-	-	-	-
Magpie	Confirmed	23	-	-	-	-
Jay	Confirmed	4	-	-	-	-
Jackdaw	Confirmed	11	-	-	-	-
Rook	Non-breeding	-	-	-	-	Amber**
Carrion crow	Confirmed	15	-	-	-	-
Starling	Confirmed	2	-	-	■	Red
House sparrow	Confirmed	4	-	-	■	Red
Chaffinch	Confirmed	8	-	-	-	-
Linnet	Confirmed	1	-	-	■	Red
Goldfinch	Confirmed	10	-	-	-	-
Greenfinch	Non-breeding	-	-	-	-	Red**
Siskin	Non-breeding	-	-	-	-	-
Bullfinch	Confirmed	1	-	-	■	Amber
Reed bunting	Confirmed	2	-	-	■	Amber

Note: 1. The native population of Greylag Goose in the UK is amber listed, however, the birds recorded during the survey are part of the introduced feral population and, as such, do not meet the criteria relating to species of conservation importance. Any BoCC statuses marked with \*\* have had an updated status within BoCC 5 since this report was first written.

Annex 2.1.37: Alphabetical List of Bird Species Recorded During the Survey in 2019

English name	Scientific name
Blackbird	<i>Turdus merula</i>
Blackcap	<i>Sylvia atricapilla</i>
Black-headed gull	<i>Chroicocephalus ridibundus</i>
Blue Tit	<i>Cyanistes caeruleus</i>
Bullfinch	<i>Pyrrhula pyrrhula</i>
Buzzard	<i>Buteo buteo</i>
Canada goose	<i>Branta canadensis</i>
Carrion crow	<i>Corvus corone</i>
Chaffinch	<i>Fringilla coelebs</i>
Chiffchaff	<i>Phylloscopus collybita</i>
Coal tit	<i>Periparus ater</i>
Collared dove	<i>Streptopelia decaocto</i>
Coot	<i>Fulica atra</i>
Cormorant	<i>Phalacrocorax carbo</i>
Dunnock	<i>Prunella modularis</i>
Feral Dove	<i>Columba livia</i>
Firecrest	<i>Regulus ignicapilla</i>
Garden warbler	<i>Sylvia borin</i>
Goldcrest	<i>Regulus regulus</i>
Goldfinch	<i>Carduelis carduelis</i>
Great crested grebe	<i>Podiceps cristatus</i>
Great spotted woodpecker	<i>Dendrocopos major</i>
Great tit	<i>Parus major</i>
Green woodpecker	<i>Picus viridis</i>
Greenfinch	<i>Chloris chloris</i>
Grey heron	<i>Ardea cinerea</i>
Grey wagtail	<i>Motacilla cinerea</i>
Greylag goose	<i>Anser anser</i>
Herring gull	<i>Larus argentatus</i>
House martin	<i>Delichon urbicum</i>
House sparrow	<i>Passer domesticus</i>
Jackdaw	<i>Coloeus monedula</i>
Jay	<i>Garrulus glandarius</i>
Kestrel	<i>Falco tinnunculus</i>
Lesser black-backed gull	<i>Larus fuscus</i>
Lesser whitethroat	<i>Sylvia curruca</i>
Linnet	<i>Linaria cannabina</i>
Little ringed plover	<i>Charadrius dubius</i>

English name	Scientific name
Long-tailed tit	<i>Aegithalos caudatus</i>
Magpie	<i>Pica pica</i>
Mallard	<i>Anas platyrhynchos</i>
Marsh tit	<i>Poecile palustris</i>
Mistle thrush	<i>Turdus viscivorus</i>
Moorhen	<i>Gallinula chloropus</i>
Nightingale	<i>Luscinia megarhynchos</i>
Nuthatch	<i>Sitta europaea</i>
Peregrine	<i>Falco peregrinus</i>
Pheasant	<i>Phasianus colchicus</i>
Pied wagtail	<i>Motacilla alba</i>
Red kite	<i>Milvus milvus</i>
Red-legged partridge	<i>Alectoris rufa</i>
Redwing	<i>Turdus iliacus</i>
Reed bunting	<i>Emberiza schoeniclus</i>
Reed warbler	<i>Acrocephalus scirpaceus</i>
Ring-necked parakeet	<i>Psittacula krameri</i>
Robin	<i>Erithacus rubecula</i>
Rook	<i>Corvus frugilegus</i>
Siskin	<i>Spinus spinus</i>
Skylark	<i>Alauda arvensis</i>
Snipe	<i>Gallinago gallinago</i>
Song thrush	<i>Turdus philomelos</i>
Sparrowhawk	<i>Accipiter nisus</i>
Starling	<i>Sturnus vulgaris</i>
Stock dove	<i>Columba oenas</i>

Wintering Bird Surveys

Annex 2.1.38: Summary Count Data of Birds Recorded During Survey - October 2018 and March 2019

Species	Peak Count	Mean Count	Species	Peak Count	Mean Count
Blackbird	54	44.4	Kestrel	4	3
Bullfinch	7	2.8	Red kite	1	0.2
Black-headed gull	110	34.6	Lapwing	240	48
Blue tit	140	98.2	Lesser black-backed gull	2	0.4
Buzzard	3	2	Long-tailed tit	58	24.4
Carrion crow	42	31.4	Mistle thrush	3	1.2
Chiffchaff	15	3	Mallard	17	14.2
Collared dove	2	0.6	Magpie	36	22.6
Canada goose	28	7.2	Moorhen	8	3.6
Chaffinch	6	2.6	Mandarin duck	2	0.4
Common gull	1	0.2	Meadow pipit	31	7.8
Coal tit	10	5	Marsh tit	3	0.6
Dunnock	15	10.4	Nuthatch	11	6.4
Feral rock dove	9	2.2	Pheasant	3	0.8
Egyptian goose	2	0.4	Pied wagtail	10	5
Firecrest	1	0.2	Robin	81	57.8
Fieldfare	19	7.8	Reed bunting	1	0.2
Green woodpecker	3	1.8	Redwing	75	20.4
Goldcrest	33	16.4	Ring-necked parakeet	2	0.6
Green sandpiper	1	0.4	Rook	27	8
Greylag goose	5	1	Skylark	13	3.2
Grey wagtail	3	1.4	Starling	55	26.8
Goldfinch	12	7.4	Sparrowhawk	2	0.4
Greenfinch	2	0.4	Siskin	23	5.2
Great spotted woodpecker	11	7.6	Snipe	7	1.4
Great tit	83	64	Song thrush	17	14.2
Grey heron	3	1.4	Treecreeper	7	4.8
Herring gull	10	3	Woodcock	1	0.2
House sparrow	3	1.2	Woodpigeon	102	62.2
Jay	15	7.2	Wren	38	21.8
Jackdaw	175	75.6			

Annex 2.1.39: Conservation Status of Birds Recorded within the Project Area - October 2018 and March 2019

Species	Annex 1 EU Birds Directive	UK BAP Priority Species	NERC Species of Principal Importance	Birds of Conservation Concern
Bullfinch		•	•	Amber
Black-headed gull				Amber
Common gull				Amber
Dunnock		•	•	Amber
Fieldfare				Red
Green sandpiper				Amber
Greylag goose				Amber
Grey wagtail				Red
Herring gull		•	•	Amber
House sparrow		•	•	Red
Kestrel				Amber
Lapwing		•	•	Red
Lesser black-backed gull				Amber
Mallard				Amber
Marsh tit		•	•	Red
Mistle thrush				Red
Meadow pipit				Amber
Red kite	•			N/A
Redwing				Amber**
Skylark		•	•	Red
Snipe				Amber
Song thrush		•	•	Amber**
Starling		•	•	Red
Woodcock				Red

Reptile Surveys

Annex 2.1.40: Reptile Survey Results

Survey	Survey Area	Date	Weather	Species recorded
1	A3, A5	17/04/19	15C, Wind F2, Cloud 3/8	None
	A6 River Mole Corridor	18/04/19	14C, Wind F3, Cloud 2/8	None
	A1	12/06/19	14C, Wind F1, Cloud 7/8	None
	A6 Field south of Brockley Wood	03/09/19	19C, Wind F1, Cloud 7/8	None
2	A3	01/05/19	10-11C, Wind F1, Cloud 1/8-3/8	None
	A5			None
	A6 River Mole Corridor	18/06/19	16C, Wind F2, Cloud 2/8	3 female grass snake, 1 sub-adult male grass snake, 3 juvenile grass snake and 3 grass snake
	A1			None

Survey	Survey Area	Date	Weather	Species recorded
	A6 Field south of Brockley Wood	05/09/19	15C, Wind F2, Cloud 3/8	None
3	A3	13/05/19	15C, Wind F1, Cloud 2/8	None
	A5			None
	A6 River Mole Corridor			4 grass snake, 1 juvenile grass snake, 3 adult grass snake and 2 sub-adult grass snake
	A1	26/06/19	18C, Wind F1, Cloud 7/8	1 grass snake, 1 juvenile grass snake
	A6 Field south of Brockley Wood	10/09/2019	16C, Wind F1, Cloud 3/8	None
4	A5	03/06/19	19C, Wind F3, Cloud 3/8	2 juvenile grass snake
	A3, A6 River Mole Corridor	13/06/19	13C, wind F3, Cloud 6/8	None
	A1	08/08/2019	18C, Wind F1, Cloud 1/8	None
	A6 Field south of Brockley Wood	16/09/2019	15C, Wind F1, Cloud 5/8	None
5	A3	26/06/19	18C, wind F1, Cloud 7/8	1 grass snake, 1 juvenile grass snake
	A5			None
	A6 River Mole Corridor			None
	A1	16/09/2019	15C, Wind F1, Cloud 5/8	None
	A6 Field south of Brockley Wood	19/09/2019	15C, Wind F1, Cloud 1/8	None
6	A5	06/08/19	20C, wind F4, Cloud 4/8	None
	A3	08/08/19	18C, Wind F3, Cloud 6/8	None
	A6 River Mole Corridor			2 grass snake slough
	A6 Field south of Brockley Wood	19/09/2019	15C, Wind F1, Cloud 1/8	None
7	A1	26/09/2019	18C, Wind F3, Cloud 3/8	None
	A3, A5, A6 River Mole Corridor, A6 Field south of Brockley Wood	02/10/2019	14C, Wind F2, Cloud 1/8	None

Great Crested Newt Surveys

Annex 2.1.41: HSI Scores for All Ponds within Project Boundary

Pond No.	Description	HSI score
<b>FCZ</b>	No access was granted to this pond.	N/S
<b>9VG</b>	No access was granted to this pond.	N/S
<b>Pond F</b>	A large man-made attenuation pond, a barrier crosses the middle running north to south and marginal vegetation was present around all the sides.	Poor
<b>SM7</b>	Small pond behind services with poor water quality and little aquatic vegetation.	Poor
<b>981</b>	Large pond within a woodland. No aquatic vegetation and little woodland ground flora. Mature trees surround pond and a film of algae across pond.	Below average
<b>Pond G</b>	Shallow pond with a silt bed that was mostly dry	Below average
<b>30Z</b>	Large pond within Horleyland Wood. Pond has shallow banks and marginal vegetation.	Average
<b>8N8</b>	Woodland pond flooded over into surrounding woodland and cutting off footpath to rest of woodland	Good
<b>W46</b>	Small man-made wildlife pond, lots of aquatic vegetation along the southern bank. Banks relatively steep sided with a fallen tree across middle.	Average
<b>Old Lagoon</b>	Y shaped lagoon, man made, amenity grass banks and steep sided.	N/S
<b>E11</b>	Long linear settlement pond, wider at eastern end, linked to the M23 spur road. Reeds and bulrush dominated.	Average
<b>Pond E</b>	No access granted to this pond,	N/S
<b>A0A</b>	Pond within Police training area, swamped with willow and surrounded by woodland.	Below average
<b>MHA</b>	Circular pond in the middle of the southern staff car park. The pond was surrounded with vegetation.	Poor

Pond No.	Description	HSI score
<b>JCT</b>	Outside Project boundary, no access granted.	N/S
<b>Pond A</b>	A pond located to the north of the runway near the fire training area. It was surrounded by dense bramble scrub and marginal vegetation such as pond sedge and bulrush	Good
<b>New Lagoon</b>	A circular sewage pond known as 'New lagoon' was a man-made, steep sided amenity grassland settlement lagoon.	N/S
<b>Pond M</b>	Settlement pond east of the biodiversity wall. A man-made structure with steep concrete walls. Semi-improved grassland surrounded the pond. Only the eastern half of the pond held water.	N/S
<b>WP9</b>	No access granted to this pond.	N/S
<b>AA20</b>	Awaiting details.	Poor
<b>AA21</b>	Awaiting details.	Poor
<b>K5F</b>	A long pond with 0.5m high banks around the northern side. The southern bank was covered with scrub and inaccessible. Around all sides there was a large amount of aquatic vegetation.	Excellent
<b>TTD</b>	A small circular man-made pond surrounded by willow and pine trees. Aquatic vegetation was present around the eastern, northern and southern sides. A concrete outflow was identified in the south east corner of the pond.	Excellent
<b>C24</b>	Large pond around 30 x 20 m. Lots of marginal vegetation mainly bulrush completely dry willow and ash growing around edge.	Good*
<b>Pond D</b>	A rectangular attenuation pond. It was concrete sided with an outflow into the Mole corridor.	Poor
<b>Pond D</b>	A triangular attenuation pond made from concrete and steep sided. The pond was surrounded by managed grassland.	Poor

Pond No.	Description	HSI score
<b>293</b>	Large open fishing lake in middle of public park. Two islands within middle of the lake densely covered with trees.	Poor
<b>FFJ</b>	A small attenuation pond for the runway. Marginal vegetation was present here with rushes being dominant.	Good
<b>29A</b>	A long thin man-made channel. 5 metre high sides with a fence line around the top of it. Water was swamped by algae and had little aquatic vegetation The banks were vegetated with tufted grass.	Average*
<b>30P</b>	Murky shallow pond, with clear animal tracks leading to it.	Poor
<b>AVF</b>	A large pond within the Land East of the Gatwick Aviation Museum Field covered with reeds and willowherb and algae topped. Nettle and willowherb ruderal surrounded the pond.	Good
<b>Dog Kennel Pond</b>	A small manmade attenuation pond with steep banks showing high levels of maintenance. A diverse mix of aquatic and marginal vegetation was found within the pond.	Average
<b>AAA4</b>	Newly created pond along Mole corridor.	N/S
<b>1WH</b>	Pond in centre of eastern part of woodland had a small amount of water in with a heavy covering of duckweed.	Average
<b>NU1</b>	Pond in centre of eastern part of woodland had a small amount of water in with a heavy covering of duckweed. Linked to Pond 1WH	Average
<b>A1 Ditch 1</b>	Ditch goes along the boundary of a small area of woodland. No water or aquatic vegetation present. Visible from adjacent parallel footpath to the south.	Good
<b>A1 Ditch 2</b>	Field boundary ditch with trees and scrub on either side. No aquatic vegetation. Dry in places. Completely shaded. Visible from adjacent parallel footpath to the south.	Average

Pond No.	Description	HSI score
<b>A1 Ditch 3</b>	Ditch on field boundary at the base of a wooded bank. Shaded by adjacent trees. No aquatic vegetation. Some stagnant water. Ditch continues west, parallel to the A23 but is mostly dry.	Average
<b>Un-named additional ditch 1</b>	Additional ditch recorded under bridge.	????
<b>A1 Ditch 5</b>	Ditch parallel to the B2036. Wet with some aquatic vegetation. Visible along the side of the road.	????
<b>A1 Ditch 4</b>	Wet ditch at the base of an earth bank for the A23. No aquatic vegetation. Quite shaded by trees either side. Visible from the B2036.	Below average
<b>A1 services ditch</b>	Dry ditch parallel to the A23/Airport Way, at the bottom of the earth bank for the A23. Surrounded by trees. Access from adjacent footpath.	????
<b>A1 Services lake</b>	Large lake behind the services area. No aquatic vegetation. Wildfowl present, kingfisher seen twice. Banks are lined with trees. Access from adjacent footpath.	????
<b>A1 additional ditch by M23</b>	Wet ditch with reeds adjacent to the M23 slip road. Trees and scrub either side. Access – can see from adjacent footpath but surrounded by dense scrub.	????
<b>A2 Ditch 5</b>	Wet ditch parallel to a railway embankment. Occasional adjacent trees and scrub. No aquatic vegetation. No direct access, viewed through fence.	Average
<b>A2 SUDS area</b>	Large SUDS area in the middle of the car park. Surrounded by trees, scrub and grassland.	????
<b>A2 Ditch 6</b>	Dry ditch along the edge of some woodland. No aquatic vegetation. Also continues along the southern edge of the car park.	Average

Pond No.	Description	HSI score
<b>A2 Ditch 7</b>	Wet ditch with some emergent vegetation. Surrounded by car parking with some adjacent trees.	Good
<b>A2 Ditch 8</b>	Ditch surrounded by car parking. Partly wet at eastern end with some aquatic vegetation. Adjacent trees along south side of ditch.	Good
<b>A2 Ditch 9</b>	Dry ditch along the edge of an access track with mature trees on its northern bank. No aquatic vegetation.	Average
<b>A2 Ditch 10</b>	Partly wet ditch along the edge of an access track. No aquatic vegetation. Mature trees present on the southern bank of the ditch.	Average
<b>A2 additional ditch south of D10</b>	Additional ditch parallel to D10, to the south, along the edge of a car park. Partly wet but no aquatic vegetation.	????
<b>A2 Ditch 11</b>	There are two ditches here, either side of the road. Both are dry with no aquatic vegetation.	Average
<b>A2 Ditch 12</b>	Wet ditch through a car park with adjacent mature trees and dense bracken. No aquatic vegetation.	Below average
<b>A2 Ditch 13 - West</b>	Dry ditch along a tree line. No aquatic vegetation.	Average
<b>A2 Ditch 13 - East</b>	Wet ditch along a tree line with scrub either side. No aquatic vegetation.	????
<b>A2 Ditch 14</b>	Wet ditch surrounded by scrub and self-seeded tree saplings. No aquatic vegetation present.	Good
<b>A2 Pond 1</b>	Large woodland pond with some emergent vegetation.	Average
<b>A2 Pond 2</b>	Small woodland pond with over-hanging trees. No aquatic vegetation present.	????
<b>A3 Pond 1</b>	New environmental pond. Lined. No aquatic vegetation visible apart from duckweed. Pond situated in an area of woodland, adjacent to a large wet meadow. Habitat connectivity to P2.	Below average

Pond No.	Description	HSI score
<b>A3 Pond 2</b>	Woodland pond surrounded by tussocky grassland and trees. No aquatic vegetation. GCN recorded in 2021 during bottle trapping.	Average
<b>A3 Green Ditch 1</b>	Wet ditch within a large field of tussocky grassland / wet meadow. Some young willow trees growing either side of the ditch. No aquatic vegetation but surrounding habitat very good, with habitat connectivity to P2 with confirmed GCN presence.	Below average
<b>A3 Purple Ditch 1</b>	Ditch along the north side of a grass access track. Wet but no aquatic vegetation. A hedge is on the north side of the ditch.	????
<b>A3 Purple Ditch 2</b>	Ditch along the south side of a grass access track. Wet but no aquatic vegetation. A hedge is on the south side of the ditch.	????
<b>A3 Purple Ditch 3</b>	Ditch along the east side of a grass access track. Wet but no aquatic vegetation. A hedge is on the east side of the ditch.	????
<b>A3 Purple Ditch 4</b>	Ditch along the west side of a grass access track. Wet but no aquatic vegetation. Woodland is adjacent to the west side of the ditch.	????
<b>A5 Ditch 1</b>	Wet ditch along the edge of a grass field, with adjacent scrub and trees. No aquatic vegetation. Ditch is only adjacent to the southern field boundary, there is no ditch along the eastern field boundary as marked on the map.	Excellent
<b>A5 Ditch 2</b>	Wet ditch with grass banks along the edges of a field. No aquatic vegetation.	Good
<b>A5 Ditch 3</b>	Dry ditch with grass banks along the edge of a field. Adjacent trees and hedge. Visible from Aviation museum field to the west. Direct access via clay shooting school.	Excellent



Pond No.	Description	HSI score
<b>A5 Pond 1</b>	No pond at the location shown on the map.	Below average
<b>A5 additional Pond 2</b>	Large pond with dense reeds surrounded by grassland.	????
<b>A5 additional Pond 3</b>	Small pond surrounded by trees and tussocky grassland. No aquatic vegetation.	????
<b>B2036 Ditch 1</b>	Unknown location, not labelled on map.	Below average
<b>B2036 Ditch 2</b>	Unknown location, not labelled on map.	Poor
<b>B2036 Ditch 3</b>	????	Good
<b>B2036 Ditch 4</b>	????	Below average
<b>Crawter's Wood- west</b>	No access available. This pond is airside so security access and a security escort need to be pre-arranged for access.	Poor
<b>Crawter's Wood- east</b>	No access available. This pond is airside so security access and a security escort need to be pre-arranged for access.	Poor
<b>Detention Centre Pond</b>	No access available.	Below average
<b>Ditch, North boundary, Car park X</b>	Partly wet ditch with bulrushes. Grass and ruderal vegetation on the banks. Surrounded by hard-standing.	Good
<b>Ditch, Southern boundary, Car park X</b>	Wet ditch along southern edge of car park. Trees and scrub adjacent to the south. No aquatic vegetation.	Good
<b>Ditch, Western boundary, Car park X</b>	Wet ditch along the edge of the car park with some aquatic vegetation and adjacent scrub.	????
<b>Pond, Car Park X</b>	Additional pond. In the centre of the car park. Surrounded by trees and grassland. No aquatic vegetation.	????

Pond No.	Description	HSI score
<b>North long stay A6/A7- Ditch 1</b>	Ditch runs through the centre of a car park. No connecting habitat. The ditch holds water but no aquatic vegetation is present. The banks are lined with self-seeded tree saplings and some scrub.	Below average
<b>North long stay A6/A7- Ditch 2</b>	Ditch runs along the southern and western boundaries of the car park. Wet, but no visible aquatic vegetation. Bordered by scrub and self-seeded tree saplings. More open along western boundary.	Excellent
<b>North long stay A6/A7- Ditch 3</b>	Ditch runs parallel to southern edge of car park. Some water but no visible aquatic vegetation. Heavily shaded by adjacent trees and scrub.	Average
<b>North long stay A6/A7- Ditch 4</b>	Ditch between two car parks. Some water but no visible aquatic vegetation. Heavily shaded by adjacent trees and scrub.	Average
<b>North long stay A6/A7- Pond 1</b>	Woodland pond surrounded by trees and grassland. Some emergent vegetation present.	Good
<b>Woodland behind Premier Inn- Ditch 1</b>	A8 South – Shallow wet ditch through an area of woodland. No aquatic vegetation present.	Below average
<b>Woodland behind Premier Inn- Riverside Garden</b>	A8 North – Ditch runs parallel to the A23 London Road, through an area of woodland. Dry in places. No aquatic vegetation. Access from within Riverside Gardens	Below average
<b>A8 Island ditch</b>	Island of land surrounded by road. There are a few metres of amenity grassland before a ditch, which encircles the western half of the island. The majority of the island is wooded. The ditch itself is deep but mostly dry, with no aquatic vegetation.	????
<b>Pond 1</b>	Large pond in the corner of an arable field with lots of emergent vegetation and adjacent trees.	Excellent

Pond No.	Description	HSI score
<b>Pond 4</b>	Large pond in the corner of a grass field surrounded by trees. No aquatic vegetation is present.	Poor
<b>Pond 5</b>	Small pond in the corner of a grass field, surrounded by trees. No aquatic vegetation is present.	Poor
<b>Pond 6</b>	Large garden pond with extensive emergent vegetation. Surrounded by amenity grassland. Mature trees line the garden boundary.	Excellent
<b>Additional pond right next to the M23</b>	Large drainage pond adjacent to the earth bank of the M23. Extensive emergent vegetation present. Trees and scrub are present along the northern edge of the pond.	Excellent

### Bat Emergence/Re-entry Surveys

#### Building JW9 (Landside)

##### Bat Emergence Survey 15 July 2019

- A2.1.1 The bat emergence survey on 15 July commenced at 21:00 hours, 15 minutes before sunset and finished at 22:45 hours.
- A2.1.2 No bats were seen emerging from the building but were detected foraging nearby. Bat activity was recorded at low levels during the survey.
- A2.1.3 The following bat activity was recorded during the survey:
- 21:59 – noctule heard but not seen;
  - 22:15 – noctule heard close by;
  - 22:19 – noctule heard close by; and
  - 22:22 – noctule heard close by.

##### Bat emergence survey 20 August 2019

- A2.1.4 The bat emergence survey on the 20<sup>th</sup> August commenced at 20:00 hours, 15 minutes before sunset and finished at 21:45 hours.
- A2.1.5 No bats were seen emerging from the building but were detected foraging and commuting nearby. Bat activity was recorded at low levels during the survey.
- A2.1.6 The following bat activity was recorded during the survey:

- 20:56 – Leisler’s bat pass and foraging soprano pipistrelle heard nearby;
- 20:58 – soprano pipistrelle heard foraging;
- 21:19 – soprano pipistrelle foraging; and
- 21:49 – brief Leisler’s bat pass, not seen.

#### Bat Emergence Survey 26 September 2019

- A2.1.7 The bat emergence survey on 26 September commenced at 18:55 hours, 15 minutes before sunset and finished at 20:30 hours.
- A2.1.8 Not bats were seen emerging from the building but were detected foraging and commuting nearby. Bat activity was recorded at moderate levels during the survey; although no bats were seen, it was presumed that bats were foraging near to the grassland area to the west of the feature.
- A2.1.9 The following bat activity was recorded during the survey.
- 19:18 – noctule heard but not seen.
  - 19:28 – distant noctule call.
  - 19:29 – brief *Myotis* call – not seen.
  - 19:30 – noctule heard but not seen – made several passes until 19:39, possibly over the grassland areas adjacent to building. Mainly foraging and social calls.
  - 19:32 – faint common pipistrelle call heard.
  - 19:39 – distant common pipistrelle call.
  - 19:45 – distant common pipistrelle call.
  - 19:46 – *Myotis* bat heard.
  - 19:48 – at least two noctule’s foraging nearby.
  - 20:03-20:14 – noctule, *Myotis* and pipistrelle heard foraging nearby; calls gradually getting quieter towards the end of the survey.

#### Building D9H (Airside)

#### Bat Emergence Survey 15 July 2019

- A2.1.10 The bat emergence survey on the 15 July commenced at 21:00 hours, 15 minutes before sunset and finished at 22:45 hours.
- A2.1.11 No bats were seen emerging from the building but were detected foraging and commuting nearby.
- A2.1.12 The following bat activity was recorded during the survey:
- 21:49 – faint common pipistrelle bat pass;
  - 22:04 – distant pass from common pipistrelle;
  - 22:11 – common pipistrelle commuting east along building;

- 22:12 – common pipistrelle heard but not seen, foraging nearby;
- 22:27 – noctule heard but not seen;
- 22:34 – brief noctule pass;
- 22:35 – brief noctule pass;
- 22:36 – brief noctule pass; and
- 22:39 – brief noctule pass.

#### Bat Emergence Survey 7 August 2019

- A2.1.13 The bat emergence survey on 7 August commenced at 20:26 hours, 15 minutes before sunset and finished at 22:11 hours.
- A2.1.14 No bats were seen emerging from the building and only a single Noctule was recorded briefly at 21:43 hours.

#### Bat Emergence Survey 2 October

- A2.1.15 The bat emergence survey on 2 October commenced at 18:22 hours, 15 minutes before sunset, and finished at 20:07 hours.
- A2.1.16 No bats were seen emerging from the building but were recorded foraging nearby. Bat activity was recorded at low levels during the survey; noctule were heard making regular, brief passes between 19:11 and 20:03 hours.

#### Bat Activity Transect Surveys

#### Annex 2.1.42: Bat Activity Transect Survey Dates, Weather Conditions and Sunset Times

Survey date	Sunset time	Survey start	Weather conditions
<b>Transect 1</b>			
09/04/19	19:46	19:31	8°C, cloudy, light breeze, no rain
24/04/19	20:10	19:57	10°C, heavy cloud cover, light wind
08/05/19	20:33	20:26	11°C, dry, light cloud, light breeze
21/05/19	20:52	20:37	17°C, no cloud, light breeze
12/06/19	21:16	21:09	13°, overcast, occasional light rain
25/06/19	21:20	21:05	22°C, humid, cloudy, light wind
09/07/19	21:15	21:00	20°C, dry, warm, overcast
23/07/19	21:01	20:46	26°C, clear, hot, humid

Survey date	Sunset time	Survey start	Weather conditions
06/08/19	20:41	20:25	18°C, cloudy, calm
28/08/19	19:56	19:40	20°C, cloudy, light breeze, no rain
03/09/19	19:41	19:36	18°C, dry, cloudy, light wind
25/09/19	18:55	18:40	16°C, patchy cloud, dry, light wind
15/10/19	18:09	19:50	16°C, clear sky, dry, light wind
30/10/19	16:41	16:25	10°C, light wind, clear, dry
<b>Transect 2</b>			
09/04/19	19:46	19:31	8°C, cloudy, light breeze, no rain
24/04/19	20:10	19:58	10°C, heavy cloud cover, light wind
08/05/19	20:33	20:18	11°C, dry, light cloud, light breeze
21/05/19	20:52	20:37	17°C, no cloud, light breeze
12/06/19	21:16	21:09	13°, overcast, occasional light rain
25/06/19	21:20	21:05	22°C, humid, cloudy, light wind
09/07/19	21:15	21:00	20°C, dry, warm, overcast
23/07/19	21:01	20:46	26°C, clear, hot, humid
06/08/19	20:41	20:25	18°C, cloudy, calm
28/08/19	19:56	19:40	20°C, cloudy, light breeze, no rain
25/09/19	18:55	18:40	16°C, patchy cloud, dry, light wind
16/10/19	18:09	19:50	16°C, clear sky, dry, light wind
30/10/19	16:41	16:25	10°C, light wind, clear, dry
<b>Transect 3</b>			
09/04/19	19:46	19:26	12°C, cloudy, no wind
24/04/19	20:10	20:00	Heavy cloud. damp
08/05/19	20:34	20:18	11°C, dry, light cloud, light breeze
21/05/19	20:52	20:43	17°C, no cloud, light breeze

Transect 7- Riverside			
Survey date	Sunset time	Survey start	Weather conditions
10/08/2020	20:31	20:16	30°C, dry, sunny, humid
25/08/2020	20:00	19:45	20°C, wind 6, rain in the day
7/09/2020	19:17	19:37	20°C, cloudy (95%), light breeze (2), dry
21/09/2020	19:00		17°C, cool, dry, little to no breeze
5/10/2020	18:28	18:13	13 °c, complete cloud cover, rain in day
27/10/2020	18:25		17 °c, light rain, little wind, cloudy
15/04/2021	19:57	19:42	10°C, clear skies, light breeze
27/04/2021	20:16	20:02	12°C, mild, wind-2, sunny in the day
18/05/2021	20:48	20:31	13°C, overcast, dry, light breeze
25/05/2021	20:58	20:43	14°C, clear sky, light breeze
8/06/2021	21:13	20:58	19°C, dry, no wind, 3/8 cloud
22/06/2021	21:20	21:05	14°C, dry, no wind, 2/8 cloud
6/07/2021	21:17	21:20	14°C, dry but rain in the day, wind-2
19/07/2021	21:05	20:40	23°C, mild, little breeze, dry
Survey date	Sunset time	Survey start	Weather conditions
18/06/19	21:19	21:00	16°C, dry, overcast, light wind
25/06/19	21:20	21:05	22°C, humid, cloudy, light wind
09/07/19	21:15	21:00	20°C, dry, warm, overcast

Survey date	Sunset time	Survey start	Weather conditions
23/07/19	21:01	20:46	26°C, clear, hot, humid
06/08/19	20:41	20:25	18°C, cloudy, calm
29/08/19	19:54	19:46	19°C, light cloud, no rain
03/09/19	19:41	19:33	18°C, dry, cloudy, light wind
25/09/19	18:55	18:40	16°C, patchy cloud, dry, light wind
16/10/19	18:09	19:50	16°C, clear sky, dry, light wind
30/10/19	16:41	16:25	10°C, light wind, clear, dry
Transect 4			
10/04/19	19:48	19:32	10°C, clear skies, light breeze
25/04/19	20:12	19:57	11°C, high cloud, light wind
13/05/19	20:42	20:27	11°C, light winds, fair
22/05/19	20:54	20:39	19°C, clear, dry, no wind
13/06/19	21:17	21:00	13°C, cloudy, occasional light rain
26/06/19	21:20	21:05	18°C, cloudy, windy
10/07/19	21:15	21:00	17°C, dry, light wind, patchy cloud
24/07/19	21:00	20:40	27°C, patchy cloud, no wind
05/08/19	20:41	20:20	20°C, light wind, no rain
29/08/19	19:54	19:46	19°C, light cloud, no rain
04/09/19	19:40	19:25	17°C, clear, breezy
24/09/19	18:55	18:40	18°C, light wind and light cloud
15/10/19	18:09	19:50	16°C, clear sky, dry, light wind
29/10/19	16:40	16:25	13°C, cloudy, light wind
Transect 5			
10/04/19	19:48	19:32	10°, clear skies, light breeze
25/04/19	20:12	19:57	11°C, high cloud, light wind
14/05/19	20:43	20:28	12°C, moderate breeze, fair
22/05/19	20:54	20:39	19°C, clear, dry, no wind
13/06/19	21:17	21:02	13°C, cloudy, occasional light rain
26/06/19	21:20	21:05	18°C, cloudy, windy

Survey date	Sunset time	Survey start	Weather conditions
10/07/19	21:15	21:00	17°C, dry, light wind, patchy cloud
24/07/19	21:00	20:40	27°C, patchy cloud, no wind
05/08/19	20:41	20:20	20°C, light wind, no rain
29/08/19	19:54	19:46	19°C, light cloud, no rain
04/09/19	19:40	19:25	17°C, clear, breezy
24/09/19	18:55	18:40	18°C, light wind and light cloud
15/10/19	18:09	19:50	18°C, clear sky, dry, light wind
29/10/19	16:40	16:25	13°C, cloudy, light wind
Transect 6-Mole corridor			
10/08/2020	20:31	20:16	29°C, cloud 20%, no wind
25/08/2020	19:58		19 °c, cloudy, windy, slight rain
7/09/2020	19:32	19:17	19 °c, little breeze, foggy
21/09/2020	19:00	19:00	19 °c, dry, little wind, cloudy 6/8
5/10/2020	18:28	18:13	15°C, complete cloud cover
14/04/2021	19:54	19:39	11°C, cloudy 70%, little wind
26/04/2021	20:14	19:59	12 °c, slight wind (2), no rain
13/05/2021	20:41	20:28	18 °c, 80% cloud cover, light breeze
26/05/2021	20:33		11°C, slight wind, cloudy
7/06/2021	21:12	21:34	17 °c, dry, clear, no wind
24/06/2021	21:20	21:05	19 °c, light breeze
6/07/2021	21:25	21:17	15 °c, moderate breeze, dry
19/07/2021	21:05	20:50	26 °c, dry, sunny

Survey date	Sunset time	Survey start	Weather conditions
<b>Transect 8- Surrey CC</b>			
11/08/2020	20:17		28°C, little cloud, light wind
25/08/2020	20:00	19:45	20°C, wind 6, rain in the day Used from other transect
8/09/2020	19:30	19:15	24°C, partial cloud (40%), no wind, dry
22/09/2020	18:58	18:43	19°C, full cloud cover, light wind, dry
6/10/2020	18:26	18:11	15 °c, partial cloud cover (50%), slight breeze (4)
28/10/2020	18:24		17 °c, light rain, little wind, cloudy
14/04/2021	19:55	19:39	10°C, 4/8 cloud, little breeze
26/04/2021	20:14	19:57	10°C, 6/8 cloud, light breeze
12/05/2021	20:41	20:26	15°C, overcast, no breeze
07/06/2021	21:12	20:51	18°C, light clouds, dry, light breeze
24/06/2021	21:20	21:05	20°C, 8/8 cloud, little wind
7/07/2021	21:23		14°C, scattered clouds, light breeze
20/07/2021	21:05	20:50	23°C, dry, no clouds, little breeze
<b>Transect 9- Church meadows</b>			
19/10/202	18:15	18:10	17°C, 8/8 cloud, light breeze
<b>Transect 10- Dairy Farm</b>			

Survey date	Sunset time	Survey start	Weather conditions
28/05/2022	20:59	20:59	15°C, 7/8 cloud, dry, little breeze
10/06/2022	21:15	21:15	17°C, no cloud, dry, slight breeze
22/06/2022	21:20	21:20	20°C, no cloud, dry, little breeze
10/07/2022	21:15		22°C, no cloud, dry, no wind
20/07/2022	21:04	21:04	24°C, 8/8 cloud, dry, little breeze
7/08/2022	20:38	20:38	21°C, 1/8 cloud, dry, no wind
23/08/2022	20:06		22°C, 8/8 cloud, dry, little wind
12/09/2022	19:22	19:22	24°C, 8/8 cloud, dry, no wind
29/09/2022	18:42	18:59	13°C, no cloud, dry, no wind
12/10/2022	18:20	18:20	17°C, 8/8 cloud, dry, little breeze

### Pre-maternity

#### Transect 1

A2.1.17 A total of four visits were undertaken for Transect 1 during the pre-maternity season in 2019: 9 April, 24 April, 8 May and 21 May.

A2.1.18 A total of 240 bat passes were recorded during the surveys. These comprised passes from:

- 217 common pipistrelles;
- 19 soprano pipistrelles;
- Three *Myotis* sp. (including two characteristic of whiskered/Brandt's *Myotis mystacinus/brandtii* bat); and
- One noctule.

A2.1.19 Figure 3.12.1a shows the transect route and the number and location of species recorded during the surveys.

#### Transect 2

A2.1.20 A total of four visits were undertaken for Transect 2 during the pre-maternity season in 2019: 9 April, 24 April, 8 May and 21 May.

A2.1.21 A total of 217 bat passes were recorded during the surveys. These comprised passes from:

- 192 common pipistrelles;
- Two soprano pipistrelles;
- Three *Myotis* sp. (including two characteristic of Daubenton's bat *Myotis daubentonii*);
- One long-eared *Plecotus* sp. bat; and
- 19 noctule.

A2.1.22 Figure 3.12.1b shows the transect route and the number and location of species recorded during the surveys

#### Transect 3

A2.1.23 A total of four visits were undertaken for Transect 3 during the pre-maternity season in 2019: 9 April, 24 April, 8 May and 21 May.

A2.1.24 A total of 286 bat passes were recorded during the surveys. These comprised passes from:

- 242 common pipistrelles;
- 30 soprano pipistrelles; and
- 14 *Myotis* sp. (including three characteristic of Natterer's bat *Myotis nattereri* and one of Daubenton's bat).

A2.1.25 Figure 3.12.1c and 3.12.3d shows the transect route and the number and location of species recorded during the surveys.

#### Transect 4

A2.1.26 A total of four visits were undertaken for Transect 4 during the pre-maternity season in 2019: 10 April, 25 April, 13 May and 22<sup>nd</sup> May.

A2.1.27 A total of 24 bat passes were recorded during the surveys. These comprised passes from:

- 21 common pipistrelles; and
- Three noctule.

A2.1.28 Figure 3.12.1e shows the transect route and the number and location of species recorded during the surveys.

#### Transect 5

A2.1.29 A total of four visits were undertaken for Transect 5 during the pre-maternity season in 2019: 10 April, 25 April, 14 May and 22 May.

A2.1.30 A total of 131 bat passes were recorded during the surveys. These comprised passes from:

- 77 common pipistrelles;
- 12 soprano pipistrelles;
- One pipistrelle species;
- 25 *Myotis* sp. (including three characteristic of whiskered/Brandt's bats, two of Daubenton's bats and four of Natterer's bats); and
- 16 noctule.

A2.1.31 Figure 3.12.1f shows the transect route and the number and location of species recorded during the surveys.

#### Transect 6

A2.1.32 A total of four visits were undertaken for Transect 6 during the pre-maternity season in 2021: 14 April, 26 April, 13 May and 26 May.

A2.1.33 A total of 313 bat passes were recorded during the surveys. These comprised passes from:

- 256 common pipistrelles;
- 2 soprano pipistrelles;
- 9 *Myotis* sp.;
- 44 noctule; and
- 2 *Plecotus* sp.

A2.1.34 Figures 3.12.4f-h shows the transect route and the number and location of species recorded during the surveys.

#### Transect 7

A2.1.35 A total of four visits were undertaken for Transect 7 during the pre-maternity season in 2021: 15 April, 27 April, 18 May and 25 May.

A2.1.36 A total of 436 bat passes were recorded during the surveys. These comprised passes from:

- 341 common pipistrelles;

- 57 soprano pipistrelles;
- 7 unidentified pipistrelle species;
- 4 *Myotis* sp.

A2.1.37 Figure 3.12.5f-h shows the transect route and the number and location of species recorded during the surveys.

#### Transect 8

A2.1.38 A total of three visits were undertaken for Transect 8 during the pre-maternity season in 2021: 14 April, 26 April, and 12 May.

A2.1.39 A total of 250 bat passes were recorded during the surveys. These comprised passes from:

- 144 common pipistrelles;
- 2 soprano pipistrelles;
- 1 unidentified pipistrelle species;
- 3 *Myotis* sp.

A2.1.40 Figure 3.12.6f-h shows the transect route and the number and location of species recorded during the surveys.

#### Transect 9

A2.1.41 One visit was undertaken for Transect 10 during the pre-maternity season in 2022: 28 May.

A2.1.42 A total of 90 bat passes were recorded during the survey. These comprised passes from:

- 77 common pipistrelles;
- 10 soprano pipistrelles;
- 2 noctule.

A2.1.43 Figure 3.12.7a shows the transect route and the number and location of species recorded during the survey.

#### Maternity

##### Transect 1

A2.1.44 A total of four visits were undertaken for Transect 1 during the maternity season in 2019: 12 June, 25 June, 9 July and 23 July.

A2.1.45 A total of 400 bat passes were recorded during the surveys. These comprised passes from:

- 301 common pipistrelles;
- 56 soprano pipistrelles;
- 16 *Myotis* sp. (including one characteristic of whiskered/Brandt's bat and one of Natterer's bat);

- 15 noctule;
- Six Leisler's bats;
- Ten *Nyctalus* sp.; and
- Six serotine bats.

A2.1.46 Figure 3.13.2a shows the transect route and the number and location of species recorded during the surveys.

#### Transect 2

A2.1.47 A total of four visits were undertaken for Transect 2 during the maternity season in 2019: 12 June, 25 June, 9 July and 23 July.

A2.1.48 A total of 218 bat passes were recorded during the surveys. These comprised passes from:

- 197 common pipistrelles;
- Two soprano pipistrelles;
- One Nathusius' pipistrelle;
- 14 noctule; and
- Four serotine bats.

A2.1.49 Figure 3.12.2b shows the transect route and the number and location of species recorded during the surveys

#### Transect 3

A2.1.50 A total of four visits were undertaken for Transect 3 during the maternity season in 2019: 18 June, 2 June, 9 July and 23 July.

A2.1.51 A total of 252 bat passes were recorded during the surveys. These comprised passes from:

- 211 common pipistrelles;
- 31 soprano pipistrelles;
- Two *Myotis* sp. (including one characteristic of Natterer's bat);
- One noctule;
- One *Nyctalus* sp.; and
- Six serotine bats.

A2.1.52 Figure 3.12.2c and 3.12.2d shows the transect route and the number and location of species recorded during the surveys.

#### Transect 4

A2.1.53 A total of four visits were undertaken for Transect 4 during the maternity season in 2019: 13 June, 26 June, 10 July and 24 July.

A2.1.54 A total of 23 bat passes from common pipistrelles were recorded during the surveys.

<p>A2.1.55 Figure 3.12.2e shows the transect route and the number and location of species recorded during the surveys.</p>	<ul style="list-style-type: none"> <li>▪ 12 <i>Myotis</i> sp.</li> <li>▪ 1 brown long-eared bat.</li> </ul>	<ul style="list-style-type: none"> <li>▪ Nine <i>Myotis</i> sp.;</li> <li>▪ One <i>Plecotus</i> sp.</li> <li>▪ 16 noctule;</li> <li>▪ One serotine bat; and</li> <li>▪ Two <i>Nyctalus</i> sp.</li> </ul>
<p>A2.1.56</p> <p><b>Transect 5</b></p>	<p>A2.1.65 Figure 3.12.5i-k shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Transect 8</b></p>	
<p>A2.1.57 A total of four visits were undertaken for Transect 5 during the maternity season in 2019: 13 June, 26 June, 10 July and 24 July.</p>	<p>A2.1.66 A total of four visits were undertaken for Transect 6 during the maternity season in 2021: 7 June, 24 June, 7 July and 20 July.</p>	<p>A2.1.74 Figure 3.12.3a shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Transect 2</b></p>
<p>A2.1.58 A total of 333 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 260 common pipistrelles;</li> <li>▪ 32 soprano pipistrelles;</li> <li>▪ 23 <i>Myotis</i> sp. (including three characteristic of whiskered/Brandt's bats and one of Daubenton's bat);</li> <li>▪ 15 noctule;</li> <li>▪ Two Leisler's bats; and</li> <li>▪ One <i>Nyctalus</i> sp.</li> </ul>	<p>A2.1.67 A total of 251 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 245 common pipistrelles;</li> <li>▪ 2 soprano pipistrelles;</li> <li>▪ 4 Nathusius' pipistrelle .</li> </ul>	<p>A2.1.75 A total of five visits were undertaken for Transect 2 during the post-maternity season in 2019: 6 August, 28 August, 25 September, 16 October and 30 October. One survey was cancelled in early September due to access constraints.</p>
<p>A2.1.59 Figure 3.12.2f shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Transect 6</b></p>	<p>A2.1.68 Figure 3.12.6i-k shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Transect 9</b></p>	<p>A2.1.76 A total of 243 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 227 common pipistrelles;</li> <li>▪ Four soprano pipistrelles;</li> <li>▪ Five <i>Myotis</i> sp. (including one characteristic of Daubenton's bat);</li> <li>▪ One <i>Plecotus</i> sp.;</li> <li>▪ Four noctule; and</li> <li>▪ Two <i>Nyctalus</i> sp.</li> </ul>
<p>A2.1.60 A total of four visits were undertaken for Transect 6 during the maternity season in 2021: 7 June, 24 June, 6 July and 19 July.</p>	<p>A2.1.69 A total of four visits were undertaken for Transect 6 during the maternity season in 2022: 10 June, 22 June, 10 July and 20 July.</p>	
<p>A2.1.61 A total of 192 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 149 common pipistrelles;</li> <li>▪ 9 soprano pipistrelles;</li> <li>▪ 2 Nathusius' pipistrelle;</li> <li>▪ 10 <i>Myotis</i> sp.</li> <li>▪ 21 noctule.</li> <li>▪ 1 brown long-eared bat</li> </ul>	<p>A2.1.70 A total of 462 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 374 common pipistrelles;</li> <li>▪ 84 soprano pipistrelles;</li> <li>▪ 1 noctule.</li> <li>▪ 1 <i>Nyctalus</i> sp.</li> <li>▪ 1 Serotine</li> <li>▪ 1 <i>Plecotus</i> sp.</li> </ul>	<p>A2.1.77 Figure 3.12.3b shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Transect 3</b></p>
<p>A2.1.62 Figure 3.12.4i-k shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Transect 7</b></p>	<p>A2.1.71 Figure 3.12.7b-e shows the transect route and the number and location of species recorded during the surveys.</p> <p><b>Post-maternity</b></p> <p><b>Transect 1</b></p>	<p>A2.1.78 A total of six visits were undertaken for Transect 3 during the post-maternity season in 2019: 6 August, 29 August, 3 September, 25 September, 16 October and 30 October.</p>
<p>A2.1.63 A total of four visits were undertaken for Transect 6 during the maternity season in 2021: 8 June, 22 June, 6 July and 19 July.</p>	<p>A2.1.72 A total of six visits were undertaken for Transect 1 during the post-maternity season in 2019: 6 August, 28 August, 3 September, 25 September, 15 October and 30 October.</p>	<p>A2.1.79 A total of 378 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 328 common pipistrelles;</li> <li>▪ 37 soprano pipistrelles;</li> <li>▪ One Nathusius' pipistrelle;</li> <li>▪ Five <i>Myotis</i> sp.;</li> <li>▪ Three Leisler's bats; and</li> <li>▪ Four noctule.</li> </ul>
<p>A2.1.64 A total of 480 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 402 common pipistrelles;</li> <li>▪ 65 soprano pipistrelles;</li> </ul>	<p>A2.1.73 A total of 508 bat passes were recorded during the surveys. These comprised passes from:</p> <ul style="list-style-type: none"> <li>▪ 433 common pipistrelles;</li> <li>▪ 46 soprano pipistrelles;</li> </ul>	<p>A2.1.80 Figure 3.12.3c and 3.12.3d shows the transect route and the number and location of species recorded during the surveys.</p>

#### Transect 4

A2.1.81 A total of six visits were undertaken for Transect 4 during the post-maternity season in 2019: 5 August, 29 August, 4 September, 24 September, 15 October and 29 October.

A2.1.82 A total of 52 passes were recorded during the surveys. These comprised passes from:

- 32 common pipistrelles;
- Four soprano pipistrelles;
- One Nathusius' pipistrelle;
- 12 noctule;
- One Leisler's bat; and
- Two *Nyctalus* sp.

A2.1.83 Figure 3.12.3e shows the transect route and the number and location of species recorded during the surveys.

#### Transect 5

A2.1.84 A total of six visits were undertaken for Transect 5 during the post-maternity season in 2019: 5 August, 29 August, 4 September, 24 September, 15 October and 29 October.

A2.1.85 A total of 297 bat passes were recorded during the surveys. These comprised passes from:

- 212 common pipistrelles;
- 16 soprano pipistrelles;
- 47 *Myotis* sp. (including 12 characteristic of whiskered/Brandt's bat, six Daubenton's bats and six Natterer's bats);
- One serotine bat;
- Five *Plecotus* sp.;
- 14 noctule; and
- Two *Nyctalus* sp.

A2.1.86 Figure 3.12.3f shows the transect route and the number and location of species recorded during the surveys.

#### Transect 6

A2.1.87 A total of five visits were undertaken for Transect 6 during the post-maternity season in 2020: 10 August, 25 August, 7 September, 21 September, and 5 October.

A2.1.88 A total of 296 bat passes were recorded during the surveys. These comprised passes from:

- 177 common pipistrelles;

- 30 soprano pipistrelles;
- 39 *Myotis* sp.
- 50 noctule;

A2.1.89 Figure 3.12.4a-e shows the transect route and the number and location of species recorded during the surveys.

#### Transect 7

A2.1.90 A total of six visits were undertaken for Transect 7 during the post-maternity season in 2020: 10 August, 25 August, 7 September, 21 September, 5 October and 27 October.

A2.1.91 A total of 408 bat passes were recorded during the surveys. These comprised passes from:

- 368 common pipistrelles;
- 18 soprano pipistrelles;
- 19 *Myotis* sp.
- 3 noctule.

A2.1.92 Figure 3.12.5a-e shows the transect route and the number and location of species recorded during the surveys.

#### Transect 8

A2.1.93 A total of six visits were undertaken for Transect 8 during the post-maternity season in 2020: 11 August, 25 August, 8 September, 22 September, 6 October and 28 October.

A2.1.94 A total of 541 bat passes were recorded during the surveys. These comprised passes from:

- 507 common pipistrelles;
- 12 soprano pipistrelles;
- 9 *Myotis* sp.
- 1 serotine bat;
- 12 noctule.

A2.1.95 Figure 3.12.6a-e shows the transect route and the number and location of species recorded during the surveys.

#### Transect 9

A2.1.96 A total of five visits were undertaken for Transect 10 during the post-maternity season in 2022: 7 August, 23 August, 12 September, 29 September, and 12 October.

A2.1.97 A total of 965 bat passes were recorded during the surveys. These comprised passes from:

- 701 common pipistrelles;

- 232 soprano pipistrelles;
- 1 Nathusius's pipistrelle sp;
- 12 *Myotis* sp.
- 4 serotine bat;
- 1 Leisler's bat
- 1 *Plecotus* sp.;
- 11 noctule; and
- 2 *Nyctalus* sp.
- 

A2.1.98 Figure 3.12.7f-j shows the transect route and the number and location of species recorded during the surveys.

Bat Static/Automated Surveys

Annex 2.1.43: Bat Records at Location 1

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
24/04/19	0	0	0	0	0	0	0	0	0	0
25/04/19	2	0	0	0	1	0	0	0	0	3
26/04/19	1	0	0	0	0	0	1	0	0	2
27/04/19	0	0	0	0	0	0	0	0	0	0
28/04/19	16	0	1	0	1	1	2	0	0	21
29/04/19	10	0	0	0	1	0	0	0	0	11
30/04/19	22	0	0	0	0	0	0	0	0	22
<b>Species total</b>	<b>51</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>3</b>	<b>1</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>59</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	14	1	0	0	0	0	0	0	0	15
11/05/19	159	4	3	2	0	0	4	0	0	172
12/05/19	114	8	0	0	0	0	3	0	0	125
13/05/19	116	0	0	0	0	0	0	0	0	116
14/05/19	64	1	1	0	0	0	0	0	0	66
15/05/19	65	5	0	0	0	2	0	0	0	72
<b>Species total</b>	<b>532</b>	<b>19</b>	<b>4</b>	<b>2</b>	<b>0</b>	<b>2</b>	<b>7</b>	<b>0</b>	<b>0</b>	<b>566</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
11/06/19	0	0	0	0	0	0	0	0	0	0
12/06/19	0	0	0	0	0	0	0	0	0	0
13/06/19	9	0	0	0	0	0	3	0	0	12
14/06/19	69	0	0	0	0	0	1	0	0	70
15/06/19	99	4	0	0	0	0	4	0	0	107
<b>Species total</b>	<b>177</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>8</b>	<b>0</b>	<b>0</b>	<b>189</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	81	1	0	1	4	1	9	3	1	101
13/07/19	171	3	1	1	3	1	36	2	1	219
14/07/19	119	0	0	2	1	0	54	0	0	176
15/07/19	80	0	0	2	2	0	4	0	0	88
16/07/19	104	4	0	0	4	1	48	0	0	161
<b>Species total</b>	<b>555</b>	<b>8</b>	<b>1</b>	<b>6</b>	<b>14</b>	<b>3</b>	<b>151</b>	<b>5</b>	<b>2</b>	<b>745</b>



Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	43	1	0	0	4	0	6	0	1	55
14/08/19	34	0	0	0	1	1	5	0	1	42
15/08/19	106	2	0	0	6	0	4	0	0	118
16/08/19	2	0	0	0	1	0	2	0	0	5
17/08/19	16	0	0	0	2	0	14	0	1	33
18/08/19	21	0	3	0	3	2	0	0	0	29
<b>Species total</b>	<b>222</b>	<b>3</b>	<b>3</b>	<b>0</b>	<b>17</b>	<b>3</b>	<b>31</b>	<b>0</b>	<b>3</b>	<b>282</b>

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	0	15	0	0	3	1	0	0	0	0	19
26/09/19	0	5	0	0	0	2	0	53	0	0	60
27/09/19	0	1	0	0	0	2	0	158	0	0	161
28/09/19	0	2	0	0	0	1	0	31	0	0	34
29/09/19	1	11	0	0	0	1	0	70	0	0	83
<b>Species total</b>	<b>1</b>	<b>34</b>	<b>0</b>	<b>0</b>	<b>3</b>	<b>7</b>	<b>0</b>	<b>312</b>	<b>0</b>	<b>0</b>	<b>357</b>

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	0	8	0	0	0	0	0	2	0	0	10
15/10/19	0	91	1	0	8	3	0	14	0	0	117
16/10/19	1	4	2	0	0	3	0	0	0	0	10
17/10/19	0	0	0	0	0	0	0	0	0	0	0
18/10/19	0	0	0	0	0	1	0	0	0	0	1
<b>Species total</b>	<b>1</b>	<b>103</b>	<b>3</b>	<b>0</b>	<b>8</b>	<b>7</b>	<b>0</b>	<b>16</b>	<b>0</b>	<b>0</b>	<b>138</b>

#### Annex 2.1.44: Bat Records at Location 2

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
24/04/19	0	0	0	0	0	0	0	0	0	0
25/04/19	4	2	0	0	1	0	0	0	0	7
26/04/19	2	0	0	0	0	0	1	0	0	3
27/04/19	13	0	0	0	1	0	0	0	0	14
28/04/19	35	2	0	0	5	0	0	0	0	42
29/04/19	28	1	0	0	1	0	0	0	0	30
30/04/19	14	0	0	0	0	0	0	0	0	14
<b>Species total</b>	<b>96</b>	<b>5</b>	<b>0</b>	<b>0</b>	<b>8</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>110</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	58	2	0	0	0	0	4	0	0	64
11/05/19	173	6	0	0	3	0	7	0	0	189
12/05/19	135	10	0	0	1	1	16	0	0	163
13/05/19	241	22	0	0	1	0	4	0	0	268
14/05/19	217	8	0	0	0	0	20	0	0	245
15/05/19	124	0	0	0	2	0	46	0	0	172
<b>Species total</b>	<b>948</b>	<b>48</b>	<b>0</b>	<b>0</b>	<b>7</b>	<b>1</b>	<b>97</b>	<b>0</b>	<b>0</b>	<b>1,101</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/06/19	7	0	0	0	0	0	25	0	0	32
13/06/19	15	3	0	0	6	0	131	0	0	155
14/06/19	16	2	0	3	3	1	224	0	0	249
15/06/19	11	0	0	0	1	0	90	0	0	102
16/06/19	17	0	0	0	3	0	172	0	0	193
<b>Species total</b>	<b>66</b>	<b>5</b>	<b>0</b>	<b>3</b>	<b>13</b>	<b>1</b>	<b>642</b>	<b>0</b>	<b>0</b>	<b>730</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	390	6	0	0	5	2	12	0	1	416
13/07/19	348	5	0	0	6	0	8	0	0	367
14/07/19	245	1	0	0	2	0	7	0	0	255
15/07/19	99	3	0	1	4	0	5	0	2	114
16/07/19	101	5	0	0	1	0	9	0	1	117
<b>Species total</b>	<b>1,183</b>	<b>20</b>	<b>0</b>	<b>1</b>	<b>18</b>	<b>2</b>	<b>41</b>	<b>0</b>	<b>4</b>	<b>1,269</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	28	7	0	1	10	9	29	5	1	90
14/08/19	0	0	0	0	0	0	0	0	0	0
15/08/19	80	4	0	1	22	1	2	8	2	120
16/08/19	2	0	0	0	1	0	2	0	0	5
17/08/19	38	3	0	0	17	6	33	0	8	105
18/08/19	1	1	0	1	3	0	3	0	1	10
<b>Species total</b>	<b>149</b>	<b>15</b>	<b>0</b>	<b>39</b>	<b>53</b>	<b>16</b>	<b>69</b>	<b>13</b>	<b>12</b>	<b>330</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	18	0	0	0	2	0	53	0	0	73
26/09/19	21	1	0	0	5	2	96	0	0	125
27/09/19	3	0	0	1	0	0	89	0	0	93
<b>Species total</b>	<b>42</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>7</b>	<b>2</b>	<b>238</b>	<b>0</b>	<b>0</b>	<b>291</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	8	3	0	0	3	0	0	0	0	14
15/10/19	5	2	0	0	2	0	0	1	0	10
16/10/19	1	0	0	0	0	0	0	0	0	1
17/10/19	0	0	0	0	0	0	0	0	0	0
18/10/19	10	0	0	0	0	0	0	0	0	10
<b>Species total</b>	<b>24</b>	<b>5</b>	<b>0</b>	<b>0</b>	<b>5</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>35</b>

**Annex 2.1.45: Bat Records at Location 3**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/04/19	61	28	0	0	2	0	0	0	0	91
26/04/19	1	1	0	0	3	0	0	0	0	5
27/04/19	0	5	0	0	0	0	0	0	0	5
28/04/19	404	37	0	0	39	0	0	0	0	480
29/04/19	585	64	0	0	48	0	3	0	0	700
30/04/19	485	53	0	0	6	0	0	0	0	544
01/05/19	525	40	0	1	19	0	0	0	0	585
<b>Species total</b>	<b>2,061</b>	<b>228</b>	<b>0</b>	<b>1</b>	<b>117</b>	<b>0</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>2,410</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	3,200	63	0	1	382	65	1	0	0	3,712
11/05/19	3,381	76	0	1	599	48	16	1	0	4,135
12/05/19	3,838	148	0	0	931	38	31	3	0	4,989
13/05/19	3,545	103	0	1	780	53	12	0	0	4,494
14/05/19	1,648	139	0	0	410	30	8	1	0	2,236
<b>Species total</b>	<b>15,612</b>	<b>529</b>	<b>0</b>	<b>3</b>	<b>3,102</b>	<b>234</b>	<b>68</b>	<b>5</b>	<b>0</b>	<b>19,553</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/06/19	0	2	0	0	0	0	2	0	0	4
13/06/19	248	58	0	0	74	1	24	0	0	406

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/06/19	604	82	0	0	159	3	24	0	0	895
15/06/19	276	84	0	0	563	0	31	0	0	662
16/06/19	174	42	0	1	143	0	28	0	0	391
<b>Species total</b>	<b>1,302</b>	<b>268</b>	<b>0</b>	<b>1</b>	<b>639</b>	<b>4</b>	<b>109</b>	<b>0</b>	<b>0</b>	<b>2,358</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	1,432	86	0	2	306	0	6	0	0	1,832
13/07/19	1,440	83	1	2	251	0	12	0	0	1,789
14/07/19	1,880	99	0	0	414	0	11	0	0	2,404
15/07/19	1,269	81	0	0	347	0	1	0	0	1,698
16/07/19	1,667	106	0	1	410	3	4	0	0	2,191
<b>Species total</b>	<b>7,688</b>	<b>455</b>	<b>1</b>	<b>5</b>	<b>1,728</b>	<b>3</b>	<b>34</b>	<b>0</b>	<b>0</b>	<b>9,914</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	630	186	0	109	137	3	2	0	0	1,067
14/08/19	487	95	0	99	107	0	0	0	0	788
15/08/19	257	166	0	92	54	0	3	0	1	573
16/08/19	14	23	0	0	5	0	0	0	0	42
17/08/19	543	250	0	154	92	0	1	0	0	1,040
18/08/19	408	186	0	81	146	0	0	0	1	820
<b>Species total</b>	<b>2,339</b>	<b>904</b>	<b>0</b>	<b>535</b>	<b>541</b>	<b>3</b>	<b>6</b>	<b>0</b>	<b>2</b>	<b>4,330</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	40	41	0	127	17	0	56	0	1	282
26/09/19	101	19	0	131	48	0	40	0	0	339
27/09/19	34	3	0	203	22	0	26	0	0	288
28/09/19	52	10	0	33	9	0	20	0	0	124
29/09/19	106	10	0	176	49	0	19	0	0	360
<b>Species total</b>	<b>333</b>	<b>83</b>	<b>0</b>	<b>670</b>	<b>145</b>	<b>0</b>	<b>161</b>	<b>0</b>	<b>1</b>	<b>1,393</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	66	11	0	0	126	0	0	0	0	203
15/10/19	330	124	0	39	679	1	1	0	0	1,174
16/10/19	19	53	0	12	182	4	0	0	0	270
17/10/19	34	0	0	0	0	0	0	0	0	34
18/10/19	6	80	0	2	18	0	0	0	0	106

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
<b>Species total</b>	<b>455</b>	<b>268</b>	<b>0</b>	<b>53</b>	<b>1,005</b>	<b>5</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1,787</b>

**Annex 21.46: Bat Records at Location 4**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/04/19	662	27	1	1	42	18	0	0	0	751
26/04/19	354	25	0	0	7	0	0	0	1	387
27/04/19	14	8	0	0	3	0	0	0	1	25
28/04/19	272	10	1	0	13	3	3	1	0	303
29/04/19	400	31	0	0	18	8	0	0	1	458
30/04/19	1,093	46	3	0	14	10	3	0	0	1,169
<b>Species total</b>	<b>2,795</b>	<b>147</b>	<b>5</b>	<b>1</b>	<b>96</b>	<b>39</b>	<b>6</b>	<b>1</b>	<b>3</b>	<b>3,093</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	31	4	1	0	6	0	0	1	0	43
11/05/19	52	15	1	0	7	0	1	0	0	76
12/05/19	142	18	0	4	2	0	0	1	0	167
13/05/19	138	512	0	6	1	0	0	0	0	657
14/05/19	1,214	375	0	7	10	0	0	0	0	1,606
15/05/19	828	386	1	12	5	0	0	0	0	1,232
<b>Species total</b>	<b>2,405</b>	<b>1,310</b>	<b>3</b>	<b>29</b>	<b>31</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>0</b>	<b>3,781</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/06/19	4	0	0	0	3	0	0	0	0	7
13/06/19	30	0	0	0	4	0	1	0	0	35
14/06/19	37	2	0	0	2	4	23	0	0	68
15/06/19	28	0	1	0	0	0	2	0	0	31
<b>Species total</b>	<b>99</b>	<b>2</b>	<b>1</b>	<b>0</b>	<b>9</b>	<b>4</b>	<b>26</b>	<b>0</b>	<b>0</b>	<b>141</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	59	9	0	1	15	2	4	0	6	96
13/07/19	56	3	0	1	25	8	5	0	1	99
14/07/19	54	0	0	1	11	4	4	0	2	76
15/07/19	62	4	1	0	8	0	5	0	0	80
16/07/19	68	7	1	2	19	4	18	0	0	119
<b>Species total</b>	<b>299</b>	<b>23</b>	<b>2</b>	<b>5</b>	<b>78</b>	<b>18</b>	<b>36</b>	<b>0</b>	<b>9</b>	<b>470</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	85	2	0	3	23	0	15	9	2	139
14/08/19	53	2	0	3	7	0	2	0	0	67
15/08/19	102	0	0	0	28	3	10	1	2	146
16/08/19	32	0	0	0	4	0	0	0	0	36
17/08/19	56	3	0	0	1	0	5	0	0	65
18/08/19	57	6	0	0	4	0	0	0	0	68
<b>Species total</b>	<b>385</b>	<b>13</b>	<b>0</b>	<b>6</b>	<b>67</b>	<b>3</b>	<b>32</b>	<b>10</b>	<b>4</b>	<b>520</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	19	1	0	0	4	0	21	0	0	45
26/09/19	10	3	0	0	5	4	21	0	0	43
27/09/19	9	0	0	0	3	2	21	0	0	35
<b>Species total</b>	<b>38</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>12</b>	<b>6</b>	<b>63</b>	<b>0</b>	<b>0</b>	<b>123</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	0	0	0	0	0	0	0	0	0	0
15/10/19	11	0	0	0	2	0	8	0	0	21
16/10/19	3	1	0	0	3	0	12	0	0	19
17/10/19	0	0	0	0	0	0	0	0	0	0
18/10/19	2	1	0	0	10	0	0	0	0	13
<b>Species total</b>	<b>16</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>15</b>	<b>0</b>	<b>20</b>	<b>0</b>	<b>0</b>	<b>53</b>

**Annex 2.147: Bat Records at Location 5**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	522	6	0	0	0	0	1	0	0	529
11/05/19	395	2	0	0	3	0	0	0	0	400
12/05/19	281	0	0	0	3	0	0	0	0	284
13/05/19	582	7	0	0	0	1	0	0	0	590
14/05/19	696	4	0	0	0	6	0	0	0	706
15/05/19	985	16	0	0	0	0	0	0	0	1,001
<b>Species total</b>	<b>3,461</b>	<b>35</b>	<b>0</b>	<b>0</b>	<b>6</b>	<b>7</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>3,694</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	1,337	47	0	1	1	0	9	0	50	14,45
13/07/19	234	6	0	0	2	0	2	0	27	271
14/07/19	878	84	0	0	2	0	5	0	5	974
15/07/19	339	6	0	1	1	0	0	0	6	353
16/07/19	272	1	0	0	2	0	0	0	3	278
<b>Species total</b>	<b>3,060</b>	<b>144</b>	<b>0</b>	<b>2</b>	<b>8</b>	<b>0</b>	<b>16</b>	<b>0</b>	<b>91</b>	<b>3,321</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	162	20	0	1	0	0	0	0	0	183
14/08/19	53	2	0	3	7	0	2	0	0	67
15/08/19	102	0	0	0	28	3	10	1	2	146
16/08/19	32	0	0	0	4	0	0	0	0	36
17/08/19	56	3	0	0	1	0	5	0	0	65
18/08/19	57	6	0	0	4	0	0	0	0	67
<b>Species total</b>	<b>462</b>	<b>31</b>	<b>0</b>	<b>4</b>	<b>44</b>	<b>3</b>	<b>17</b>	<b>1</b>	<b>2</b>	<b>564</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	1	0	0	0	0	0	1	0	0	2
26/09/19	31	3	0	14	3	0	2	0	0	53
27/09/19	24	6	0	26	1	1	5	0	0	63
28/09/19	20	2	0	3	0	0	1	0	0	26
29/09/19	92	17	0	36	7	3	6	0	0	161
<b>Species total</b>	<b>168</b>	<b>28</b>	<b>0</b>	<b>79</b>	<b>11</b>	<b>4</b>	<b>15</b>	<b>0</b>	<b>0</b>	<b>305</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	0	0	0	0	0	0	0	0	0	0
15/10/19	25	3	0	2	3	1	0	0	0	34
16/10/19	14	2	0	4	2	2	0	0	0	24
17/10/19	0	0	0	0	0	0	0	0	0	0
18/10/19	8	1	0	0	0	1	0	0	0	10
<b>Species total</b>	<b>47</b>	<b>6</b>	<b>0</b>	<b>6</b>	<b>5</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>68</b>

Annex 2.148: Bat Records at Location 6

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/04/19	0	0	0	0	0	0	0	0	0	0
26/04/19	239	0	1	0	0	0	0	0	0	240
27/04/19	27	2	0	0	0	0	0	0	0	29
<b>Species total</b>	<b>266</b>	<b>2</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>269</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	2,728	124	2	0	4	0	2	0	0	2,860
11/05/19	1,746	64	0	0	20	1	0	0	0	1,831
12/05/19	365	35	0	0	2	0	0	0	0	402
<b>Species total</b>	<b>4,839</b>	<b>223</b>	<b>2</b>	<b>0</b>	<b>26</b>	<b>1</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>5,093</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/06/19	10	0	0	0	2	0	0	0	0	10
13/06/19	2,588	1	9	0	4	0	7	0	0	2,607
14/06/19	1,791	2	4	1	0	2	22	0	0	1,826
15/06/19	1,752	1	30	0	2	0	3	0	1	1,787
16/06/19	1,613	9	8	0	0	0	14	0	0	1,646
<b>Species total</b>	<b>7,754</b>	<b>13</b>	<b>51</b>	<b>1</b>	<b>8</b>	<b>2</b>	<b>46</b>	<b>0</b>	<b>1</b>	<b>7,876</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	2,074	12	8	0	1	3	25	0	1	2,124
13/07/19	581	1	0	0	2	0	7	0	1	592
14/07/19	1,061	4	0	0	7	1	10	0	0	1,083
15/07/19	866	1	1	0	5	0	18	0	0	891
16/07/19	1	0	0	0	0	0	0	0	0	1
<b>Species total</b>	<b>4,583</b>	<b>18</b>	<b>9</b>	<b>0</b>	<b>15</b>	<b>4</b>	<b>60</b>	<b>0</b>	<b>2</b>	<b>4691</b>



Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	0	0	0	0	0	0	0	0	0	0
14/08/19	1,839	1	0	0	3	0	11	1	0	1,855
15/08/19	1,560	0	0	0	10	2	69	1	1	1,643
16/08/19	51	0	0	0	0	0	0	0	0	51
17/08/19	2,173	1	0	0	1	0	12	0	0	2,187
18/08/19	2,149	3	0	0	5	0	4	0	0	2,161
<b>Species total</b>	<b>7,772</b>	<b>5</b>	<b>0</b>	<b>0</b>	<b>19</b>	<b>2</b>	<b>96</b>	<b>2</b>	<b>1</b>	<b>7,897</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
24/09/19	0	0	0	0	0	0	0	0	0	0
25/09/19	1,429	15	0	2	1	0	7	0	0	1,454
26/09/19	1,411	3	0	0	3	0	2	0	0	1,419
27/09/19	11	3	0	0	0	0	4	0	0	18
28/09/19	21	0	0	0	2	0	6	0	0	29
<b>Species total</b>	<b>2,872</b>	<b>21</b>	<b>0</b>	<b>2</b>	<b>6</b>	<b>0</b>	<b>19</b>	<b>0</b>	<b>0</b>	<b>2,920</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	54	7	0	0	0	0	0	0	0	61
15/10/19	85	3	0	0	0	0	1	0	0	89
16/10/19	26	0	0	0	1	0	1	0	0	28
17/10/19	0	0	0	0	0	0	0	0	0	0
18/10/19	126	19	0	0	1	0	0	0	0	146
19/10/19	55	0	0	0	0	0	0	0	0	55
<b>Species total</b>	<b>346</b>	<b>29</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>379</b>

**Annex 2.1.49: Bat Records at Location 7**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/04/19	1,966	38	0	0	38	1	0	0	0	2,043
26/04/19	559	18	0	0	4	0	0	0	0	581
27/04/19	201	98	0	0	1	0	0	0	0	300
28/04/19	1,815	81	0	0	34	2	0	0	0	1,932
29/04/19	1,577	72	0	0	14	0	0	0	0	1,663
30/04/19	1,903	30	0	0	7	0	0	0	0	1,940
<b>Species total</b>	<b>8,021</b>	<b>337</b>	<b>0</b>	<b>0</b>	<b>98</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>8,459</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	3,674	79	0	0	7	2	0	0	0	3,762
11/05/19	3,897	75	1	0	1	0	2	0	0	3,976
12/05/19	3,596	80	0	0	2	2	0	0	0	3,680
13/05/19	1,403	56	0	0	1	0	0	0	0	1,460
<b>Species total</b>	<b>12,570</b>	<b>290</b>	<b>1</b>	<b>0</b>	<b>11</b>	<b>4</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>12,878</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/06/19	615	11	0	0	0	0	8	0	0	634
13/06/19	2,037	58	0	0	10	0	0	0	0	2,105
14/06/19	2,883	118	0	0	10	1	3	0	0	3,015
15/06/19	1,952	61	0	0	40	5	0	0	0	2,058
16/06/19	396	2	0	0	1	1	9	0	0	409
<b>Species total</b>	<b>7,883</b>	<b>250</b>	<b>0</b>	<b>0</b>	<b>61</b>	<b>7</b>	<b>20</b>	<b>0</b>	<b>0</b>	<b>8,221</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	855	3	0	1	1	3	0	0	15	878
13/07/19	1,075	7	0	2	5	6	4	0	34	1,133
14/07/19	1,900	12	0	0	2	6	0	0	11	1,931
15/07/19	1,274	16	0	2	4	4	4	0	4	1,308
<b>Species total</b>	<b>5,104</b>	<b>38</b>	<b>0</b>	<b>5</b>	<b>12</b>	<b>19</b>	<b>8</b>	<b>0</b>	<b>64</b>	<b>5,250</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	645	10	0	9	9	1	0	0	0	674
14/08/19	28	0	0	1	0	0	0	0	0	29
15/08/19	443	9	0	5	12	4	0	0	0	479
16/08/19	35	1	0	0	0	0	0	0	0	36
17/08/19	559	2	0	33	95	9	0	0	0	702
18/08/19	444	5	0	24	0	2	0	0	25	501
<b>Species total</b>	<b>2,154</b>	<b>27</b>	<b>0</b>	<b>72</b>	<b>116</b>	<b>16</b>	<b>0</b>	<b>0</b>	<b>25</b>	<b>2,421</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	48	0	0	47	0	0	5	0	0	100
26/09/19	98	2	0	28	1	5	2	0	0	136
27/09/19	2	0	0	9	0	0	3	0	0	14
<b>Species total</b>	<b>148</b>	<b>2</b>	<b>0</b>	<b>84</b>	<b>1</b>	<b>5</b>	<b>10</b>	<b>0</b>	<b>0</b>	<b>250</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
15/10/19	21	2	0	0	0	0	0	0	0	23
16/10/19	23	1	0	0	1	0	0	0	0	25
17/10/19	0	0	0	0	0	0	0	0	0	0
18/10/19	152	4	0	2	2	0	0	0	0	160
19/10/19	29	19	0	0	0	0	1	0	0	49
20/10/19	211	16	0	0	4	0	0	0	0	231
<b>Species total</b>	<b>436</b>	<b>42</b>	<b>0</b>	<b>2</b>	<b>7</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>488</b>

**Annex 2.1.50: Bat Records at Location 8**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
24/04/19	0	0	0	0	0	0	0	0	0	0
25/04/19	173	0	0	0	0	0	0	0	0	173
26/04/19	23	0	0	0	0	0	0	0	0	25
27/04/19	13	0	0	0	0	0	0	0	0	13
28/04/19	280	0	1	0	0	0	3	0	0	284
29/04/19	367	0	0	4	0	0	0	0	0	371
30/04/19	267	0	0	0	0	0	10	0	0	277
01/05/19	603	2	0	3	0	0	7	0	0	615
<b>Species total</b>	<b>1,728</b>	<b>2</b>	<b>1</b>	<b>7</b>	<b>0</b>	<b>0</b>	<b>20</b>	<b>0</b>	<b>0</b>	<b>1,758</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/05/19	71	0	0	0	0	0	0	0	0	71
13/05/19	1,118	1	0	0	0	0	0	0	0	1,119
14/05/19	787	2	0	0	0	0	0	0	0	789
15/05/19	142	0	0	0	0	0	0	0	0	142
<b>Species total</b>	<b>2,118</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>2,121</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
11/07/19	0	0	0	0	0	0	0	0	0	0
12/07/19	5	0	0	0	0	0	7	0	0	12
13/07/19	7	0	0	0	0	0	39	0	0	46
14/07/19	14	0	0	0	0	0	101	0	0	114
15/07/19	8	0	0	0	0	0	0	0	0	8
16/07/19	3	0	0	0	0	0	20	0	0	23
<b>Species total</b>	<b>37</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>164</b>	<b>0</b>	<b>0</b>	<b>203</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/08/19	1	0	0	0	0	0	0	0	0	1
15/08/19	2	0	0	0	0	0	0	0	0	2
16/08/19	0	0	0	0	0	0	0	0	0	0
17/08/19	7	0	0	0	0	0	0	0	0	7
18/08/19	4	0	0	0	0	0	0	0	0	4
<b>Species total</b>	<b>14</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>14</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	227	4	0	31	0	0	113	0	0	375
26/09/19	138	6	0	4	0	0	175	2	0	325
27/09/19	9	0	0	0	0	0	39	0	0	48
28/09/19	125	1	9	6	0	0	46	0	0	187
29/09/19	180	8	0	2	0	0	642	6	1	840
<b>Species total</b>	<b>679</b>	<b>19</b>	<b>9</b>	<b>43</b>	<b>1</b>	<b>0</b>	<b>1,015</b>	<b>8</b>	<b>1</b>	<b>1,775</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	21	3	0	0	0	0	11	0	0	35
15/10/19	332	16	22	2	1	0	14	1	0	388
16/10/19	38	0	0	0	0	0	2	0	0	40
17/10/19	1	0	0	0	0	0	0	0	0	1
18/10/19	217	3	0	0	7	0	5	0	0	232
19/10/19	184	2	0	1	0	0	6	0	0	193
<b>Species total</b>	<b>793</b>	<b>24</b>	<b>22</b>	<b>3</b>	<b>8</b>	<b>0</b>	<b>38</b>	<b>1</b>	<b>0</b>	<b>889</b>

**Annex 2.1.51: Bat Records at Location 9**

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
24/04/19	0	0	0	0	0	0	0	0	0	0	0
25/04/19	0	4	0	0	0	0	0	0	0	0	4
26/04/19	0	0	0	0	0	0	0	0	0	0	0
27/04/19	0	0	0	0	0	0	0	0	0	0	0
28/04/19	0	1	0	0	0	0	0	0	0	0	1
29/04/19	0	4	0	0	0	0	0	0	0	0	4
30/04/19	0	12	0	0	0	0	0	0	0	0	12
01/05/19	0	1	0	0	0	0	0	0	0	0	1

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
<b>Species total</b>	<b>0</b>	<b>22</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>22</b>

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/05/19	0	197	0	0	0	0	0	0	0	0	197
11/05/19	0	214	0	0	0	0	0	0	0	0	214
12/05/19	0	305	1	0	0	0	0	0	0	0	306
13/05/19	0	352	0	0	0	0	0	0	0	0	352
14/05/19	0	453	0	0	0	1	0	0	0	0	454
15/05/19	0	565	0	0	0	1	0	0	0	0	566
<b>Species total</b>	<b>0</b>	<b>2,086</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>2,089</b>

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/06/19	1	428	7	8	0	2	0	0	0	0	446
14/06/19	0	950	6	1	0	0	0	3	0	0	960
15/06/19	0	653	6	0	0	0	0	0	0	0	659
16/06/19	0	763	0	0	0	0	0	0	0	0	763
<b>Species total</b>	<b>1</b>	<b>2,794</b>	<b>19</b>	<b>9</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>2,828</b>

Survey Date	Bb	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	0	91	2	0	0	0	0	16	0	0	109
13/07/19	0	31	0	0	0	0	0	2	0	0	33
14/07/19	0	46	0	0	0	0	0	0	0	0	46
15/07/19	0	60	0	0	0	0	0	0	0	0	60
16/07/19	0	10	0	0	0	0	0	1	0	0	11
<b>Species total</b>	<b>0</b>	<b>238</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>19</b>	<b>0</b>	<b>0</b>	<b>259</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	18	1	0	0	0	0	0	0	0	19

14/08/19	3	0	0	0	0	0	0	0	0	3
15/08/19	10	0	0	0	2	0	0	0	0	12
16/08/19	7	0	0	0	0	0	0	0	0	7
17/08/19	26	0	0	0	1	0	0	0	0	27
18/08/19	40	0	0	0	0	0	0	0	0	40
<b>Species total</b>	<b>104</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>108</b>
<b>Survey Date</b>	<b>Pp</b>	<b>Ppy</b>	<b>Pn</b>	<b>Psp</b>	<b>Msp</b>	<b>Plsp</b>	<b>Nn</b>	<b>NI</b>	<b>Es</b>	<b>Total</b>
25/09/19	52	0	0	1	0	0	0	0	0	53
26/09/19	45	0	0	5	0	0	0	0	0	50
27/09/19	19	0	0	0	0	0	0	0	0	19
28/09/19	6	0	0	0	0	0	0	0	0	6
29/09/19	4	0	0	0	0	0	0	0	0	4
<b>Species total</b>	<b>126</b>	<b>0</b>	<b>0</b>	<b>6</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>132</b>

<b>Survey Date</b>	<b>Pp</b>	<b>Ppy</b>	<b>Pn</b>	<b>Psp</b>	<b>Msp</b>	<b>Plsp</b>	<b>Nn</b>	<b>NI</b>	<b>Es</b>	<b>Total</b>
15/10/19	2	0	0	0	0	0	0	0	0	2
16/10/19	0	0	0	0	1	0	0	0	0	1
<b>Species total</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>3</b>

#### Annex 2.1.52: Bat Records at Location 10

<b>Survey Date</b>	<b>Pp</b>	<b>Ppy</b>	<b>Pn</b>	<b>Psp</b>	<b>Msp</b>	<b>Plsp</b>	<b>Nn</b>	<b>NI</b>	<b>Es</b>	<b>Total</b>
10/05/19	33	0	0	1	0	0	0	0	0	34
11/05/19	118	2	0	2	1	0	1	0	0	124
12/05/19	133	23	0	0	0	0	0	0	0	156
13/05/19	670	1	0	0	2	0	0	0	0	673
14/05/19	736	112	0	1	0	0	0	0	0	849
15/05/19	603	207	0	0	0	0	0	0	0	810
<b>Species total</b>	<b>2,293</b>	<b>345</b>	<b>0</b>	<b>4</b>	<b>3</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>2,646</b>

<b>Survey Date</b>	<b>Pp</b>	<b>Ppy</b>	<b>Pn</b>	<b>Psp</b>	<b>Msp</b>	<b>Plsp</b>	<b>Nn</b>	<b>NI</b>	<b>Es</b>	<b>Total</b>
12/07/19	289	20	0	0	0	0	6	0	0	315
13/07/19	324	9	0	1	2	0	5	0	0	341
14/07/19	1,243	68	0	0	3	1	3	0	0	1,318
15/07/19	369	39	0	0	1	0	9	0	0	418
16/07/19	431	0	0	0	0	0	0	0	0	431
<b>Species total</b>	<b>2,656</b>	<b>136</b>	<b>0</b>	<b>1</b>	<b>6</b>	<b>1</b>	<b>23</b>	<b>0</b>	<b>0</b>	<b>2,823</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	459	93	0	1	6	2	4	2	2	569
14/08/19	305	1	0	0	1	0	2	0	0	309
15/08/19	463	31	0	11	12	5	6	0	1	529
<b>Species total</b>	<b>1,227</b>	<b>125</b>	<b>0</b>	<b>12</b>	<b>19</b>	<b>7</b>	<b>12</b>	<b>2</b>	<b>3</b>	<b>1,407</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	248	17	0	2	2	0	57	2	0	328
26/09/19	109	13	1	0	2	1	12	0	0	138
27/09/19	1	0	0	0	0	0	2	0	0	3
28/09/19	25	8	0	0	2	0	37	0	0	72
29/09/19	108	36	0	0	3	1	9	0	0	157
<b>Species total</b>	<b>491</b>	<b>74</b>	<b>1</b>	<b>2</b>	<b>9</b>	<b>2</b>	<b>117</b>	<b>2</b>	<b>0</b>	<b>698</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	30	1	0	0	1	0	4	1	0	37
15/10/19	6	0	0	0	2	0	3	0	0	11
16/10/19	6	1	0	0	1	0	4	0	0	12
17/10/19	10	0	0	0	0	0	0	0	0	10
18/10/19	26	2	0	0	1	0	0	0	0	29
<b>Species total</b>	<b>78</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>5</b>	<b>0</b>	<b>11</b>	<b>1</b>	<b>0</b>	<b>99</b>

**Annex 2.1.53: Bat Records at Location 11**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
24/04/19	0	0	0	0	0	0	0	0	0	0
25/04/19	585	3	2	1	1	0	0	0	0	592
26/04/19	94	0	0	0	0	0	0	0	0	94
27/04/19	1	0	0	0	0	0	0	0	0	1
28/04/19	357	2	0	1	0	1	1	0	0	362
29/04/19	166	1	0	3	0	0	0	0	0	170
30/04/19	626	1	0	0	3	1	1	0	0	632
01/05/19	182	0	0	0	0	0	4	0	0	186
<b>Species total</b>	<b>2,011</b>	<b>7</b>	<b>2</b>	<b>5</b>	<b>4</b>	<b>2</b>	<b>6</b>	<b>0</b>	<b>0</b>	<b>2,037</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
11/05/19	5	0	0	0	0	0	0	0	0	5
12/05/19	9	0	0	0	0	0	0	0	0	9
13/05/19	21	0	0	0	0	0	0	0	0	21
14/05/19	2	0	0	0	0	0	0	0	0	2
15/05/19	23	0	0	0	0	0	0	0	0	23
<b>Species total</b>	<b>60</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>60</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/06/19	196	0	0	0	0	0	6	0	0	202
14/06/19	97	0	0	0	0	0	5	0	0	102
15/06/19	155	0	0	0	0	0	3	0	0	158
16/06/19	480	0	0	0	0	0	3	0	0	483
<b>Species total</b>	<b>928</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>17</b>	<b>0</b>	<b>0</b>	<b>945</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
12/07/19	865	0	1	3	26	1	6	0	0	902
13/07/19	1,376	0	1	2	19	0	3	0	0	1,401
14/07/19	1,207	1	2	3	46	0	5	0	0	1,264
15/07/19	482	1	1	0	23	1	9	0	0	517
16/07/19	431	1	2	3	7	0	10	0	0	454
<b>Species total</b>	<b>4,361</b>	<b>3</b>	<b>7</b>	<b>11</b>	<b>121</b>	<b>2</b>	<b>33</b>	<b>0</b>	<b>0</b>	<b>4,538</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
13/08/19	57	0	0	0	1	0	5	0	0	63
14/08/19	663	0	0	25	1	0	5	0	1	695
15/08/19	57	1	0	0	6	1	6	0	0	71
16/08/19	210	0	0	1	5	0	7	0	0	223
17/08/19	223	0	0	11	0	0	4	0	0	238
<b>Species total</b>	<b>1,210</b>	<b>1</b>	<b>0</b>	<b>37</b>	<b>13</b>	<b>1</b>	<b>27</b>	<b>0</b>	<b>1</b>	<b>1,290</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/19	2,003	2	0	165	35	1	322	0	0	2,528
26/09/19	875	1	0	80	22	1	212	0	0	1,191
27/09/19	17	1	0	1	1	1	5	0	0	26



Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
<b>Species total</b>	<b>2,895</b>	<b>4</b>	<b>0</b>	<b>246</b>	<b>58</b>	<b>3</b>	<b>539</b>	<b>0</b>	<b>0</b>	<b>3,745</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
14/10/19	252	5	0	0	17	0	0	0	0	274
15/10/19	183	0	0	0	19	0	16	0	0	218
16/10/19	26	1	0	0	5	0	4	0	0	36
17/10/19	0	0	0	0	0	0	0	0	0	0
18/10/19	986	3	0	2	18	0	0	0	0	1,009
19/10/19	9	0	0	0	0	0	0	0	0	9
<b>Species total</b>	<b>1,456</b>	<b>9</b>	<b>0</b>	<b>2</b>	<b>59</b>	<b>0</b>	<b>20</b>	<b>0</b>	<b>0</b>	<b>1,546</b>

**Annex 2.1.34: Bat Records at Location 12**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
28/05/2022	51	1	0	0	1	0	8	0	0	61
29/05/2022	3	0	0	0	0	0	5	0	0	8
30/05/2022	4	0	0	0	1	0	6	0	0	11
31/05/2022	1	0	0	0	0	0	2	0	0	3
01/06/2022	7	0	0	0	0	0	4	0	0	11
02/06/2022	22	2	0	0	0	0		0	0	24
<b>Species total</b>	<b>88</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>25</b>	<b>0</b>	<b>0</b>	<b>118</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
27/07/2022	18	0	0	0	1	0	1	0	1	21
28/07/2022	23	2	0	0	3	0	1	1	1	31
29/07/2022	85	2	0	0	8	0	10	1	0	106
30/07/2022	44	1	0	0	5	0	11	0	2	63
31/07/2022	57	2	0	0	0	0	50	1	5	115
<b>Species total</b>	<b>227</b>	<b>7</b>	<b>0</b>	<b>0</b>	<b>17</b>	<b>0</b>	<b>73</b>	<b>3</b>	<b>9</b>	<b>336</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
18/08/2022	3,064	1,018	0	0	6	0	0	0	4	4,092
19/08/2022	1,661	491	0	0	13	0	0	0	2	2,167
20/08/2022	2,613	632	0	0	4	0	0	0	6	3,255
21/08/2022	1,829	445	0	0	9	1	2	0	4	2,290
22/08/2022	2,736	503	0	0	6	0	1	0	8	3,254

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
23/08/2022	1,111	153	0	0	7	0	0	0	7	1,278
<b>Species total</b>	<b>13,014</b>	<b>3,242</b>	<b>0</b>	<b>0</b>	<b>45</b>	<b>1</b>	<b>3</b>	<b>0</b>	<b>31</b>	<b>16,336</b>

**Annex 2.1.35: Bat Records at Location 13**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
27/05/2022	31	0	0	0	0	0	1	2	0	34
28/05/2022	112	2	0	0	0	0	0	0	0	114
29/05/2020	3	0	0	0	0	0	1	0	0	4
30/05/2022	2	0	0	0	0	0	0	0	0	2
01/06/2022	87	0	0	0	0	0	2	1	0	90
02/06/2022	6	0	0	0	0	0	1	0	0	7
<b>Species total</b>	<b>241</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>5</b>	<b>3</b>	<b>0</b>	<b>251</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
22/06/2022	715	36	0	0	0	0	8	1	0	760
23/06/2022	892	16	0	0	0	0	14	0	0	922
24/06/2022	798	6	0	0	0	0	5	0	1	810
25/06/2022	1,100	12	0	0	4	0	2	0	0	1,118
26/06/2022	1,144	16	0	0	5	0	2	0	0	1,167
27/06/2022	1,095	18	0	0	1	0	6	0	0	1,120
<b>Species total</b>	<b>5,744</b>	<b>104</b>	<b>0</b>	<b>0</b>	<b>10</b>	<b>0</b>	<b>37</b>	<b>0</b>	<b>1</b>	<b>5,897</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
23/08/2022	1,117	5	0	8	0	0	0	0	0	1,130
24/08/2022	954	24	0	0	3	0	20	2	6	1,009
25/08/2022	647	10	0	0	1	0	4	0	2	664
26/08/2022	787	6	0	0	1	0	10	1	1	806
27/08/2022	314	12	0	0	1	0	6	0	2	335
<b>Species total</b>	<b>3,819</b>	<b>57</b>	<b>0</b>	<b>8</b>	<b>6</b>	<b>0</b>	<b>40</b>	<b>3</b>	<b>11</b>	<b>3,944</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
21/09/2022	62	9	0	0	0	0	13	4	0	88
22/09/2022	132	1	0	0	0	0	1	0	0	134
23/09/2022	349	6	0	0	0	0	13	0	0	368
24/09/2022	354	24	0	0	2	1	11	0	0	392

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
25/09/2022	771	45	1	0	1	1	5	0	1	825
<b>Species total</b>	<b>1,668</b>	<b>85</b>	<b>1</b>	<b>0</b>	<b>3</b>	<b>2</b>	<b>43</b>	<b>4</b>	<b>1</b>	<b>1,807</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
18/10/2022	74	11	0	0	0	0	0	0	0	85
19/20/2022	36	2	0	4	0	0	1	0	0	43
20/10/2022	47	0	0	0	0	0	2	0	0	49
21/10/2022	4	0	0	0	0	0	0	0	0	4
22/10/2022	12	1	0	0	0	0	0	0	0	13
<b>Species total</b>	<b>173</b>	<b>14</b>	<b>0</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>194</b>

Annex 2.1.34: Bat Records at Location 14

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
21/07/2022	30	1	0	0	0	0	2	0	0	33
22/07/2022	31	0	0	0	0	0	7	0	0	38
23/07/2022	3	1	0	0	0	0	9	0	0	13
24/07/2022	6	0	0	0	0	0	1	0	0	7
25/07/2022	15	0	0	0	0	0	0	0	0	15
26/07/2022	3	1	0	0	0	0	11	0	0	15
27/07/2022	1	1	0	0	0	0	2	0	0	4
28/07/2022	8	0	0	0	0	0	1	0	0	9
29/07/2022	58	0	0	0	0	0	8	0	0	66
30/07/2022	6	0	0	0	0	0	1	0	0	7
<b>Species total</b>	<b>161</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>42</b>	<b>0</b>	<b>0</b>	<b>207</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
23/08/2022	1,834	1	0	0	0	0	2	0	0	1,837
24/08/2022	2,679	204	0	39	0	0	27	0	0	2,949
25/08/2022	2,548	31	1	0	0	0	9	0	0	2,589
26/08/2022	1,215	18	0	0	4	0	8	0	0	1,245
27/08/2022	1,530	21	0	0	0	0	6	0	0	1,558
<b>Species total</b>	<b>9,806</b>	<b>275</b>	<b>1</b>	<b>0</b>	<b>4</b>	<b>0</b>	<b>52</b>	<b>0</b>	<b>0</b>	<b>10,178</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
08/09/2022	47	1	0	0	0	0	5	1	1	55
09/09/2022	5	1	0	0	0	0	0	0	0	6

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/09/2022	675	102	0	0	0	0	1	0	0	778
11/09/2022	103	37	0	0	0	0	0	0	0	140
12/09/2022	832	166	0	0	0	0	0	0	0	998
14/09/2022	1283	187	1	0	0	0	0	0	0	1,478
<b>Species total</b>	<b>2,945</b>	<b>494</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>6</b>	<b>1</b>	<b>1</b>	<b>3,455</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
18/10/2022	188	5	0	0	0	0	0	0	0	193
19/10/2022	279	6	0	0	0	0	0	0	0	285
20/10/2022	85	0	0	0	0	0	0	0	0	85
21/10/2022	14	0	0	0	0	0	0	0	0	14
22/10/2022	201	0	0	0	10	0	0	0	0	211
23/10/2022	0	1	0	0	0	0	0	0	0	1
<b>Species total</b>	<b>767</b>	<b>12</b>	<b>0</b>	<b>0</b>	<b>10</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>789</b>

**Annex 2.1.34: Bat Records at Location 15**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
18/05/2022	701	272	0	0	0	1	0	0	22	996
19/05/2022	899	347	0	0	1	0	0	0	1	1,248
20/05/2022	677	58	0	0	1	0	0	0	0	736
21/05/2022	308	192	0	0	0	0	0	0	0	500
22/05/2022	609	278	0	0	0	0	0	0	29	916
23/05/2022	1,358	307	0	0	1	0	0	0	1	1,667
24/05/2022	470	249	0	0	0	0	0	0	0	719
<b>Species total</b>	<b>5,022</b>	<b>1,703</b>	<b>0</b>	<b>0</b>	<b>3</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>53</b>	<b>6,782</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/06/2022	1,147	638	0	0	3	0	0	0	0	1,788
11/06/2022	1,386	1,541	1	0	1	0	0	1	7	2,937
12/06/2022	1,454	1,141	4	0	1	0	0	0	1	2,601
13/06/2022	1,040	460	2	0	1	0	0	0	2	1,505
14/06/2022	1,051	194	2	0	2	1	0	0	0	1,250
15/06/2022	1,554	255	3	0	3	0	0	0	1	1,816
16/06/2022	1,502	303	0	0	1	0	1	8	1	1,816
19/06/2022	1,299	2,024	0	0	9	0	0	0	0	3,322
<b>Species total</b>	<b>10,433</b>	<b>6,556</b>	<b>12</b>	<b>0</b>	<b>21</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>12</b>	<b>17,045</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
07/07/2022	2,624	218	0	0	6	1	0	0	3	2,852
08/07/2022	1,472	485	0	0	6	0	6	0	5	1,974
09/07/2022	2,271	610	0	0	8	0	1	3	3	2,896
10/07/2022	176	8	0	0	0	0	0	0	0	184
<b>Species total</b>	<b>6,543</b>	<b>1,321</b>	<b>0</b>	<b>0</b>	<b>20</b>	<b>1</b>	<b>7</b>	<b>3</b>	<b>11</b>	<b>7,906</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
07/08/2022	1,821	1,416	0	0	13	0	12	0	3	3,265
08/08/2022	895	1,980	0	0	5	0	0	2	8	2,890
09/08/2022	1,173	1,716	0	0	7	0	2	2	2	2,902
10/08/2022	2,146	1,654	0	0	5	1	2	0	2	3,810
11/08/2022	500	225	0	0	3	0	3	0	0	731
<b>Species total</b>	<b>6,535</b>	<b>6,991</b>	<b>0</b>	<b>0</b>	<b>33</b>	<b>1</b>	<b>19</b>	<b>4</b>	<b>15</b>	<b>13,598</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
21/09/2022	1,631	1,214	0	0	24	13	3	0	0	2,885
22/09/2022	2,286	2,014	0	0	16	0	7	0	1	4,324
23/09/2022	240	556	0	0	22	5	4	0	1	828
24/09/2022	75	168	0	0	14	15	0	0	0	272
25/09/2022	23	903	0	0	7	1	0	0	0	934
<b>Species total</b>	<b>4,255</b>	<b>4,855</b>	<b>0</b>	<b>0</b>	<b>83</b>	<b>34</b>	<b>14</b>	<b>0</b>	<b>2</b>	<b>9,243</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
20/10/2022	1,304	396	0	0	6	0	0	0	0	1,706
21/10/2022	894	88	0	0	3	0	0	0	0	985
22/10/2022	748	1,636	0	0	15	0	1	0	0	2,400
23/10/2022	64	28	0	0	1	0	0	0	0	93
27/10/2022	6	2	0	0	0	0	0	0	0	8
<b>Species total</b>	<b>3,016</b>	<b>2,150</b>	<b>0</b>	<b>0</b>	<b>25</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>5,192</b>

**Annex 2.1.34: Bat Records at Location 16**

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
19/05/2022	18	0	0	0	0	0	0	0	0	18
21/05/2022	3	0	0	0	0	0	0	0	0	3
22/05/2022	1	0	0	0	0	0	0	0	0	1
<b>Species total</b>	<b>22</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>22</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/06/2022	876	32	0	0	0	0	1	0	1	909
11/06/2022	604	64	0	0	1	0	1	0	0	670
12/06/2022	632	24	0	0	0	0	0	0	0	656
13/06/2022	370	33	0	0	3	0	0	1	0	407
14/06/2022	348	54	0	0	0	0	0	0	0	402
15/06/2022	468	36	0	0	1	0	0	0	0	505
<b>Species total</b>	<b>3,297</b>	<b>243</b>	<b>0</b>	<b>0</b>	<b>5</b>	<b>0</b>	<b>2</b>	<b>1</b>	<b>1</b>	<b>3,550</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
10/07/2022	715	174	0	0	0	0	4	0	1	894
11/07/2022	542	213	0	0	0	0	0	0	4	759
12/07/2022	388	211	0	0	2	0	3	1	2	607
13/07/2022	514	205	0	0	0	0	0	0	2	721
14/07/2022	521	269	0	0	0	0	1	1	2	794
15/07/2022	512	353	0	0	0	0	4	3	4	876
<b>Species total</b>	<b>3,192</b>	<b>1,425</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>12</b>	<b>5</b>	<b>15</b>	<b>4,651</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
18/08/2022	351	95	0	0	0	0	0	0	0	446
19/08/2022	283	105	0	0	0	0	0	0	0	388
20/08/2022	267	62	0	0	1	0	0	0	0	330
21/08/2022	198	60	0	0	0	0	1	0	0	259
22/08/2022	170	42	0	4	1	0	0	0	0	217
23/08/2022	172	11	0	0	0	0	0	0	1	184
<b>Species total</b>	<b>1,441</b>	<b>375</b>	<b>0</b>	<b>4</b>	<b>2</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>1,824</b>

Survey Date	Pp	Ppy	Pn	Psp	Msp	Plsp	Nn	NI	Es	Total
18/10/2022	677	156	0	0	8	0	0	0	0	841
19/10/2022	645	252	0	0	0	0	0	0	0	897
20/10/2022	366	134	0	0	2	0	0	0	0	502
21/10/2022	1,525	168	0	0	2	0	0	0	1	1,696
22/10/2022	794	633	0	0	1	0	0	0	0	1,428
23/10/2022	314	43	0	0	0	0	0	0	0	5,721
<b>Species total</b>	<b>4,321</b>	<b>1,386</b>	<b>0</b>	<b>0</b>	<b>13</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>5,721</b>

## A2.2 Preliminary Bat Roost Assessment Trees

Tree ref.	Species	Tree Height (m)	Tree condition	Features / Description	Roost potential	Further action required	Hibernation potential?
1	Ash	10	Semi-mature	Knot hole with staining	Moderate	Climbing or emergence/ re-entry surveys	Yes
2	Aspen	15	Mature	Mature tree by river with lots of ivy	Moderate	Climbing or emergence/ re-entry surveys	Yes
3	Poplar	12	Semi-mature	Two woodpecker holes, next to two trees with dense ivy	High	Climbing or emergence/ re-entry surveys	Yes
4	Lime	10	Semi-mature	????	Moderate	Emergence/ re-entry surveys	Yes
5	Poplar	10	Semi-mature	Rot hole with staining	Moderate	Climbing or emergence/ re-entry surveys	Yes
6	Ash	14	Semi-mature	Woodpecker hole with various splits and cavities	High	Climbing or emergence/ re-entry surveys	Yes
7	Ash	12	Semi-mature	Various rot holes	High	Climbing or emergence/ re-entry surveys	Yes
8	Poplar	15	Mature	Multiple rot holes with staining and a woodpecker hole	High	Climbing or emergence/ re-entry surveys	Yes
9	Willow	6	Semi-mature	????	Moderate	Emergence/ re-entry surveys	Yes
10	Poplar	14	Mature	????	Moderate	Emergence/ re-entry surveys	Yes
11	Lime	10	Semi-mature	Various rot holes	Moderate	Climbing or emergence/ re-entry surveys	Yes
12	Poplar	12	Semi-mature	Multiple trunk cavities	Moderate	Climbing or emergence/ re-entry surveys	Yes
13	Elder	10	Mature	Group of three elder with lots of flaking bark all over	Moderate	Climbing or emergence/ re-entry surveys	Yes

Tree ref.	Species	Tree Height (m)	Tree condition	Features / Description	Roost potential	Further action required	Hibernation potential?
14	Oak	10	Semi-mature	Two rot cavities from cut branches	Moderate	Climbing or emergence/ re-entry surveys	Yes
15	Oak	12	Mature	????	Moderate	Emergence/ re-entry surveys	Yes
16	Oak	10	Mature	Multiple features on all sides	High	Climbing or emergence/ re-entry surveys	Yes
17	Oak	10	Mature	Group of 8 trees with ivy	Low	Supervised soft-fell	No
18	Oak	10	Semi-mature	????	Moderate	Emergence/ re-entry surveys	Yes
19	Ash	10	Mature	Tree on edge of car park with rot hole	Moderate	Climbing or emergence/ re-entry surveys	Yes
20	Ash	10	Semi-mature	????	Moderate	Emergence/ re-entry surveys	Yes
21	Ash	8	Immature	????	Moderate	Emergence/ re-entry surveys	Yes
23.1	Ash	14	Mature	Dense ivy all over	Low	Supervised soft-fell	No
23.2	Willow	10	Mature	Dead branch overhanging fence with multiple holes	Moderate	Climbing or emergence/ re-entry surveys	Yes
24	Willow	10	Mature	Rot hole on south side of northern branch by street lamp	Moderate	Climbing or emergence/ re-entry surveys	Yes
25	Ash	12	Semi-mature	Tree by fence with multiple holes	Moderate	Climbing or emergence/ re-entry surveys	Yes
26	Elder	10	Mature	Next to fence, rot hole with staining	High	Climbing or emergence/ re-entry surveys	Yes
27	Sycamore	14	Semi-mature	Group of ten trees with dense ivy	Low	Supervised soft-fell	No
28	Unknown	6	Dead	Standing dead tree with multiple holes	High	Climbing or emergence/ re-entry surveys	Yes
29	Sycamore	8	Immature	Just inside fence line, two rot holes	Moderate	Climbing or emergence/ re-entry surveys	Yes



Tree ref.	Species	Tree Height (m)	Tree condition	Features / Description	Roost potential	Further action required	Hibernation potential?
30	Cherry	10	Mature	South side of path on top of bank	Low	Supervised soft-fell	No
31	Mixed	10	Semi-mature	Large group of mixed species trees with dense ivy	Low	Supervised soft-fell	No
32	Unknown	8	Dead	Multiple woodpecker holes	High	Climbing or emergence/ re-entry surveys	Yes

## Annex 3

### Evaluation

### A3.1 Evaluation

#### Breeding Bird Surveys

**Annex 3.1.1: Species of Conservation Interest, Number of Territories, National, Regional and County Status and Geographical Importance of Survey Area Population**

Species	No. of pairs	UK Breeding Population	Regional Breeding Population	County Status	Geographical Importance
<b>Mallard</b>	9	61,000-146,000	-	Surrey: common breeding resident. Sussex: common resident and winter visitor.	Local
<b>Little ringed plover<sup>1</sup></b>	1	1,115	123	Surrey: summer visitor breeding annually in small numbers and passage migrant (estimated at 10 pairs in 2016). Sussex: scarce breeding summer visitor and passage migrant (14 pairs in 2016).	County
<b>Stock dove</b>	3	260,000	-	Surrey: common breeding resident and passage migrant. Sussex: common resident and possible winter visitor.	Local
<b>Kestrel</b>	4	46,000	-	Surrey: moderately common breeding resident. Sussex: Fairly common resident and passage migrant.	Local
<b>Peregrine<sup>1</sup></b>	1	1,731	93	Surrey: increasing breeding resident, passage migrant and winter visitor (14 pairs in 2016). Sussex: scarce breeding resident (33 pairs in 2016).	Regional
<b>Marsh tit</b>	1	41,000	-	Surrey: uncommon and declining breeding resident. Sussex: scarce resident.	County
<b>Skylark</b>	12	1,500,000	-	Surrey: common but declining breeding resident, passage migrant and winter visitor. Sussex: very common but declining resident; and probably common passage migrant and winter visitor.	Local
<b>Starling</b>	2	1,900,000	-	Surrey: common breeding resident. Sussex: common but declining resident; and very common to abundant winter visitor.	Local
<b>Song thrush</b>	19	1,200,000	-	Surrey: common breeding resident. Sussex: very common but decreasing resident and partial migrant; common passage migrant and winter visitor.	Local
<b>Mistle thrush</b>	2	170,000	-	Surrey: common breeding resident. Sussex: common resident and partial migrant.	Local
<b>Firecrest<sup>1</sup></b>	1	4,000+	c.250	Surrey: moderately common breeding resident, passage migrant and winter visitor (estimated at 150 singing males in 2016). Sussex: scarce or possibly fairly common breeding resident; passage migrant; and winter visitor (estimated at 100 singing males in 2015).	County
<b>House sparrow</b>	4	5,300,000	-	Surrey: common breeding resident. Sussex: very common but possibly declining resident.	Local
<b>Dunnock</b>	18	2,500,000	-	Surrey: common breeding resident. Sussex: very common resident.	Local
<b>Grey wagtail</b>	1	38,000	-	Surrey: moderately common breeding resident and passage migrant. Sussex: scarce resident and fairly common passage migrant and winter visitor.	Local
<b>Bullfinch</b>	1	220,000	-	Surrey: moderately common breeding resident. Sussex: fairly common or common resident.	Local
<b>Linnet</b>	1	430,000	-	Surrey: moderately common resident, passage migrant and winter visitor. Sussex: common but decreasing resident and partial migrant.	Local
<b>Reed bunting</b>	2	250,000	-	Surrey: moderately common breeding resident. Sussex: fairly common resident; passage migrant and winter visitor.	Local

## Annex 4

### Invertebrate Scoping Survey

# COLIN PLANT ASSOCIATES (UK)

## CONSULTANT ENTOMOLOGISTS

30a Alexandra Road, London, N8 0PP

Tel: [REDACTED] Mobile telephone: [REDACTED]  
E-mail: [REDACTED]@btinternet.com

---

FAO Nick Betson  
RPS Group Plc  
Lakesbury House  
Hiltingbury Road  
Chandlers Ford  
Southampton  
SO53 5SS

25<sup>th</sup> June 2019

Our Reference: CPA - 19110

### **Land bordering Gatwick Airport: Appraisal of invertebrate habitats outside the Biodiversity Areas**

Dear Nick,

Further to your instruction of 24<sup>th</sup> May 2019, we have now visited the above site; the surveyors on this occasion were Marcel Ashby and Tristan Bantock. This letter is our formal report of that visit.

#### **Statement of impartiality**

Please note that this report presents our surveyors' impartial and unbiased opinion on the existing invertebrate ecology of the site at the date of examination. Unless otherwise stated, our findings and any conclusions drawn or recommendations made are independent of the detail of any proposed development to the site and are wholly independent of any third party opinions where these may exist.

If this report contains suggestions or recommendations relating to mitigating losses, these have been made without specific consideration of the details of the proposed development works and are offered on the assumption that the entire area inside the red line would be lost.


#### **Introduction and scope of visit**

The purpose of the visit was to appraise the invertebrate habitats present on site and to advise whether or not it is likely that a proposed development would have an impact on invertebrate ecology. Of particular concern was the potential for the site to support Species of Principal Importance in England, as defined within Section 41 of the *Natural Environment and Rural Communities (NERC) Act 2006*, although species included in other conservation categories were also considered.


It was previously agreed that the scope of the appraisal should focus solely on land outside the two existing Biodiversity Areas. These two areas comprise (1) the River Mole corridor and Brockley Wood, located close to the north west perimeter and (2) land east of the railway line, including the Gatwick Stream, Horleyland Wood, Lower Picketts Wood, Upper Picketts Wood, Goat Meadow, Rolls Field and Ashleys Field. Both

Biodiversity Areas are managed for nature conservation and are known to support important invertebrate assemblages on the basis of recent records gathered by Gatwick Airport Biodiversity Consultant Rachel Bicker. These include various species of conservation significance such as Dingy Skipper *Erynnis tages*, Grizzled Skipper *Pyrgus malvae*, Brown Hairstreak *Thecla betulae*, Long-horned Bee *Eucera longicornis* and Black-headed Mason Wasp *Odynerus melanocephalus*, all of which are Section 41 species (Bicker, 2018).

In the light of the above, eight areas were selected which were considered to be of potential importance for invertebrates. The locations of these are shown in Figures 1 and 2. The site visit was undertaken on 17<sup>th</sup> June in sunny and warm conditions.



**Fig. 1** Areas examined in the eastern sector (Sites A - C).



**Fig. 2** Areas examined in the western sector (Sites D - H).

## INVERTEBRATE HABITATS PRESENT IN JUNE 2019

### Site A: Pentagon Field and adjacent ditches

This area lies directly north of Lower Picketts Wood and covers approximately 10 ha. The habitats present comprise an expanse of dry semi-improved neutral grassland bounded by a wooded hedgerow and a ditch on the western margin. A strip of immature plantation woodland is present along the southern boundary.

The grassland is rather uniform in nature and very few herbaceous species are represented in the sward. This lack of structural variation combined with its low floristic diversity predicts a species-poor invertebrate assemblage dominated by those with more generalist ecological requirements, which are usually of lower conservation value.

The hedgerow includes numerous overmature oaks, some of which contain obvious amounts of standing dead wood and aerial wood decay features, including a large red rotten cavity at the base of one tree. These trees offer a range of potential niches for invertebrates which are both phytophagous and saproxylic.

In Britain alone, there are at least 700 native species of beetle (Coleoptera) and over 700 species of two-winged fly (Diptera) which appear to be dependent on decaying wood at some stage in their life cycles. Many of these are of high conservation value and are listed as Section 41 species.

The ditch on the western margin is open and unshaded along much of its length and held a high water level on the day of examination. The ditch profile is gently shelving which has allowed a diverse riparian and emergent flora to develop, including Hemlock Water-dropwort and Meadowsweet, as well as sedges and stands of bulrush. These are positive features that suggest its potential importance for invertebrates is likely to be raised.

Overall we consider that Site A has a **moderate** intrinsic invertebrate interest.

#### **Site B: Pond F**

This area covers approximately 1.7 ha and lies between the A23 to the north, railway lines to the west and airport car parks to the south.

It comprises a single waterbody which is rather deep and steep sided and was probably originally constructed as a balancing pond. The pond is surrounded by a narrow zone of alder and willow scrub, but there is only minimal emergent vegetation at the margins and the bankside vegetation is largely dominated by dense bramble scrub. Several mats of White Water-lily are present. The water is subject to nutrient enrichment by wildfowl and presumably there is also some runoff from surrounding roads.

Overall we consider that Site B has a **low** intrinsic invertebrate interest.

#### **Site C: Riverside Garden Park and ponds**

This area covers approximately 11 ha and lies between the A23 to the south and the urban edge of Horley to the north.

The site presents as a mosaic of dense mature woodland interspersed with open areas of grassland. A range of tree species are represented including oak, hawthorn and elder which offer numerous niches for phytophagous invertebrates, but only a minimal standing or fallen dead wood resource is apparent with the exception of a single dead barkless oak. The grassland is highly improved in nature and a minimal herbaceous flora is present in the open areas. In places the woodland understorey is dominated by dense stands of stinging nettles, indicating high soil fertility.

A stocked fishing lake is present close to the southern boundary. The water column appeared turbid and is presumably subject to extensive nutrient enrichment from the large numbers of feral Greylag Geese present. Areas of marginal vegetation are minimal and emergent macrophytes are represented only by small stands of Yellow Flag and some cover by White Water-lily.

The Gatwick Stream runs through the Riverside Garden Park but the channel is very eroded and steep-sided, supporting minimal riparian vegetation and dominated by dense bramble cover.

Overall we consider that Site C has a **low** intrinsic invertebrate interest.

#### **Site D: Pond D**

This area covers approx. 2 ha and is located between the northern airport perimeter and the River Mole.



The eastern half of the site comprises a pond adjacent to a water management facility which makes up the western section. The pond margins were bare and muddy indicating recent fluctuation and the surrounding banks dominated by an improved grass sward which had recently been mown. It was not possible to physically access the pond edges but the marginal flora appeared to be impoverished and represented solely by small stands of rushes and sedges.

Overall we consider that Site D has a **low** intrinsic invertebrate interest.

#### **Site E: Dog Kennel Wood, pond and ditch**

This area covers approximately 2.7 ha and is located very close to the northern airport perimeter and largely surrounded by the built environment of the airport on all sides.

Despite this it contains a range of habitats, comprising a water body and a small copse of mature woodland which encloses the dog kennels. The pond is set in a deep and roughly triangular depression and is almost entirely vegetated, with only minimal standing water apparent on the day of examination. The western half is dominated by Common Reed and the remainder by bulrush, Hemlock Water-dropwort, Meadowsweet, rushes and sedges, with a number of small willows also present.

The bankside flora comprises dry semi-improved neutral grassland with a range of herbaceous species including Creeping Cinquefoil, Common Bird's-foot Trefoil, Self-heal and tall ruderal species such as ragworts, docks, teasel and Perforate St John's Wort. The sloping nature of the bank presents a warm south-facing aspect across the northern section, a positive feature for invertebrates requiring a warm microclimate at the ground surface.

The relatively diverse flora which includes a range of host plants in combination with the transition from wet to dry soils provides a large range of potential niches for invertebrates. The presence of a range of mature trees in the adjoining woodland contributes to the overall interest.

Overall we consider that Site E has a **moderate** intrinsic invertebrate interest.

#### **Site F: Pond M and top of environment bund**

This area covers approximately 6 ha and is located between the northern airport perimeter and the River Mole.

Various habitats are present around a concrete-sided water body which is split into two halves. This pond is presumably used in silt extraction as the western half was almost entirely dry on the day of examination and the bed entirely covered by silt deposits. This area has some potential for invertebrates which require very fine-grained sediments, although is unlikely to support a rich fauna.

The surrounding area comprises dry semi-improved neutral grassland with a range of herbaceous species including Creeping Cinquefoil, Common Bird's-foot Trefoil, Meadow Vetchling, Grass Vetchling and Tufted Vetch, as well as tall ruderal flora in the form of docks and thistles. The structural variation within the grassland, combined with its floristic diversity, predicts that various plant-feeding groups of invertebrates such as phytophagous beetles and true bugs may have rich faunas. During the visit a single Section 41 species were noted, the Small Heath *Coenonympha pamphilus*. This area possibly lies within the foraging range of the Long-horned Bee *Eucera longicornis* which is known to nest along the adjacent River Mole corridor and use legumes such as vetchlings and trefoils as its principal forage plants.

Overall we consider that Site F has a **moderate** intrinsic invertebrate interest.

### Site G: Gatwick Aviation Museum and Brook Farm

This large area covers approximately 35 ha and presents as a network of hedgerows dominated by mature oaks surrounded by dry grassland.

The grassland is rather improved in nature around the Aviation Museum and has been recently mown to produce amenity areas. Further east the sward is more diverse and presents as dry semi-improved neutral grassland with a range of herbaceous species including Common Bird's-foot Trefoil, Meadow Vetchling, Grass Vetchling and Foxglove, offering a range of niches for phytophagous invertebrates. During the visit a single Section 41 species were noted, the Small Heath *Coenonympha pamphilus*. The eastern boundary of the site possibly lies within the foraging range of the Long-horned Bee *Eucera longicornis* which is known to nest along the adjacent River Mole corridor and use legumes such as vetchlings and trefoils as its principal forage plants.

Several ponds are present along the southern margin although the water column is entirely covered by duckweed. Stands of bulrush and Hemlock Water-dropwort are also apparent.

Numerous open grown overmature oaks are present in the hedgerows which contain a significant standing dead wood resource and may support a range of saproxylic invertebrates of conservation significance.

Overall we consider that Site G has a **moderate** intrinsic invertebrate interest.

### Site H: Westfield Stream site

This area covers approximately 3 ha and is located between the southern boundary of Site G and the airport perimeter.

The site presents as a mosaic of wet and dry habitats with elements of wet woodland grading through to dry, sparsely-vegetated areas. The transitional nature of the habitats present on this site ensure that numerous potential niches for invertebrates are represented.

The Westfield stream runs along the western margin and was almost dry on the day of examination. The channel contains stands of bulrush and Hemlock Water-dropwort, while the tops and sides of the bank support a community of ephemeral short perennial vegetation, including Creeping Cinquefoil, Common Bird's-foot Trefoil, Common Mallow and Meadow Vetchling, as well as numerous alder saplings which are presumably rather heat-stressed. An area of damp woodland containing alder, willow and White Poplar is present in the southwest sector of the site, while the areas of woodland along the northern edge are drier and contain more oak. In freely draining parts of the site extensive stands of gorse are apparent, an important plant for invertebrates, while the areas that retain a wetter influence throughout the year support numerous *Juncus* tussocks.

Overall we consider that Site H has a **moderate** intrinsic invertebrate interest

### Conclusions and recommendations

Several of the sites under discussion presents features of potential value to invertebrates and in our opinion, have a moderate invertebrate interest that is likely to be raised above the expected regional background level.

## References

Bicker, R. (2018) Gatwick Biodiversity Action Plan: Five Year Review 2012-2017.

*\* \* end of formal report \* \* \**

I hope that you will find this report adequate for your client's current needs.

With all best wishes,



Tristan Bantock  
Partner

## Annex 5

### Terrestrial Invertebrate Survey



# **Gatwick Airport Northern Runway Project –**

## **Assessment of Terrestrial Invertebrate Interest**

**RPS Group Plc**

*Report prepared by:*  
Ecus Ltd.  
Unit 1  
Woodlands Business Village  
Coronation Road  
Basingstoke  
RG21 4JX

01256 224588



**June 2021**


## Ecus Ltd


Report to: **RPS Group Plc**

Report Title: **Gatwick Airport Northern Runway Project - Assessment of Invertebrate Interest**

Version: **V1.0**  
Issue Date: **June 2021**  
Report Ref: **14864**

Originated By:    
**Alex Hellyar** **Richard Jones**  
**Consultant Ecologist** **Entomologist** Date: March 2021

Reviewed By:   
**Abel Drewett**  
**Technical Director**  
**(Ecology)** Date: June 2021

Approved By:   
**Paul White**  
**Regional Director** Date: June 2021

Version	Author	Description	Date
0.1	AH/RJ	Initial draft	19/03/2021
0.2	AD	QA1 Review	16/06/2021
0.3	AD	Edits	18/06/2021
1.0	PAW	Approve and issue to client	21/06/2021

The report and the site assessments carried out by Ecus on behalf of the client in accordance with the agreed terms of contract and/or written agreement form the agreed Services. The Services were performed by Ecus with the skill and care ordinarily exercised by a reasonable Environmental Consultant at the time the Services were performed. Further, and in particular, the Services were performed by Ecus taking into account the limits of the scope of works required by the client, the time scale involved and the resources, including financial and manpower resources, agreed between Ecus and the client.

Other than that expressly contained in the paragraph above, Ecus provides no other representation or warranty whether express or implied, in relation to the services.

This report is produced exclusively for the purposes of the client. Ecus is not aware of any interest of or reliance by any party other than the client in or on the services. Unless expressly provided in writing, Ecus does not authorise, consent or condone any party other than the client relying upon the services provided. Any reliance on the services or any part of the services by any party other than the client is made wholly at that party's own and sole risk and Ecus disclaims any liability to such parties.

This report is based on site conditions, regulatory or other legal provisions, technology or economic conditions at the time of the Service provision. These conditions can change with time and reliance on the findings of the Services under changing conditions should be reviewed.

Ecus accepts no responsibility for the accuracy of third party data used in this report.

## Contents

<b>SUMMARY</b> .....	<b>IV</b>
<b>1. INTRODUCTION</b> .....	<b>1</b>
1.1 BACKGROUND.....	1
<b>2. CONSERVATION STATUS AND LEGAL PROTECTION</b> .....	<b>2</b>
2.1 CONSERVATION STATUS.....	2
2.2 LEGISLATION.....	3
<b>3. METHODOLOGY</b> .....	<b>4</b>
3.1 SURVEY VISITS.....	4
3.2 SITE COMPARTMENTS.....	4
3.3 LOCATION AND COLLECTION OF SPECIMENS.....	4
3.4 TAXONOMIC COVERAGE.....	4
3.5 LIMITATIONS.....	4
<b>4. RESULTS</b> .....	<b>5</b>
4.1 GENERAL.....	5
4.2 NOTEWORTHY SPECIES.....	5
<b>5. SITE ASSESSMENT</b> .....	<b>13</b>
5.1 DISCUSSION.....	13
<b>6. CONCLUSION</b> .....	<b>14</b>
<b>7. REFERENCES</b> .....	<b>15</b>
<b>FIGURE 1. SITE LOCATIONS (THREE SEPARATE PARCELS - GATWICK AIRPORT)</b> .....	<b>17</b>
<b>APPENDIX 1. FULL SPECIES LIST</b> .....	<b>18</b>

## Summary

Ecus Ltd was commissioned by RPS Group Plc in May 2020 to undertake terrestrial invertebrate surveys of three separate land parcels/compartments adjoining Gatwick Airport as part of the surface water management and flood alleviation for the Northern Runway Project development. The three survey areas comprised Compartment P: Riverside Park (National Grid Reference (NGR): TQ 28055 42020, Compartment M: River Mole Corridor NGR: TQ 25772 40623 and Compartment G: Gatwick Brook Grasslands NGR: TQ 29000 39799, hereafter referred to as 'the Site'. The three land parcel locations are shown on **Figure 1**.

Although Gatwick Airport is now nominally in the County of West Sussex, for the purposes of biological recording it is in Vice-County 17, Surrey.

Six site field visits were made during 2020: 27<sup>th</sup> May, 19<sup>th</sup> June, 22<sup>nd</sup> June, 30<sup>th</sup> June, 10<sup>th</sup> September and 14<sup>th</sup> September. Six visits were made to cover all sites in total, however all three parcels were not covered on all six visits, but more on a rotation basis to cover each one at appropriate points across the 2020 season.

A list of 303 terrestrial invertebrate species was recorded in total (see **Appendix 1**). This total was considered a relatively diverse list for such a site.

Numerous unusual and scarce insects were found, including:

- *Acinia corniculata*, a nationally rare fly that breeds in the seed-heads of knapweed.
- *Catoplatus fabricii*, a nationally scarce lacebug that breeds on oxeye daisy.
- *Dioxya bidentis*, a nationally scarce fly that breeds on trifid bur-marigold.
- *Dorycera graminum*, a nationally rare fly of damp meadows and floodplains.
- *Ectobius lapponicus*, the dusky cockroach, a species of rough heathy grasslands.
- *Hylaeus cornutus*, a nationally scarce yellow-faced bee.
- *Merzomyia westermanii*, a nationally scarce picture-winged fly that breeds on ragwort.
- *Paraclusia tigrina*, a nationally rare fly of old broad-leaved woodlands.
- *Podagrica fuscicornis*, a nationally scarce leaf beetle that feeds on mallows.
- *Reptalus quinquicostatus*, a nationally scarce frog-hopper of dry grasslands.
- *Rhinocyllus conicus*, a nationally scarce weevil that feeds on thistles.
- *Squamapion vicinum*, a nationally scarce weevil that feeds on water mint.
- *Tachys bistriatus*, a nationally scarce ground beetle of damp muddy pond edges.
- *Uleiota planata*, a nationally scarce beetle that breeds under fungoid tree bark.
- *Variimorda villosa*, a nationally scarce flower beetle associated with ancient woods.

The diverse species list and numerous scarce and unusual insects recorded, reflect a diversity of habitat types present at the Site: river edge, flowery grassland, hedgerows, disturbed ground, woodland and scrub.

The individual parcels are suggested as having the following invertebrate interest:

- Riverside Park — low.
- River Mole Corridor — medium/high.
- Gatwick Brook Grasslands — low/medium.



# 1. Introduction

## 1.1 Background

- 1.1.1 Ecus Ltd was commissioned by RPS Group Plc in April 2020 to undertake terrestrial invertebrate surveys of three land parcels adjoining Gatwick Airport as part of the surface water management and flood alleviation for the Northern Runway Project development. The three land parcels (compartments), hereafter referred to as 'the Site', are described below and the location is shown in **Figure 1**.
- 1.1.2 **Compartment P: Riverside Park.** This is a public open space, mainly wooded, although likely more open in the past. It is heavily managed with plentiful short amenity grassland. Of little interest in terms of entomology, apart from a narrow corridor alongside the A23 which was the focus for survey here. This area is centred on National grid reference TQ 28055 42020.
- 1.1.3 **Compartment M: River Mole Corridor.** This is an irregular plot to the west of the airport, comprising brownfield zones, rough grassy areas, the River Mole and riverbanks, hedgerows and woodland edges. This area is centred on National grid reference TQ 25772 40623.
- 1.1.4 **Compartment G: Gatwick Brook Grasslands.** This is an irregular plot to the east of the airport, comprising rough grassy areas, a few mature trees, scrub, Gatwick Brook, hedgerows and woodland edges. This area is centred on National grid reference TQ 29000 39799.
- 1.1.5 This survey is part of a wider ecological assessment of the site to be undertaken by RPS Group Plc.

## 2. Conservation Status and Legal Protection

### 2.1 Conservation Status

- 2.1.1 The national significance of species recorded in this survey is assessed here with regard to the Red Data Book Species. IUCN guidelines are used to give rare insects a status. It is, however, dependent on the degree of threats that they face (originally published in Shirt, 1987; Hyman & Parsons, 1992 and recently updated in a series of ongoing reviews).
- 2.1.2 Statuses continue to be assessed and reassessed over time. **Table 1** details the statuses that can be applied at this current time.

**Table 1: Categories and criteria used to assess invertebrate statuses**

Status	Description
Endangered (RDB-1)	The rarest taxa. Taxa in danger of extinction in Great Britain; species with very few recorded localities or living in especially vulnerable habitats.
Vulnerable (RDB-2)	Very rare species. Taxa likely to move into the RDB1 category; species declining in their range.
Rare (RDB-3)	Rare species. Taxa with small populations and which are at risk; species estimated to occur in 15 or fewer of the 10-km squares in the national Ordnance Survey grid since 1970, or nominated later date if applicable.
Insufficiently known (RDB-K)	Species thought to be very rare in Britain, recorded from less than 15 of the 10-km squares of the national Ordnance Survey grid since 1970 or later date, and which warrant RDB classification of some sort, but for which there is a recognized lack of accurate information.
Nationally scarce (notable A)	Very local species, thought to occur in 16 to 30 of the 10-km squares of the national Ordnance Survey grid since 1970, or later date.
Nationally scarce (notable B)	Very local species, thought to occur in 31 to 100 of the 10-km squares of the national Ordnance Survey grid since 1970 or later date.
Nationally scarce	Status is sometimes not subdivided into categories A and B, (notable, occurring in 16 to 100 10-km squares).
Very local	Status is a much more subjective, but nevertheless useful, measure of scarcity and is based on personal experience, published and unpublished records. It is applied to species that are very limited in distribution or confined to very limited specialist habitats. This group includes species previously considered nationally rare or scarce, but which have had statuses reviewed following more recent study.

## 2.2 Legislation

- 2.2.1 Three invertebrate species occurring in the UK are European protected species (EPS), so are therefore protected under European law. These species are Fisher's Estuarine Moth (*Gortyna borelii lunata*), Large Blue Butterfly (*Phengaris arion*) and Lesser Whirlpool Ramshorn Snail (*Anisus vorticulus*).
- 2.2.2 It is an offence to capture, kill, disturb or injure these species. As well as to damage or destroy their breeding or resting places or to obstruct access to any such place (either deliberately or accidentally).
- 2.2.3 Forty invertebrate species occurring in the UK are included on Schedule 5 Section 9.1 of the Wildlife & Countryside Act 1981 (as amended). This it makes it an offence to kill, injure or take any of the species.
- 2.2.4 Four species are listed under Schedule 5 Section 9.4 of the Wildlife & Countryside Act 1981 (as amended). This makes it illegal to damage or destroy their breeding or resting places or to obstruct access to any such place.
- 2.2.5 Twenty seven species are listed under Schedule 5 Section 9.5 of the Wildlife & Countryside Act 1981 (as amended), which prevents them from being sold or transported.

## **3. Methodology**

### **3.1 Survey Visits**

3.1.1 The surveys were undertaken by an experienced entomologist to the Site six times during 2020: 27<sup>th</sup> May, 19<sup>th</sup> June, 22<sup>nd</sup> June, 30<sup>th</sup> June, 10<sup>th</sup> September and 14<sup>th</sup> September. Six visits were made to cover all sites in total, however all three sites were not covered on all six visits, but more on a rotation basis to cover each one at appropriate points across the 2020 season. A walk-over assessment of the Site, taking note of habitats and features in relation to invertebrates was complemented by the collection of specimens for either Site or subsequent laboratory analysis.

### **3.2 Site Compartments**

3.2.1 Three broad compartments/parcels were identified which equated to three separate sites. These sites are described in Section 1.1 and illustrated in Figure 1. Each of these compartments was visited on at least three occasions during 2020. In the following species descriptions, reference is made of these compartments and 10-figure National grid references are also provided where appropriate.

### **3.3 Location and Collection of Specimens**

3.3.1 A walk-over type survey was carried out. Invertebrates were located and collected by general methods using sweep net, beating tray and a stout trowel. Flowers, leaf surfaces, rocks, bare ground, logs and tree trunks were examined by visual searching. Others were found by finger-tip grubbing in loose soil, rubble, and plant roots, logs, stumps and animal dung. Squares of roofing felt left out for reptile monitoring were also examined. Voucher specimens of all but the most common and characteristic species were collected for examination later under the microscope.

### **3.4 Taxonomic Coverage**

3.4.1 The survey concentrated on the following major groups (orders): Coleoptera (beetles), Diptera (flies), Hemiptera (bugs, froghoppers, etc), Hymenoptera (bees, wasps and ants) and Lepidoptera (butterflies and moths). Some examples of other groups were noted if found.

3.4.2 These are hugely numerous and diverse orders of insects, and identification to species is not always possible, especially in many of the microscopically small specimens. Consequently on any given site, there is much subjective selection of which families or genera are worth taking as sample specimens, for later study. This is often influenced by a knowledge of the groups for which useable identification keys are available, and for which the individual entomologist has particular experience. Nevertheless, a wide coverage of insect orders allows some assessment of just how important any given site may be for its invertebrate biodiversity.

### **3.5 Limitations**

3.5.1 The surveys were undertaken during the optimal season for terrestrial invertebrates (April – September inclusive) and all areas of the land parcels could be accessed safely. It is therefore considered that the findings of the survey provide an accurate representation of the insect assemblages present.

## 4. Results

### 4.1 General

4.1.1 A list of 303 terrestrial invertebrate species was recorded across the 2020 season. They represent a range of different groups of insects as set out in **Table 2** below.

**Table 2: Orders of invertebrates recorded**

Order & Group Common Name	Total of Species
Coleoptera (beetles)	119
Dermaptera (earwigs)	1
Diptera (flies)	65
Hemiptera (bugs)	48
Hymenoptera (bees, wasps, etc)	24
Lepidoptera (butterflies & moths)	23
Dictyoptera (cockroaches)	1
Odonata (dragonflies)	7
Orthoptera (grasshoppers)	6
Aranaea (spiders)	3
Opiliones (harvestmen)	1
Isopoda (woodlice & hoglice)	3
Mollusca (slugs & snails)	2
<b>TOTAL</b>	<b>303</b>

4.1.2 303 species is a relatively high number and reflects a diversity produced by several different habitat types spread over three separate sites. Several interesting and unusual species are included in this list as discussed below.

4.1.3 The full list of species recorded is provided in **Appendix 1**.

### 4.2 Noteworthy Species

4.2.1 Most of the insects recorded were common examples, which might be expected to occur in any open area in southern England. However, a number are uncommon or otherwise unusual and merit highlighting.

4.2.2 Common or garden species occur commonly in gardens, or indeed almost anywhere; they are often mobile, adaptable, fast-reproducing with quick generation times, feeding on common and widespread plants or occurring in a wide variety of diverse habitat types. They tell us very little about a site since they often occur in almost every bit of open space available.

4.2.3 Scarce species, however, are scarce because they have very particular habitat requirements — they feed on scarce plants which only occur in limited habitat pockets, they have very narrow

toleration of climate, including daily or yearly temperature or rainfall minima or maxima, or they only occur in niches where they can avoid serious competition, predation or parasitism from abundant and widespread species. The occurrence of these scarcer species gives a much clearer picture of the environmental health or conservation biodiversity status of a particular site.

4.2.4 The following species are highlighted as being especially noteworthy. Most are uncommon nationally. Criteria for allocation of accepted 'nationally rare' (previously red data book) and 'nationally scarce' (previously notable) statuses are varied and complex (originally published in Shirt, 1987; Hyman & Parsons, 1992 and recently updated in a series of ongoing reviews). Statuses continue to be assessed, reassessed and altered over time and a JNCC database is available giving an up to date summary overview. Every time a rare insect is found there are more records added to the scoring system (based on grid squares in which an insect is found) and it becomes less rare. These statuses are useful to gauge relative rarity, but despite the apparent objectivity of counting numerical records, there is still a subjective element of exactly how rare an organism may be. Those that are relevant to this Site are listed in brief here:

- **Endangered (RDB-1).** The rarest taxa. Taxa in danger of extinction in Great Britain; species with very few recorded localities or living in especially vulnerable habitats.
- **Vulnerable (RDB-2).** Very rare species. Taxa likely to move into the RDB1 category; species declining in their range.
- **Rare (RDB-3).** Rare species. Taxa with small populations and which are at risk; species estimated to occur in 15 or fewer of the 10-km squares in the national Ordnance Survey grid since 1970, or nominated later date if applicable.
- **Insufficiently known (RDB-K).** Species thought to be very rare in Britain, recorded from less than 15 of the 10-km squares of the national Ordnance Survey grid since 1970 or later date, and which warrant RDB classification of some sort, but for which there is a recognized lack of accurate information.
- **Nationally scarce (notable A).** Very local species, thought to occur in 16 to 30 of the 10-km squares of the national Ordnance Survey grid since 1970, or later date.
- **Nationally scarce (notable B).** Very local species, thought to occur in 31 to 100 of the 10-km squares of the national Ordnance Survey grid since 1970 or later date.
- **Nationally scarce** status is sometimes not subdivided into categories A and B, (notable, occurring in 16 to 100 10-km squares).
- **Very local** status is a much more subjective, but nevertheless useful, measure of scarcity and is based on personal experience, published and unpublished records. It is applied to species that are very limited in distribution or confined to very limited specialist habitats. This group includes species previously considered nationally rare or scarce, but which have had statuses reviewed following more recent study.

4.2.5 The following is a description of the more interesting and noteworthy species taken at the Site. Where possible a nominal 10-figure National grid reference is given to indicate the exact location(s) where they were found.

***Acinia corniculata* (Zetterstedt)**

4.2.6 A small pink picture-winged fly, family Tephritidae. Status: endangered (red data book category 1, Shirt, 1987; Falk, 1991b). At the time of the national review this very rare fly was known only from a handful of localities in southern England, all National Nature Reserves. It breeds in the seed heads of common knapweed, but despite the abundance of its host plant, it remains very elusive. Until the late 1990s, it seemed to be primarily associated with a few East Anglian fens, but has recently been recorded from several localities in Sussex, London, Surrey, Hampshire, Dorset and

Gloucestershire (Clemons, 1997, 2004, 2015). It may be increasing its range after a series of warm summers and mild winters. It remains, nevertheless, very elusive, but its status may need revision. Several specimens were found by sweeping knapweed, Gatwick Brook Grassland, dated 22.6.2020, at TQ 28941 40057.

***Athous campyloides* (Newman)**

- 4.2.7 A medium-sized brown click beetle, family Elateridae. Status: nationally scarce (notable B, Hyman & Parsons, 1992). This very local species is associated with rough grassy places in south-east England (Mendel & Clarke, 1996). The larvae are thought to feed at the roots of grass and herbs. It was once regarded as an extremely rare species, but appears to have colonized Britain in the early 19th century, and is still spreading (Jones, 2001). Several specimens were found by sweeping, River Mole Corridor and Gatwick Brook Grassland, dated 27.5.2020 and 19.6.2020.

***Bruchidius imbricornis* (Panzer)**

- 4.2.8 A tiny mottled bean weevil, family Chrysomelidae. Status: very local. A recent colonist to Britain, this small but distinctive beetle was first found, in Essex, in 2012. Its food plant is nominally goat's rue, *Galega officinalis*, a widespread alien vetch that has become widely established in brownfield sites in England. Several specimens were found by sweeping, River Mole Corridor, dated 19.6.2020 and 10.9.2020.

***Camarota curvipennis* (Latreille)**

- 4.2.9 A minute black 'frit' fly, family Chloropidae. Status: very local. Once much more widespread, breeding the heads of wheat, rye and barley, this species has declined dramatically in the last 50 years following 'advances' in agriculture. Although not accorded notable status in the national review by Falk (1991b), the decline of this fly has alerted dipterists to suggest that this species be monitored for possible future nationally scarce notification. Two specimens were found by sweeping, Gatwick Brook Grassland, dated 14.9.2020, at TQ 28976 39926.

***Campiglossa malaris* (Seguy)**

- 4.2.10 A minute pink and grey picture-winged fly, family Tephritidae. Status: very local. Originally suggested to be nationally rare, but insufficiently known (red data book category K, Clemons, 1997), this status has not been official agreed by JNCC. This scarce fly is thought to feed on ragworts, but whether it forms stem or leaf galls, or breeds in the seed heads is not known. In the 1970s and 80s it was known from only two UK sites, both on the Kent coast, and was accorded endangered status (red data book category 1) by Shirt (1987) and Falk (1991b), but this was later revised to RDB-K by Clemons (1997, 2004) when further Kent localities were discovered. Since then further reports appear to document a spread into England and there are now numerous records from inland Kent, East Sussex and other Home Counties and outliers beyond into central England. This species often occurs in rough grassy places and disturbed ground where the food-plants grow. The fly remains very rare, but its status may need another revision. It is currently not accorded any conservation status by JNCC. Several specimens were found by sweeping, River Mole Corridor, dated 19.6.2020, at TQ 26064 40625.

***Catoplatus fabricii* (Stal)**

- 4.2.11 A minute pale lacebug, family Tingidae. Status: nationally scarce (notable B, Kirby, 1992). This scarce lacebug feeds on ox-eye daisy, *Leucanthemum vulgare* and although the food-plant grows commonly and widely on disturbed ground, chalk downland, verges, railway cuttings and rough meadows, the bug is extremely local. It occurs in widely scattered localities in central and southern England, but its precise ecological requirements are unclear. A single specimen was swept from the roadside verge of Riverside Park, dated 27.5.2020, at TQ 28105 41966.

***Coccidula scutellata* (Herbst)**

- 4.2.12 A small red and black ladybird, family Coccinellidae. Status: very local. Although widespread across much of England and parts of Wales, the localities for this beetle are widely scattered. It is

confined to freshwater sites, stream banks, pond-sides, and marshes (Roy *et al.*, 2011). Several specimens were swept from waterside vegetation, River Mole Corridor, dated 27.5.2020, at TQ 25573 40591.

***Dioxya bidentis* (Ronbinaeu-Desvoidy)**

- 4.2.13 A small grey picture-winged fly, family Tephritidae. Status: nationally scarce (Notable, Falk, 1991b). This very scarce fly occurs in a few widely scattered localities in England and South Wales, with an old record from Scotland (Clemons, 1997, 2004, 2015). It breeds in the flower heads of various composites (Asteraceae), particularly *Bidens tripartita*, the trifid bur-marigold. Several specimens were swept from vegetation bordering the stream, River Mole Corridor, dated 10.9.2020, at TQ 25565 40797.

***Dorycera graminum* (Fabricius)**

- 4.2.14 A small mottled fly, family Ulidiidae. Status: nationally rare (red data book category 3, Shirt, 1987; Falk, 1991b). *Dorycera* is usually associated with herb-rich unimproved meadows, often in association with umbellifers and broad-leaved trees. The life history is unknown, but the larvae probably develop in decaying vegetable matter, possibly in the dead or dying roots of hogweed, *Heracleum sphondylium* or a near relative. It was once regarded as a fairly frequent insect, but appears to have declined dramatically in recent years. Threats are thought to come from loss of unimproved flowery meadows through drainage or lack of grazing leading to scrub invasion. Although there are old records from Hampshire and Worcestershire, most of the recent records are from the Thames Estuary where it regularly occurs on brownfield sites (Ismay, 2000; Jones, 2003, 2007). On a personal note, however, I have recorded this fly in many widely spread localities in the Home Counties in the last 5 years, indicating that it may be increasing and spreading in some areas at least. Two specimens were found sweeping, River Mole Corridor, dated 27.5.2020, at TQ 25825 40566.

***Ectobius lapponicus* (Linnaeus)**

- 4.2.15 The 'dusky cockroach', family Blatellidae. Status: nationally scarce (notable B, Haes & Harding, 1997). This is one of Britain's native cockroach species, not to be confused with the many domestic pest species that have been introduced into buildings. It is very uncommon, and in Britain it is confined to southern England and most colonies are either in the Sussex Weald or the New Forest (Marshall & Haes, 1988) or west Surrey (Baldock, 1999). It is an omnivorous scavenger, living in grass litter. Several specimens were found by sweeping, River Mole Corridor, dated 27.5.2020, at TQ 25570 40654.

***Hippodamia* (formerly *Adonia*) *variegata* (Goeze)**

- 4.2.16 The Adonis ladybird, family Coccinellidae. Status: nationally scarce (notable B, Hyman & Parsons, 1992), but status may need revision. Until about 30 years ago, this species was always regarded as having a coastal distribution, occurring in warm sheltered locations such as chalk downs, dunes, undercliffs and other disturbed areas (Majerus *et al.*, 1997). However, it is now known to be fairly widespread in England, especially in the London area and Thames Estuary, where it is associated with sparsely vegetated post-industrial brownfield sites, and it has also spread across central England (Roy *et al.*, 2011). Several specimens were found by general sweeping of sparse herbage, River Mole Corridor, dated 10.9.2020, at TQ 25872 40560.

***Hylaeus cornutus* (Curtis)**

- 4.2.17 A small black white-faced bee, family Colletidae. Status: nationally scarce (notable A, Falk, 1991a). This uncommon bee occurs in a variety of habitats, including woodland and fenland, but is mainly found in dry chalky areas, particularly in the Thames Estuary and Thames Valley (Edwards & Telfer, 2001) and Surrey (Baldock, 2008). It visits a variety of flowers after nectar and pollen and nests in the tough hollow stems of various dead plants such as dock and bramble. Several specimens were found visiting flowers, River Mole Corridor and Gatwick Brook Grassland, dated 27.5.2020 and 22.6.2020.



***Lasius brunneus* (Latreille)**

4.2.18 A small brown ant, family Formicidae. Status: nationally scarce (notable A, Falk, 1991a). This is a very local species restricted mainly to central and southern England from Essex to Shropshire. It seems to be centred on the Thames and Severn Valleys (Edwards, 1998), but appears to be spreading. It nests exclusively in dead wood (logs and standing timber) where it excavates its galleries, and it is particularly associated with ancient woodlands. Several were found crawling up tree trunks or under bark of dead trunks, Gatwick Brook Grassland, dated 22.6.2020, at TQ 28976 39926 and TQ 28874 39869.

***Magdalis armigera* (Geoffroy)**

4.2.19 A small black weevil, family Curculionidae. Status: very local. This widespread, but scarce species breeds in the twigs and branches of elm trees. It is currently increasing again, after becoming extremely scarce following the disappearance of elm trees from the landscape after the ravages of Dutch elm disease in the 1970s. One was beaten from small elm trees, Riverside Park, dated 27.5.2020, at TQ 27924 42151.

***Malthodes pumillus* (Brebisson)**

4.2.20 A minute soldier beetle, family Cantharidae. Status: very local. This beetle is usually found on herbage in old woodlands, and although recorded from numerous localities across much of the British Isles, it is very local and seldom seen. A single specimen was found by sweeping, Gatwick Brook Grassland, dated 22.6.2020, at TQ 29195 40057.

***Merzomyia* (formerly *Ictericia*) *westermanni* (Meigen)**

4.2.21 A medium-sized brown and orange picture-winged fly, family Tephritidae. Status: nationally scarce (notable, Falk, 1991b). This very local fly is known from an area south-east of a line from The Wash, to Gloucester to Weymouth. It breeds in the heads of ragwort, *Senecio* species, but despite the widespread abundance of its foodplant, it remains a scarce fly (Clemons, 1997, 2004, 2015). Several specimens were found by sweeping, River Mole Corridor, dated 10.9.2020, at TQ 26064 40625.

***Metopoplax ditomoides* (Costa)**

4.2.22 A small black and white ground bug, family Lygaeidae. Status: very local, but spreading. A single specimen of this bug was first found in Britain on a rubbish tip in Hounslow in 1953, after its spread had been monitored across Europe. Regarded as a vagrant or adventitious species, it was not included in the review of British Hemiptera (Kirby, 1992), but was rediscovered in Britain, in Oxfordshire, shortly after publication. It feeds on various species of mayweed, *Matricaria*. It has since been found on a number of occasions on brownfield sites in south-east England and appears to be spreading (Jones, 2008). Several specimens found by sweeping, River Mole Corridor, dated 17.5.2020 and 19.6.2020.

***Microlestes minutulus* (Goeze)**

4.2.23 A small black ground beetle, family Carabidae. Status: very local. This recent discovery in Britain was first found on the Suffolk coast in 1976, but was not recognized until 1995. It was later found in a few scattered coastal sites in Suffolk, Essex and Kent, usually in coastal litter (Luff, 1998). Since then it has continued to spread inland and is now known from numerous localities in southern and Eastern England. It seems to be associated with warm, well-drained soils with sparse vegetation. One specimen was found under rubble and broken bricks, Gatwick Brook Grassland, dated 14.9.2020, at TQ 29114 40000.

***Neottiglossa pusilla* (Gmelin)**

4.2.24 A tiny brown shieldbug, family Pentatomidae. Status: very local. This is a scarce species of rough grassy places in central and south-eastern England (Bantock, 2018). It is a secretive, ground-dwelling species, and easily overlooked. A single specimen was found by sweeping, Gatwick Brook

Grasslands, dated 22.6.2020, at TQ 29104 40030.

***Ophonus ardosiacus* (Luts)**

- 4.2.25 A medium-sized blue-black ground beetle, family Carabidae. Status: very local. Although given nationally scarce (notable B) status by Hyman & Parsons (1992), this was not confirmed by Telfer (2016) after recent records show it to be more widespread, even increasing in numbers and geographic range. This is still an uncommon species of southern England, south of the Severn/Wash line, and most localities are coastal or estuarine, with a large series of localities on the north Kent and South Essex coast of the Thames Estuary and London area (Luff, 1998). Several specimens were found under rubble and by finger-tip grubbing, Gatwick Brook Grassland, dated 27.5.2020, at TQ 29114 40000.

***Orchesia undulata* (Kraatz)**

- 4.2.26 A small mottled fungus beetle, family Melandryidae. Status: very local. A widespread, but rather local species found under the rotten bark of fungoid logs and trees. Originally listed as nationally scarce (notable B) by Hyman (1985), this was not confirmed by Hyman & Parsons (1992) or Alexander *et al.* (2014). A single specimen was found under the fungoid bark of a fallen/ cut tree trunk, Riverside Park, dated 14.9.2020, at TQ 28001 42089.

***Paraclusia tigrina* (Fallen)**

- 4.2.27 A small pink fly, family Clusiidae. Status: vulnerable (red data book category 2, Shirt, 1987, Falk, 1991b). This small fly is thought to breed in dead and decaying timber and is associated with woodlands and parklands. Since the review of Diptera was published (Falk, 1991b), there have been many more records of this species, suggesting that it is either increasing in abundance and range, or was previously overlooked. Its status probably needs to be reviewed. One specimen was seen running on a dead tree trunk, Riverside Park, dated 14.9.2020, at TQ 27716 42294.

***Podagrica fuscicornis* (Linnaeus)**

- 4.2.28 A small pink and blue leaf beetle, family Chrysomelidae. Status: nationally scarce (notable B, Hyman & Parsons, 1992; Hubble, 2014). A very local species, mainly of east and south-eastern England where it feeds on mallows, *Malva* species (Cox, 2007). Several specimens were swept from the foodplant, River Mole Corridor, dated 19.6.2020, at TQ 25987 40641.

***Polydrusus formosus* (Mayer)**

- 4.2.29 A small metallic green weevil, family Curculionidae. Status: nationally scarce (notable A, Hyman & Parsons, 1992). This very local weevil occurs on various broad-leaved trees, including hazel, oak, beech, apple and sallow in southern England. Until recently (about 2000) it was only recorded from Sussex, Hampshire and Kent, but has apparently started to spread and is now widely recorded in southern England, London and Thames Gateway area, with a scatter of records throughout much of England and Wales and outliers in Scotland. Its status may need revision. One was beaten from bushes, River Mole Corridor, dated 27.5.2020, at TQ 25678 40692.

***Reptalus quinquicostatus* formerly *Oliarus panzeri* (Low)**

- 4.2.30 A small brown plant hopper, family Cixiidae. Status: nationally scarce (notable, Kirby, 1992). This scarce bug has a very restricted south-eastern distribution and is thought to have declined dramatically in the last 50 years (Kirby, 1992). It has recently only been found in the extreme south-east, London, Thames Estuary, Sussex and Kent (Jones & Hodge, 1999). It seems to be associated with areas of rough ground, particularly where there are areas of bare soil, or where there is regular cracking in the ground during periods of drought. It may be a root-feeder during its nymph stage. A single specimen was found by sweeping, River Mole Corridor, dated 19.6.2020, at TQ 25650 40606.

***Rhinocyllus conicus* (Frohlich)**

- 4.2.31 A small mottled brown weevil, family Curculionidae. Status: nationally scarce (notable A, Hyman &

Parsons, 1992). Historically, this very scarce beetle was only known from a few scattered localities in south and south-west England, usually on disturbed ground. It was usually regarded as a coastal species, but appears to have been spreading in recent years, occurring at many inland sites right across central England. Its status may need revision. Several specimens were swept from thistles, River Mole Corridor, dated 27.5.2020 and 19.6.2020, at TQ 26064 40625.

***Sermylassa halensis* (Linnaeus)**

4.2.32 A small pink and green leaf beetle, family: Chrysomelidae. Status: very local. This beetle feeds on bedstraws, in rough grassy places such as verges, heathland, downs and chalk pits, usually in warm dry places (Cox, 2007). Two specimens were found by sweeping, River Mole Corridor, dated 10.9.2020, at TQ 25936 40624.

***Squamapion vicinum* (Kirby)**

4.2.33 A minute grey weevil, family Apionidae. Status: nationally scarce (notable B, Hyman & Parsons, 1992). This very local weevil breeds in the stems of water mint, *Mentha aquatica*, and although the food-plant is very common and widespread, the beetle seems very restricted. It is recorded widely, but sporadically, across England and Wales. Several specimens were found by sweeping water mint, River Mole Corridor, dated 27.5.2020, at TQ 25549 40781.

***Stictopleurus abutilon* (Rossi)**

4.2.34 A medium-sized brown leaf bug (family Rhopalidae). Status: extinct (Kirby, 1992), but now recolonized (Bantock, 2016). At the time of the national review of the British Hemiptera in 1992, this bug had only been found in the UK on a handful of occasions, the last being in 1948 and it was regarded as being extinct. During 1996 it was found in several localities in southern England and appeared to have successfully recolonized Britain. Since then it has been recorded on many occasions. Like the following species it has become a species typical of dry, well-drained and sparsely vegetated rough grassy places and brownfield sites in southern England, but remains relatively scarce and localized (Bantock, 2018). Several specimens were found by sweeping, River Mole Corridor, dated 30.6.2020.

***Stictopleurus punctatonervosus* (Goeze)**

4.2.35 A medium-sized brown leaf bug, family Rhopalidae. Status: extinct (Shirt, 1987, Kirby, 1992), but now recolonized and spreading across Britain (Bantock, 2016). At the time of the national review of British Hemiptera, this species was regarded as being extinct. It had been recorded from only two localities in Britain, the last in 1870. It appears to have successfully recolonized Britain since it was recorded in Essex in 1997. It has now become a species typical of the dry, well-drained and sparsely vegetated brownfield sites in and around urban London and the Thames Estuary (Jones, 2008) and is spreading widely across England (Bantock, 2018). Several specimens were found by sweeping, River Mole Corridor and Gatwick Brook Grasslands, dated 27.5.2020, 19.6.2020, 22.6.2020 and 10.9.2020.

***Tachys bistriatus* (Duftschmid)**

4.2.36 A minute brown ground beetle, family Carabidae. Status: nationally scarce (notable B, Hyman & Parsons, 1992; Telfer, 2016). This scarce beetle occurs on damp clay or sand by freshwater pools and ditches; it is more or less confined to southern England, and most records are from near the coast (Luff, 1998). Several specimens were found running about on the muddy edges of the river, River Mole Corridor, dated 19.6.2020, at TQ 25528 40714.

***Temnocerus (Rhynchites) nanus* (Paykull)**

4.2.37 A very small blue-black weevil, family Attelabidae. Status: very local. This scarce beetle breeds in the leaf buds of birch trees. Although recorded widely in England and Wales, records are scattered. One specimen was beaten from trees, Gatwick Brook Grassland, dated 27.5.2020, at TQ 29133 39913.

***Uleiota planata* (Linnaeus)**

4.2.38 A small flat bark beetle, family Cucujidae. Status: nationally scarce (notable A, Hyman & Parsons, 1992). This rare beetle occurs under the fungoid bark of broad-leaved trees, usually beech, elm, oak or birch, in ancient woodlands. It is listed in ancient woodland saproxylic fauna group 1 by Harding & Rose, 1986. Although recorded from Wales and Lancashire, most records are from central southern England: Hampshire, Surrey, Sussex and Berkshire. Several specimens were found under the bark of a large fallen/felled tree, Riverside Park, dated 14.6.2020, at TQ 28001 42089.

***Variimorda villosa* (Schrank)**

4.2.39 A small grey and black flower beetle, family Mordellidae. Status: nationally scarce (notable B, Hyman & Parsons, 1992; Alexander *et al.*, 2014). This scarce southern beetle is mostly found in Hampshire, Sussex and Kent. It is usually associated with old broadleaved woodland where it is thought to breed in dead fungoid wood, or wood mould, though it is most often found visiting flowers. Two specimens were found resting on hogweed flower heads, River Mole Corridor, dated 19.6.2020, at TQ 25937 40597.

## 5. Site Assessment

### 5.1 Discussion

5.1.1 The terrestrial invertebrate value of the three Gatwick sites, as discussed below, is mixed. Each has its own contribution to local biodiversity for the different habitat types they represent.

#### **Compartment P: Riverside Park**

5.1.2 Much of this linear habitat is heavily shaded, secondary woodland, although several scarce species were recorded associated with living trees, fallen logs and rotten tree stumps: *Magdalis armigera*, *Orchesia undulata*, *Uleiota planata* and *Paraclusia tigrina*. The most unusual insect recorded was the scarce lace-bug, *Catoplatus fabricii*, swept from the narrow overgrown verge of the busy A23 on the south western perimeter.

#### **Compartment M: River Mole Corridor**

5.1.3 This large river corridor area contained a variety of habitat types: brownfield/disturbed ground, river edge, rough grassland, woodland edge and scrub. Many of the scarce and unusual insects were recorded in this area, highlighting its diversity and ecological value. The most interesting species were: *Campiglossa malaris*, *Coccidula scutellata*, *Dioxyna bidentis*, *Dorycera graminum*, *Ectobius lapponicus*, *Hylaeus cornutus*, *Merzomyia westermanni*, *Reptalus quinquecostatus*, *Rhinocyllus conicus*, *Squamapion vicinum*, *Tachys bistratus* and *Variimorda villosa*. Several of these (*Coccidula*, *Dioxyna*, *Squamapion*, *Tachys*) are closely associated with the riverside habitat. Apart from the old woodland *Variimorda*, all the others are species of rough flowery, grassy places or disturbed ground.

#### **Compartment G: Gatwick Brook Grasslands**

5.1.4 This large rough grassland area belies the fact that it was recently re-profiled for surface water management and flood alleviation with ground levels significantly lowered, leaving the large mature trees (mostly oak) standing on tumulus-like hummocks. The unusual species found here — *Acinia corniculata*, *Camarota curvipennis*, *Hylaeus cornutus*, *Microlestes minutulus*, *Neottiglossa pusilla*, *Ophonus ardosiacus* — are mainly species of disturbed flowery and grassy land. However, *Lasius brunneus* is solely an old woodland species and must still occur on the mature trees in relic colonies, even though the land between the undisturbed tree mounds has been completely altered in approximately the last decade. *Temnocerus nanus* and *Malthodes pumillus* are mature hedgerow species.

5.1.5 The 303 species recorded across the combined Gatwick survey area is a relatively diverse overall assemblage. Considering the unusual and scarce species found, the biodiversity values of each of the three compartments is recommended as follows:

- Compartment P: Riverside Park - Low. It should be noted that the survey effort here focussed on the narrow densely flowering zone alongside the A23 and its environs and not the heavily shaded amenity park.
- Compartment M: River Mole Corridor - Medium/High.
- Compartment G: Gatwick Brook Grasslands - Low/Medium.

## 6. Conclusion

- 6.1.1 In total the three survey parcels associated with Gatwick Airport as part of this study, have provided a diverse and high value species list for the biodiversity of the area. All three parcels have varying diverse habitat types which provide for differing species and species groups. Numerous scarce, unusual and higher value species were recorded and should be taken into account during any development proposals.

## 7. References

- Alexander, K.N.A., Dodd, S. & Denton, J. 2014. *A review of the beetles of Great Britain. The darkling beetles and their allies. Species Status Number 18.* Natural England.
- Baldock, D.W. 1999. *Grasshoppers and crickets of Surrey.* Woking: Surrey Wildlife Trust.
- Baldock, D.W. 2008. *Bees of Surrey.* Woking: Surrey Wildlife Trust.
- Baldock, D.W. 2010. *Wasps of Surrey.* Woking: Surrey Wildlife Trust.
- Bantock, T.A. 2016. *A review of the Hemiptera of Great Britain: the shieldbugs and allied families.* Natural England/JNCC.
- Bantock, T.A. 2018. *Provisional atlas of shieldbugs and allies.* Terrestrial Heteroptera Recording Scheme.
- Clemons, L. 1997. A provisional atlas of the Tephritidae (Diptera) of Britain and Ireland. *British Tephritidae Newsletter* **6**: 1-42.
- Clemons, L. 2004. A provisional atlas of the Tephritidae (Diptera) of Britain and Ireland. Version 2. *British Tephritidae Newsletter* **8**: 1-25.
- Clemons, L. 2015. *The Tephritidae of Britain and Ireland.*
- Cox, M.L. 2007. *Atlas of the seed and leaf beetles of Britain and Ireland.* Newbury: Pisces Publications.
- Ecus Ltd. (2021). *Gatwick Airport Northern Runway Project – Aquatic Ecology Surveys Report.* Ecus Ltd, Basingstoke.
- Edwards, R. 1998. *Provisional atlas of the aculeate Hymenoptera of Britain and Ireland. Part 2.* Abbots Ripton: Biological Records Centre.
- Edwards, R. & Telfer, M. 2001. *Provisional atlas of the aculeate Hymenoptera of Britain and Ireland. Part 3.* Abbots Ripton: Biological Records Centre.
- Falk, S. 1991a. *A review of the scarce and threatened bees, wasps and ants of Great Britain.* Peterborough: Nature Conservancy Council.
- Falk, S. 1991b. *A review of the scarce and threatened flies of Great Britain (part 1).* Peterborough: Nature Conservancy Council.
- Haes, E.C.M. & Harding, P.T. 1997. *Atlas of grasshoppers, crickets and allies insects in Britain and Ireland.* London: The Stationery Office.
- Harding, P.T. and Rose, F. 1986. *Pasture-woodlands in lowland Britain: a review of their importance for wildlife conservation.* Abbots Ripton: Institute of Terrestrial Ecology.
- Hubble, D.S. 2014. *A review of the scarce and threatened beetles of Great Britain. The leaf beetles and their allies: Chrysomelidae, Megalopodidae, Orsodacnidae. Species Status Number 19.* JNCC/Natural England.
- Hyman, P.S. 1985. *A provisional review of the status of British Coleoptera. Invertebrate Site Register Report Number 60.* Peterborough: Nature Conservancy Council.
- Hyman, P.S. & Parsons, M.S. 1992. *A review of the scarce and threatened Coleoptera of Great Britain. Part 1.* Peterborough: Joint Nature Conservation Committee.
- Ismay, J.W. 2000. *The status, distribution and biology of Dorycera graminum (Fabricius) (Diptera, Ulidiidae).* English Nature Research Report Number 395.
- Kirby, P. 1992. *A review of the scarce and threatened Hemiptera of Great Britain.* Peterborough: Joint Nature Conservation Committee.

Luff, M.L. 1998. *Provisional atlas of the ground beetles (Coleoptera, Carabidae) of Britain*. Abbots Ripton: Biological Records Centre.

Majerus, M.E.N., Majerus, T.O., Bertrand, D. & Walker, L.E. 1997. *The geographic distribution of ladybirds (Coleoptera: Coccinellidae) in Britain (1984–1994)*. *Entomologist's Monthly Magazine* **133**: 181-203.

Mendel, H. & Clarke, R.E. 1996. *Provisional atlas of the click beetles (Coleoptera: Elateroidea) of Britain and Ireland*. Ipswich: Ipswich Borough Council Museums.

Roy, H., *et al.* 2011. *The ladybirds (Coccinellidae) of Britain and Ireland*. Biological Recording Centre.


Shirt, D.B. (ed.) 1987. *British red data books: 2. Insects*. Peterborough: Nature Conservancy Council.

Telfer, M. 2016. *A review of the beetles of Great Britain: ground beetles (Carabidae)*. Natural England Commissioned Report 189. Natural England.



## Figure 1. Site Locations (Three separate parcels - Gatwick Airport)

Image taken from MAGIC (<https://magic.defra.gov.uk/home.htm>)



## Appendix 1. Full Species List

Locality name:		Gatwick Airport (3 separate sites)	
Nominal National grid reference:		TQ 256406	
Vice-County		17, Surrey	
<b>Order &amp; Species</b>	<b>Status</b>	<b>Site Notes</b>	<b>Survey Date* &amp; Land Parcel (G = Gatwick Brook; P = Riverside Park; M = River Mole)</b>
			* month as Roman numeral
<b>COLEOPTERA, Beetles</b>			
<b>Ptinidae (formerly Anobiidae) woodworm beetles</b>			
Anobium (Hemicoelus) fulvicorne (Sturm)	common	Various dead timber	22.vi.2020 (G)
Ptilinus pectinicornis (Lin.)	local	On various dead timbers	27.v.2020 (P)
Stegobium paniceum (Lin.)	local	Indoors, in stored food products	
<b>Anthicidae, 'ant' beetles</b>			
Anthicus antherinus L.	local	Compost heaps, grass cuttings	14.ix.2020 (G)
<b>Apionidae, Minute weevils</b>			
Apion frumentarium (Lin.) (formerly A. miniatum)	local	On docks, Rumex species	27.v, 19.vi, 14.ix.2020 (G, M)

Aspidapion radiolus (Mars.)	common	On mallows, Malva species.	22.vi.2020 (P)
Ceratapion gibbirostre (Gyll.)	local	On thistles, Cirsium and Carduus	22.vi.2020 (G)
Eutrichapion ervi Kirby	common	On vetches, Viccia, grassy places	19.vi.2020 (M)
Exapion ulicis (For.)	common	On gorse, Ulex europeus, etc.	10.ix.2020 (M)
Nanophyes marmoratus (Goeze)	local	On Lythrum salicaria	30.vi, 10.ix.2020 (M)
Perapion curtirostre Germ.	common	On docks, Rumes species	27.v.2020 (M, P)
Squamapion vicinum (Kirby)	Nb	On water mint	27.v.2020 (M)
Trichapion simile (Kirby)	common	On birch	10.ix.2020 (M)
<b>Attelabidae, weevils</b>			
Temnocerus (Rhynchites) nanus (Payk.)	v. local	On birch, willow and alder.	27.v.2020 (G)
<b>Biphylidae, fungus beetles</b>			
Biphyllus lunatus (Fab.)	local	In Daldinia fungus on ash, ancient woodland indicator	30.vi.2020 (M)
<b>Cantharidae, Soldier beetles</b>			
Cantharis lateralis (Lin.)	local	Woods, larvae predatory in rotten wood	27.v, 19.vi, 22.vi.2020 (M, G)
Cantharis flavilabris (formerly nigra (Deg.))	local	Larvae predatory in rotten wood, soil, etc	27.v, 19.vi.2020 (G, M)
Cantharis nigricans Mull.	common	Larvae predatory in rotten wood, soil, etc	22.vi.2020 (G)
Cantharis pallida Goez	common	Larvae predatory in rotten wood, soil, etc	22.vi.2020 (G)

<i>Cantharis rustica</i> Fallen	common	Larvae predatory in rotten wood, soil, etc	19.vi.2020 (M)
<i>Malthodes marginatus</i> Latr.	local	Woods and meadows	22.vi.2020 (G)
<i>Malthodes minimus</i> (Lin.)	common	Woods, larvae predatory in rotten wood	22.vi.2020 (G)
<i>Malthodes pumilus</i> (Breb.)	v. local	Woods and grassland	22.vi.2020 (G)
<i>Rhagonycha fulva</i> (Scop.)	common	Adults on flowers, larvae predators in soil layer	27.v, 22.vi.2020 (M, G)
<i>Rhagonycha nigriventris</i> (formerly <i>limbata</i> )	local	Adults on flowers, larvae predators in soil layer	27.v.2020 (M)
<b>Carabidae, Ground beetles</b>			
<i>Abax parallelepipedus</i> (P. & M.)	common	Under logs, stones, etc, in woods	22.vi.2020 (G)
<i>Acupalpus dubius</i>	local	Damp grassy places	27.v.2020 (M, G)
<i>Amara plebeja</i> Gyll.	common	Various habitats	27.v.2020 (M)
<i>Bembidion articulatum</i> Panz.	local	River and stream banks	19.vi.2020 (M)
<i>Bembidion biguttatum</i> (Fab.)	local	Damp places, stream and pond sides	30.vi.2020 (M)
<i>Bembidion mannerheimi</i> Sahl.	local	Damp grasslands	27.v.2020 (M)
<i>Demetrias atricapillus</i> (Lin.)	common	Long grass	27.v.2020 (G)
<i>Paradromius linearis</i> Ol.	common	Dry grassy areas	27.v, 19.vi, 10.ix.2020 (M)
<i>Harpalus ardosiacus</i> Luts.	v. local	Chalk or limestone, usually coastal	27.v.2020 (G)
<i>Microlestes minutulus</i>	v. local	Dry sandy places, mostly East Anglia and London	14.ix.2020 (G)

Poecilus (Pterostichus) cupreus (Lin.)	common	Open fields, bare ground	27.v.2020 (M)
Poecilus (Pterostichus) versicolor Sturm	local	Open ground, bare soil.	27.v, 14.ix.2020 (G)
Tachys bistriatus (Dufts.)	Nb	On damp clay soils.	19.vi.2020 (M)
<b>Cerambycidae, Longhorn beetles</b>			
Pseudovadonia (Leptura) livida Fab.	local	Larvae in fungal hyphae in soil	19.vi.2020 (M)
Stenurella (Leptura) melanura (Lin.)	local	Larvae in dead timber or roots	19.vi, 22.vi.2020 (M, G)
Rutpela (Strangalia) maculata (Poda)	common	Larvae in dead wood of various trees	27.v.2020 (G)
<b>Chrysomelidae, Leaf and flea beetles</b>			
Bruchidius varius (Ol.)	local	On red clover, recent colonist	27.v.2020 (M)
Bruchidius imbricornis Panz.	v. local	New colonist in Britain	19.vi, 10.ix.2020 (M)
Bruchidius villosus (Fab.)	local	Rough grassy places	27.v.2020 (P)
Bruchus atomarius (Lin.)	local	Various habitats	27.v, 10.ix.2020 (M)
Bruchus loti Payk.	common	On Lotus corniculata	19.vi.2020 (M)
Cassida flaveola Thunb.	local	Dry grassy places	27.v.2020 (M)
Cassida rubiginosa Mull.	common	On thistles, Cirsium species.	22.vi.2020 (G)
Cassida vibex Fab.	local	On Centaurea species	27.v, 22.vi.2020 (G)
Cassida viridis (Lin.)	common	On water mint etc, Mentha species	19.vi.2020 (M)

<i>Crepidodera (Chalcoides) aurea</i> (Fourc.)	common	On willows, sallows, poplars etc	22.vi.2020 (P)
<i>Crepidopdera (Chalcoides) fulvicornis</i> (Fab.)	common	On willows, sallows, poplars etc	27.v.2020 (G)
<i>Chalcoides plutus</i> (Latr.)	local	On willows, sallows, poplars etc	19.vi, 10.ix.2020 (M)
<i>Neocrepidodera transversa</i> (Marsh.)	common	On thistles	22.vi.2020 (G)
<i>Cryptocephalus fulvus</i> Goeze	local	Dry grassy areas.	19.vi.2020 (M)
<i>Cryptocephalus pusillus</i> Fab.	local	On birch, sandy and chalky areas	14.ix, 19.vi.2020 (M, P)
<i>Galerucella lineola</i> Fab.	common	On alders, willows etc	19.vi.2020 (M)
<i>Gastrophysa viridula</i> Deg.	local	Wet meadows, marshes	19.vi, 22.vi.2020 (M, G)
<i>Lilioceris lili</i> (Scop.)	local	On cultivated lilies, gardens and parks	14.ix.2020 (P)
<i>Oulema obscura</i> Steph.	common	Meadows and fields	22.vi.2020 (G)
<i>Podagrica fuscicornis</i> (Lin.)	Nb	Grassland, rough ground, on mallows.	19.vi.2020 (M)
<i>Psylliodes chrysocephala</i> (Lin.)	common	On crucifers	27.v.2020 (P)
<i>Sermylassa halensis</i> (Lin.)	v. local	On bedstraws	10.ix.2020 (M)
<i>Sphaeroderma rubidum</i> (Gra.)	common	On thistles, <i>Cirsium</i> species	19.vi, 22.vi.2020 (M, G)
<i>Sphaeroderma testaceum</i> Fab.	common	On thistles, <i>Cirsium</i> species	19.vi, 14.ix.2020 (M, G)
<b>Coccinellidae, Ladybirds</b>			
<i>Anisosticta 19-punctata</i> (Lin.)	local	Water ladybird, near ponds	19.vi.2020 (M)

<i>Coccidula rufa</i> (Herbst)	local	Marshy places, reed and sedge beds	19.vi.2020 (M)
<i>Coccidula scutellata</i> (herbst)	v. local	Marshes, reed beds, stream sides	27.v.2020 (M)
<i>Coccinella 7-punctata</i> Lin.	common	7-spot. Wide variety of habitats.	27.v, 19.vi, 22.vi, 10.ix, 14.ix.2020 (G, M, P)
<i>Harmonia axyridis</i> Pallas	common	Recent arrival in Britain	27.v, 19.vi, 22.vi, 30.vi.2020 (M, G, P)
<i>Hippodamia variegata</i>	Nb	Adonis ladybird. Mainly coastal and London basin	10.ix.2020 (M)
<i>Micraspis 16-punctata</i> (Lin.)	common	16-spot, mildew feeder, grassy places	27.v, 19.vi, 22.vi, 10.ix, 14.ix.2020 (G, M)
<i>Propylea 14-punctata</i> (Lin.)	common	14-spot. Wide variety of habitats	19.vi, 22.vi, 10.ix, 14.ix.2020 (M, G, P)
<i>Psyllobora 22-punctata</i> (Lin.)	common	22-spot. Wide variety of habitats, mildew-feeder.	10.ix, 14.ix.2020 (M, P, G)
<i>Rhyzobius litura</i> (Fab.)	common	Rough grassy places	27.v, 14.ix.2020 (G, P)
<i>Subcoccinella 24-punctata</i> (Lin.)	common	24-spot. On false-oat grass	27.v, 19.vi, 10.ix, 14.ix.2020 (M, G)
<b>Colydiidae, fungus beetles</b>			
<i>Bitoma crenata</i>	local	Under fungoid bark of broadleaved trees	14.ix.2020 (P)
<b>Cryptophagidae, Fungus beetles, etc</b>			
<i>Micrambe ulicis</i> Steph.	local	General grassy places	27.v.2020 (G, P)
<b>Cucujidae (fungus beetles)</b>			
<i>Uleiota planata</i> (L.)	Na	Under fungoid bark	14.ix.2020 (P)
<b>Curculionidae (weevils)</b>			

<i>Ceutorhynchus obstrictus</i> Marsh. (assimilis)	common	On various crucifers	27.v.2020 (M)
<i>Ceutorhynchus pallidactylus</i> = <i>quadridens</i> (Pz.)	common	On aliaria and other crucifers, woods and hedges	27.v.2020 (M)
<i>Nedyus</i> ( <i>Cidnorhynchus</i> ) <i>4-maculatus</i> (L.)	common	On stinging nettles	22.vi.2020 (G)
<i>Cionus alauda</i> (Herbst)	common	On water figwort, <i>Scrophulria aquatica</i>	19.vi.2020 (M)
<i>Curculio glandium</i> (L.)	common	On oaks	27.v, 22.vi.2020 (P, G)
<i>Archarius</i> ( <i>Curculio</i> ) <i>pyrrhoceras</i> Mars.	local	On oaks	27.v, 22.vi.2020(P, G)
<i>Gymnetron pascuorum</i> Gyll.	common	On plantains, <i>Plantago</i> species	27.v, 19.vi, 22.vi.2020 (G, M, P)
<i>Hypera nigrirostris</i> 9Fab.)	common	Grassy places, on clovers	19.vi.2020 (M)
<i>Hypera rumicis</i> (Lin.)	common	Rough grassy places, on docks, <i>Rumex</i> species	19.vi.2020 (M)
<i>Magdalis armigera</i> (Geoff.)	v. local	Breeds in the twigs and branches of elm trees	27.v.2020 (P)
<i>Miccotrogus picirostris</i> (Fab.)	common	On clovers, <i>Trifolium</i> species	22.vi.2020 (P)
<i>Polydrusus formosus</i> ( <i>splendidus</i> ) (Mayer)	Na	On hazel, oak and other trees, spreading	27.v.2020 (M)
<i>Rhamphus pulicarius</i>	local	On sallow, birch, poplars, boggy places	19.vi.2020 (M)
<i>Rhinocyllus conicus</i> Froh.	Na	On thistles, southern England	27.v, 19.vi.2020 (M)
<i>R. pericarpus</i> (Lin.)	common	On <i>Rumex</i> species	22.vi, 10.ix.2020 (G, M)
<i>Sitona hispidulus</i> (Fab.)	common	On clovers and other legumes	10.ix.2020 (M)
<i>Sitona humeralis</i> Steph.	common	Dry grassy areas, on clover etc.	10.ix.2020 (M)



<i>Sitona lineatus</i> (Lin.)	common	On clovers, and many other legumes	19.vi, 10.ix.2020 (M)
<i>Trichosirocalus troglodytes</i> (Fab.)	common	On ribwort plantain, <i>Plantago lanceolata</i>	27.v, 19.vi, 14.ix.2020 (G, M)
<b>Dermestidae, Hide &amp; larder beetles</b>			
<i>Anthrenus verbasci</i> (Lin.)	common	Museum beetle. Indoors in kitchens, carpets, outdoors on flowers	22.vi.2020 (P)
<b>Elateridae, Click beetles</b>			
<i>Agriotes acuminatus</i> Steph.	common	Larvae in grass roots etc	22.vi.2020 (G)
<i>Agriotes obscurus</i> (Lin.)	common	Larvae in grass roots etc	19.vi.2020 (M)
<i>Athous campyloides</i> Newm.	Nb	Larvae in grass roots, rotten wood etc	27.v, 19.vi.2020 (G, M)
<b>Hydrophilidae, Water beetles</b>			
<i>Helophorus minutus</i> F.	common	Wet areas, ponds, streams, marshes	22.vi.2020 (G, M)
<b>Melandryidae, Fungus beetles</b>			
<i>Orchesia undulata</i> Kr.	v. local	Under rotten wood, in fungus	14.ix.2020 (P)
<b>Melyridae, False soldier beetles</b>			
<i>Axinotarsus marginalis</i> (Lap.)	common	Various habitats, larvae probably in rotten wood or soil.	19.vi, 22.vi.2020 (M, G)
<i>Malachius bipustulatus</i> (Lin.)	common	Open grassy areas, on flowers, larvae predatory	27.v, 19.vi, 22.vi.2020 (P, M, G)
<i>Malachius viridis</i> Fab.	common	Open grassy areas, on flowers, larvae predatory	27.v, 19.vi.2020 (G, M)

<b>Mordellidae, Flower beetles</b>			
<i>M. pumila</i> (Gyll.)	common	Flowery places	27.v.2020 (G)
<i>Variimorda villosa</i> Schr.	Nb	Broad-leaved woodland, larvae in dead wood.	19.vi.2020 (M)
<b>Nitidulidae, Pollen beetles</b>			
<i>Brachypterus glaber</i> (Steph.)	common	On stinging nettles	27.v.2020 (P)
<b>Oedemeridae, Flower beetles</b>			
<i>Oedemera lurida</i> (Marsh.)	common	On flowers, leaves, etc.	27.v, 19.vi, 22.vi, 30.vi.2020 (G, P, M)
<i>Oedemera nobilis</i> (Scopoli)	local	On flowers	27.v, 19.vi, 22.vi.2020 (G, P, M)
<b>Scarabaeidae, chafers and dung beetles</b>			
<i>Onthophagus coenobita</i> Herbst	local	In mammalian dung	19.vi.2020 (M)
<b>Scirtidae, marsh beetles</b>			
<i>Microcara testacea</i> (L).	local	Marshy places	27.v.2020 (M)
<b>Scraptiidae, Flower beetles</b>			
<i>Anaspis maculata</i> Fourc.	common	Adults on flowers, larvae in rotten wood	27.v, 22.vi.2020 (M, P)
<b>Silvanidae, fungus beetles</b>			
<i>Silvanus unidentatus</i> (Ol.)	local	Under fungoid bark	14.ix.2020 (P)
<b>Staphylinidae, Rove beetles</b>			

Anotylus rugosus (Fab.)	common	Damp grassy places	19.vi, 30.vi.2020 (M)
Platystethus cornutus/ degener	local	Muddy places, stream and pond banks	19.vi.2020 (M)
Stenus cicindeloides (Sch.)	local	Marshy places	19.vi.2020 (M)
Stenus junco Payk.	common	Rough grassy places	27.v.2020 (G)
<b>DERMAPTERA, Earwigs</b>			
<b>Forficulidae, Earwigs</b>			
Forficula auricularia L.	common	Variety of habitats, woods, gardens etc.	22.vi.2020 (G, P)
<b>DIPTERA, True flies</b>			
<b>Asilidae, robberflies</b>			
Dioctria atricapilla Meig.	local	Dry grasslands	27.v.2020 (M)
Dioctria baumhaueri Meigen	common	Grassy places, predatory	22.vi.2020 (G)
Leptogaster cylindrica (Deg.)	local	Grassy places in southern England	27.v, 19.vi, 22.vi, 30.vi.2020 (G, M)
<b>Chloropidae, fruit flies</b>			
Camarota curvipennis Lat.	v. local	Breeds in heads of wheat, rye, barley	14.ix.2020 (G)
<b>Clusiidae, 'druid' flies</b>			

Paraclusia tigrina Fallen	RDB2	Breeds in dead timber	14.ix.2020 (P)
<b>Conopidae, Thick-headed flies</b>			
Sicus ferrugineus (Lin.)	common	Parasitoid of various bumblebee species.	19.vi, 22.vi.2020 (M, G)
<b>Dolichopodidae, long-footed flies</b>			
Poecilobothrus nobilitatus Lin.	common	Associated with wet areas, mud, etc	19.vi.2020 (M)
Scellus notatus Fab.	local	Damp woods and meadows	27.v, 19.vi, 22.vi, 30.vi.2020 (M, G)
<b>Lauxaniidae, acalyprate flies</b>			
Peplomyza litura (Meig.)	common	Hedgerows, woodland edges etc	22.vi.2020 (G)
<b>Opomyzidae, minute flies</b>			
Geomyza tripunctata (Fall.)	common	Grassy places	22.vi.2020 (G)
Opomyza germinationis (Lin.)	common	Grassy places	22.vi, 10.ix, 14.ix.2020 (G)
<b>Sciomyzidae, snail-killing flies</b>			
Coremacera marginata (Fab.)	common	Biology unknown, probably snail parasitoid	27.v, 19.vi, 10.ix, 14.ix.2020 (M G)
Dichetophora obliterated (Fab.)	local	Biology unknown, probably snail parasitoid	10.ix.2020 (M)
Ilione albiseta (Scop.)	local	Attacks snails, moist places	19.vi, 30.vi.2020 (M)
Limnia unguicornis (Scop.)	local	Attacks snails, moist places	19.vi.2020 (M)
Pherbina coryleti (Scop.)	common	Attacks snails, moist places	10.vi, 10.ix.2020 (M)

<i>Sepedon spegea</i> (Fab.)	local	Parasitoid of snails	19.vi.2020 (M)
<i>Tetanocera elata</i> (Fab.)	common	Probably predator/parasitoid of land snails	19.vi.2020 (M)
<b>Stratiomyidae, Soldier flies</b>			
<i>Beris chalybata</i> (Fors.)	common	Larvae in decaying organic matter	27.v.2020 (P)
<i>Chloromyia formosa</i> (Scop.)	common	Larvae in dung and compost.	27.v, 19.vi.2020 (G)
<i>Chorisops tibialis</i> Meigen	common	Woodland edges, breeds in wood mould etc	22.vi.2020 (G)
<i>Pachygaster atra</i> (Panz.)	local	Larvae in fungus, soil, rotten wood	19.vi.2020 (M)
<i>Pachygaster leachii</i> (Curtis)	common	Larvae in fungus, soil, rotten wood	19.vi, 22.vi.2020 (M, G)
<b>Syrphidae, Hoverflies</b>			
<i>Baccha elongata</i> (Fab.)	common	Woodland and hedgerows	19.vi, 14.ix.2020 (M)
<i>Chrysotoxem bicinctum</i> (Lin.)	local	Grassland, hedgerows, woodland edges.	30.vi.2020 (M)
<i>Episyrphus balteatus</i> (Lin.)	common	Wide variety of habitats, gardens etc.	19.vi, 22.vi.2020 (M, G)
<i>Eristalinus sepulchralis</i> (Lin)	local	Larvae in rot holes in trees, ditches, ponds	10.ix.2020 (M)
<i>Eristalis arbustorum</i> (Lin.)	common	Larvae in rot holes in trees, ditches, ponds	19.vi, 22.vi, 30.vi.2020 (M, G)
<i>Eristalis nemorum</i> (Lin.)	local	Larvae in rot holes, ditches, stagnant ponds	10.ix.2020 (M)
<i>Eristalis pertinax</i> (Scop.)	common	Larvae in rot holes, ditches, stagnant ponds	19.vi.2020 (M)
<i>Eristalis tenax</i> (Lin.)	common	Larvae in rot holes in trees, ditches, ponds	27.v, 19.vi, 22.vi.2020 (M, G)

<i>Eupeodes luniger</i> (Meig.)	common	Wide variety of habitats, gardens.	22.vi.2020 (G)
<i>Helophilus pendulus</i> (Lin.)	common	Breeds in ditches and stagnant ponds.	27.v, 19.vi, 22.vi, 10.ix, 14.ix.2020 (G, M)
<i>Myathropa florea</i> (Lin.)	common	Larvae in rot holes in trees	27.v, 19.vi.2020 (G)
<i>Pipizella viduata</i> (Lin.)	local	Dry grassland, chalk and coastal	19.vi.2020 (M)
<i>Platycheirus albimanus</i> Fab.	common	Woods and fields	22.vi.2020 (G)
<i>Rhingia campestris</i> (Meig.)	common	Woodlands, hedgerows, in cow and horse dung	10.ix.2020 (M)
<i>Scaeva pyrastris</i> (Lin.)	common	Wide variety of grassy habitats	22.vi.2020 (G)
<i>Sphaerophoria scripta</i> (Lin.)	common	Wide variety of grassy habitats	19.vi, 22.vi, 30.vi, 10.ix.2020 (M, G)
<i>Syrirta pipiens</i> (Lin.)	common	Wide variety of habitats, gardens etc.	27.v, 19.vi, 22.vi.2020 (M, G)
<i>Volucella pellucens</i> (Lin.)	common	Variety of habitats, breeds in wasp nests	22.vi.2020 (P)
<i>Xylota sylvarum</i> (Lin.)	local	Woodlands	22.vi.2020 (G)
<b>Tabanidae, horseflies</b>			
<i>Haematopota pluvialis</i> (Lin.)	local	Adults suck blood, wet meadows	19.vi.2020 (M)
<b>Tachinidae, parasitic flies</b>			
<i>Eriothrix rufomaculata</i> (Deg.)	common	Hosts unknown, even though fairly common	22.vi.2020 (G)
<i>Exorista</i> species	—	Several species, impossible to separate females	19.vi.2020 (M)
<i>Lydella grisescens</i> R.-D.	common	Parasitoid of various moth caterpillars	10.ix.2020 (M)

<i>Phasia obesa</i> (Fab.)	local	Parasitoid of bugs	27.v, 22.vi, 10.ix.2020 (P, G, M)
<i>Phasia pusilla</i> Meig.	local	Parasitoid of shieldbugs	19.vi, 10.ix.2020 (M)
<i>Siphona geniculata</i> Degeer	common	Parasitoid of various insect larvae	19.vi, 22.vi.2020 (M, G)
<i>Tachina fera</i> (Lin.)	common	Parasitoid of various common moth caterpillars	10.ix.2020 (M)
<b>Tephritidae, picture-winged flies</b>			
<i>Acinia corniculata</i> (Zett.)	RDB1	Breed in heads of <i>Centaurea nigra</i>	22.vi.2020 (G)
<i>Campiglossa malaris</i> Seguy	RDB1	Larvae in heads of ragworts, spreading	19.vi.2020 (M)
<i>Chaetorellia jaceae</i> (R.-D.)	local	In heads of <i>Centaurea</i>	27.v, 19.vi, 22.vi.2020 (M, G, P)
<i>Chaetostomella cylindrica</i> (R.-D.)	local	Breeds in heads of <i>Centaurea</i> etc	22.vi.2020 (G)
<i>Dioxya bidentis</i>	N	Breeds in flower heads of <i>Bidens tripartita</i>	10.ix.2020 (M)
<i>Merzomyia (Icteric) westermanni</i> (Meig.)	N	In heads of ragwort	10.ix.2020 (M)
<i>Tephritis formosa</i> (Loew)	local	Larvae in the heads of <i>Sonchus</i> species	27.v.2020 (M)
<i>Terellia colon</i> (Meig.)	local	In heads of <i>Centaurea scabiosa</i>	22.vi.2020 (G)
<i>Terellia ruficauda</i> (Fab.)	common	Larvae in heads of <i>Cirsium arvense</i>	19.vi, 22.vi, 30.vi.2020 (M, G)
<i>Terellia serratulae</i> (Lin.)	local	Breeds in heads of thistles	19.vi.2020 (M)
<i>Urophora cardui</i> (Lin.)	common	Larvae in galls in stems of <i>Cirsium arvense</i>	27.v.2020 (M)
<i>Urophora jaceana</i> (Her.)	common	Larvae in galls in knapweed heads	19.vi, 19.vi.2020 (M, P)

<i>Urophora quadrifasciata</i> (Meig.)	local	Larvae in galls in knapweed heads	27.v, 19.vi, 22.vi.2020 (M, G)
<i>Urophora stylata</i> (Fab.)	common	Larvae in heads of <i>Cirsium arvense</i>	19.vi, 22.vi.2020 (M, G)
<b>Ulidiidae, picture-winged flies</b>			
<i>Dorycera graminum</i> (Fab.)	RDB3	Rough meadows, southern England, Thames Estuary	27.v.2020 (M)
<b>HEMIPTERA, True bugs</b>			
<b>Anthocoridae, flower bugs</b>			
<i>Orius niger</i> (Woolf)	common	Predatory on small insects, on flowers and grass	14.ix.2020 (P)
<b>Berytinidae, stilt bugs or thread bugs</b>			
<i>Cymus melanocephalus</i> Fieb.	local	Marshy places	27.v, 10.ix.2020 (G, M)
<b>Cercopidae, Frog hoppers</b>			
<i>Aphrophora alni</i> (Fallen)	common	On willows, sallows, etc.	22.vi.2020 (G)
<i>Philaenus spumarius</i> (Lin.)	common	Nymphs on various herbs, variety of habitats	19.vi, 22.vi, 10.ix, 14.ix.2020 (M, G, P)
<b>Cicadellidae, leafhoppers</b>			
<i>Aphrodes bicinctus</i> (Schr.)	common	Various grassy habitats	22.vi.2020 (G)
<i>Cicadella viridis</i> (Lin.)	local	Damp grassy places	10.ix.2020 (M)
<i>Eupterycyba jucunda</i> (H.-S.)	Local	On alder	22.vi.2020 (G)



Eupteryx aurata L.	common	On stinging nettles	14.ix.2020 (P)
Iassus scutellaris (Fieb.)	local	On elms	14.ix.2020 (P)
<b>Cixiidae, froghoppers</b>			
Cixius cunicularius (L.)	Local	Rough grassy places	10.ix.2020 (M)
Reptalus quinquicostatus (formerly Oliarus) panzeri Low	N	Dry grassy places	19.vi.2020 (M)
<b>Coreidae, Leaf bugs</b>			
Coreus marginatus (Lin.)	common	Woods, meadows, gardens, on docks, Rumex species.	27.v, 19.vi, 22.vi, 10.ix, 14.ix.2020 (P, M, G)
Coriomeris denticulatus (Scop.)	local	On medicks, trefoils and melilots, dry areas.	27.v, 14.ix.2020 (G, M)
<b>Delphacidae, Ground hoppers</b>			
Allygus mixtus (Fab.)	common	Usually on trees	22.vi.2020 (G)
Ditropis pteridis (Spin.)	common	On bracken, Pteridium aquilinum	22.vi.2020 (G)
Stenocranus minutus (Fab.)	common	Grassy places	10.ix.2020 (M)
<b>Lygaeidae, Ground bugs</b>			
Heterogaster urticae (Fab.)	common	On stinging nettles	27.v, 19.vi, 22.vi, 14.ix.2020 (M, G, P)
Ischnodemus sabuleti (Fall.)	common	Grassy places, meadows and marshes	27.v, 19.vi, 10.ix.2020 (M)
Kleidocerys resedae (Panz.)	common	On wide variety of trees and shrubs	14.ix.2020 (P)

<i>Metopoplax ditomoides</i> (Costa)	v. local	On chamomile and scentless mayweed	27.v, 19.vi.2020 (M)
<i>Nysius senecionis</i> (Schill.)	local	On Guernsey fleabane and ragworts	10.ix.2020 (M)
<i>Peritrechus geniculatus</i> (Hahn)	common	In leaf litter and grass roots	27.v, 14.ix.2020 (G, M)
<i>Scolopostethus thomsoni</i> Reut.	common	Under stones, bare ground, sparse vegetation	19.vi.2020 (M)
<b>Miridae, Leaf bugs</b>			
<i>Capsus ater</i> (Lin.)	common	Various habitats on various plants	27.v, 19.vi, 22.vi.2020 (G, M)
<i>Charagochilus gyllenhali</i> (Fall.)	common	Dry places, on bedstraws	30.vi, 10.ix.2020 (M)
<i>Deraeocoris flavilinea</i> (Costa)	local	On maples and sycamores	19.vi, 22.vi.2020 (M, G, P)
<i>Deraeocoris ruber</i> (Lin.)	common	On stinging nettles and various other plants	22.vi.2020 (G)
<i>Deraeocoris lutescens</i>	common	On various low plants and trees	22.vi, 10.ix.2020 (G, P)
<i>Dryophilocoris flavoquadrimaculatus</i>	common	On oaks	27.v.2020 (G, P)
<i>Heterotoma planicornis</i> (Fab.)	common	On stinging nettles	19.vi, 22.vi.2020 (M, G)
<i>Leptopterna dolobrata</i> (Lin.)	common	Grassy places	19.vi, 22.vi.2020 (M, G, P)
<i>Notostira elongata</i> (Geoff.)	common	Grassy places	14.ix.2020 (G)
<i>Orthops campestris</i> (L.)	common	On umbellifers and other plants	30.vi.2020 (M)
<i>Phytocoris varipes</i> Boh.	common	Grassy places	10.ix.2020 (M)
<i>Pithanus maerkeli</i> (H.-S.)	common	Grassy places	19.vi.2020 (M)

<b>Nabidae, Damsel bugs</b>			
Nabis flavomarginatus Sch.	common	Rough grassland, damp areas	19.vi.2020 (M)
Nabis rugosus (Lin.)	common	Grassy areas, bare ground, predatory.	19.vi, 10.ix.2020 (M)
<b>Pentatomidae, Shield bugs</b>			
Aelia acuminata (Lin.)	local	Various grassy habitats	27.v, 19.vi, 22.vi, 10.ix, 14.ix.2020 (G, M)
Dolycoris baccarum (Lin.)	local	Woodland edges and hedges, on variety of plants	27.v, 19.vi, 22.vi, 30.vi, 10.ix, 14.ix.2020 (G, M)
Eurydema oleracea (Lin.)	common	On wild and garden brassicas and other crucifers	19.vi, 30.vi.2020 (M)
Neottiglossa pusilla (Gmel.)	v.local	Grassy places	22.vi.2020 (G)
Palomena prasina (Lin.)	common	On a variety of plants	22.vi, 10.ix, 14.ix.2020 (G, P, M)
Pentatoma rufipes Lin.	common	On a variety of trees, mainly oak	27.v, 22.vi.2020 (P, G)
<b>Rhopalidae, Leaf bugs</b>			
Myrmus miriformis (Fall.)	local	Grassy places, chalk, sand and marshes	19.vi, 10.ix.2020 (M)
Stictopleurus abutilon (Rossi)	v. local	Open, sunny localities	30.vi.2020 (M)
Stictopleurus punctatonervosus	v. local	Open, sunny localities	27.v, 19.vi, 22.vi, 10.ix.2020 (M, G)
<b>Scutelleridae, tortoise bugs</b>			
Eurygaster testudinaria (Geoff.)	local	Grassy and marshy places	27.v, 19.vi.2020 (M)
<b>Tingidae, Lace bugs</b>			

Catoplatus fabricii	Nb	On ox-eye daisy	27.v.2020 (P)
<b>HYMENOPTERA, Bees, wasps, etc</b>			
<b>Andrenidae, solitary bees</b>			
Andrena bimaculata (Kirby)	Local	Various habitats	27.v.2020 (M)
<b>Anthophoridae, solitary bees</b>			
Nomada flavoguttata Kirby	common	Cleptoparasite of Andrena species	22.v.2020 (G)
<b>Apidae, bees</b>			
Bombus lapidarius (Lin.)	common	Wide variety of habitats	19.vi, 10.ix.2020 (M)
Bombus pascuorum (Scop.)	common	Wide variety of habitats	10.ix.2020 (M)
Bombus terrestris L.	common	Wide variety of habitats	19.vi.2020 (M)
<b>Bethylidae, aculeate wasps</b>			
Bethylus cephalotes (Forster)	local	Parasitoid of moth caterpillars	22.vi, 14.ix.2020 (G)
Bethylus fuscicornis (Jurine)	Local	Parasitoid of moth caterpillars	19.vi.2020 (M)
<b>Chrysididae, rubytails</b>			
Omalus auratus (Lin.)	common	Woods, gardens, parks, parasitoid of solitary wasps in dead stems	22.v.2020 (G)

<b>Colletidae, solitary bees</b>			
<i>Hylaeus cornutus</i> Curt.	Na	Woodland edges, chalky and sandy places	27.v, 22.vi.2020 (M, G)
<b>Cynipidae, gall wasps</b>			
<i>Andricus kollari</i>	common	Makes marble galls on oak twigs	10.ix.2020 (M)
<i>Diplolepis rosae</i> (L.)	common	Robin pin cushion or bedeguar galls on wild roses	19.vi, 10.ix.2020 (M)
<i>Neuroterus numismalis</i> (Geoff.)	common	silk button galls on oak leaves	14.ix.2020 (P)
<i>Neuroterus quercusbaccarum</i> (Sch.)	common	spangle galls on oak leaves	14.ix.2020 (P)
<b>Eumenidae, Potter wasps</b>			
<i>Ancistrocerus parietum</i> (Lin.)	common	Builds mud nest in cavities in walls, tree trunks, rocks etc	19.vi, 22.vi.2020 (M, G)
<b>Formicidae, Ants</b>			
<i>Formica fusca</i> Lin.	common	England and Wales, widespread	19.vi, 22.vi.2020 (M, G)
<i>Lasius brunneus</i> (Latr.)	Na	Central England, Severn Valley, local, spreading	22.vi.2020 (G)
<i>Lasius niger</i> Lin.	common	Ubiquitous	22.vi.2020 (P)
<i>Myrmica rubra</i> (Lin.)	common	Various habitats, nests under stones, logs, etc.	27.v.2020 (G, M)
<b>Ichneumonidae, ichneumon wasp</b>			
<i>Amblyteles armatorius</i> (Fab.)	common	Parasitoid of moth caterpillars	22.vi.2020 (G)
<b>Melittidae, solitary bees</b>			

Lasioglossum morio (F.)	common	Various localities, on flowers	19.vi.2020 (M)
Lasioglossum fum (Schenck)	local	sandy of dry soils, flowery places	19.vi.2020 (M)
<b>Sphecidae, Solitary wasps</b>			
Pemphredon inornata (Say)	common	Nests in hollow stems.	27.v.2020 (M)
<b>Tenthredinidae, Sawflies</b>			
Rhogogaster viridis (L.)	common	Larvae on alder	27.v.2020 (P)
<b>Vespidae, social wasps</b>			
Vespa crabro L.	local	Hornet. Woodlands and parks	10.ix.2020 (M)
<b>LEPIDOPTERA, Butterflies &amp; moths</b>			
<b>Erebidae, tiger moths, etc</b>			
Tyria jacobaeae (Lin.)	common	Cinnabar moth, caterpillars on ragwort	27.v, 19.vi, 22.vi.2020 (G, M)
<b>Hesperidae, skippers</b>			
Ochlodes venata (L.)	common	Large skipper, grassy places	27.v, 19.vi, 22.vi.2020 (G, M)
Thymelicus lineola (Ochs.)	common	Essex skipper, grassy places, larvae on grasses	19.vi.2020 (M)
Thymelicus sylvestris (Poda)	common	Small skipper. Grassy places, larvae on various grasses	19.vi, 22.vi.2020 (M, G)
<b>Lycaenidae, Blues</b>			

Lycaena phlaeas (L.)	common	Small copper, larvae on trefoils and medics	14.ix.2020 (G)
Polyommatus icarus Rott.	common	Common blue. Grassy places, larvae of trefoils, clovers and medicks.	14.ix.2020 (G)
<b>Lymantridae, tussock moths</b>			
Orygia antiqua (Lin.)	common	Vapourer, larvae on wide variety of plants	27.v.2020 (M)
<b>Noctuidae, moths</b>			
Acronicta rumicis	common	Knotgrass moth, larvae on various food plants	22.vi.2020 (G)
<b>Notodontidae, prominent moths</b>			
Cerura vinula	common	Puss moth, on willows and poplars. Cocoon.	30.vi.2020 (M)
<b>Nymphalidae, Brush-footed butterflies</b>			
Aglais urticae (Lin.)	common	Small tortoiseshell. Larvae on stinging nettles.	22.vi.2020 (G)
Aphantopus hyperanthus (L.)	local	Ringlet, woods and woodland edges	30.vi.2020 (M)
Coenonympha pamphilus (L.)	local	Small heath, larvae on grasses	27.v, 19.vi, 22.vi, 10.ix, 14.ix.2020 (G, M)
Inachis io (Lin.)	common	Peacock, larvae on stinging nettles	27.v, 22.vi.2020 (M, G)
Maniola jurtina (Lin.)	common	Meadow brown. Grassy places, on various grasses.	19.vi, 22.vi, 30.vi, 10.ix.2020 (M, G)
Melanargia galathea (Lin.)	Local	Marbled white. Chalk and limestone downs	19.vi.2020 (M)
Pararge aegeria (Lin.)	common	Speckled wood. Woodland edges and rides, larvae on grasses	27.v, 22.vi, 14.ix.2020 (G, P)

Polygonia c-album (Lin.)	common	Comma, larvae on stinging nettles	27.v, 22.vi.2020 (M, G, P)
Pyronia tithonus (L.)	common	Gatekeeper, hedges and grassy places	22.vi.2020 (G)
Vanessa atalanta (Lin.)	common	Red admiral. Larvae on stinging nettles. Migrant.	19.vi.2020 (M)
<b>Pieridae, cabbage whites</b>			
Pieris brassicae (Lin.)	common	Large white, on brassicas, wild and garden species	22.vi, 14.ix.2020 (P)
Pieris napi (Lin.)	common	Green-veined white, on brassicas	10.ix.2020 (M)
<b>Tortricidae, micromoths</b>			
Tortrix viridana (L.)	common	Green tortrix, on oak, larvae can defoliate trees	27.v.2020 (P)
<b>Zygaenidae, burnets</b>			
Zygaena filipendulae (L.)	common	6-spot burnet, dry grassy places	22.vi.2020 (G)
<b>DICTYOPTERA, Cockroaches</b>			
<b>Blattellidae, Cockroaches</b>			
Ectobius lapponicus (L.)	Nb	Southern, heathland, sand or chalk soils	27.v.2020 (M)
<b>ODONATA, Dragonflies</b>			
<b>Aeshnidae, hawkers</b>			



Aeshna juncea (L.)	common	Ponds, streams, rivers, lakes	10.ix.2020 (M)
Anax imperator Leach	local	Ponds, lakes and canals	27.v, 19.vi.2020 (G, M)
Calopteryx splendens (Har.)	common	Slow-moving streams and ditches	27.v, 22.vi.2020 (M, G)
<b>Cordulegastridae, hawkers</b>			
Cordulegaster boltoni (Don.)	local	Heaths and moors, mostly western in UK	10.ix.2020 (M)
<b>Coenagrionidae, Damselflies</b>			
Enallagma cyathigerum (Ch.)	common	Ponds, streams and lakes.	19.vi, 22.vi.2020 (M, G)
<b>Libellulidae, darters</b>			
Libellula depressa (Lin.)	common	Lakes and ponds	27.v.2020 (M)
Sympetrum striolatum (Charp.)	common	Ponds, lakes and streams	27.v, 10.ix.2020 (M)
<b>ORTHOPTERA, Grasshoppers</b>			
<b>Acrididae, grasshoppers</b>			
Chorthippus brunneus (Thunb.)	common	Wide variety of grassy habitats.	10.ix.2020 (M)
Chorthippus parallelus (Zett.)	common	Wide variety of grassy habitats.	19.vi, 22.vi, 30.vi, 10.ix.2020 (M, G)
<b>Conocephalidae, coneheads</b>			
Conocephalus fuscus (Fab.) = discolor (Thunb.)	Local	Grassy places, spreading recently	19.vi, 10.ix.2020 (M)

<b>Tettigoniidae, bush crickets</b>			
Leptophyes punctatissima (Bos.)	common	Various habitats, woodlands and gardens	19.vi, 22.vi.2020 (M, G)
Meconema thalassinum Deg.	common	On trees and shrubs	22.vi.2020 (G)
Metrioptera roeselii (Hag.)	local	Dry grassy places, Essex, Kent, London spreading west	27.v, 19.vi.2020 (M)
<b>ARANAEA, Spiders</b>			
<b>Araneidae, orb-web spiders</b>			
Agalenatea redii (Scop.)	local	Fields and meadows	10.ix.2020 (M)
<b>Pisauridae, nursery web spiders</b>			
Pisaura mirabilis (Clerck)	common	Wide variety of habitats	22.vi, 10.ix.2020 (G)
<b>Thomisidae, crab spiders</b>			
Misumena vatia (Cl.)	local	Southern England, on flowers	19.vi.2020 (M)
<b>OPILIONES, Harvestmen</b>			
<b>Leiobunidae, harvestmen</b>			
Dicranopalpus ramosus (Sim.)	local	On trees	14.ix.2020 (G)

<b>ISOPODA, Woodlice and hoglice</b>			
<b>Armadillidiidae, pill woodlice</b>			
Armadillidium vulgare (Latr.)	common	Under logs and stones etc, mainly dry places	27.v, 19.vi, 30.vi.2020 (G, M)
<b>Philosciidae, striped woodlice</b>			
Philoscia muscorum (Scop.)	common	Under logs, stones, leaf litter etc	27.v, 22.vi, 30.vi.2020 (G, M)
<b>Porcellionidae, Rough woodlice</b>			
Porcellio scaber (Latr.)	vc	Under logs, stones, leaf litter etc	27.v, 22.vi.2020 (G)
<b>MOLLUSCA, Slugs and snails</b>			
<b>Helicidae, snails</b>			
Cornu aspersum (formerly Helix aspersa)	common	Gardens, parks, fields and woods	14.ix.2020 (G)
Monacha cantiana	common	Various roughly vegetated habitats	22.vi.2020 (G)

Annex 6

Aquatic Ecology Survey



# **Gatwick Airport Northern Runway Project –**

## **Aquatic Ecology Surveys Report**

**RPS Group PLC**

*Report prepared by:*  
Ecus Ltd.  
Unit 1  
Woodlands Business Village  
Coronation Road  
Basingstoke  
RG21 4JX

01256 224588


**June 2021**


## Ecus Ltd


Report to: **RPS Group PLC**

Report Title: **Gatwick Airport Northern Runway Project – Fish and Aquatic Macroinvertebrate Surveys**

Version: **1.0**  
Issue Date: **June 2021**  
Report Ref: **14864**

Originated By:   
**Claire Evans**  
**Consultant Ecologist** Date: 25/05/2021

Reviewed By:   
**Abel Drewett**  
**Technical Director**  
**(Terrestrial Ecology)** Date: 16/06/2021

Approved By:   
**Paul White**  
**Regional Director** Date: 21/06/2021

Version	Author	Description	Date
0.1	CE	First Draft	25/05/2021
0.2	AD	QA1 Review	16/06/2021
0.3	AD	Edits	17/06/2021
1.0	PAW	Approve and issue to client	21/06/2021

The report and the site assessments carried out by Ecus on behalf of the client in accordance with the agreed terms of contract and/or written agreement form the agreed Services. The Services were performed by Ecus with the skill and care ordinarily exercised by a reasonable Environmental Consultant at the time the Services were performed. Further, and in particular, the Services were performed by Ecus taking into account the limits of the scope of works required by the client, the time scale involved and the resources, including financial and manpower resources, agreed between Ecus and the client.

Other than that expressly contained in the paragraph above, Ecus provides no other representation or warranty whether express or implied, in relation to the services.

This report is produced exclusively for the purposes of the client. Ecus is not aware of any interest of or reliance by any party other than the client in or on the services. Unless expressly provided in writing, Ecus does not authorise, consent or condone any party other than the client relying upon the services provided. Any reliance on the services or any part of the services by any party other than the client is made wholly at that party's own and sole risk and Ecus disclaims any liability to such parties.

This report is based on site conditions, regulatory or other legal provisions, technology or economic conditions at the time of the Service provision. These conditions can change with time and reliance on the findings of the Services under changing conditions should be reviewed.

Ecus accepts no responsibility for the accuracy of third party data used in this report.

## Contents

<b>SUMMARY AND KEY RECOMMENDATIONS.....</b>	<b>1</b>
SUMMARY.....	1
CONCLUSIONS.....	2
KEY RECOMMENDATIONS.....	2
<b>1. INTRODUCTION.....</b>	<b>4</b>
1.1 DEVELOPMENT BACKGROUND.....	4
1.2 THE OBJECTIVES.....	4
1.3 BACKGROUND TO THE WATERCOURSES.....	4
<b>2. METHODOLOGY.....</b>	<b>5</b>
2.1 AQUATICS DESK STUDY.....	5
2.2 SURVEY: AQUATIC MACROINVERTEBRATES.....	5
2.3 AQUATIC MACROINVERTEBRATE DATA ANALYSIS.....	6
2.4 SURVEY: FISH.....	7
2.5 LIMITATIONS.....	8
<b>3. RESULTS.....</b>	<b>9</b>
3.1 AQUATICS DESK STUDY.....	9
3.2 FIELD DATA.....	12
<b>4. DISCUSSION.....</b>	<b>18</b>
4.1 RIVER MOLE.....	18
4.2 GATWICK STREAM.....	19
<b>5. CONCLUSIONS AND RECOMMENDATIONS.....</b>	<b>21</b>
5.1 CONCLUSIONS.....	21
5.2 RECOMMENDATIONS.....	21
5.3 FURTHER SURVEY.....	22
<b>6. REFERENCES.....</b>	<b>23</b>
<b>APPENDIX 1: AQUATICS TEAM REPORT WITH FIGURES.....</b>	<b>24</b>

## Summary and Key Recommendations

### Summary

ECUS Ltd was commissioned by RPS Group Plc in May 2020 to undertake fish and aquatic invertebrate surveys of two water courses adjoining Gatwick Airport as part of the surface water management and flood alleviation for the Northern Runway Project development. This will affect the two sites close to Gatwick Airport, one on the River Mole and the other on the Gatwick Stream (Gatwick Brook). The River Mole may be re-meandered and land close to the river may be re-profiled to increase flood storage. The Gatwick Stream has already had surrounding land re-profiled for flood storage in recent years, however this area may be expanded to encompass both sides of the river. The aquatics team at Thomson Environmental were commissioned to assist ECUS in undertaking aquatic macroinvertebrate and fish baseline surveys and a targeted aquatics desk study, to inform this report and the proposals.

The study area encompasses two watercourses; a 1.3km stretch of the River Mole immediately downstream of where it emerges from under the runway at Gatwick Airport and a 750m stretch of the Gatwick Stream (a tributary of the River Mole) upstream of the Crawley sewage works. A 100m survey section was identified on each watercourse from an initial walkover survey conducted in June 2020. Three survey visits during 2020 were undertaken for aquatic macroinvertebrates (spring, summer and autumn) and two for fish (spring and autumn). The spring survey visit was delayed until early July due to restrictions related to the Covid 19 outbreak. Desk study data was obtained from the Sussex Biological Records Centre and the Environment Agency on behalf of Ecus.

The desk study returned one record from 2013 of the shining ram's-horn snail *Segmentina nitida* within the study section on the River Mole (TQ 25623 40908). The species is nationally scarce, a UK Priority Species under the UK Post 2010 Biodiversity Framework and listed on the Sussex Rare Species Inventory. It was not recorded during the surveys for this study, although the survey section did not coincide with the reach in which the snail was recorded. The desk study returned records of two fish species for the River Mole; bullhead *Cottus gobio* and brown trout *Salmo trutta subsp. fario*.

**River Mole** - A mean of 19.3 macroinvertebrate taxa were recorded at the River Mole site across the three survey visits. Biotic indices measuring water quality (Biological Monitoring Working Party (BMWP) score and Average Score per Taxon) indicate moderately polluted conditions in the River Mole. Lotic invertebrate Index for Flow Evaluation (LIFE) scores suggest that the aquatic macroinvertebrate community is characteristic of sluggish flow conditions and low Proportion of Sediment intolerant Invertebrates (PSI) scores indicate heavily sedimented conditions.

A total of 415 fish were caught on the River Mole in spring after three survey 'runs' compared with only 28 fish caught in autumn with the same level of effort. Roach *Rutilus rutilus* were the most abundant fish species identified (252) in spring and in autumn (13). The study stretch on the River Mole lies within open floodplain grassland with no shading which means that water temperature, and therefore dissolved oxygen (DO), fluctuated considerably. Extensive stands of submerged and emergent macrophyte plants occur through the study section and their decomposition are likely to be contributing to low DO in the autumn. These dissolved oxygen conditions coupled with organic pollution from within the catchment is considered to be influencing the composition and abundance of both the macroinvertebrate and fish community. Predatory fish such as pike *Esox lucius* are able to exploit the dense macrophyte stands and are further reducing populations of cyprinid fish.

**Gatwick Stream** - Environment Agency data from 3 sites on the Gatwick Stream indicate that the study section is of moderate to poor water quality, with sluggish flow and sedimented condition. Fewer macroinvertebrate taxa were recorded at the two Gatwick Stream sites compared with River Mole (mean of 10 taxa per visit). BMWP and ASPT scores indicate moderate water quality conditions at the upstream and poor to very poor at the downstream site. A high LIFE score for the upstream site during the spring visit suggests that velocities are high in the early part of the season and decline through the summer and



autumn. PSI scores for the upstream site fluctuated considerably across the three seasons, from only slightly sedimented conditions in spring to sedimented conditions in autumn.

A total of 300 and 317 fish were caught in spring and autumn respectively at the Gatwick Stream site after three survey 'runs'. Chub *Squalis cephalus* was the most abundant species in the spring survey and dace *Leuciscus leuciscus* in the autumn. Shading of the channel by overhanging trees meant that both water temperature and dissolved oxygen remained high throughout the three seasons.

## Conclusions

Both watercourses supported macroinvertebrate communities indicative of moderately polluted conditions, exacerbated by relatively low flow conditions and high levels of sedimentation. Dense macrophyte growth on the River Mole is contributing to acute reductions in dissolved oxygen which are impacting on the macroinvertebrate assemblage.

The presence of one record from 2013 of shining ram's-horn snail, an IUCN Red List species and UK species of principal importance under the 2006 NERC Act has implications for the design of the River Mole scheme. Although not recorded during the survey, there remains a possibility that the species may occur at the site of the 2013 record at the downstream end of the desk study area. A targeted survey is required to determine its potential presence.

The Gatwick Stream appears to be impacted by both organic pollution and silt deposition, possibly from a storm water discharge outlet from a nearby industrial area. Consistently high populations of fish caught in spring and in autumn are likely to be a consequence of stable temperature and DO conditions caused by shading and potentially high abundances of pollution tolerant macroinvertebrates such as *Oligochaete* worms as a food source.

## Key Recommendations

The main recommendations are set out below:

### ***River Mole***

- Undertake survey to establish presence/absence of shining ramshorn snail. Survey to focus initially on the site of the 2013 record and if found to be present, extended to incorporate the whole study section. Surveys should be scoped and undertaken by a specialist mollusc ecologist.
- If shining ramshorn snail found to be absent from main channel, undertake vegetation removal/dredging from the central channel/thalweg of the River Mole in selected sections. Marginal berms should be retained on alternate sides of the channel throughout the dredged section for the re-establishment of emergent vegetation. Brushwood faggots or fascines anchored with wooden stakes can be used to maintain the riverward edge of the berm and prevent silt from slumping into the main channel.
- Off-line scrapes and shallow pond could be created within the floodplain grassland area to provide habitat for wetland birds. If shining ramshorn snail is found to be present this recommendation can be adapted to incorporate new, permanently wet ditches supporting dense emergent reed vegetation.
- If shining ramshorn snail is found to be absent it is advised that some level of routine maintenance of macrophyte and bankside vegetation is undertaken annually under an appropriate management plan.
- Before any in-channel works begin it is advised that a fish rescue and exclusion or translocation is undertaken to safeguard fish populations.
- Stop nets should be installed at either end of the site proposed for in-channel works to prevent access by any fish species whilst the works are on-going.

***Gatwick Stream (Brook)***

- Identify point sources of pollution from industrial area associated with Crawley STW, including storm drains and surface water discharges points from roads and urban areas.
- Consider a Sustainable Drainage System (SuDS) scheme to address these discharges, including settlement ponds and reedbed treatment systems which would have additional biodiversity benefit.

Before any in-channel works begin on either watercourse it is advised that a fish rescue survey is undertaken to safeguard fish populations in the affected area. It will also be necessary to install stop nets at either end of the reach where in-channel works will be undertaken to prevent access by any other fish species whilst the works are ongoing.

Hydrometric surveys should be undertaken at various points along both rivers to better understand present hydrological conditions and inform plans to modify the channels.

# 1. Introduction

## 1.1 Development Background

1.1.1 Two watercourses, the River Mole and Gatwick Stream (Brook), will be directly affected by proposals for a surface water management and flood alleviation scheme to the east and west of Gatwick Airport. The scheme may include proposals to re-meander the River Mole close to where it emerges from beneath the airport runway and create new flood attenuation areas to the west of the watercourse. New flood storage has already been created to the west of the Gatwick Stream, with further areas likely planned within the floodplain to the east of the watercourse.

1.1.2 The study area encompasses two watercourses; a 1.3 km stretch of the River Mole immediately downstream of where it emerges from under the runway at Gatwick Airport and a 750 m stretch of the Gatwick Stream (a tributary of the River Mole) upstream of the Crawley sewage works.

## 1.2 The Objectives

1.2.1 RPS commissioned ECUS in May 2020 to undertake fish and aquatic macroinvertebrate surveys of the two rivers within the proposed study area. The objectives were to:

- Determine baseline populations for both fish and aquatic macroinvertebrates in these two watercourses over the course of a year (2020).
- Carry out a targeted desk study for the surrounding areas of both sites including a 1 km perimeter.
- Provide a report on the surveys giving the methods and results of the surveys, with recommendations, including opportunities for enhancement, mitigation and further survey recommendations.

## 1.3 Background to the Watercourses

1.3.1 The River Mole rises in Baldhorns Copse in West Sussex and discharges into the River Thames at the town of Molesey in Surrey. The Mole catchment flows over the Wealden and London clays, however, between Dorking and Leatherhead, the river cuts its way through the North Downs chalk. In this area part of the river water disappears through holes in the underlying chalk feeding into the groundwater aquifers before flowing back into the river near to Leatherhead. This action has been suggested as the origin to the name of this river, but is more likely attributed to the fact it meets the Thames at Molesey.

1.3.2 Approximately 7 miles downstream of the source, the River Mole reaches the boundary of Gatwick Airport where it passes beneath the runway in a culvert. The reach that will be affected by the proposed flood alleviation scheme extends 1.3 km downstream from where the river emerges from beneath the airport runway (see Figure 1a, in **Appendix 1**). The survey stretch on the Gatwick Stream surveyed (TQ 291 398) lies upstream of the Crawley sewage works (see Figure 1b, in **Appendix 1**).

## 2. Methodology

### 2.1 Aquatics Desk Study

- 2.1.1 A study area was defined as an area that encompassed the site and all land within 1 km of the perimeter of each of the sites, (see Figures 2a and 2b, in **Appendix 1**). Records of designated sites and protected and/or otherwise notable species were then sought for both study areas.
- 2.1.2 Sources of information were as state in Error! Reference source not found..

**Table 3-1: Sources of data**

Data Type	Source
Statutory sites for nature conservation related to the river environment	Multi-Agency Geographical Information for the Countryside (MAGIC) <a href="https://magic.defra.gov.uk/magicmap.aspx">https://magic.defra.gov.uk/magicmap.aspx</a>
Non-statutory sites for nature conservation, protected and notable species and invasive and non-native species (fish and macroinvertebrates only)	Sussex Biodiversity Records Centre
Background information on Water Framework Directive status	<a href="https://environment.data.gov.uk/catchment-planning">https://environment.data.gov.uk/catchment-planning</a>
Aquatic macroinvertebrate, fish and invasive and non-native species data	Environment Agency data request (EA Analysis and Reporting)

- 2.1.3 A request for information was sent to the Sussex Biological Records Centre in October 2020. The boundaries of any designated site and records of species were sought for part of the study area encompassing the site and within 1 km of the perimeter of each of the sites.
- 2.1.4 The records included in this report are those relating to fish and macroinvertebrates. Records over 10 years old have been excluded.

### 2.2 Survey: Aquatic Macroinvertebrates

- 2.2.1 A representative 100m section on each watercourse was identified from a walkover survey conducted prior to the spring sampling visit. Two sampling locations were identified on the Gatwick Stream, one at the upstream and one at the downstream end of the 100 m section (Figure 1a and 1b, **Appendix 1**). Only one sampling at the upstream end of the reach was safely accessible on the River Mole.
- 2.2.2 Samples were collected at each of the sites using the Whalley Hawkes Paisley Trigg (WHPT) method comprising of a standard three-minute kick sample using a long-handled pond net with 1 mm mesh size, which was supplemented by a one-minute hand search (Environment Agency, 2017). Sampling of habitats within the three-minute kick/sweep sampling were in proportion to their occurrence. Samples were then preserved in industrial methylated spirits (IMS) for processing in the laboratory to the requirements outlined in EA Operational Instruction 024\_08 Freshwater macroinvertebrate analysis of riverine samples (Environment Agency, 2014).
- 2.2.3 Macroinvertebrates were identified to Mixed Taxon Level 5, to enable evaluation of the macroinvertebrate community and calculation of the relevant biotic indices including Biological Monitoring Working Party (BMWP), Average Score Per Taxon (ASPT) and Lotic-Invertebrate index for Flow Evaluation (LIFE). Proportion of Sediment-sensitive Invertebrates (PSI) and Community Conservation Index (CCI).

2.2.4 Aquatic Macroinvertebrate sampling was undertaken in spring, summer and autumn on the dates presented in **Table 3-2**.

**Table 3-2: Aquatic Macroinvertebrate survey dates.**

Aquatic Macroinvertebrate survey visit	Date
Spring	04/06/2020
Summer	29/07/2020
Autumn	29/09/2020

### 2.3 Aquatic Macroinvertebrate Data Analysis

2.3.1 The macroinvertebrate abundance data collected during the field surveys and background data from the Environment Agency has been analysed using a range of biotic indices. Each of the indices used in the analyses are summarised below.

#### **Biological Monitoring Working Party (BMWP) Score**

2.3.2 The BMWP score is a method for indexing river water quality in England and Wales using macroinvertebrate families. Originally published in the early 1980's, the system was updated in 2013 based on a more robust baseline data set (Paisley *et al*, 2013). A score of between 1 and 10 is assigned to families found within a sample based on their tolerance to organic pollution, with a score of 1 indicating high tolerance, and 10 indicating low tolerance. Low scoring families include worms (*Oligochaeta*) and midge larvae (*Chironimidae*), whilst the presence of mayfly (*Ephemeroptera*) and stonefly (*Plecoptera*) larvae is indicative of clean water conditions. The scores for each family recorded in the sample are summed to give the overall BMWP site score. Since the overall site score is influenced by the number of families as well as the scores of the individual families in the sample, an average is taken by dividing the overall BMWP score by the number of families/taxa in the sample. This is termed the Average Score per Taxon (ASPT).

2.3.3 **Table 3-3** provides an interpretation of the BMWP scoring system.

**Table 3-3: BMWP scoring system**

BMWP score	Category	Interpretation
0 – 10	Very poor	Heavily polluted
11 – 40	Poor	Polluted or impacted
41 – 70	Moderate	Moderately impacted
71 – 100	Good	Clean but slightly impacted
> 100	Very good	Unpolluted, un-impacted

#### **River Invertebrate Classification Tool (RICT)**

2.3.4 BMWP and ASPT has largely be superseded by the River Invertebrate Classification Tool (RICT), which is one of the parameters used for classifying rivers according to their ecological status under the Water Framework Directive (WFD). The scores derived for an individual site under RICT are compared with those expected under unpolluted conditions (known as reference conditions) in order to give an Environmental Quality Ratio (EQR). This aims to take account of the variability of macroinvertebrate families in rivers resulting from environmental parameter such as altitude,

underlying geology and proximity to the river source.

**Lotic invertebrate Index for Flow Evaluation (LIFE) Score**

2.3.5 The LIFE score system links flow conditions in rivers, and specifically flow velocity, with commonly identified macroinvertebrate species and families (Extence *et al.*, 1999). Macroinvertebrates are assigned to one of 6 groups depending on their tolerance to low flow conditions. The groups range from ‘I’ comprising taxa associated with rapid flow conditions (>100 cm s<sup>-1</sup>) to ‘VI’ including those associated with drying or drought impacted sites. A flow score is obtained for each species/taxon by combining the flow category with an estimated abundance score as described by Extence *et al.* (1999). The LIFE score for a sample is obtained by summing the individual flow scores for each taxon by the number of taxa in the sample. LIFE scores range from 1 to 12, with scores of 8 or above indicating moderate to high flow conditions, and scores of 7 or below indicating sluggish conditions.

**Proportion of Sediment sensitive Invertebrates (PSI)**

2.3.6 The PSI index provides an indication of the extent to which watercourses have been impacted by the deposition of fine sediment (Extence *et al.*, 2017). Following the same principle as the LIFE score system, invertebrates are assigned to one of four groups depending on their sensitivity to fine sediment, with Group A comprising highly sensitive taxa, and Group D those that are highly insensitive. The method also requires a log abundance category to be estimated for all taxa identified in a sample (1–9, 10–99, 100–999 and 1000+ individuals present). Scores range from 80 -100 for un-sedimented sites down to 0-20 for highly sedimented sites (Table 3-4).

**Table 3-4: Interpretation of PSI scores**

PSI score	Riverbed condition
81 – 100	Minimally sedimented/un-sedimented
61 – 80	Slightly sedimented
41 – 60	Moderately sedimented
21 – 40	Sedimented
0 – 20	Heavily sedimented

**Community Conservation Index (CCI)**

2.3.7 The CCI combines the rarity of constituent species in a sample with the diversity of the community, or community richness, to give a single integrated score which can be used as the basis for site evaluating (Chadd & Extence, 2004). Species identified from a survey site or area are given a Conservation Score (CS), based on standard rarity categories, with Red Data Book 1 (Endangered) species scoring 10, and very common species scoring 1. The sum of each of the conservation scores in the sample is then divided by the number of contributing species to give the overall CCI score.

**2.4 Survey: Fish**

2.4.1 The surveys were undertaken using the catch depletion method in order to assess species composition, age structure and to estimate population size. Surveys were undertaken by an accredited electric fishing team comprising three members of staff. Surveys and analysis conformed to the relevant guidance outlined in BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity (British Standards, 2003). An FR2 consent (application to use fishing instruments other than rod and line) was sought from the Environment Agency prior to conducting the survey.

- 2.4.2 The survey was undertaken over a 100 m reach and there was one survey reach per watercourses, coinciding with the macroinvertebrate survey locations on both watercourses. Stop nets were installed across the channel at either end of the reach to prevent fish entering or leaving the survey area. Holding containers for captured fish were established in a small boat with an aerator installed to provide oxygen to captured fish.
- 2.4.3 The survey was undertaken using an electrofishing box alternating between a single anode and two anodes depending on the width of the river in order to maximise catch efficiency. One surveyor, operating the electrofishing anode waded from downstream to upstream and a second surveyor netted any stunned fish. In areas where the rivers was wider the second surveyor also operated an anode. The operatives were followed by an additional surveyor pulling a small boat with the electrofishing box and holding tank on board, and also equipped with a hand net to maximise the catch rate. At the end of each run all caught fish were identified, measured and placed in a submerged holding net to facilitate their recovery and prevent re-capture.
- 2.4.4 Two survey visits were undertaken, one in spring (04/06/2020) and one in autumn (29/09/2020) to establish a baseline of the species composition on the two watercourses. Undertaking the autumn visit in September ensured that air temperatures are above the minimum of 10 °C and minimise the risk of high flow conditions. It would also avoid risk of disturbance to salmonid spawning habitat, should it be present.

## 2.5 Limitations

- 2.5.1 Only one macroinvertebrate sample could be retrieved from the downstream River Mole site, due to various access issues, such as dense bankside vegetation creating a barrier to the river and steep banks, which prevented safe access and egress to the river.
- 2.5.2 The River Mole has exceptionally high coverage of aquatic plants, which made electrofishing difficult. In spring the filamentous algae blanket weed *Cladophora agg* was found in dense clumps making progress slow, as the anode became blanketed by the filamentous algae each time it was placed in to the water and needed regular clearing in order to progress. In addition to this, the macroinvertebrate surveys were difficult due to the dense macrophyte growth and deep waters preventing more than one macroinvertebrate sample being taken using the WHPT method.
- 2.5.3 The timing of the spring survey was delayed by a nationwide lockdown related to the COVID-19 outbreak.

## 3. Results

### 3.1 Aquatics Desk Study

#### *River Mole*

##### Environment Agency: Water Framework Directive Status

- 3.1.1 Under the Water Framework Directive (WFD) rivers and standing waters are termed waterbodies and are classified according to their ecological status. Ecological status is classified using five categories of high, good, moderate, poor and bad and is measured and classified via a range of inter-linked biological, physico-chemical and physical (morphological) parameters. The classification process is based primarily on the biological quality elements of the water body but considered alongside support elements covering physico-chemical standards and hydromorphological quality elements. Each of these supporting elements is assigned to a status category (i.e. high to bad). The overall status of the waterbody is based on the status category of the worst supporting element.
- 3.1.2 The affected reach of the River Mole falls within the WFD waterbody named 'Mole Upstream of Horley (GB106039017481)'. There is little information relating to the stretch of the River Mole. It was first classified as good under the WFD classification system in 2015, although the most recent classification in 2019 designates it as moderate. Although the biological quality elements are classified as good (based on fish data only), one of the physico-chemical quality elements (phosphorous) is classified at moderate status, and therefore the overall waterbody status is classed as moderate.
- 3.1.3 No Environment Agency background records were received for the River Mole.

##### Sussex Biological Records Centre Data

- 3.1.4 A total of 3 records of fish species were returned from the Sussex Biological Records Centre for the River Mole within 1 km of the study section, comprising one record of bullhead approximately 0.5 km downstream from the study section in 2014, and a record of 2 adult brown trout within the survey section in 2016. Bullhead is listed as a non-priority species under Annex 2 of the EU Habitats Directive and listed on the Sussex Rare Species Inventory. Brown trout is a UK Priority Species under the UK Post 2010 Biodiversity Framework, and Section 41 of the Natural Environment and Rural Communities Act 2006.
- 3.1.5 There is one record of the shining ram's-horn snail within the study section (TQ 25623 40908; Figure 3, **Appendix 1**) from February 2013. The species was recorded as being 'u/s Pond M and Tributary'. The shining ram's-horn snail is nationally scarce, a UK Priority Species under the UK Post 2010 Biodiversity Framework and listed on the Sussex Rare Species Inventory.
- 3.1.6 One notable dragonfly species, common darter *Sympetrum striolatum* was recorded within 1 km of the site. The species is listed in the UK Red Data Book. A total of 44 observations were made of the species in the vicinity of the Gatwick airport, with a number of them within the study section. There are no records of the larvae in the River Mole, either from the Sussex Biological Records Centre or the Environment Agency and therefore breeding sites are unclear.
- 3.1.7 A number of invasive and non-native invertebrate and aquatic plant species occur within 1 km of the River Mole study section. Three records of signal crayfish *Pacifastacus leniusculus* were returned from within 1 km of the study section between 2011 and 2013. Two of the sites were on the River Mole within the study section and the third north of the Gatwick runway within a tributary of the River Mole. Signal crayfish is listed on Schedule 9 of the Wildlife and Countryside Act. Under Section 14 of the Act it is an offence 'to release or allow to escape into the wild' any species listed under Schedule 9'.



- 3.1.8 Records of several invasive aquatic/ riparian plant species were also recorded within 1 km of the site including Nuttall's pond-weed *Elodea nuttallii*, Japanese knotweed *Fallopia japonica* and Himalayan balsam *Impatiens glandulifera*.

### **Gatwick Stream**

#### Environment Agency: Water Framework Directive Status

- 3.1.9 The Gatwick Stream is a tributary of the River Mole and is approximately 12km in length. It rises near Three Bridges and joins the River Mole near the centre of Horley. It falls within the Tilgate Stream and Gatwick Stream at Crawley WFD waterbody (GB106039017500). The overall waterbody status has remained moderate since 2013, although the biological quality elements are assigned bad status on the basis of fish data. This is a deterioration from poor status in 2016. Macroinvertebrates were classified at poor status in 2019 and have remained at that classification since 2013. Sewage discharges and the invasive signal crayfish are given by the Environment Agency as the reasons for poor biological quality in the brook.

#### Environment Agency: Macroinvertebrate Data

- 3.1.10 Macroinvertebrate data was received from the Environment Agency for 3 sites on the Gatwick Stream that lie within the study area (U/S Crawley STW (TQ 29160 39780); Downstream Tinsley Bridge (Flylife site) (TQ 29129 39864) and at Tinsley Bridge, Tinsley Green (TQ 29150 39800)). One sample was collected at U/S Crawley STW in October 2017, and duplicate samples were collected at Downstream Tinsley Bridge (Flylife site) and at Tinsley Bridge, Tinsley Green) in March 2019 (Figure 1b, **Appendix 1**). The Environment Agency have provided feedback that the 2019 samples were taken in response to a pollution incident and that the duplicate sample from both sites with lower number of taxa recorded was sorted on the bank. In comparing the data with this study only the laboratory sorted sample has been considered, although the results for both samples are presented in **Table 4-1**.
- 3.1.11 A total of 13 families were recorded during the survey at the U/S Crawley STW site in October 2017. The freshwater shrimp *Gammarus pulex*, a species indicative of moderate water quality, was the most numerous. However, the site also supported relatively high numbers of *Oligochaeta* worms, a family highly tolerant of low oxygen conditions. BMWP and ASPT scores have been calculated since none were provided by the Environment Agency (**Table 4-1**). This site had a BMWP score of 43 with an ASPT of 4.3 indicating moderate to poor water quality.
- 3.1.12 Of the two duplicate samples taken at the Downstream Tinsley Bridge (Flylife site) (TQ 29129 39864) on 14th March 2019 a total of 22 families were recorded in one sample and 7 in the other. Midge larvae (*Chironomidae*) and *Oligochaeta* worms were present in relatively high numbers in the second sample indicating poor water quality. Based on the second sample the site had a BMWP of 47 and ASPT of 3.92.
- 3.1.13 At the most upstream site at Tinsley Bridge (TQ 29150 39800) site a total of 8 families were recorded in one of the duplicate samples and 21 in the second. In general, the samples supported pollution tolerant families and species such as *Oligochaeta* (20 and 40 individuals respectively) and the water louse *Asellus aquaticus* (30 individuals in the second sample). However, the site also supported the damselfly larvae *Calopteryx* sp., a relatively pollution sensitive family. The second sample at this site had a BMWP of 48 and ASPT of 4.

**Table 4-1: EA Macroinvertebrate Biotic Scores for Gatwick Stream**

Site	U/S Crawley STW	Downstream Tinsley Bridge (Flylife Site)	Downstream Tinsley Bridge (Flylife Site)	At Tinsley Bridge, Tinsley Green	At Tinsley Bridge, Tinsley Green
Date	12/10/2107	14/03/2019	14/03/2019	14/03/2019	14/03/2019
BMWP (TL1)	43	15	47	20	48
ASPT	4.3	3.00	3.92	3.33	4
LIFE	7.5	7.0	7.11	7.00	7.00
PSI	40.00	28.57	36.00	36.36	32.14
CCI	1.00	N/A	1.00	N/A	1.00

3.1.14 LIFE scores for each of the 3 sites ranged from 7.0 to 7.5 indicating sluggish to moderate flow conditions. PSI scores for all three sites indicate sedimented conditions, although the U/S Crawley STW site is close to moderately sedimented with a score of 40. CCI scores of 1 indicate low conservation value.

Sussex Biological Records Centre Data

3.1.15 Records of two fish species, bullhead and brown trout were returned for Gatwick Stream, from the Sussex Biological Records Centre. One adult bullhead was recorded within the study section in October 2015 and a brown trout in a similar location in July 2016.

3.1.16 A total of 15 records of adult common darter dragonflies were returned for the study section on the Gatwick Stream between 2012 and 2017, although there are no records of the larvae. Six records of the downy emerald dragonfly *Cordulia aenea* and 2 of the brilliant emerald dragonfly *Somatochlora metallica* were returned from within the past 10 years. Downy emerald dragonfly is a Red List species on the IUCN Red List and a Priority Species on the UK Post 2010 Biodiversity Framework. The downy emerald dragonfly is listed on the Sussex Rare Species Inventory. None of the records were on the Gatwick Stream and there are no records of the larvae.

3.1.17 There were three records of the invasive signal crayfish from within the study section on the Gatwick Stream in 2017. The invasive aquatic plant, Nuttall's pond weed was recorded within the study section in 2016 and there are records of Japanese knotweed and Himalayan balsam.

Environment Agency: Fish data (River Mole and Gatwick Stream)

3.1.18 Data provided by the Environment Agency indicates that both the Gatwick Stream and River Mole were stocked in 2018 and 2019 with Roach, Barbel *Barbus barbus*, Dace and Chub. In 2018, 3200 fish were added to the lower Gatwick Stream in response to a pollution event which occurred in 2017.

3.1.19 In 2019, 3600 fish were stocked in the River Mole in response to a prolonged dry weather event in 2018, which occurred as a result of low flows and first flush effect, which was estimated to have affected approximately 2000 fish.

### 3.2 Field Data

#### River Mole

##### Water Quality

3.2.1 A maximum temperature of 17.6°C was recorded at the sampling site on the River Mole during the summer visit on 29th July 2020 (**Table 4-2**). The temperature was only slightly lower on the first visit (16.4°C on 1st July), which was delayed due to Covid 19 restrictions. Water temperature dropped to 13.8°C by the autumn visit on 29th September 2020. DO concentrations dropped sharply between the first and second visits, from 60.8% in early July to 17% on 29th July, before recovering slightly to 28.7% by end of September. Both conductivity and turbidity increased progressively through the season. Conductivity increased from 358.5 to 471µS/cm, whilst turbidity increased from 3.78 to 4.3NTU.

**Table 4-2: Water Quality Data Recorded at River Mole Sampling Site**

Season	Temperature (°C)	Dissolved oxygen (%)	Dissolved oxygen (mg/l)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Spring	16.4	60.8	5.94	7.30	358.5	3.78
Summer	17.6	17.0	1.60	7.19	341.0	4.02
Autumn	13.8	28.2	3.03	8.38	471.0	4.30

##### Aquatic Macroinvertebrates

3.2.2 A mean of 19.3 taxa were recorded at the River Mole site across the three visits. There was relatively little variation in the number of taxa recorded on each visit, with the maximum of 21 in the spring/early summer sample, and a minimum of 17 in the summer sample (**Table 4-3**). Of these, 12 taxa/species occurred in all three samples, including the water shrimp, the pea mussel *Sphaereum corneum* and the mayfly larvae *Cloeon dipterum*. However, abundances of individual taxa within the samples varied considerably across the 3 visits, with the crustacean *Cladocera* the most abundant in the early summer samples 01/07/20, replaced by the water boatman, *Coroxidae* one month later. The most abundance taxa in the autumn samples was the Isopod *Asellus aquatica* (waterlouse). These changes in abundance are likely to be driven by seasonal changes in life stage from early to later (larger, and therefore more readily sampled) larval instars as well as the availability of food resources.

**Table 4-3: Number of Macroinvertebrate Species/Taxa Recorded at River Mole and Gatwick Stream Sites**

Site	Spring	Summer	Autumn
River Mole	21	17	20
Gatwick Stream upstream	12	10	13
Gatwick Stream downstream	8	8	9

3.2.3 The consistent occurrence of low BMWP scoring (i.e. 3 or below) species and taxa such as the waterlouse, *Chironimidae* and *Oligochaeta* on all three visits, suggest that the watercourse is

affected by organic pollution. This is confirmed by BMWP scores of 44, 46 and 49 and ASPT of 3.73, 3.45 and 3.43 in the spring, summer and autumn samples respectively indicating moderately polluted conditions.

- 3.2.4 LIFE scores for the River Mole ranged from 6.25 in the spring/early summer sample, 6.1 in the summer sample and 5.87 in the autumn sample, indicating sluggish flow conditions (**Table 4-4**). The decline in LIFE scores over the summer period are likely to be primarily a result of low flow conditions due to low summer rainfall, although extensive macrophyte beds in the channel may also be impeding flow. Low PSI scores of less than 20 also indicate heavily sedimented conditions. This correlates with low flow velocities in the channel indicated by the LIFE scores and is likely to be exacerbated by the extensive macrophyte plant beds.
- 3.2.5 CCI scores of between 5 and 10 indicate that the macroinvertebrate community is of moderate conservation value. The presence of *Sigara limitata*, a species of water boatman, contributed to a slightly higher score of 9.62 in the autumn sample.

**Table 4-4: Macroinvertebrate Biotic Indices**

Biotic Index	Spring			Summer			Autumn		
	U/S River Mole	Gatwick Brook U/S	Gatwick Brook D/S	River Mole U/S	Gatwick Brook U/S	Gatwick Brook D/S	U/S River Mole	Gatwick Brook U/S	Gatwick Brook D/S
	01/07/2020	01/07/2020	01/07/2020	27/07/2020	27/07/2020	27/07/2020	29/09/2020	29/09/2020	29/09/2020
BMWP (TL1)	44	46	14	46	37	29	49	41	20
LIFE (TL5)	6,25	8,17	7,5	6,10	7,40	7,75	5,87	6,75	8
ASPT (TL2)	3,73	4,92	3,50	3,45	4,53	3,91	3,43	4,13	2,88
PSI (TL5)	10,00	66,67	14,29	5,00	41,67	50,00	6,25	21,05	33,33
CCI (TL5)	5,50	4,50	0	4,00	5,00	1,00	9,62	1,20	1

Fish

- 3.2.6 A total of 415 fish were caught on the River Mole in spring after three survey ‘runs’ compared with only 28 fish caught in autumn with the same level of effort. Roach were the most abundant fish species identified (252) in spring and in autumn (13).
- 3.2.7 The size range of species caught on the electrofishing surveys in spring (**Table 4-5**) suggests that there are multiple age classes of each species, ranging from juveniles to mature adults. The stretch of the River Mole sampled in this study appears to be a good breeding and spawning environment for roach and perch *Perca fluviatilis*, due to its slow flow environment and dense vegetation. The mean size data in spring would suggest that this stretch also appears to be a good environment for juvenile and sub-adult chub and dace as well as providing optimal foraging habitat for predatory fish species such as pike.

**Table 4-5: River Mole Fish Survey Data**

River Mole					
Spring					
Species	Latin name	Abundance	Mean size (mm)	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	45	166.55	77	386
Roach	<i>Rutilus rutilus</i>	252	106.91	45	256
Dace	<i>Leuciscus lueciscus</i>	37	127.86	59	203
Pike	<i>Esox lucius</i>	14	344.86	108	595
Perch	<i>Perca fluviatilis</i>	46	130.00	73	258
Bream	<i>Abramis brama</i>	3	72.33	62	79
Tench	<i>Tinca tinca</i>	2	89.0	85	93
Gudgeon	<i>Gobio gobio</i>	13	93.1	82	109
Rudd	<i>Scardinius erythrophthalmus</i>	2	138.50	81	196
Roach/ Bream Hybrid		1	143	143	143
Autumn					
Species	Latin name	Abundance	Mean size (mm)	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	3	217.67	181	289
Roach	<i>Rutilus rutilus</i>	13	123	64	200
Tench	<i>Tinca tinca</i>	7	198	153	248
Pike	<i>Esox lucius</i>	4	121	110	127
Perch	<i>Perca fluviatilis</i>	1	86	86	86

**Gatwick Stream**

Water quality

3.2.8 Water temperature at the two Gatwick Stream sites remained relatively consistent across the three seasons (**Table 4-6**), peaking at 16.7°C at the downstream site during the summer visit on 29th July. The lowest temperature was recorded at the upstream site (14.8°C) at the end of September. The sites are moderately shaded by overhanging trees, which will help to buffer water temperature. DO concentrations also remained relatively high at over 70% throughout the three seasons, reaching a maximum of 78.7% at the downstream site in autumn. Turbidity was relatively high compared with the River Mole site, with a minimum of 5.95 NTU at the upstream site in autumn and a maximum of 11.85 NTU at the upstream site in summer.

**Table 4-6: Water Quality Data for Gatwick Stream**

Spring						
Site	Temperature (°C)	Dissolved oxygen (%)	Dissolved oxygen (mg/l)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Gatwick Stream US	15.4	73.5	7.34	7.53	276.2	11.21
Gatwick Stream DS	16.7	71.7	7.37	7.76	333.9	10.74
Summer						
Site	Temperature (°C)	Dissolved oxygen (%)	Dissolved oxygen (mg/l)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Gatwick Stream US	16.5	72.0	7.04	7.68	280.1	11.85
Gatwick Stream DS	16.7	73.5	7.13	8.00	269.1	10.92
Autumn						
Site	Temperature (°C)	Dissolved oxygen (%)	Dissolved oxygen (mg/l)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Gatwick Stream US	14.8	76.8	7.73	7.46	413.9	5.95
Gatwick Stream DS	16.0	78.7	7.73	8.20	387.8	11.64

### Aquatic Macroinvertebrates

- 3.2.9 Fewer taxa were recorded at the two Gatwick Stream sites compared with River Mole (mean of 10 taxa per visit, compared with 19.3) (**Table 4-3**). As with the River Mole, the number of taxa recorded per visit remained relatively consistent, with a maximum of 13 in the autumn sample and a minimum of 10 in the summer sample at the upstream site. Eight taxa were recorded at the downstream site during spring and summer and 9 in the autumn.
- 3.2.10 BMWP scores indicate moderate water quality conditions for the upstream site at the Gatwick Stream in spring and autumn (46 and 41 respectively) but were classed as poor in summer (37) (**Table 4-4**). The boundary between moderate and poor lies at 40 and therefore the difference between the three visits is unlikely to be significant and is due to the smaller number of taxa recorded. However, an additional three species were recorded in the autumn sample, including the coloniser species *Asellidae* (isopod crustaceans) and the caddisfly *Polycentropus flavomaculatus*, suggesting an increase in water quality at this location, although both species were found in low abundance. The ASPT for the upstream site is similar across the three visits and is lowest in the

autumn sample (4.92, 4.53 and 4.13 for the spring, summer and autumn visit respectively).

- 3.2.11 At the downstream site of the Gatwick Stream the BMWP scores are classified as poor across all three visits, with the score of 14 for the spring visit being close to very poor (**Table 4-4**). The ASPT is also consistently lower for this site than the upstream site (3.50, 3.91 and 2.88 for the spring, summer and autumn visit respectively) over all three visits indicating the presence of pollution tolerant taxa only.
- 3.2.12 The PSI scores for the upstream site fluctuated considerably across the three season, with the maximum score of 66.67 in the spring indicating only slightly sedimented conditions (**Table 4-4**). However, the scores dropped progressively at this site through the season to 41.67 in the summer (moderately sedimented conditions) and then to 21.05 (sedimented conditions) in the autumn. Assuming no changes in the inputs of sediment upstream of the site, this suggests that flow velocity dropped through the season, leading to increased sediment deposition. A high LIFE score for the upstream site of 8.17 during the spring visit also suggests that velocities are high in the early part of the season.
- 3.2.13 The PSI scores for the downstream site indicated heavily sedimented conditions during the spring season (score of 14.29), with a change to moderately sedimented conditions (score of 50) in the summer and a return to sedimented conditions in the autumn (score of 33.33). LIFE scores remained relatively high and consistent across the three seasons at the downstream site (7.5, 7.75 and 8 at the spring, summer and autumn visit respectively) suggesting relatively consistent flow velocities (**Table 4-4**).
- 3.2.14 CCI scores for both of the Gatwick Stream sites were relatively low indicating that rare and/or notable species are absent from the macroinvertebrate assemblage. Although scores for both sites were below 5 on all sampling occasions, the upstream site had scores of 4.5 and 5 in the spring and summer respectively, whilst the scores for the downstream site was either 1 or 0 on all occasions. This indicates that the assemblage at the upstream site is of marginally higher conservation value.

#### Fish

- 3.2.15 A total of 300 and 317 fish were caught in spring and autumn respectively in the Gatwick Stream after three survey 'runs'. Chub were the most abundant fish species identified (111) in spring on the Gatwick Stream, whereas dace were the most abundant fish species identified (137) in autumn (**Table 4-7**).
- 3.2.16 The size range of species caught during the electrofishing surveys carried out on the Gatwick Stream in spring suggests that there are multiple age classes of each species, ranging from juveniles to mature adults all year round.

**Table 4-7: Gatwick Stream Fish Survey Data**

Gatwick Brook					
Spring					
Species	Latin name	Abundance	Mean size (mm)	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	111	194.50	52	360
Dace	<i>Leucisus luecisus</i>	74	145.35	63	220
Perch	<i>Perca fluviatilis</i>	36	85.05	65	156
Roach	<i>Rutilus rutilus</i>	11	105.45	72	153
Bream	<i>Abramis brama</i>	6	146	92	279
Gudgeon	<i>Gobio gobio</i>	57	107.24	75	197
Stone Loach	<i>Barbatula barbatula</i>	3	127.33	97	179
Autumn					
Species	Latin name	Abundance	Mean size	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	85	211.56	71	436
Dace	<i>Leucisus luecisus</i>	137	149.38	50	204
Roach	<i>Rutilus rutilus</i>	28	111.32	71	156
Perch	<i>Perca fluviatilis</i>	21	113.14	80	213
Bream	<i>Abramis brama</i>	10	158	132	284
Gudgeon	<i>Gobio gobio</i>	36	118.55	52	146
Stone Loach	<i>Barbatula barbatula</i>	3	86	65	98



## 4. Discussion

### 4.1 River Mole

- 4.1.1 The study stretch on the River Mole lies within open floodplain grassland with no shading from trees. This means that water temperatures and therefore DO, fluctuate considerably since oxygen is less soluble in warm water. Bacterial activity associated with organic pollution also depletes DO levels and therefore macroinvertebrate taxa which occur in organically polluted conditions are tolerant of low DO conditions. Both factors are likely to be influencing the macroinvertebrate community at the River Mole site.
- 4.1.2 Extensive stands of macrophyte plants covered approximately 90% of the channel surface, including submerged species such as water crowfoot *Ranunculus aquatilis* and the invasive non-native Canadian pondweed *Elodea canadensis*. Emergent species such as branched bur-reed *Sparganium erectum*, old world arrowhead *Sagittaria sagittifolia* and reed sweet-grass *Glyceria maxima* also dominated the channel. Although this channel vegetation will have contributed DO to the water during the summer through photosynthesis, their decay in autumn will contribute to organic pollution in reducing DO (28.2% and 3.03 mg/l during the autumn visit). Significant increases in conductivity such as those seen on the River Mole from spring to autumn (358 – 471  $\mu\text{S/cm}$ ) (**Table 4-2**) are likely attributed to the decay of macrophytes and the release of ions such as phosphorous.
- 4.1.3 Submerged and emergent macrophyte stands are also contributing to reduced flow velocity and increased sedimentation, reflected in the low LIFE and PSI scores for this reach.
- 4.1.4 The presence of one record from 2013 of shining ram's-horn snail, an IUCN Red List species and UK species of principal importance under the 2006 NERC Act has implications for the design of the scheme (Figure 2, **Appendix 1**). Once abundant in ditch networks in the UK, the species has declined steeply and now only occurs in a restricted number of sites in Norfolk Broads, Pevensey Levels, Lewis Levels and East Kent (Clarke, 2011). The reasons for its decline are not fully understood, but are thought to be over-frequent ditch clearance, eutrophication due to fertiliser run-off and conversion of grazing levels to arable farming with associated water table lowering (Suffolk Biological Information Service, 2003).
- 4.1.5 In a study of the associations of the species with ditch vegetation communities Clarke (2011) only found the species in ditches supporting the *Carex-Juncus-Eleocharis-Oenanthe* community of emergent vegetation. Although a full macrophyte survey was not undertaken during this study, incidental recording of macrophytes at the sampling location was undertaken and this community type was not present. However, the entire stretch from the boundary with Gatwick airport to the end of the study reach is heavily vegetated and largely impenetrable. More suitable habitat may therefore exist further downstream towards the location where it was recorded in 2013. Recommendations for further survey to determine the potential presence of the species within the study section are presented in Section 6.
- 4.1.6 The extensive macrophyte growth on the River Mole throughout the year made electrofishing difficult. In spring the filamentous algae, *Cladophora* created dense mats, which surrounded the anode each time it was placed into the water, making progress slow as the anodes regularly needed to be brought to the surface and cleared of the algae. In some cases, *Cladophora* can be beneficial to an ecosystem by providing a food source to aquatic organisms and providing a buffer to nitrification. However, excessive growth of *Cladophora* prevents aeration of deeper waters as the dense mats prevent circulation of water, which is detrimental to an ecosystem.
- 4.1.7 The high variability and remarkably low concentration of DO in the waters of the River Mole, likely contributed to the low catch in autumn where only 28 fish were caught in comparison to 415 in spring. The slow/sluggish flow of the River Mole, in combination with higher water temperatures in

summer (17.6°C) could be causing DO to disassociate faster from the water. The increased presence of tench *Tinca tinca* in the River Mole in autumn acted as an in-field indicator of low DO conditions, as tench are able to tolerate much lower DO conditions than most other UK fish species.

4.1.8 The abundance of predatory fish in summer such as pike and perch, may have been having a disproportionate impact on prey species on the River Mole. The prevalence of these predators has likely contributed to the significant decline in the fish population from 417 in summer to 28 in autumn. In total 14 pike were caught in summer ranging in size from 108 mm – 595 mm indicating the full range of age classes. Pike are very effective freshwater hunters and as ambush predators are aided by the abundant macrophyte growth. In addition to this 46 perch were caught in summer and ranged in size from 73 mm – 258 mm, also suggesting a full range of age classes. Perch also utilise macrophytes to aid in their hunting techniques, however, they are more temperature sensitive, retreating to deeper waters throughout the autumn and winter months, which has likely contributed to their decline in the area to one individual in autumn on the River Mole.

## 4.2 Gatwick Stream

4.2.1 The downstream site of the Gatwick Stream appears to be suffering from poorer biological water quality than the upstream site, with the LIFE and PSI scores indicating an influx of organic pollution somewhere between these sites. This is supported by the absence of *Asellidae* (isopod crustaceans), which suggests that organic pollution is chronic and there has been no recovery between Spring and Autumn. Crawley sewage treatment works lies immediately east of the Gatwick Stream and although the discharge is directly into the River Mole, it is possible that storm water discharges from the associated industrial area enter the Gatwick Stream between the two sites. Relatively high turbidity levels of between 5.95 and 11.85 NTU compared with a maximum of 4.3 NTU at the River Mole site.

4.2.2 Differences in habitat quality and diversity between the two Gatwick Stream sites may also have influenced the macroinvertebrate community. Both sites were moderately shaded by overhanging trees, but the upstream sites was located on a tight bend with a small riffle section on the outer side of the bend and a shallow berm on the inside edge. These microhabitats are likely to support distinct macroinvertebrate communities, with the more pollution sensitive species present in the riffle section.

4.2.3 The considerable variation in PSI score between the three seasonal visits at the upstream sites (maximum of 66.67 in spring compared to a minimum of 21.05 in autumn) may indicate that the macroinvertebrate community at this site is sensitive to changes on sediment deposition. Equally, it may have resulted from small differences in sampling effort in each of the microhabitats leading to a higher number of sediment sensitive taxa in the spring sample. Limited conclusions can be drawn with only one sample per visit and data from a single visit and further sampling would be required to determine any trends in the data. Overall, both sites are moderately to heavily sedimented with likely potential storm water discharges resulting in greater sedimentation at the downstream site.

4.2.4 The invasive New Zealand mud snail *Potamopyrgus antipodarum* was identified at both sites except for the Gatwick Stream downstream site in autumn. The New Zealand pond snail is now one of the most common gastropods in the UK, its ability to avoid desiccation and its tolerance for a range of conditions enables it to dominate native gastropods, which may lead to disruptions in the food chain and effect native fish species. Currently the Gatwick Stream upstream site hosts the largest population of New Zealand mud snail, where abundances increased from 12 to 40 from spring to autumn in the samples collected. Signal crayfish were observed in relatively high numbers at both the Gatwick Stream sites during each of the visits.

4.2.5 The macroinvertebrate results from this study compare favourably with the Environment Agency data collected in 2017 and 2019 (**Table 4-1**). A slightly higher ASPT score of 4.92 was obtained for the upstream site in early July compared with values of 3.92 and 4.0 for the 'At Tinsley Bridge,

Tinsley Green' and 'Downstream Tinsley Bridge (Flylife Site)' in March 2019. However, this may reflect seasonal changes in the macroinvertebrate community between March and July. LIFE and PSI scores for both data sets indicate relatively sluggish and sedimented conditions.

- 4.2.6 The Gatwick Stream on first appearances seemed to be poor for fish species, but surprisingly a consistently healthy population of fish were caught in spring (300) and in autumn (317). This is likely due to the Gatwick Stream maintaining a relatively consistent water temperature (14.8-16.7°C) across all three seasons and dissolved oxygen concentrations >71%. Furthermore, although the macroinvertebrate community is poor on the Gatwick Stream, the abundance of *Chironomids*, *Oligochaetes* and *Gastropods* provide an excellent food source. There is also a diverse range of microhabitats present, such as shaded pools and undercut banks, interspersed with roots providing shelter for fish.
- 4.2.7 As a point of interest, a roach – bream *Abramis brama* hybrid was identified in spring. Hybridisation between these two species is not uncommon as hybridisation between members of cyprinids is more widespread than in any other group of freshwater fish.

## 5. Conclusions and Recommendations

### 5.1 Conclusions

- 5.1.1 There are no Environment Agency WFD monitoring sites on this stretch of the River Mole and therefore no background data to compare the field data collected in this study with. Data from a single site on a single year, albeit across three seasons, does not enable a comprehensive assessment of trends in the macroinvertebrate assemblage. However, based on the analysis of macroinvertebrate data collected for this study, the River Mole exhibits moderate biological water quality. Dense macrophyte growth within the channel, exacerbated by organic pollution are causing acute reductions in DO are likely to be impacting on the macroinvertebrate assemblage.
- 5.1.2 The record from 2013 of shining ramshorn snail, an IUCN Red List species and UK species of principal importance under the 2006 NERC Act has implications for the design of the surface water management and flood alleviation scheme. The species was not recorded during the surveys for this study, although the survey section did not coincide with the reach in which the snail was recorded.
- 5.1.3 A targeted survey for the species is required to determine its presence or absence (Section 6.2). If the species is found to be present the marginal and channel macrophyte vegetation and flow conditions will need to be preserved in the section of the river in which the population occurs. Creation of new habitat, possibly in the form of off-line ditches supporting dense emergent vegetation is likely to be a requirement of the scheme if the species is found to be present.
- 5.1.4 The structure and abundance of the cyprinid fish community in the River Mole appears to be driven by sluggish flow conditions and high summer water temperatures which favour species such as tench. The dense stands of submerged and emergent macrophytes provide foraging habitat for predatory species such as pike. Periodic dredging of the macrophyte beds would help to establish larger areas of open and deeper water thus providing refuges for prey species, improving flow conditions and creating areas of deeper, cooler water.
- 5.1.5 Based on macroinvertebrate biotic scores the Gatwick Stream has biological quality ranging from moderate at the upstream site to poor at the downstream site. Nevertheless, it retains a natural sinuous course with a variety of microhabitats supporting a range of macroinvertebrate and fish species. However, the watercourse appears to be impacted by both organic pollution and silt deposition, possibly from a storm water discharge from a nearby industrial area.
- 5.1.6 The invasive New Zealand mud snail was identified at the River Mole and Gatwick Stream sites, and signal crayfish were observed at both the Gatwick Stream sites during each visit.

### 5.2 Recommendations

- 5.2.1 Both the Gatwick Stream and the River Mole retain natural sinuous channels characteristic of lowland rivers. It will be important to maintain and enhance this characteristic in both watercourses. The following recommendations for each watercourse are based on the findings from this study and will need refinement in light of the design of the surface water management and flood alleviation scheme and in the case of the River Mole, the findings of the survey for shining ramshorn snail. However, the habitat improvement measures recommended below are largely consistent with the requirements for this species.

#### ***River Mole***

- Undertake survey to establish presence/absence of shining ramshorn snail. Survey to focus initially on the site of the 2013 record and if found to be present, extended to incorporate the whole study section. Surveys should be scoped and undertaken by a specialist mollusc ecologist.

- If shining ramshorn snail found to be absent from main channel, undertake vegetation removal/dredging from the central channel/thalweg of the River Mole in selected sections. Marginal berms should be retained on alternate sides of the channel throughout the dredged section for the re-establishment of emergent vegetation. Brushwood faggots or fascines anchored with wooden stakes can be used to maintain the riverward edge of the berm and prevent silt from slumping into the main channel.
- Off-line scrapes and shallow pond could be created within the floodplain grassland area to provide habitat for wetland birds. If shining ramshorn snail is found to be present this recommendation can be adapted to incorporate new, permanently wet ditches supporting dense emergent reed vegetation.

### ***Gatwick Stream***

- Identify point sources of pollution from industrial area associated with Crawley STW including storm drains and surface water discharge points from roads and urban areas. Consider a SuDS scheme to address these discharges, including settlement ponds and reedbed treatment systems which would have additional biodiversity benefit.

5.2.2 Before any in-channel works begin it is advised that a fish rescue survey is undertaken to safeguard fish populations in the affected area. It will also be necessary to install stop nets at either end of the reach where in-channel works will be undertaken to prevent access by any other fish species whilst the works are ongoing.

5.2.3 Currently the River Mole is choked with submerged and emergent macrophyte plant growth, which is impeding flow, increasing deposition of sediment and reducing the circulation of deeper waters preventing aeration and creating low DO conditions. It is therefore advised that there is some level of routine maintenance of macrophyte and bankside vegetation to aid in reducing the effects of flooding and contribute to increasing the biological water quality.

## **5.3 Further Survey**

5.3.1 It is recommended that further macroinvertebrate and fish surveys are carried out on both the River Mole and the Gatwick Stream to provide a more robust baseline of community assemblage and therefore better advice on any schemes in the future.

5.3.2 To provide additional insight into the hydrological conditions of these rivers, it is recommended that further investigations are carried out to monitor the flow velocity and the discharge rates in order to better advice on any schemes in the future, which could include the installation of level loggers.

## 6. References

- British Standards (2003). BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity.
- Chadd, R. Extence, C. (2004). The conservation of freshwater macroinvertebrate populations: A community based classification scheme. *Aquatic Conservation Marine and Freshwater Ecosystems*. Vol 14. Pp. 597-624.
- Clarke, R. (2011). Hydroseral habitat requirement of the endangered Shining Ramshorn Snail *Segmentina nitida*. *Aquatic Conserv: Mar. Freshw. Ecosyst.* 14: 597–624.
- Ecus Ltd. (2021). Gatwick Airport Northern Runway Project – Assessment of Terrestrial Invertebrate Interest. Ecus Ltd, Basingstoke.
- Extence, C.A, Balbi, D.M. Chadd, R.P. (1999). River Flow Indexing Using British Macroinvertebrates: A Framework For Setting Hydroecological Objectives. *Regulated Rivers: Research and Management* 15. pp543-574.
- Extence, C.A. Chadd, R.P. England, J. Naura, M. Pickwell, A.G.G. (2017). Application of the Proportion of Sediment-Sensitive Invertebrates (psi) Biomonitoring Index. *River Research and Applications*. Vol 33. Issue 10. December 2017. Pp. 1596 – 1605.
- Paisley, M.F. Trigg, D.J. Walley, W.J. (2013). Revision of the Biological Monitoring Working Party (BMWP) score system: Derivation of present-only and abundance-related scores from field data. <https://onlinelibrary.wiley.com/doi/abs/10.1002/rra.2686>.
- Suffolk Biological Information Service (2003). Shining Ram's Horn Snail (*Segmentina nitida*) Biodiversity Action Plan. <https://www.suffolkbis.org.uk/sites/default/files/biodiversity/priorityspecieshabitats/actionplans/shiningramshornsnail.pdf>
- Water Framework Directive – United Kingdom Advisory Group. UKTAG River Assessment Method. Benthic Invertebrate Fauna. Invertebrates (General Degradation): Whalley, Hawkes, Paisley & Trigg (WHPT) metric in River Invertebrate Classification Tool (RICT). July 2014. ISBN: 978-1-906934-49-1.

---

## **Appendix 1: Aquatics Team Report with Figures**



**Electrofishing and Aquatic  
Macroinvertebrates Surveys**

**Gatwick Fish and  
Aquatic  
Macroinvertebrate  
Surveys**

**Final Report**

For

**ECUS Ltd**

Project No.: A-ECU-101/001

June 2021



**London & South East**  
Compass House  
Surrey Research Park  
Guildford  
GU2 7AG . UK  
t: +44 (0)1483 466 000

**North & Borders**  
The Tannery  
91 Kirkstall Road  
Leeds  
LS3 1HS. UK  
t: +44 (0)113 247 3780

**Wales & South West**  
Sophia House  
28 Cathedral Road  
Cardiff  
CF11 9LJ UK  
t: +44 (0) 2920 660180

**Midlands**  
Edmund House  
12-22 Newhall Street  
Birmingham  
B3 3AS  
t: +44 (0) 121 726 3494

**Scotland**  
20-23 Woodside Place  
Glasgow  
G3 7QF . UK  
t: +44 (0)141 582 1333

**Enquiries**

e: [enquiries@thomsonec.com](mailto:enquiries@thomsonec.com)

w: [www.thomsonec.com](http://www.thomsonec.com)



---

Project Number	Report No.
A-ECU-101/001	001

Revision No.	Date of Issue	Author	Reviewer	Approver
001	05/01/2021	Alex Charlesworth	Tessa Harding	Tessa Harding
002	16/06/2021	Alex Charlesworth	Tessa Harding	Tessa Harding
003				

**Disclaimer:**

Copyright Thomson Environmental Consultants Limited. All rights reserved.

No part of this report may be copied or reproduced by any means without prior written permission from Thomson Environmental Consultants Limited. If you have received this report in error, please destroy all copies in your possession or control and notify Thomson Environmental Consultants Limited.

This report has been prepared for the exclusive use of the commissioning party and unless otherwise agreed in writing by Thomson Environmental Consultants Limited, no other party may use, make use of or rely on the contents of the report. No liability is accepted by Thomson Environmental Consultants Limited for any use of this report, other than for the purposes for which it was originally prepared and provided.

Opinions and information provided in the report are on the basis of Thomson Environmental Consultants Limited using due skill, care and diligence in the preparation of the same and no explicit warranty is provided as to their accuracy. It should be noted and it is expressly stated that no independent verification of any of the documents or information supplied to Thomson Environmental Consultants Limited has been made.

---

## Contents

1. Summary and Main Recommendations .....	5
1.1 Summary .....	5
1.2 Conclusions .....	6
1.3 Main Recommendations .....	7
2. Introduction .....	8
2.1 Development Background .....	8
2.2 The Brief and Objectives .....	8
2.3 Background to Watercourses .....	8
3. Methodology .....	12
3.1 Desk study .....	12
3.2 Survey: Macroinvertebrates .....	12
3.3 Macroinvertebrate data analysis .....	13
3.4 Survey: Fish .....	15
3.5 Limitations .....	16
4. Results .....	18
4.1 Desk study .....	18
4.2 Field data .....	22
<i>Fish</i> .....	27
5. Discussion .....	28
6. Conclusions and recommendations .....	31
6.1 Conclusions .....	31
6.2 Recommendations .....	32
6.3 Further Survey .....	33
7. References .....	34
Appendix 1 .....	35

Figure 1a: Sampling locations on R Mole

Figure 1b: Sampling locations on Gatwick Stream

Figure 2a: River Mole 1km desk study search area and priority habitats

Figure 2b: Gatwick Stream 1km desk study search area and priority habitats

Figure 3: Location of shining ramshorn snail (*Segmentina nitida*) record

---

## 1. Summary and Main Recommendations

### 1.1 Summary

- 1.1.1** ECUS Ltd are undertaking Ecological Assessment work for a proposed surface water management and flood alleviation scheme that will affect two sites close to Gatwick Airport, one on the River Mole and the other on the Gatwick Stream. The River Mole may be re-meandered and land close to the river may be re-profiled to increase flood storage. The Gatwick Stream has already had some surrounding land re-profiled for flood storage, however this area may be expanded to encompass both sides of the river. Thomson Environmental Consultants Aquatics Team were commissioned by ECUS to undertake aquatic macroinvertebrate and fish baseline surveys and a desk study, to inform the proposals.
- 1.1.2** The study area encompasses two watercourses; a 1.3km stretch of the River Mole immediately downstream of where it emerges from under the runway at Gatwick Airport and a 750m stretch of the Gatwick Stream (a tributary of the River Mole) upstream of the Crawley sewage works. A 100m survey section was identified on each watercourse from an initial walkover survey conducted in June 2020. Three survey visits during 2020 were undertaken for aquatic macroinvertebrates (spring, summer and autumn) and two for fish (spring and autumn). The spring survey visit was delayed until early July due to restrictions related to the Covid 19 outbreak. Desk study data was obtained from the Sussex Biological Records Centre and the Environment Agency on behalf of Ecus.
- 1.1.3** The desk study returned one record from 2013 of the shining ram's-horn snail (*Segmentina nitida*) within the study section on the River Mole (TQ 25623 40908). The species is nationally scarce<sup>1</sup>, a UK Priority Species under the UK Post 2010 Biodiversity Framework and listed on the Sussex Rare Species Inventory. It was not recorded during the surveys for this study, although the survey section did not coincide with the reach in which the snail was previously recorded. The desk study returned records of two fish species for the River Mole; bullhead and brown trout.
- 1.1.4** **River Mole** - A mean of 19.3 macroinvertebrate taxa were recorded at the River Mole site across the three survey visits. Biotic indices measuring water quality (Biological Monitoring Working Party (BMWP) score and Average Score Per Taxon) indicate moderately polluted conditions in the River Mole. Lotic invertebrate Index for Flow Evaluation (LIFE) scores suggest that the macroinvertebrate community is characteristic of sluggish flow conditions and low Proportion of Sediment intolerant Invertebrates (PSI) scores indicate heavily sedimented conditions.
- 1.1.5** A total of 415 fish were caught on the River Mole in spring after three survey 'runs' compared with only 28 fish caught in autumn with the same level of effort. Roach (*Rutilus rutilus*) were the most abundant fish species identified (252) in spring and in autumn (13). The study stretch on

---

<sup>1</sup> Occurring in 16-100 hectads in Great Britain. Excludes rare species qualifying under the main IUCN criteria. This category replaces Notable, Notable A and Notable B.

the River Mole lies within open floodplain grassland with no shading, which means that water temperature and therefore dissolved oxygen (DO), fluctuated considerably. Extensive stands of submerged and emergent macrophyte plants occur through the study section and their decomposition are likely to be contributing to low DO in the autumn. These DO conditions coupled with organic pollution from within the catchment is considered to be influencing the composition and abundance of both the aquatic macroinvertebrate and fish communities present. Predatory fish such as pike are able to exploit the dense macrophyte stands and are further reducing populations of cyprinid fish.

**1.1.6 Gatwick Stream** - Environment Agency data from 3 sites on the Gatwick Stream indicate that the study section is of moderate to poor water quality, with sluggish flow and sedimented condition. Fewer macroinvertebrate taxa were recorded at the two Gatwick Stream sites compared with River Mole (mean of 10 taxa per visit). BMWP and ASPT scores indicate moderate water quality conditions at the upstream and poor to very poor at the downstream site. A high LIFE score for the upstream site during the spring visit suggests that velocities are high in the early part of the season and decline through the summer and autumn. PSI scores for the upstream site fluctuated considerably across the three seasons, from only slightly sedimented conditions in spring to sedimented condition in autumn.

**1.1.7** A total of 300 and 317 fish were caught in spring and autumn respectively at the Gatwick Stream site after three survey 'runs'. Chub was the most abundant species in the spring survey and dace in the autumn. Shading of the channel by overhanging trees meant that water temperature was relatively consistent and dissolved oxygen remained high throughout the three seasons.

## **1.2 Conclusions**

**1.2.1** Both watercourses supported macroinvertebrate communities indicative of moderately polluted conditions, exacerbated by relatively low flow conditions and high levels of sedimentation. Dense macrophyte growth on the River Mole is contributing to acute reductions in dissolved oxygen which are impacting on the macroinvertebrate assemblage.

**1.2.2** The presence of one record from 2013 of shining ram's-horn snail, an IUCN Red List species and UK species of principal importance under the 2006 NERC Act, has implications for the design of the River Mole scheme. Although not recorded during the survey, there remains a possibility that the species may occur at the site of the 2013 record at the downstream end of the desk study area. A targeted survey is required to determine its potential presence.

**1.2.3** The Gatwick Stream appears to be impacted by both organic pollution and silt deposition, possibly from a storm water discharge outlet from a nearby industrial area. Consistently high populations of fish caught in spring and in autumn are likely to be a consequence of stable temperature and dissolved oxygen conditions caused by shading and potentially high abundances of pollution tolerant macroinvertebrates such as Oligochaete worms as a food source.

---

### 1.3 Main Recommendations

#### 1.3.1 The main recommendations are set out below:

##### **River Mole**

- Undertake survey to establish presence/absence of shining ramshorn snail. Survey to focus initially on the site of the 2013 record and if found to be present, extended to incorporate the whole study section. Surveys should be scoped and undertaken by a specialist mollusc ecologist.
- If shining ramshorn snail found to be absent from main channel, undertake vegetation removal/dredging from the central channel/thalweg of the River Mole in selected sections. Marginal berms should be retained on alternate sides of the channel throughout the dredged section for the re-establishment of emergent vegetation. Brushwood faggots or fascines anchored with wooden stakes can be used to maintain the riverward edge of the berm and prevent silt from slumping into the main channel.
- Off-line scrapes and shallow ponds could be created within the floodplain grassland area to provide habitat for wetland birds. If shining ramshorn snail is found to be present this recommendation can be adapted to incorporate new, permanently wet ditches supporting dense emergent reed vegetation.
- If shining ramshorn snail is found to be absent it is advised that some level of routine maintenance of macrophyte and bankside vegetation is undertaken annually under an appropriate management plan.
- Before any in-channel works begin, it is advised that a fish rescue and exclusion or translocation is undertaken to safeguard fish populations.
- Stop nets should be installed at either end of the site proposed for in-channel works to prevent access by any fish species whilst the works are on-going.

##### **Gatwick Stream**

- Identify point sources of pollution from industrial area associated with Crawley STW, including storm drains and surface water discharge points from roads and urban areas.
  - Consider SUDS scheme to address these discharges including settlement ponds and reedbed treatment systems which would have additional biodiversity benefit.
- 1.3.2 Before any in-channel works begin it is advised that a fish rescue survey is undertaken to safeguard fish populations in the affected area. It will also be necessary to install stop nets at either end of the reach where in-channel works will be undertaken to prevent access by any other fish species whilst the works are ongoing.
- 1.3.3 Hydrometric surveys should be undertaken at various points along both rivers to better understand present hydrological conditions and inform plans to modify the channels.

## 2. Introduction

### 2.1 Development Background

**2.1.1** Two watercourses, the River Mole and Gatwick Stream will be directly affected by proposals for a surface water management and flood alleviation scheme to the east and west of Gatwick Airport. The scheme may include proposals to re-meander the River Mole close to where it emerges from beneath the airport runway and create new flood attenuation areas to the west of the watercourse. New flood storage has already been created to the west of the Gatwick Stream, with further areas likely planned within the floodplain to the east of the watercourse.

**2.1.2** The study area encompasses two watercourses; a 1.3km stretch of the River Mole immediately downstream of where it emerges from under the runway at Gatwick Airport and a 750m stretch of the Gatwick Stream (a tributary of the River Mole) upstream of the Crawley sewage works.

### 2.2 The Brief and Objectives

**2.2.1** ECUS Ltd commissioned Thomson Environmental Consultants Aquatic Team in May 2020 to undertake fish and aquatic macroinvertebrate surveys of the two rivers within the proposed site. The brief was to:

- To determine baseline populations for both fish and aquatic macroinvertebrates in these two watercourses over the course of a year.
- Carry out a desk study for the surrounding areas of both sites including a 1km perimeter.
- Provide a report on the surveys giving the methods and results of the surveys, with recommendations, including opportunities for enhancement, mitigation and further surveys.

### 2.3 Background to Watercourses

**2.3.1** The River Mole rises in Baldhorns Copse in West Sussex and discharges into the River Thames at the town of Molesey in Surrey. The Mole catchment flows over the Wealden and London clays, however, between Dorking and Leatherhead, the river cuts its way through the North Downs chalk. In this area part of the river water disappears through holes in the underlying chalk feeding into the groundwater aquifers before flowing back into the river near to Leatherhead. This action has been suggested as the origin to the name of this river, but is more likely attributed to the fact it meets the Thames at Molesey,

**2.3.2** Approximately 7 miles downstream of the source, the River Mole reaches the boundary of Gatwick Airport where it passes beneath the runway in a culvert. The reach that will be affected by the proposed scheme extends 1.3km downstream from where the river emerges

from beneath the airport runway (Figure 1a). The survey stretch of the Gatwick Stream surveyed (TQ291398) lies upstream of the Crawley sewage works (Figure 1b).







Figure 1a: Thomson Environmental Consultants - Ordnance Survey Data and Thomson Environmental Consultants - Thomson Ecology Sampling Sites - LF\_041220.mxd



## 3. Methodology

### 3.1 Desk study

- 3.1.1** A study area was defined as an area that encompassed the site and all land within 1 km of the perimeter of each of the sites, (Figures 2a and 2b). Records of designated sites and protected or otherwise notable species were then sought for both study areas.
- 3.1.2** Sources of information were as state in Table 3-1.

**Table 3-1: Sources of data**

Data type	Source
Statutory sites for nature conservation related to the river environment	Multi-Agency Geographical Information for the Countryside (MAGIC) <a href="https://magic.defra.gov.uk/magicmap.aspx">https://magic.defra.gov.uk/magicmap.aspx</a>
Non-statutory sites for nature conservation, protected and notable species and invasive and non-native species (fish and macroinvertebrates only)	Sussex Biodiversity Records Centre
Background information on Water Framework Directive status	<a href="https://environment.data.gov.uk/catchment-planning">https://environment.data.gov.uk/catchment-planning</a>
Macroinvertebrate, fish and invasive and non-native species data	Environment Agency data request (EA Analysis and Reporting)

- 3.1.3** A request for information was sent to the Sussex Biological Records Centre on 07/10/2020 with responses requested by 20/10/2020. The boundaries of any designated site and records of species were sought for part of the study area encompassing the site and within 1km of the perimeter of each of the sites.
- 3.1.4** The records included in this report are those relating to fish and macroinvertebrates. Records over 10 years old have been excluded.

### 3.2 Survey: Macroinvertebrates

- 3.2.1** A representative 100m section on each watercourse was identified from a walkover survey conducted prior to the spring sampling visit. Two sampling locations were identified on the Gatwick Stream, one at the upstream and one at the downstream end of the 100m section (Figure 1a and 1b). Only one sampling at the upstream end of the reach was safely accessible on the River Mole.

- 3.2.2** Samples were collected at each of the sites using the Whalley Hawkes Paisley Trigg (WHPT) method comprising of a standard three-minute kick sample using a long-handled pond net with 1mm mesh size, which was supplemented by a one-minute hand search (Environment Agency, 2017). Sampling of habitats within the three-minute kick/sweep sampling were in proportion to their occurrence. Samples were then preserved in industrial methylated spirits (IMS) for processing in the laboratory to the requirements outlined in EA Operational Instruction 024\_08 Freshwater macroinvertebrate analysis of riverine samples (Environment Agency, 2014).
- 3.2.3** Macroinvertebrates were identified to Mixed Taxon Level 5, to enable evaluation of the macroinvertebrate community and calculation of the relevant biotic indices including Biological Monitoring Working Party (BMWP), Average Score Per Taxon (ASPT) and Lotic-Invertebrate index for Flow Evaluation (LIFE). Proportion of Sediment-sensitive Invertebrates (PSI) and Community Conservation Index (CCI).
- 3.2.4** Macroinvertebrate sampling was undertaken in spring, summer and autumn on the dates presented in Table 3-2.

**Table 3-2: Macroinvertebrate survey dates.**

Macroinvertebrate survey visit	Date
Spring	04/06/20
Summer	29/07/20
Autumn	29/09/20

### 3.3 Macroinvertebrate data analysis

- 3.3.1** The macroinvertebrate abundance data collected during the field surveys and background data from the Environment Agency has been analysed using a range of biotic indices. Each of the indices used in the analyses are summarised below.

#### **Biological Monitoring Working Party (BMWP) score**

- 3.3.2** The BMWP score is a method for indexing river water quality in England and Wales using macroinvertebrate families. Originally published in the early 1980's, the system was updated in 2013 based on a more robust baseline data set (Paisley et al, 2013). A score of between 1 and 10 is assigned to families found within a sample based on their tolerance to organic pollution, with a score of 1 indicating high tolerance, and 10 indicating low tolerance. Low scoring families include worms (Oligochaeta) and midge larvae (Chironimidae), whilst the presence of mayfly (Ephemeroptera) and stonefly (Plecoptera) larvae is indicative of clean water conditions. The scores for each family recorded in the sample are summed to give the overall BMWP site score. Since the overall site score is influenced by the number of families as well

as the scores of the individual families in the sample, an average is taken by dividing the overall BMWP score by the number of families/taxa in the sample. This is termed the Average Score Per Taxon (ASPT).

Table 3-3 provides an interpretation of the BMWP scoring system.

**Table 3-3: BMWP Scoring System**

BMWP score	Category	Interpretation
0-10	Very poor	Heavily polluted
11-40	Poor	Polluted or impacted
41-70	Moderate	Moderately impacted
71-100	Good	Clean but slightly impacted
>100	Very good	Unpolluted, unimpacted

### River Invertebrate Classification Tool (RICT)

- 3.3.3** BMWP and ASPT has largely be superseded by the River Invertebrate Classification Tool (RICT), which is one of the parameters used for classifying rivers according to their ecological status under the Water Framework Directive (WFD). The scores derived for an individual site under RICT are compared with those expected under unpolluted conditions (known as reference conditions) in order to give an Environmental Quality Ratio (EQR). This aims to take account of the variability of macroinvertebrate families in rivers resulting from environmental parameter such as altitude, underlying geology and proximity to the river source.

### Lotic invertebrate Index for Flow Evaluation (LIFE) Score

- 3.3.4** The LIFE score system links flow conditions in rivers, and specifically flow velocity, with commonly identified macroinvertebrate species and families (Extence *et al.* 1999). Macroinvertebrates are assigned to one of 6 groups depending on their tolerance to low flow conditions. The groups range from I comprising taxa associated with rapid flow conditions (>100cm s<sup>-1</sup>) to VI including those associated with drying or drought impacted sites. A flow score is obtained for each species/taxon by combining the flow category with an estimated abundance score as described by Extence *et al.* (1999). The LIFE score for a sample is obtained by summing the individual flow scores for each taxon by the number of taxa in the sample. LIFE scores range from 1 to 12, with scores of 8 or above indicating moderate to high flow conditions, and scores of 7 or below indicating sluggish conditions.

### Proportion of Sediment sensitive Invertebrates (PSI)

- 3.3.5** The PSI index provides an indication of the extent to which watercourses have been impacted by the deposition of fine sediment (Extence et al, 2017). Following the same principle as the LIFE score system, invertebrates are assigned to one of four groups depending on their sensitivity to fine sediment, with Group A comprising highly sensitive taxa, and Group D those

that are highly insensitive. The method also requires a log abundance category to be estimated for all taxa identified in a sample (1-9, 10-99, 100-999 and 1000+ individuals present). Scores range from 80 -100 for unsedimented sites down to 0-20 for highly sedimented sites (Table 3-4).

**Table 3-4:- Interpretation of PSI scores**

PSI score	River bed condition
81-100	Minimally sedimented/unsedimented
61-80	Slightly sedimented
41-60	Moderately sedimented
21-40	Sedimented
0-20	Heavily sedimented

### Community Conservation Index (CCI)

The CCI combines the rarity of constituent species in a sample with the diversity of the community, or community richness, to give a single integrated score which can be used as the basis for site evaluating (Chadd and Extence, 2004). Species identified from a survey site or area are given a Conservation Score (CS), based on standard rarity categories, with Red Data Book 1 (Endangered) species scoring 10, and very common species scoring 1. The sum of each of the conservation scores in the sample is then divided by the number of contributing species to give the overall CCI score.

### 3.4 Survey: Fish

**3.4.1** The surveys were undertaken using the catch depletion method in order to assess species composition, age structure and to estimate population size. Surveys were undertaken by an accredited electric fishing team comprising three members of staff. Surveys and analysis conformed to the relevant guidance outlined in BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity (British Standards, 2003). An FR2 consent (application to use fishing instruments other than rod and line) was sought from the Environment Agency prior to conducting the survey.

**3.4.2** The survey was undertaken over a 100m reach and there was one survey reach per watercourses, coinciding with the macroinvertebrate survey locations on both watercourses. Stop nets were installed across the channel at either end of the reach to prevent fish entering

or leaving the survey area. Holding containers for captured fish were established in a small boat with an aerator installed to provide oxygen to captured fish.

- 3.4.3** The survey was undertaken using an electrofishing box alternating between a single anode and two anodes depending on the width of the river in order to maximise catch efficiency. One surveyor, operating the electrofishing anode waded from downstream to upstream and a second surveyor netted any stunned fish. In areas where the rivers was wider the second surveyor also operated an anode. The operatives were followed by an additional surveyor pulling a small boat with the electrofishing box and holding tank on board, and also equipped with a hand net to maximise the catch rate. At the end of each run all caught fish were identified, measured and placed in a submerged holding net to facilitate their recovery and prevent re-capture.
- 3.4.4** Two survey visits were undertaken, one in spring (04/06/20) and one in autumn (29/09/20) to establish a baseline of the species composition on the two watercourses. Undertaking the autumn visit in September ensured that air temperatures are above the minimum of 10 degrees and minimise the risk of high flow conditions. It would also avoid risk of disturbance to salmonid spawning habitat, should it be present.

### **3.5** Limitations

- 3.5.1** Only one macroinvertebrate sample could be retrieved from the downstream River Mole site, due to various access issues, such as, dense bankside vegetation creating a barrier to the river and steep banks, which prevented safe access and egress to the river.
- 3.5.2** The River Mole has exceptionally high coverage of aquatic plants, which made electrofishing difficult. In spring the filamentous algae blanket weed (*Cladophera* agg.) was found in dense clumps making progress slow, as the anode became blanketed by the filamentous algae each time it was placed in to the water and needed regular clearing in order to progress. In addition to this, the macroinvertebrate surveys were difficult due to the dense macrophyte growth and deep waters preventing more than one macroinvertebrate sample being taken using the WHPT method.
- 3.5.3** The timing of the spring survey was delayed by a nationwide lockdown related to the COVID-19 outbreak.



**Figure 2a. River Mole 1km desk study search area and priority habitats**



**Figure 2b. Gatwick stream 1km desk study search area and habitats priority habitats**



---

## 4. Results

### 4.1 Desk study

#### River Mole

##### *Environment Agency: Water Framework Directive status*

**4.1.1** Under the Water Framework Directive (WFD) rivers and standing waters are termed waterbodies and are classified according to their ecological status. Ecological status is classified using five categories of high, good, moderate, poor and bad and is measured and classified via a range of inter-linked biological, physico-chemical and physical (morphological) parameters. The classification process is based primarily on the biological quality elements of the water body but considered alongside support elements covering physico-chemical standards and hydromorphological quality elements. Each of these supporting elements is assigned to a status category (i.e. high to bad). The overall status of the waterbody is based on the status category of the worst supporting element.

**4.1.2** The affected reach of the R Mole falls within the WFD waterbody named 'Mole Upstream of Horley (GB106039017481)'. There is little information relating to the stretch of the River Mole. It was first classified as good under the WFD classification system in 2015, although the most recent classification in 2019 designates it as moderate. Although the biological quality elements are classified as good (based on fish data only), one of the physico-chemical quality elements (phosphorous) is classified at moderate status, and therefore the overall waterbody status is classed as moderate.

**4.1.3** No Environment Agency background records were received for the River Mole.

##### *Sussex Biological Records Centre data*

**4.1.4** A total of 3 records of fish species were returned from the Sussex Biological Records Centre for the River Mole within 1km of the study section, comprising one record of bullhead (*Cottus gobio*) approximately 0.5km downstream from the study section in 2014, and a record of 2 adult brown trout (*Salmo trutta subsp. Fario*) within the survey section in 2016. Bullhead is listed as a non-priority species under Annexe 2 of the EU Habitats Directive and listed on the Sussex Rare Species Inventory. Brown trout is a UK Priority Species under the UK Post 2010 Biodiversity Framework, and Section 41 of the Natural Environment and Rural Communities Act 2006.

**4.1.5** There is one record of the shining ram's-horn snail (*Segmentina nitida*) within the study section (TQ2562340908; Figure 3) from February 2013. The species was recorded as being 'u/s Pond M and Tributary'. The shining ram's-horn snail is nationally scarce<sup>2</sup>, a UK Priority Species

---

<sup>2</sup> Occurring in 16-100 hectads in Great Britain. Excludes rare species qualifying under the main IUCN criteria. This category replaces Notable, Notable A and Notable B.

under the UK Post 2010 Biodiversity Framework, and listed on the Sussex Rare Species Inventory.

- 4.1.6** One notable dragonfly species, common sympetrum (*Sympetrum striolatum*) was recorded within 1km of the site. The species is listed in the UK Red Data Book. A total of 44 observations were made of the species in the vicinity of the Gatwick airport, with a number of them within the study section. There are no records of the larvae in the River Mole, either from the Sussex Biological Records Centre or the Environment Agency, and therefore breeding sites are unclear.
- 4.1.7** A number of invasive and non-native invertebrate and aquatic plant species occur within 1km of the River Mole study section. Three records of signal crayfish (*Pacifastacus leniusculus*) were returned from within 1km of the study section between 2011 and 2013. Two of the sites were on the River Mole within the study section and the third north of the Gatwick runway within a tributary of the R Mole. Signal crayfish is listed on Schedule 9 of the Wildlife and Countryside Act. Under Section 14 of the Act it is an offence 'to release or allow to escape into the wild' any species listed under Schedule 9'.
- 4.1.8** Records of several invasive aquatic plant species were also recorded within 1km of the site including Nuttall's pond-weed (*Elodea nuttallii*), Japanese knotweed (*Fallopia japonica*) and Himalayan balsam (*Impatiens glandulifera*).


### **Gatwick Stream**

#### *Environment Agency: Water Framework Directive status*

- 4.1.9** The Gatwick Stream is a tributary of the River Mole and is approximately 12km in length. It rises near Three Bridges and joins the River Mole near the centre of Horley. It falls within the Tilgate Stream and Gatwick Stream at Crawley WFD waterbody (GB106039017500). The overall waterbody status has remained moderate since 2013, although the biological quality elements are assigned bad status on the basis of fish data. This is a deterioration from poor status in 2016. Macroinvertebrates were classified at poor status in 2019 and have remained at that classification since 2013. Sewage discharges and the invasive signal crayfish are given by the Environment Agency as the reasons for poor biological quality in the brook.

#### *Environment Agency: Macroinvertebrate data*

- 4.1.1** Macroinvertebrate data was received from the Environment Agency for 3 sites on the Gatwick Stream that lie within the study area (U/S Crawley STW (TQ 29160 39780); Downstream Tinsley Bridge (Flylife site) (TQ 29129 39864) and At Tinsley Bridge, Tinsley Green (TQ-29150-39800)). One sample was collected at U/S Crawley STW in October 2017, and duplicate samples were collected at Downstream Tinsley Bridge (Flylife site) and At Tinsley Bridge, Tinsley Green) in March 2019 (Figure 1b). The Environment Agency have provided feedback that the 2019 samples were taken in response to a pollution incident and that the duplicate sample from both sites with lower number of taxa recorded was sorted on the bank. In comparing the data with this study only the laboratory sorted sample has been considered, although the results for both samples are presented in Table 4-1.



- 4.1.2** A total of 13 families were recorded during the survey at the U/S Crawley STW site in October 2017. The freshwater shrimp *Gammarus pulex*, a species indicative of moderate water quality, was the most numerous. However, the site also supported relatively high numbers of Oligochaete worms, a family highly tolerant of low oxygen conditions. BMWP and ASPT scores have been calculated since none were provided by the Environment Agency (Table 4-1). This site had a BMWP score of 43 with an ASPT of 4.3 indicating moderate to poor water quality.
- 4.1.3** Of the two duplicate samples taken at the Downstream Tinsley Bridge (Flylife site) (TQ-29129-39864) on 14<sup>th</sup> March 2019 a total of 22 families were recorded in one sample and 7 in the other. Midge larvae (Chronomidae) and Oligochaete worms were present in relatively high numbers in the second sample indicating poor water quality. Based on the second sample the site had a BMWP of 47 and ASPT of 3.92.
- 4.1.4** At the most upstream site at Tinsley Bridge (TQ-29150-39800) site a total of 8 families were recorded in one of the duplicate samples and 21 in the second. In general, the samples supported pollution tolerant families and species such as Oligochaeta (20 and 40 individuals respectively) and the water louse (*Asellus aquaticus*) (30 individuals in the second sample). However, the site also supported the damselfly larvae (*Calopteryx* sp.), a relatively pollution sensitive family. The second sample at this site had a BMWP of 48 and ASPT of 4.

**Table 4-1: EA macroinvertebrate biotic scores for Gatwick Stream**

Site	U/S Crawley STW	Downstream Tinsley Bridge (Flylife Site)	Downstream Tinsley Bridge (Flylife Site)	At Tinsley Bridge. Tinsley Green	At Tinsley Bridge. Tinsley Green
<b>Date</b>	12/10/2107	14/03/2019	14/03/2019	14/03/2019	14/03/2019
<b>BMWP (TL1)</b>	43	15	47	20	48
<b>ASPT</b>	4.3	3.00	3.92	3.33	4
<b>LIFE</b>	7.5	7.0	7.11	7.00	7.00
<b>PSI</b>	40.00	28.57	36.00	36.36	32.14
<b>CCI</b>	1.00	N/A	1.00	N/A	1.00

- 4.1.5** LIFE scores for each of the 3 sites ranged from 7.0 to 7.5 indicating sluggish to moderate flow conditions. PSI scores for all three sites indicate sedimented conditions, although the U/S

Crawley STW sites is close to moderately sedimented with a score of 40. CCI scores of 1 indicate low conservation value.

*Sussex Biological Records Centre data*

- 4.1.6** Records of two fish species, bullhead and brown trout were returned for the Gatwick Stream. Bullhead from Sussex Biological Records Centre. One adult bullhead was recorded within the study section in October 2015, and a brown trout in a similar location in July 2016.
- 4.1.7** A total of 15 records of adult common sympetrum dragonflies were returned for the study section on the Gatwick Stream between 2012 and 2017, although there are no records of the larvae. Six records of the downy emerald dragonfly (*Cordulia aenea*), and 2 of the brilliant emerald dragonfly (*Somatochlora metallica*) were returned from within the past 10 years. Downy emerald dragonfly is a Red List species on the IUCN Red List, and a Priority Species on the UK Post 2010 Biodiversity Framework. The downy emerald dragonfly is listed on the Sussex Rare Species Inventory. None of the records were on the Gatwick Stream and there are no records of the larvae.
- 4.1.8** There were three records of the invasive signal crayfish from within the study section on the Gatwick Stream in 2017. The invasive aquatic plant, Nuttall's pond weed was recorded within the study section in 2016, and there are records of Japanese knotweed and Himalayan balsam.

*Environment Agency: Fish data (R Mole and Gatwick Stream)*

- 4.1.9** Data provided by the Environment Agency indicates that both the Gatwick Stream and River Mole were stocked in 2018 and 2019 with Roach, Barbel, Dace, and Chub. In 2018, 3200 fish were added to the lower Gatwick Stream in response to a pollution event which occurred in 2017.
- 4.1.10** In 2019, 3600 fish were stocked in the River Mole in response to a prolonged dry weather event in 2018, which occurred as a result of low flows and first flush effect, which was estimated to have affected approximately 2000 fish.

**4.2 Field data**

**River Mole**

*Water Quality*

- 4.2.1** A maximum temperature of 17.6°C was recorded at the sampling site on the River Mole during the summer visit on 29<sup>th</sup> July 2020 (Table 4-2). The temperature was only slightly lower on the first visit (16.4°C on 1<sup>st</sup> July), which was delayed due to Covid 19 restrictions. Water temperature dropped to 13.8°C by the autumn visit on 29<sup>th</sup> September 2020. Dissolved oxygen concentrations dropped sharply between the first and second visits, from 60.8% in early July to 17% on 29<sup>th</sup> July, before recovering slightly to 28.7% by end of September. Both

conductivity and turbidity increased progressively through the season. Conductivity increased from 358.5 to 471  $\mu\text{S}/\text{cm}$ , whilst turbidity increased from 3.78 to 4.3 NTU.

**Table 4-2: Water quality data recorded at River Mole sampling site**

Season	Temperature (°C)	DO (%)	DO (mg/L)	pH	Conductivity ( $\mu\text{S}/\text{cm}$ )	Turbidity (NTU)
Spring	16.4	60.8	5.94	7.3	358.5	3.78
Summer	17.6	17.0	1.6	7.19	341.0	4.02
Autumn	13.8	28.2	3.03	8.38	471.0	4.3

### Macroinvertebrates

**4.2.2** A mean of 19.3 taxa were recorded at the River Mole site across the three visits. There was relatively little variation in the number of taxa recorded on each visit, with the maximum of 21 in the spring/early summer sample, and a minimum of 17 in the summer sample (Table 4-34-3). Of these, 12 taxa/species occurred in all three samples, including the water shrimp *Gammarus pulex*, the pea mussel, *Sphaereum corneum* and the mayfly larvae *Cloeon dipterum*. However, abundances of individual taxa within the samples varied considerably across the 3 visits, with the crustacean Cladocera the most abundant in the early summer samples 01/07/20, replaced by the water boatman, Corixidae one month later. The most abundance taxa in the autumn samples was the Isopod *Asellus aquaticus* (waterlouse). These changes in abundance are likely to be driven by seasonal changes in life stage from early to later (larger, and therefore more readily sampled) larval instars as well as the availability of food resources.

**Table 4-3 Number of macroinvertebrate species/taxa recorded at River Mole and Gatwick Stream sites**

	Spring	Summer	Autumn
River Mole	21	17	20
Gatwick Stream upstream	12	10	13
Gatwick Stream downstream	8	8	9

**4.2.3** The consistent occurrence of low BMWP scoring (i.e. 3 or below) species and taxa such as the waterlouse *A. aquaticus*, Chironimidae and Oligochaeta on all three visits suggest that the watercourse is affected by organic pollution. This is confirmed by BMWP scores of 44, 46 and 49, and ASPT of 3.73, 3.45 and 3.43 in the spring, summer and autumn samples respectively indicating moderately polluted conditions.

**Table 4-4. Macroinvertebrate biotic indices**

Biotic Index	Spring			Summer			Autumn		
	U/S River Mole	Gatwick Brook U/S	Gatwick Brook D/S	River Mole U/S	Gatwick Brook U/S	Gatwick Brook D/S	U/S River Mole	Gatwick Brook U/S	Gatwick Brook D/S
	01/07/2020	01/07/2020	01/07/2020	27/07/2020	27/07/2020	27/07/2020	29/09/2020	29/09/2020	29/09/2020
BMWP (TL1)	44	46	14	46	37	29	49	41	20
LIFE (TL5)	6,25	8,17	7,5	6,10	7,40	7,75	5,87	6,75	8
ASPT (TL2)	3,73	4,92	3,50	3,45	4,53	3,91	3,43	4,13	2,88
PSI (TL5)	10,00	66,67	14,29	5,00	41,67	50,00	6,25	21,05	33,33
CCI (TL5)	5,50	4,50	0	4,00	5,00	1,00	9,62	1,20	1

**4.2.4** LIFE scores for the R Mole ranged from 6.25 in the spring/early summer sample, 6.1 in the summer sample and 5.87 in the autumn sample, indicating sluggish flow conditions (Table 4-44-4). The decline in LIFE scores over the summer period are likely to be primarily a result of low flow conditions due to low summer rainfall, although extensive macrophyte beds in the channel may also be impeding flow. Low PSI scores of less than 20 also indicate heavily sedimented conditions. This correlates with low flow velocities in the channel indicated by the LIFE scores, and is likely to be exacerbated by the extensive macrophyte beds.

**4.2.5** CCI scores of between 5 and 10 indicate that the macroinvertebrate community is of moderate conservation value. The presence of *Sigara limitata*, a species of water boatman, contributed to a slightly higher score of 9.62 in the autumn sample.

#### Fish

**4.2.6** A total of 415 fish were caught on the River Mole in spring after three runs compared with only 28 fish caught in autumn with the same level of effort. Roach (*Rutilus rutilus*) were the most abundant fish species identified (252) in spring and in autumn (13).

**4.2.7** The size range of species caught on the electrofishing surveys in spring (Table 4-54-5) suggests that there are multiple age classes of each species, ranging from juveniles to mature adults. The stretch of the River Mole sampled in this study appears to be a good breeding and spawning environment for Roach and Perch, due to its slow flow environment and dense vegetation. The mean size data in spring would suggest that this stretch also appears to be a good environment for juvenile and sub-adult Chub and Dace as well as providing optimal foraging habitat for predatory fish species such as Pike.

**Table 4-5: River Mole Fish Survey Data**

River Mole					
Spring					
Species	Latin name	Abundance	Mean size (mm)	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	45	166.55	77	386
Roach	<i>Rutilus rutilus</i>	252	106.91	45	256
Dace	<i>Leuciscus leuciscus</i>	37	127.86	59	203
Pike	<i>Esox lucius</i>	14	344.86	108	595
Perch	<i>Perca fluviatilis</i>	46	130.00	73	258
Bream	<i>Abramis brama</i>	3	72.33	62	79
Tench	<i>Tinca tinca</i>	2	89.0	85	93
Gudgeon	<i>Gobio gobio</i>	13	93.1	82	109
Rudd	<i>Scardinius erythroplthalmus</i>	2	138.50	81	196
Roach/ Bream Hybrid		1	143	143	143
Autumn					
Species	Latin name	Abundance	Mean size (mm)	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	3	217.67	181	289
Roach	<i>Rutilus rutilus</i>	13	123	64	200
Tench	<i>Tinca tinca</i>	7	198	153	248
Pike	<i>Esox lucius</i>	4	121	110	127
Perch	<i>Perca fluviatilis</i>	1	86	86	86

## Gatwick Stream

### Water quality

- 4.2.8** Water temperature at the two Gatwick Stream sites remained relatively consistent across the three seasons (Table 4-74-6), peaking at 16.7°C at the downstream site during the summer visit on 29<sup>th</sup> July. The lowest temperature was recorded at the upstream site (14.8°C) at the end of September. The sites are moderately shaded by overhanging trees, which will help to buffer water temperature. Dissolved oxygen concentrations also remained relatively high at over 70% throughout the three seasons, reaching a maximum of 78.7% at the downstream site in autumn. Turbidity was relatively high compared with the River Mole site, with a minimum of 5.95NTU at the upstream site in autumn and a maximum of 11.85NTU at the upstream site in summer.



**Table 4-6: Water quality data for Gatwick Stream**

Spring						
Site	Temperature (°C)	DO (%)	DO (mg/L)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Gatwick Stream US	15.4	73.5	7.34	7.53	276.2	11.21
Gatwick Stream DS	16.7	71.7	7.37	7.76	333.9	10.74
Summer						
Site	Temperature (°C)	DO (%)	DO (mg/L)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Gatwick Stream US	16.5	72.0	7.04	7.68	280.1	11.85
Gatwick Stream DS	16.7	73.5	7.13	8.00	269.1	10.92
Autumn						
Site	Temperature (°C)	DO (%)	DO (mg/L)	pH	Conductivity (µS/cm)	Turbidity (NTU)
Gatwick Stream US	14.8	76.8	7.73	7.46	413.9	5.95
Gatwick Stream DS	16.0	78.7	7.73	8.2	387.8	11.64

### Macroinvertebrates

- 4.2.9** Fewer taxa were recorded at the two Gatwick Stream sites compared with River Mole (mean of 10 taxa per visit, compared with 19.3)(Table 4-34-3). As with the R Mole, the number of taxa recorded per visit remained relatively consistent, with a maximum of 13 in the autumn sample and a minimum of 10 in the summer sample at the upstream site. Eight taxa were recorded at the downstream site during spring and summer, and 9 in the autumn.
- 4.2.10** BMWP scores indicate moderate water quality conditions for the upstream site at the Gatwick Stream in spring and autumn (46 and 41 respectively) but were classed as poor in summer (37) (Table 4-4). The boundary between moderate and poor lies at 40, and therefore the difference between the three visits is unlikely to be significant, and is due to the smaller number of taxa recorded. However, an additional three species were recorded in the autumn sample, including the coloniser species *Asellidae*, and the caddisfly, *Polycentropus flavomaculatus*, suggesting an increase in water quality at this location, although both species were found in low abundance. The ASPT for the upstream site is similar across the three visits and is lowest in the autumn sample (4.92, 4.53 and 4.13 for the spring, summer and autumn visit respectively).

- 
- 4.2.11** At the downstream site of the Gatwick Stream the BMWP scores are classified as poor across all three visits, with the score of 14 for the spring visit being close to very poor (Table 4-4). The ASPT is also consistently lower for this site than the upstream site (3.50, 3.91 and 2.88 for the spring, summer and autumn visit respectively) over all three visits indicating the presence of pollution tolerant taxa only.
- 4.2.12** The PSI scores for the upstream site fluctuated considerably across the three season, with the maximum score of 66.67 in the spring indicating only slightly sedimented conditions (Table 4-4). However, the scores dropped progressively at this site through the season to 41.67 in the summer (moderately sedimented conditions) and then to 21.05 (sedimented conditions) in the autumn. Assuming no changes in the inputs of sediment upstream of the site, this suggests that flow velocity dropped through the season, leading to increased sediment deposition. A high LIFE score for the upstream site of 8.17 during the spring visit also suggests that velocities are high in the early part of the season.
- 4.2.13** The PSI scores for the downstream site indicated heavily sedimented conditions during the spring season (score of 14.29), with a change to moderately sedimented conditions (score of 50) in the summer and a return to sedimented conditions in the autumn (score of 33.33). LIFE scores remained relatively high and consistent across the three seasons at the downstream site (7.5, 7.75 and 8 at the spring, summer and autumn visit respectively) suggesting relatively consistent flow velocities (Table 4-4).
- 4.2.14** CCI scores for both of the Gatwick Stream sites were relatively low indicating that rare and/or notable species are absent from the macroinvertebrate assemblage. Although scores for both sites were below 5 on all sampling occasions, the upstream site had scores of 4.5 and 5 in the spring and summer respectively, whilst the scores for the downstream site was either 1 or 0 on all occasions. This indicates that the assemblage at the upstream site is of marginally higher conservation value.

#### *Fish*

- 4.2.15** A total of 300 and 317 fish were caught in spring and autumn respectively in the Gatwick Stream after three runs. Chub (*Squalius cephalus*) were the most abundant fish species identified (111) in spring on the Gatwick Stream, whereas Dace (*Leuciscus leuciscus*) were the most abundant fish species identified (137) in autumn (Table 4-7).
- 4.2.16** The size range of species caught during the electrofishing surveys carried out on the Gatwick Stream in spring suggests that there are multiple age classes of each species, ranging from juveniles to mature adults all year round.

**Table 4-7: Gatwick Stream Fish Survey Data**

Gatwick Brook					
Spring					
Species	Latin name	Abundance	Mean size (mm)	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	111	194.50	52	360
Dace	<i>Leucisus luecisus</i>	74	145.35	63	220
Perch	<i>Perca fluviatilis</i>	36	85.05	65	156
Roach	<i>Rutilus rutilus</i>	11	105.45	72	153
Bream	<i>Abramis brama</i>	6	146	92	279
Gudgeon	<i>Gobio gobio</i>	57	107.24	75	197
Stone Loach	<i>Barbatula barbatula</i>	3	127.33	97	179
Autumn					
Species	Latin name	Abundance	Mean size	Min size (mm)	Max size (mm)
Chub	<i>Squalis cephalus</i>	85	211.56	71	436
Dace	<i>Leucisus luecisus</i>	137	149.38	50	204
Roach	<i>Rutilus rutilus</i>	28	111.32	71	156
Perch	<i>Perca fluviatilis</i>	21	113.14	80	213
Bream	<i>Abramis brama</i>	10	158	132	284
Gudgeon	<i>Gobio gobio</i>	36	118.55	52	146
Stone Loach	<i>Barbatula barbatula</i>	3	86	65	98

## 5. Discussion

### River Mole

- 5.1.1** The study stretch on the River Mole lies within open floodplain grassland with no shading from trees. This means that water temperatures, and therefore dissolved oxygen, fluctuate considerably, since oxygen is less soluble in warm water. Bacterial activity associated with organic pollution also depletes dissolved oxygen levels, and therefore macroinvertebrate taxa which occur in organically polluted conditions are tolerant of low dissolved oxygen conditions. Both factors are likely to be influencing the macroinvertebrate community at the River Mole site.
- 5.1.2** Extensive stands of macrophytes covered approximately 90% of the channel surface, including submerged species such as water crowfoot (*Ranunculus aquatilis*) and the invasive non-native Canadian pondweed (*Elodea canadensis*). Emergent species such as branched bur-reed (*Sparganium erectum*), old world arrowhead (*Sagittaria sagittifolia*) and reed sweet-grass (*Glyceria maxima*) also dominated the channel. Although this channel vegetation will have contributed dissolved oxygen to the water during the summer through photosynthesis, their decay in autumn will contribute to organic pollution in reducing dissolved oxygen (28.2% and 3.03 mg/L during the autumn visit). Significant increases in conductivity such as those seen on the River Mole from spring to autumn (358 - 471  $\mu$ S/cm) (Table 4-2) are likely attributed to the decay of macrophytes and the release of ions such as phosphorous.

- 
- 5.1.3** Submerged and emergent macrophyte stands are also contributing to reduced flow velocity and increased sedimentation, reflected in the low LIFE and PSI scores for this reach.
- 5.1.4** The presence of one record from 2013 of shining ram's-horn snail (*S. nitida*), an IUCN Red List species and UK species of principal importance under the 2006 NERC Act has implications for the design of the scheme (Figure 2). Once abundant in ditch networks in the UK, the species has declined steeply, and now only occurs in a restricted number of sites in Norfolk Broads, Pevensey Levels, Lewis Levels and East Kent (Clarke, 2011). The reasons for its decline are not fully understood but are thought to be over-frequent ditch clearance, eutrophication due to fertiliser run-off, and conversion of grazing levels to arable farming with associated water table lowering (Suffolk Biological Information Service, 2003).
- 5.1.5** In a study of the associations of the species with ditch vegetation communities Clarke (2011) only found the species in ditches supporting the *Carex-Juncus-Eleocharis-Oenanthe* community of emergent vegetation. Although a full macrophyte survey was not undertaken during this study, incidental recording of macrophytes at the sampling location was undertaken and this community type was not present. However, the entire stretch from the boundary with Gatwick airport to the end of the study reach is heavily vegetated and largely impenetrable. More suitable habitat may therefore exist further downstream towards the location where it was recorded in 2013. Recommendations for further survey to determine the potential presence of the species within the study section are presented in Section 6.
- 5.1.6** The extensive macrophyte growth on the River Mole throughout the year made electrofishing difficult. In spring the filamentous algae, *Cladophora* created dense mats, which surrounded the anode each time it was placed into the water, making progress slow as the anodes regularly needed to be brought to the surface and cleared of the algae. In some cases, *Cladophora* can be beneficial to an ecosystem by providing a food source to aquatic organisms and providing a buffer to nitrification. However, excessive growth of *Cladophora* prevents aeration of deeper waters as the dense mats prevent circulation of water, which is detrimental to an ecosystem.
- 5.1.7** The high variability and remarkably low concentration of DO in the waters of the River Mole, likely contributed to the low catch in autumn where only 28 fish were caught in comparison to 415 in spring. The slow/sluggish flow of the River Mole, in combination with higher water temperatures in summer (17.6°C) could be causing DO to disassociate faster from the water. The increased presence of Tench (*Tinca tinca*) in the River Mole in autumn acted as an in-field indicator of low DO conditions, as Tench are able to tolerate much lower DO conditions than most other UK fish species.
- 5.1.8** The abundance of predatory fish in summer such as Pike (*Esox Lucius*) and Perch (*Perca fluviatilis*), may have been having a disproportionate impact on prey species on the River Mole. The prevalence of these predators has likely contributed to the significant decline in the fish population from 417 in summer to 28 in autumn. In total 14 Pike were caught in summer ranging in size from 108mm - 595mm indicating the full range of age classes. Pike are very effective freshwater hunters and as ambush predators are aided by the abundant macrophyte growth, In addition to this 46 Perch were caught in summer and ranged in size from 73mm -

258mm, also suggesting a full range of age classes. Perch also utilise macrophytes to aid in their hunting techniques, however, they are more temperature sensitive, retreating to deeper waters throughout the autumn and winter months, which has likely contributed to their decline in the area to one individual in autumn on the River Mole.

### **Gatwick Stream**

- 5.1.9** The downstream site of the Gatwick Stream appears to be suffering from poorer biological water quality than the upstream site, with the LIFE and PSI scores indicating an influx of organic pollution somewhere between these sites. This is supported by the absence of *Asellidae*, which suggests that organic pollution is chronic and there has been no recovery between Spring - Autumn. Crawley sewage treatment works lies immediately east of the Gatwick Stream, and although the discharge is directly into the River Mole, it is possible that storm water discharges from the associated industrial area enter the Gatwick Stream between the two sites. Relatively high turbidity levels of between 5.95 and 11.85NTU compared with a maximum of 4.3NTU at the River Mole site.
- 5.1.10** Differences in habitat quality and diversity between the two Gatwick Stream sites may also have influenced the macroinvertebrate community. Both sites were moderately shaded by overhanging trees, but the upstream sites was located on a tight bend with a small riffle section on the outer side of the bend, and a shallow berm on the inside edge. These microhabitats are likely to support distinct macroinvertebrate communities, with the more pollution sensitive species present in the riffle section.
- 5.1.11** The considerable variation in PSI score between the three seasonal visits at the upstream sites (maximum of 66.67 in spring compared to a minimum of 21.05 in autumn) may indicate that the macroinvertebrate community at this site is sensitive to changes on sediment deposition. Equally, it may have resulted from small differences in sampling effort in each of the microhabitats leading to a higher number of sediment sensitive taxa in the spring sample. Limited conclusions can be drawn with only one sample per visit and data from a single visit and further sampling would be required to determine any trends in the data. Overall, both sites are moderately to heavily sedimented with likely potential storm water discharges resulting in greater sedimentation at the downstream site.
- 5.1.12** The invasive New Zealand mud snail (*Potamopyrgus antipodarum*) was identified at both sites except for the Gatwick Stream downstream site in Autumn. The New Zealand pond snail is now one of the most common gastropods in the UK, its ability to avoid desiccation and its tolerance for a range of conditions enables it to dominate native gastropods, which may lead to disruptions in the food chain and effect native fish species. Currently the Gatwick Stream upstream site hosts the largest population of New Zealand mud snail, where abundances increased from 12 to 40 from spring to autumn in the samples collected. Signal crayfish (*Pacifastacus leniusculus*) were observed in relatively high numbers at both the Gatwick Stream sites during each of the visits.
- 5.1.13** The macroinvertebrate results from this study compare favourably with the Environment Agency data collected in 2017 and 2019 (Table 4-1). A slightly higher ASPT score of 4.92 was

obtained for the upstream site in early July compared with values of 3.92 and 4.0 for the 'At Tinsley Bridge, Tinsley Green' and 'Downstream Tinsley Bridge (Flylife Site)' in March 2019. However, this may reflect seasonal changes in the macroinvertebrate community between March and July. LIFE and PSI scores for both data sets indicate relatively sluggish and sedimented conditions.

- 5.1.14** The Gatwick Stream on first appearances seemed to be poor for fish species but surprisingly a consistently healthy population of fish were caught in spring (300) and in autumn (317). This likely due to the Gatwick Stream maintaining a relatively consistent water temperature (14.8-16.7°C) across all three seasons and dissolved oxygen concentrations >71%. Furthermore, although the macroinvertebrate community is poor on the Gatwick Stream, the abundance of *Chironomids*, *Oligochaetes* and *Gastropods* provide an excellent food source. There is also a diverse range of microhabitats present, such as shaded pools and undercut banks, interspersed with roots providing shelter for fish.
- 5.1.15** As a matter of interest, a Roach - Bream hybrid was identified in spring. Hybridisation between these two species is not uncommon as hybridisation between members of cyprinids is more widespread than in any other group of freshwater fish.

## 6. Conclusions and recommendations

### 6.1 Conclusions

- 6.1.1** There are no Environment Agency WFD monitoring sites on this stretch of the River Mole and therefore no background data to compare the field data collected in this study with. Data from a single site on a single year, albeit across three seasons, does not enable a comprehensive assessment of trends in the macroinvertebrate assemblage. However, based on the analysis of macroinvertebrate data collected for this study, the River Mole exhibits moderate biological water quality. Dense macrophyte growth within the channel, exacerbated by organic pollution are causing acute reductions in dissolved oxygen are likely to be impacting on the macroinvertebrate assemblage.
- 6.1.2** The record from 2013 of shining ramshorn snail (*S. nitida*), an IUCN Red List species and UK species of principal importance under the 2006 NERC Act has implications for the design of any surface water management and flood alleviation scheme. The species was not recorded during the surveys for this study, although the survey section did not coincide with the reach in which the snail was recorded.
- 6.1.3** A targeted survey for the species is required to determine its presence or absence (Section 6.2). If the species is found to be present the marginal and channel macrophyte vegetation, and flow conditions will need to be preserved in the section of the river in which the population occurs. Creation of new habitat, possibly in the form of off-line ditches supporting dense emergent vegetation is likely to be a requirement of the scheme if the species is found to be present.

- 
- 6.1.4** The structure and abundance of the cyprinid fish community in the River Mole appears to be driven by sluggish flow conditions and high summer water temperatures which favour species such as tench. The dense stands of submerged and emergent macrophytes provide foraging habitat for predatory species such as pike. Periodic dredging of the macrophyte beds would help to establish larger areas of open and deeper water thus providing refuges for prey species, improving flow conditions and creating areas of deeper, cooler water.
- 6.1.5** Based on macroinvertebrate biotic scores the Gatwick Stream has biological quality ranging from moderate at the upstream site to poor at the downstream site. Nevertheless, it retains a natural sinuous course with a variety of microhabitats supporting a range of macroinvertebrate and fish species. However, the watercourse appears to be impacted by both organic pollution and silt deposition, possibly from a storm water discharge from a nearby industrial area.
- 6.1.6** The invasive New Zealand mud snail was identified at the River Mole and Gatwick Stream sites, and signal crayfish were observed at both the Gatwick Stream sites during each visit.

## **6.2 Recommendations**

- 6.2.1** Both the Gatwick Stream and the River Mole retain natural sinuous channels characteristic of lowland rivers. It will be important to maintain and enhance this characteristic in both watercourses. The following recommendations for each watercourse are based on the findings from this study and will need refinement in light of the design of any surface water management and flood alleviation scheme and in the case of the River Mole, the findings of the survey for shining ramshorn snail. However, the habitat improvement measures recommended below are largely consistent with the requirements for this species.

### **River Mole**

1. Undertake survey to establish presence/absence of shining ramshorn snail. Survey to focus initially on the site of the 2013 record and if found to be present, extended to incorporate the whole study section. Surveys should be scoped and undertaken by a specialist mollusc ecologist.
2. If shining ramshorn snail found to be absent from main channel, undertake vegetation removal/dredging from the central channel/thalweg of the River Mole in selected sections. Marginal berms should be retained on alternate sides of the channel throughout the dredged section for the re-establishment of emergent vegetation. Brushwood faggots or fascines anchored with wooden stakes can be used to maintain the riverward edge of the berm and prevent silt from slumping into the main channel.
3. Off-line scrapes and shallow ponds could be created within the floodplain grassland area to provide habitat for wetland birds. If shining ramshorn snail is found to be present this recommendation can be adapted to incorporate new, permanently wet ditches supporting dense emergent reed vegetation.

### **Gatwick Stream**

4. Identify point sources of pollution from industrial area associated with Crawley STW including storm drains and surface water discharge points from roads and urban areas. Consider SUDS scheme to address these discharges including settlement ponds and reedbed treatment systems which would have additional biodiversity benefit.

**6.2.2** Before any in-channel works begin it is advised that a fish rescue survey is undertaken to safeguard fish populations in the affected area. It will also be necessary to install stop nets at either end of the reach where in-channel works will be undertaken to prevent access by any other fish species whilst the works are ongoing.

**6.2.3** Currently the River Mole is choked with submerged and emergent macrophyte growth, which is impeding flow, increasing deposition of sediment and reducing the circulation of deeper waters preventing aeration and creating low DO conditions. It is therefore advised that there is some level of routine maintenance of macrophyte and bankside vegetation to aid in reducing the effects of flooding and contribute to increasing the biological water quality.

### **6.3 Further Survey**

**6.3.1** It is recommended that further macroinvertebrate and fish surveys are carried out on both the River Mole and the Gatwick Stream to provide a more robust baseline of community assemblage and therefore better advise on any schemes in the future.

**6.3.2** To provide additional insight into the hydrological conditions of these rivers, it is recommended that further investigations are carried out to monitor the flow velocity and the discharge rates in order to better advise on any schemes in the future, which could include the installation of level loggers.



---

## 7. References

- 7.1.1** British Standards (2003). BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity.
- 7.1.2** Chadd, R. Extence, C. (2004). The conservation of freshwater macroinvertebrate populations: A community based classification scheme. *Aquatic Conservation Marine and Freshwater Ecosystems*. Vol 14. Pp. 597-624.
- 7.1.3** Clarke, R. (2011). Hydroseral habitat requirement of the endangered Shining Ramshorn Snail *Segmentina nitida*. *Aquatic Conserv: Mar. Freshw. Ecosyst*. 14: 597-624.
- 7.1.4** Extence, C.A, Balbi, D.M. Chadd, R.P. (1999). River Flow Indexing Using British Macroinvertebrates: A Framework For Setting Hydroecological Objectives. *Regulated Rivers: Research and Management* 15. pp543-574.
- 7.1.5** Extence, C.A. Chadd, R.P. England, J. Naura, M. Pickwell, A.G.G. (2017). Application of the Proportion of Sediment-Sensitive Invertebrates (psi) Biomonitoring Index. *River Research and Applications*. Vol 33. Issue 10. December 2017. Pp. 1596 - 1605.
- 7.1.6** Paisley, M.F. Trigg, D.J. Walley, W.J. (2013). Revision of the Biological Monitoring Working Party (BMWP) score system: Derivation of present-only and abundance-related scores from field data. <https://onlinelibrary.wiley.com/doi/abs/10.1002/rra.2686>
- 7.1.7** Suffolk Biological Information Service (2003). Shining Ram's Horn Snail (*Segmentina nitida*) Biodiversity Action Plan.  
<https://www.suffolkbis.org.uk/sites/default/files/biodiversity/priorityspecieshabitats/actionplans/shiningramshornsnail.pdf>
- 7.1.8** Water Framework Directive - United Kingdom Advisory Group. UKTAG River Assessment Method. Benthic Invertebrate Fauna. Invertebrates (General Degradation): Whalley, Hawkes, Paisley & Trigg (WHPT) metric in River Invertebrate Classification Tool (RICT). July 2014. ISBN: 978-1-906934-49-1.

## Appendix 1

Table 1: Species records for Gatwick Stream derived from the desk study

Common Name	Scientific Name	HSR Sch <sup>3</sup> 2 or 5	WCA <sup>4</sup> Sch1, 5 or 8	National Priority Species <sup>5</sup>	Local priority/ BAP species	Red Data Book/ BoCC <sup>6</sup>	Other	Grid Ref.	Distance from site	Source
<b>Birds</b>										
Goldeneye	<i>Bucephala clangula</i>		✓			Amber				
Reed Bunting	<i>Emberiza schoeniclus</i>				✓	Amber	NERC Act			
Bearded Tit	<i>Panurus biarmicus</i>		✓							
Kingfisher	<i>Alcedo atthis</i>		✓			Amber	Annex 1 Birds Directive			
Yellow Wagtail	<i>Motacilla flava</i>				✓	Red	NERC Act			

<sup>3</sup> Conservation of Habitats and Species Regulations 2010, as amended

<sup>4</sup> Wildlife and Countryside Act 1981, as amended

<sup>5</sup> Species of Principal Importance within the relevant country of the United Kingdom

<sup>6</sup> Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man

Common Name	Scientific Name	HSR Sch <sup>7</sup> 2 or 5	WCA <sup>8</sup> Sch1, 5 or 8	National Priority Species <sup>9</sup>	Local priority/ BAP species	Red Data Book/ BoCC <sup>10</sup>	Other	Grid Ref.	Distance from site	Source
<b>Amphibians</b>										
Common Toad	<i>Bufo bufo</i>		✓		✓					
Common Frog	<i>Rana temporaria</i>		✓							
Palmate Newt	<i>Lissotriton helveticus</i>		✓							
Smooth Newt	<i>Lissotriton vulgaris</i>		✓							
Great Crested Newt	<i>Triturus cristatus</i>	✓	✓		✓		NERC Act			
<b>Fish</b>										
Bullhead	<i>Cottus gobio</i>	✓								
Brown/Sea Trout	<i>Salmo trutta</i>				✓		NERC Act			
<b>Mammals (excluding bats)</b>										
<b>Reptiles</b>										
Slow Worm	<i>Anguis fragilis</i>		✓		✓		NERC Act			
Grass Snake	<i>Natrix helvetica</i>		✓		✓		NERC Act			
<b>Bats</b>										
Serotine Bat	<i>Eptesicus serotinus</i>	✓	✓							
Brandt's Bat	<i>Myotis brandtii</i>	✓	✓							

<sup>7</sup> Conservation of Habitats and Species Regulations 2010, as amended

<sup>8</sup> Wildlife and Countryside Act 1981, as amended

<sup>9</sup> Species of Principal Importance within the relevant country of the United Kingdom

<sup>10</sup> Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man

---

Brown Long-eared Bat	<i>Plecotus auritus</i>	✓	✓		✓					
----------------------	-------------------------	---	---	--	---	--	--	--	--	--

**Table 2: Species records for River Mole derived from the desk study**

Common Name	Scientific Name	HSR Sch <sup>11</sup> 2 or 5	WCA <sup>12</sup> Sch1, 5 or 8	National Priority Species <sup>13</sup>	Local priority/ BAP species	Red Data Book/ BoCC <sup>14</sup>	Other	Grid Ref.	Distance from site	Source
<b>Birds</b>										
Kingfisher	<i>Alcedo atthis</i>		✓			Amber	Annex 1 Birds Directive			
Song Thrush	<i>Turdus philomelos</i>				✓	Red	NERC Act			
Song Thrush (subspecies)	<i>Turdus philomelos clarkei</i>				✓	Red	NERC Act			
Redwing	<i>Turdus iliacus</i>		✓			Red				
Fieldfare	<i>Turdus pilaris</i>		✓			Red				
Skylark	<i>Aladua arvensis</i>				✓	Red	NERC Act			
Yellow Wagtail					✓	Red	NERC Act			
Dunnock	<i>Prunella modularis</i>				✓	Amber	NERC Act			
Black Redstart	<i>Phoenicurus ochruros</i>		✓			Red				
Nightingale	<i>Luscinia megarhynchos</i>					Red				
Marsh Tit	<i>Poecile palustris</i>				✓	Red	NERC Act			
Starling	<i>Sturnus vulgaris</i>				✓	Red	NERC Act			
House Sparrow	<i>Passer domesticus</i>				✓	Red	NERC Act			
Bullfinch	<i>Pyrrhula pyrrhula</i>				✓	Amber	NERC Act			
Hawfinch	<i>Coccothraustes coccothraustes</i>				✓	Red	NERC Act			
Yellowhammer	<i>Emberiza citrinella</i>				✓	Red	NERC Act			

<sup>11</sup> Conservation of Habitats and Species Regulations 2010, as amended

---

<sup>12</sup> Wildlife and Countryside Act 1981, as amended

<sup>13</sup> Species of Principal Importance within the relevant country of the United Kingdom

<sup>14</sup> Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man

Common Name	Scientific Name	HSR Sch <sup>15</sup> 2 or 5	WCA <sup>16</sup> Sch1, 5 or 8	National Priority Species <sup>17</sup>	Local priority/ BAP species	Red Data Book/ BoCC <sup>18</sup>	Other	Grid Ref.	Distance from site	Source
Reed Bunting	<i>Emberiza schoeniclus</i>				✓	Amber	NERC Act			
Bearded Tit	<i>Panurus biarmicus</i>		✓							
<b>Amphibians</b>										
Smooth Newt	<i>Lissotriton vulgaris</i>		✓							
<b>Invertebrates - Molluscs</b>										
Shining Ram's- Horn	<i>Segmentina nitida</i>				✓		NERC Act			
<b>Fish</b>										
Bullhead	<i>Cottus gobio</i>	✓								
Brown Trout	<i>Salmo trutta</i>				✓		NERC Act			
<b>Mammals (excluding bats)</b>										
Harvest Mouse	<i>Micromys minutus</i>				✓		NERC Act			
<b>Reptiles</b>										
Grass Snake	<i>Natrix helvetica</i>		✓		✓		NERC Act			
Common Name	Scientific Name	HSR Sch <sup>19</sup> 2 or 5	WCA <sup>20</sup> Sch1, 5 or 8	National Priority Species <sup>21</sup>	Local priority/ BAP species	Red Data Book/ BoCC <sup>22</sup>	Other	Grid Ref.	Distance from site	Source
<b>Bats</b>										
Serotine Bat	<i>Eptesicus serotinus</i>	✓	✓				Annex 4 Habitats Directive			



---

<sup>15</sup> Conservation of Habitats and Species Regulations 2010, as amended

<sup>16</sup> Wildlife and Countryside Act 1981, as amended

<sup>17</sup> Species of Principal Importance within the relevant country of the United Kingdom

<sup>18</sup> Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man

<sup>19</sup> Conservation of Habitats and Species Regulations 2010, as amended

<sup>20</sup> Wildlife and Countryside Act 1981, as amended

<sup>21</sup> Species of Principal Importance within the relevant country of the United Kingdom

<sup>22</sup> Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man

Bechsteins Bat	<i>Myotis bechsteinii</i>	✓	✓		✓	NERC Act	Annex 4 Habitats Directive Annex 2 Habitats Directive			
Brown Long-eared Bat	<i>Plecotus auritus</i>	✓	✓		✓					
Daubenton's Bat	<i>Myotis daubentonii</i>	✓	✓				Annex 4 Habitats Directive			
Whiskered Bat	<i>Myotis mystacinus</i>	✓	✓				Annex 4 Habitats Directive			
<b>Common Name</b>	<b>Scientific Name</b>	<b>HSR Sch<sup>23</sup> 2 or 5</b>	<b>WCA<sup>24</sup> Sch1, 5 or 8</b>	<b>National Priority Species<sup>25</sup></b>	<b>Local priority/ BAP species</b>	<b>Red Data Book/ BoCC<sup>26</sup></b>	<b>Other</b>	<b>Grid Ref.</b>	<b>Distance from site</b>	<b>Source</b>
Natterer's Bat	<i>Myotis nattereri</i>	✓	✓				Annex 4 Habitats Directive			
Noctule Bat	<i>Nyctalus noctula</i>	✓	✓		✓	NERC Act	Annex 4 Habitats Directive			
Nathusius's Pipistrelle Bat	<i>Pipistrellus nathusii</i>	✓	✓				Annex 4 Habitats Directive			
Common Pipistrelle Bat	<i>Pipistrellus pipistrellus</i>	✓	✓		✓	NERC Act	Annex 4 Habitats Directive			

---

Soprano Pipistrelle Bat	<i>Pipistrellus pygmaeus</i>	✓	✓		✓	NERC Act	Annex 4 Habitats Directive			
----------------------------	----------------------------------	---	---	--	---	----------	----------------------------------	--	--	--

---

<sup>23</sup> Conservation of Habitats and Species Regulations 2010, as amended

<sup>24</sup> Wildlife and Countryside Act 1981, as amended

<sup>25</sup> Species of Principal Importance within the relevant country of the United Kingdom

<sup>26</sup> Birds of Conservation Concern 4: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man



**Electrofishing and  
Macroinvertebrates Survey**

**Gatwick Fish and  
Macroinvertebrate  
surveys**

For

**RPS Ltd**

Project No.: RPS001-022

January 2023

**London & South East**

Compass House  
Surrey Research Park  
Guildford  
GU2 7AG . UK  
t: +44 (0)1483 466 000

**North & Borders**

The Tannery  
91 Kirkstall Road  
Leeds  
LS3 1HS. UK  
t: +44 (0)113 247 3780

**Wales & South West**

Sophia House  
28 Cathedral Road  
Cardiff  
CF11 9LJ UK  
t: +44 (0) 2920 660180


**Midlands**

Edmund House  
12-22 Newhall Street  
Birmingham  
B3 3AS  
t: +44 (0) 121 726 3494

**Enquiries**

e: [enquiries@thomsonec.com](mailto:enquiries@thomsonec.com)

w: [www.thomsonec.com](http://www.thomsonec.com)



Project Number	Report No.
RPS001-022	001

Revision No.	Date of Issue	Author	Reviewer	Approver
001	31/01/23	Alex Charlesworth Lucy Grinstead	Tessa Harding	Tessa Harding

**Disclaimer:**

Copyright Thomson Environmental Consultants Limited. All rights reserved.

No part of this report may be copied or reproduced by any means without prior written permission from Thomson Environmental Consultants Limited. If you have received this report in error, please destroy all copies in your possession or control and notify Thomson Environmental Consultants Limited.

This report has been prepared for the exclusive use of the commissioning party and unless otherwise agreed in writing by Thomson Environmental Consultants Limited, no other party may use, make use of or rely on the contents of the report. No liability is accepted by Thomson Environmental Consultants Limited for any use of this report, other than for the purposes for which it was originally prepared and provided.

Opinions and information provided in the report are on the basis of Thomson Environmental Consultants Limited using due skill, care and diligence in the preparation of the same and no explicit warranty is provided as to their accuracy. It should be noted and it is expressly stated that no independent verification of any of the documents or information supplied to Thomson Environmental Consultants Limited has been made.

## Contents

1. Summary and Main Recommendations .....	6
1.1 Summary .....	6
1.2 Conclusions .....	7
1.3 Recommendations.....	7
2. Introduction .....	8
2.1 Development Background .....	8
2.2 The Brief and Objectives .....	8
3. Methodology .....	13
3.1 Survey site selection.....	13
3.2 Survey methods: Macroinvertebrates.....	13
3.3 Macroinvertebrate data analysis.....	14
3.4 Survey: Fish.....	16
3.5 Limitations .....	17
4. Results .....	18
4.1 Background to waterbodies and survey sites .....	18
4.2 Environment Agency: Water Framework Directive status .....	18
4.3 <i>Gatwick Stream</i> .....	19
4.4 River Mole.....	22
4.5 Pond F .....	25
4.6 River Mole confluence .....	27
5. Discussion .....	29
6. Conclusions and recommendations .....	31
6.1 Conclusions .....	31
6.2 Recommendations.....	31
7. References .....	32
Appendix 1 - Taxa lists .....	33
Table 1: Macroinvertebrate survey dates.....	14
Table 2: BMWP Scoring System.....	15
Table 3: Interpretation of LIFE score.....	16
Table 4: Interpretation of PSI scores.....	16
Table 5: Fish survey dates .....	17
Table 6: Water quality at Gatwick Stream.....	19
Table 7: Macroinvertebrate biotic indices at Gatwick Stream.....	19
Table 8: Gatwick Stream fish species data collected through summer and autumn electrofishing surveys.....	21
Table 9: Water quality at River Mole .....	22

Table 10: Macroinvertebrate biotic indices at River Mole .....	23
Table 11: River Mole fish species data collected during the autumn electrofishing survey.....	24
Table 12: Water quality at Pond F.....	25
Table 13: Macroinvertebrate biotic indices at Pond F.....	26
Table 14: Water quality at the River Mole confluence .....	27
Table 15: Macroinvertebrate biotic indices at the River Mole confluence.....	27



## 1. Summary and Main Recommendations

### 1.1 Summary

- 1.1.1** RPS commissioned Thomson Environmental Consultants (TEC) to update baseline fish and macroinvertebrate surveys on the Gatwick Stream and River Mole to support the ecological impact assessment for expansion proposals at Gatwick airport. The scheme will entail re-meandering a stretch of the River Mole immediately north of the airport and creation of a new attenuation area to the north east of the airport.
- 1.1.2** The study area encompasses two watercourses and a pond; a 1.3km stretch of the River Mole immediately downstream of where it emerges from under the runway at Gatwick Airport, and a 750m stretch of the Gatwick Stream (a tributary of the River Mole) upstream of the Crawley sewage works. Baseline macroinvertebrate surveys were also undertaken at Pond F, a surface drainage pond located east of the airport. Surveys were previously undertaken on the River Mole and Gatwick Stream in 2020. In 2022, whilst the survey site remained the same for the River Mole survey, the Gatwick Stream site was taken further down-stream to avoid access issues with the equipment for electro-fishing.
- 1.1.3** Macroinvertebrate surveys were carried out in summer and autumn 2022. A mean of 12 species were recorded on the Gatwick Stream across the sampling site over the two visits. Biotic indices measuring water quality (Biological Monitoring Working Party (BMWP) score and Average Score Per Taxon) indicate moderately polluted conditions in the stream in summer to poor in the autumn. Lotic invertebrate Index for Flow Evaluation (LIFE) scores suggest that the macroinvertebrate community is characteristic of slow/sluggish flow conditions. The Proportion of Sediment intolerant Invertebrates (PSI) scores indicate slightly variable conditions across the watercourse and between summer and autumn, with a greater proportion of slightly sedimented conditions in summer to moderately sedimented conditions in autumn.
- 1.1.4** A mean of 22.5 species were recorded on the River Mole across the sampling site over the two visits. Biotic indices measuring water quality (Biological Monitoring Working Party (BMWP) score and Average Score Per Taxon) indicate poor pollution conditions in the River Mole. Lotic invertebrate Index for Flow Evaluation (LIFE) scores suggest that the macroinvertebrate community is characteristic of sluggish flow conditions. The Proportion of Sediment intolerant Invertebrates (PSI) scores indicate heavily sedimented conditions over both the seasons.
- 1.1.5** A mean of 16.50 species were recorded on Pond F across the sampling site over the two visits. Biotic indices measuring water quality (Biological Monitoring Working Party (BMWP) score and Average Score Per Taxon) indicate moderately polluted conditions in the summer moving to poor in the autumn. Lotic invertebrate Index for Flow Evaluation (LIFE) scores suggest that the macroinvertebrate community is characteristic of sluggish flow conditions. The Proportion of Sediment intolerant Invertebrates (PSI) scores indicate heavily sedimented conditions in both seasons.

**1.1.6** In the Gatwick Stream, a total of 302 fish were caught in summer after one run compared with 304 in autumn over three electrofishing runs. Chub (*Squalius cephalus*) was the most abundant species over both visits. A total of 71 fish were recorded from the River Mole during three runs in the autumn survey, with Roach (*Rutilus rutilus*) being the most abundant. The species composition in the Gatwick Stream was more diverse overall than the River Mole with a total of 6 species in the summer and 8 in the autumn compared to 6 species in the river Mole in the autumn survey.

**1.1.7** The fish assemblages in both watercourses were similar to the 2020 surveys with chub dominating the community in the Gatwick Stream and roach in the River Mole. Although fish abundances in the Gatwick Stream were similar to that recorded during the 2020 surveys abundances in the River Mole were over double that of 2020.

## **1.2 Conclusions**

**1.2.1** Both watercourses and Pond F supported macroinvertebrate communities indicative of moderate to poor water quality, exacerbated by relatively low flow conditions and high levels of sedimentation. Dense macrophyte growth on the R Mole is contributing to acute reductions in dissolved oxygen which are impacting on the macroinvertebrate assemblage. The findings for R Mole and Gatwick Stream are therefore consistent with the WFD classification.

**1.2.2** The PSI and LIFE scores over the 2020 and 2022 surveys indicate that the River Mole has problems with siltation. This should be considered during the design process of the river channel in order to maximise flow.

**1.2.3** Hot and dry weather conditions in summer 2022 combined with a pollution incident prior to the summer survey for the River Mole appeared to show no chronic impacts in the macroinvertebrate communities. The increase in biotic scores may be attributed to the recovery in flow conditions in the autumn sample.

**1.2.4** The presence of bullhead in the Gatwick Stream enhances the need for safeguarding to be involved in the mitigation strategy for the Gatwick airport expansion development.

## **1.3 Recommendations**

**1.3.1** The main recommendations are set out below:

- The presence of bullhead should be considered when planning the new outfall from the water treatment plant into the Gatwick Stream in order to minimise impact to the species.
- To increase habitat heterogeneity through variable flow patterns and the creation of new channel features such as pools, side bars and points bars, avoiding dominating vegetation.
- Management of the invasive, non-native species on site such as the crayfish, mink and Himalayan balsam. Particularly in areas where it has the potential to contribute to the decline of water quality and the stability of the banksides.
- The creation of a management plan for the diversion and floodplain of the River Mole. Management would have to consider the presence and protection of water voles.

---

## 2. Introduction

### 2.1 Development Background

**2.1.1** Two watercourses, the River Mole and Gatwick Stream will be directly affected by proposals to increase flood storage to the east and west of Gatwick Airport. The scheme includes proposals to re-meander the River Mole close to where it emerges from beneath the airport runway, create new flood attenuation areas to the west of the watercourse and lengthen existing culverts where the River Mole flows beneath the A23. A new outfall will be created into the Gatwick Stream from a new treatment plant treating surface water from the aprons and runway areas. The discharge point will be close to the existing Crawley waste water treatment works.

**2.1.2** The study area for the fish and macroinvertebrate surveys encompasses two watercourses; a 1.3km stretch of the River Mole immediately downstream of where it emerges from under the runway at Gatwick Airport, and a 750m stretch of the Gatwick Stream (a tributary of the River Mole) upstream of the Crawley sewage works. Macroinvertebrate sampling was also undertaken at the confluence of the River Mole and Gatwick Stream, and in Pond F, a balancing pond immediately south of the A23. The sites are described in Section 3.1.

### 2.2 The Brief and Objectives

**2.2.1** RPS commissioned Thomson Environmental Consultants on 16/06/2022 to undertake fish and macroinvertebrate surveys of the Gatwick Stream, River Mole and Pond F, a surface water balancing pond located within the industrialised area of Gatwick. The brief was to:

- To determine baseline populations for both fish, eels and macroinvertebrates in these watercourses over the course of a year.
- Provide a report on the survey giving the methods and results of the survey, with recommendations, including opportunities for enhancement, mitigation and further surveys.

Filepath: R:\Live Projects\Water Ecology\IRPS001-022\_1 - Aquatic surveys at Gatwick Airport\Mapping\Working\IRPS001\_022\_Fig1\_GatwickStream\_SC\_091222.mxd  
 Contains Ordnance Survey data © Crown copyright and database right 2022. This map must not be copied or reproduced by any means without prior written permission from Thomson Ecology Ltd.



Legend

**Sampling Points**

- Electro-Fishing Sampling Point
- Macroinvertebrate Sampling Point
- Gatwick Stream


Map Centre Grid Reference: 528,221 142,095

Client	RPS Ltd.		Drawing Ref	RPS001-022/401771/1		
Figure Number	1a		Scale at A4	1:1,000		
Figure Title	Gatwick Stream		Drawn	SC	Checked	JB
			Date	09/12/2022	Date	09/12/2022

**Thomson**  
 environmental  
 consultants

www.thomsonec.com  
 enquiries@thomsonec.com

**DRAFT**




Filepath: R:\Live Projects\Water Ecology\RPS001-022\_1 - Aquatic surveys at Gatwick Airport\Mapping\Working\RPS001\_022\_Fig1b\_RiverMole\_SC\_091222.mxd  
Contains Ordnance Survey data © Crown copyright and database right 2022. This map must not be copied or reproduced by any means without prior written permission from Thomson Ecology Ltd.

Legend

**Sampling Point Type**

- Electro-Fishing Sampling Point
- Macroinvertebrate Sampling Point
- River Mole

Map Centre Grid Reference: 525,634 140,520




Client	RPS Ltd.		Drawing Ref	RPS001-022/401772/1	
Figure Number	1b		Scale at A4	1:2,000	
Figure Title	River Mole Sampling Points		Drawn	SC	Checked
					AS
		Date	09/12/2022	Date	09/12/2022

**Thomson**  
environmental  
consultants

www.thomsonec.com  
enquiries@thomsonec.com

Filepath: S:\Live Projects\Water Ecology\IRPS001-022\_1 - Aquatic surveys at Gatwick Airport\Working\IRPS001\_022\_Fig1c\_SamplingPoint\_SC\_091222.mxd  
 Contains Ordnance Survey data © Crown copyright and database right 2023. This map must not be copied or reproduced by any means without prior written permission from Thomson Ecology Ltd.



**Legend**

- Macroinvertebrate Sampling Point
- Waterbody

Map Centre Grid Reference: 528,744 141,535


Client	RPS Ltd.
Figure Number	1c
Figure Title	Macroinvertebrate Sampling Point

Drawing Ref	RPS001-022/401773/1	
Scale at A4	1:5,000	
Drawn	JT	Checked JB
Date	10/01/2023	Date 10/01/2023

**Thomson**  
 environmental  
 consultants

www.thomsonec.com  
 enquiries@thomsonec.com


Filepath: S:\Live Projects\Water Ecology\RPS001-022\_1 - Aquatic surveys at Gatwick Airport\Working\RPS001\_022\_Fig1d\_SamplingPoint\_JT\_100123.mxd  
 Contains Ordnance Survey data © Crown copyright and database right 2023. This map must not be copied or reproduced by any means without prior written permission from Thomson Ecology Ltd.



**Legend**

- Macrolnvertebrate Sampling Point
- Waterbodies

Map Centre Grid Reference: 527,584 142,376



Client	RPS Consulting Services Ltd		Drawing Ref	RPS001-022/401910/1	
Figure Number	1d		Scale at A4	1:8,000	
Figure Title	Macrolnvertebrate Sampling Location		Drawn	JT	Checked
			Date	10/01/2023	Date

**Thomson**  
 environmental  
 consultants

www.thomsonec.com  
 enquiries@thomsonec.com

---

## 3. Methodology

### 3.1 Survey site selection

- 3.1.1** Survey sites for the Gatwick Stream and the River Mole were identified from a walkover survey undertaken prior to the 2020 surveys. One 100m electrofishing survey section was identified on the River Mole and a second on the Gatwick Stream. The survey stretch on the River Mole (TQ255405 to TQ 25681 40489) lies approximately 7 miles downstream of the source, immediately north of where the R Mole passes beneath the Gatwick airport runway in a culvert. (Figure 1b). The survey site is in approximately the same location as in 2020 with a slight adjustment made to avoid dense macrophyte growth.
- 3.1.2** The survey stretch on the Gatwick Stream (TQ 28194 42146 to TQ 28254 42033) lies within Riverside Park, Crawley approximately 650m upstream of the confluence with the River Mole. (Figure 1a). The site is approximately 2.5km downstream of the 2020 sampling location. The same sampling site was not used due to access issues regarding the electrofishing equipment and was therefore substituted with the Riverside Park site which had safer access.
- 3.1.3** Pond F was not sampled in 2020 and therefore there was no pre-existing survey site. Access to the pond was limited due to an extensive marginal macrophyte fringe and fish surveys were not considered feasible. The pond was sampled for macroinvertebrates towards the southwest corner where there was a moderately accessible point (TQ 28755 41520), illustrated in Figure 1c.
- 3.1.4** The River Mole confluence was also not surveyed in 2020 and therefore there was no pre-existing survey site to replicate. As with Pond F, access to the site was limited due to the high-water levels. It was therefore not surveyed for fish at all, and surveyed for macroinvertebrates on only the summer visit. The sample for invertebrates was taken from TQ 27592 42448 (Figure 1d).

### 3.2 Survey methods: Macroinvertebrates

- 3.2.1** 3 samples were collected, one on the Gatwick Stream and two on the River Mole using the standard method for sampling macroinvertebrates in rivers to support WHPT metric calculations (British Standard, 1994; UKTAG 2021). The method comprises a three-minute kick sample using a long-handled pond net with 1mm mesh size, which was supplemented by a one-minute hand search (Environment Agency, 2017). Sampling of habitats within the three-minute kick/sweep sampling were in proportion to their occurrence. Samples were then preserved in industrial methylated spirits (IMS) for processing in the laboratory to the requirements outlined in EA Operational Instruction 024\_08 Freshwater macroinvertebrate analysis of riverine samples (Environment Agency, 2014).
- 3.2.2** The Pond F sample was collected using the methods described in the PSYM methodology document (Environment Agency, 2002) . The procedure involves sweeping beneath the water surface using a long-handled pond net after disturbing the sediment, then agitating marginal and



submerged vegetation to dislodge macroinvertebrates. Sampling of habitats within the three minute period was in proportion to their occurrence. Deep accumulations of sediment were avoided, since these areas are typically species poor. Samples were then preserved in industrial methylated spirits (IMS) for processing in the laboratory to the requirements outlined in EA Operational Instruction 024\_08 Freshwater macroinvertebrate analysis of riverine samples (Environment Agency, 2014).

- 3.2.3** Macroinvertebrates were identified to Mixed Taxon Level 5, to enable evaluation of the macroinvertebrate community and calculation of the relevant biotic indices including Biological Monitoring Working Party (BMWP), Average Score Per Taxon (ASPT) and Lotic-Invertebrate index for Flow Evaluation (LIFE). Proportion of Sediment-sensitive Invertebrates (PSI) and Community Conservation Index (CCI).
- 3.2.4** Macroinvertebrate sampling was undertaken in summer and autumn on the dates presented in **Error! Reference source not found.**

**Table 1: Macroinvertebrate survey dates**

Macroinvertebrate survey visit	Date
Summer	02/08/2022
Autumn	11/10/2022

### 3.3 Macroinvertebrate data analysis

- 3.3.1** The macroinvertebrate abundance data collected during the field surveys and background data from the Environment Agency has been analysed using a range of biotic indices. Each of the indices used in the analyses are summarised below.

#### *Biological Monitoring Working Party (BMWP) score*

- 3.3.2** The BMWP score is a method for indexing river water quality in England and Wales using macroinvertebrate families. Originally published in the early 1980's, the system was updated in 2013 based on a more robust baseline data set (Paisley et al, 2013). A score of between 1 and 10 is assigned to families found within a sample based on their tolerance to organic pollution, with a score of 1 indicating high tolerance, and 10 indicating low tolerance. Low scoring families include worms (Oligochaeta) and non-biting midge larvae (Chironimidae), whilst the presence of mayfly (Ephemeroptera) and stonefly (Plecoptera) larvae is indicative of clean water conditions. The scores for each family recorded in the sample are summed to give the overall BMWP site score. Since the overall site score is influenced by the number of families as well as the scores of the individual families in the sample, an average is taken by dividing the overall BMWP score by the number of families/taxa in the sample. This is termed the Average Score Per Taxon (ASPT).

Table 2 provides an interpretation of the BMWP scoring system.

**Table 2: BMWP Scoring System**

BMWP score	Category	Interpretation
0-10	Very poor	Heavily polluted
11-40	Poor	Polluted or impacted
41-70	Moderate	Moderately impacted
71-100	Good	Clean but slightly impacted
>100	Very good	Unpolluted, unimpacted

*River Invertebrate Classification Tool (RICT)*

**3.3.3** BMWP and ASPT have been updated by the Walley, Hawkes, Paisley and Trigg (WHPT) method for River Invertebrate Classification Tool (RICT) (UK TAG, 2021). RICT is used to classify the aquatic macroinvertebrate sub-element according to their ecological status under the Water Framework Directive (WFD). The scores derived for an individual site under RICT are compared with those expected under unpolluted conditions (known as reference conditions) in order to give an Environmental Quality Ratio (EQR). This aims to take account of the variability of macroinvertebrate families in rivers resulting from environmental parameters such as altitude, underlying geology and proximity to the river source. For the purposes of monitoring and reporting under the WFD, the quality elements which are used to assess ecological status for surface waterbodies, are assigned to one of the following five categories: High, Good, Moderate, Poor, Bad. Macroinvertebrates are one of the biological quality elements, together with fish and phytobenthos.

*Lotic invertebrate Index for Flow Evaluation (LIFE) Score*

**3.3.4** The LIFE score system links flow conditions in rivers, and specifically flow velocity, with commonly identified macroinvertebrate species and families (Extence *et al.* 1999). Macroinvertebrates are assigned to one of 6 groups depending on their tolerance to low flow conditions. The groups range from I comprising taxa associated with rapid flow conditions (>100cm s<sup>-1</sup>) to VI including those associated with drying or drought impacted sites (Table 3). A flow score is obtained for each species/taxon by combining the flow category with an estimated abundance score as described by Extence *et al.* (1999). The LIFE score for a sample is obtained by summing the individual flow scores for each taxon by the number of taxa in the sample. LIFE scores range from 1 to 12, with scores of 8 or above indicating moderate to high flow conditions, and scores of 7 or below indicating sluggish conditions.

**Table 3: Interpretation of LIFE score**

Category	Flow group
I	Rapid
II	Moderate/fast
III	Slow/sluggish
IV	Sluggish/standing
V	Standing
VI	Drought resistant

*Proportion of Sediment sensitive Invertebrates (PSI)*

**3.3.5** The PSI index provides an indication of the extent to which watercourses have been impacted by the deposition of fine sediment (Extence et al, 2017). Following the same principle as the LIFE score system, invertebrates are assigned to one of four groups depending on their sensitivity to fine sediment, with Group A comprising highly sensitive taxa, and Group D those that are highly insensitive. The method also requires a log abundance category to be estimated for all taxa identified in a sample (1-9, 10-99, 100-999 and 1000+ individuals present). Scores range from 80 -100 for unsedimented sites down to 0-20 for highly sedimented sites.

**Table 4: Interpretation of PSI scores**

PSI score	River bed condition
81-100	Minimally sedimented/unsedimented
61-80	Slightly sedimented
41-60	Moderately sedimented
21-40	Sedimented
0-20	Heavily sedimented

*Community Conservation Index (CCI)*

The CCI combines the rarity of constituent species in a sample with the diversity of the community, or community richness, to give a single integrated score which can be used as the basis for site evaluating (Chadd and Extence, 2004). Species identified from a survey site or area are given a Conservation Score (CS), based on standard rarity categories, with Red Data Book 1 (Endangered) species scoring 10, and very common species scoring 1. The sum of each of the conservation scores in the sample is then divided by the number of contributing species to give the overall CCI score.

**3.4 Survey: Fish**

**3.4.1** The surveys were undertaken using the catch depletion method in order to assess species composition, age structure and to estimate population size. Surveys were undertaken by an accredited electric fishing team comprising four members of staff. Surveys and analysis

conformed to the relevant guidance outlined in BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity (British Standards, 2003). An FR2 consent (application to use fishing instruments other than rod and line) was sought from the Environment Agency prior to conducting the survey.

- 3.4.2** The survey was undertaken over two 100m reaches that coincided with the macroinvertebrate survey locations on the Gatwick Stream and River Mole. Stop nets were installed across the channel at either end of the reach to prevent fish entering or leaving the survey area. Holding containers for captured fish were established on the riverbank with an aerator installed to provide oxygen to captured fish. On the bankside the captured fish were immediately identified, measured and returned to another holding container after surveying each reach.
- 3.4.3** The survey was undertaken using an electrofishing box and two anodes. Two surveyors, operating the electrofishing anodes waded from downstream to upstream and a third surveyor netted any stunned fish. The operatives were supported by a fourth operative who monitored the electrofishing box and holding tank. At the end of each run all caught fish were identified, measured and placed in a submerged holding net to facilitate their recovery and prevent re-capture.
- 3.4.4** Three survey visits were undertaken, one in summer (02/08/2022) and two in autumn (11-12/10/2022) (Table 5) to establish a baseline of the species composition on the watercourses.

**Table 5: Fish survey dates**

Fish survey visit	Date
Summer	02/08/2022
Autumn	11-12/10/2022

### 3.5 Limitations

- 3.5.1** Comparison between fish populations is limited due to the incomplete runs in summer for the Gatwick Stream and no data for the River Mole due to a pollution event. Comparisons are therefore made between the complete 2020 surveys conducted by Thomson of the sites.
- 3.5.2** Electro-fishing surveys at Pond F were not possible due to the dense fringe of emergent macrophytes which created unsafe access conditions for launching a vessel. Access was possible for macroinvertebrate sampling.
- 3.5.3** The River Mole had exceptionally high coverage of aquatic plants, which made electrofishing difficult. The filamentous algae and blanket weed (*Cladophera* agg.), was found in dense clumps making progress slow, as the anode became blanketed by the filamentous algae each time it was placed into the water and needed regular clearing in order to progress. Due to the depth and access to the River Mole confluence, electro-fishing during summer and autumn and macroinvertebrate sampling in winter was not possible at this site.

---

## 4. Results

### 4.1 Background to waterbodies and survey sites

**4.1.1** The River Mole rises in Baldhorns Copse in West Sussex and flows over the Wealden and London clays, cutting through the North Downs chalk between Dorking and Leatherhead, before discharging into the River Thames at the town of Molesey in Surrey. The Gatwick Stream is a tributary of the River Mole. It rises in Worth Forest in West Sussex before flowing north to meet the River Mole at the West Sussex and Surrey border. The site surveyed for fish and macroinvertebrates on the Gatwick Stream lies immediately upstream of the confluence with the River Mole north of Gatwick Airport (Figure 1d).

**4.1.2** Pond F is a surface water balancing pond for Gatwick Airport, located to the south of Airport Way and immediately east of the rail line within the catchment of the Gatwick Stream (Figure 1c).

### 4.2 Environment Agency: Water Framework Directive status

**4.2.1** The surveyed reach of the River Mole falls within the WFD waterbody named 'Mole Upstream of Horley' (GB106039017481). The most recent classification in 2019 under the WFD classification system marks this stretch as moderate ecological status. Although it is classified as good for biological quality elements (fish only) it is moderate for physico-chemical quality elements and 'bad' for dissolved oxygen. The reasons for not achieving good have been identified as continuous discharge from the water industry (DO and phosphate).

**4.2.2** The reach of Gatwick Stream falls within the 'Tilgate Brook and Gatwick Stream at Crawley Water' (GB106039017500). Similarly to the River Mole stretch, it is of moderate ecological status under the WFD classification. However, the stretch was classified as 'bad' for biological quality elements due to bad score for fish and poor for invertebrates. The physico-chemical quality elements received a good whilst chemical failed. The reason for the Gatwick Stream not achieving good status has been attributed to continuous sewage discharge and urban and transport through diffuse pollution.

**4.2.3** The sampled area of the River Mole confluence falls within the 'Horley to Dorking' waterbody under the WFD (GB106039017625). The most recent classification (2019) under the WFD classification system identifies this stretch as moderate ecological status. All the biological quality elements achieved 'moderate' apart from fish which achieved 'good'. The physico-chemical quality elements also achieved 'moderate'. The waterbody is rated as 'fail' for chemical status due to the presence of mercury and its compounds. The reasons for not achieving good status are 'Technically infeasible' 'Disproportionately expensive'.

### 4.3 Gatwick Stream

#### Water Quality

Table 6: Water quality at Gatwick Stream

Season	Temperature (°C)	DO (%)	DO (mg/L)	Conductivity (µS/cm)	pH	ORP	Turbidity (NTU)
Summer	19.7	74.9	6.75	931	7.4	88.4	4.9
Autumn	15.9	87.3	8.6	835	7.08	233.1	3.29

4.3.1 Water temperature in the Gatwick Stream ranged from 19.7°C in summer to 15.9°C in autumn. The dissolved oxygen content remained over 70% in both the summer and autumn. Conductivity remained high over both the summer and autumn at 931mg/L and 835 mg/L (Table 6). However, the turbidity appears to be low overall with little variation between summer and autumn.

4.3.2 The oxidation reduction potential was high at the Gatwick Stream site (88.4 and 233.1 in summer and autumn respectively) reflecting the capacity of the system to break down pollutants, contaminants or detritus.

4.3.3 There was an overall increase in water temperature of about 3 degrees in the summer when compared to the 2020 data. Additionally, there was a large increase in the conductivity from 280-269µS/cm in 2020 to 931µS/cm in 2022. DO concentrations in the autumn of 2022 exceeded the levels of the Gatwick data in 2021 by almost 10%.

#### Macroinvertebrates

Table 7: Macroinvertebrate biotic indices at Gatwick Stream

Biotic Index	Summer 02/08/2022	Autumn 11/10/2022
BMWP (TL1)	27	50
NTAXA (TL1)	7	10
ASPT (TL1)	3.86	5.00
LIFE (TL5)	7.67	7.11
PSI (TL5)	66.67	47.37
EPSI (TL5)	72.99	78.53
CCI (TL5)	1.00	9.29
WHPT (TL2)	30.9	54.0
NTAXA (TL2)	7	11
ASPT (TL2)	4.41	4.91

- 
- 4.3.4** There were fewer species recorded at the Gatwick Stream than the River Mole (mean of 12 taxa per visit compared with 22.5). In the summer months the assemblage was predominantly comprised of worms, *Oligochaeta*, net spinners (caddis) *Hydropsyche angustipennis* and non-biting midges *Chironomidae*. The dominant species was the same for the autumn months with the addition of the freshwater shrimp *Crangonyx pseudogracilis/floridanus agg.* Various caddisflies and damselfly nymphs were also found in the autumn samples such as *Calopteryx virgo* and *Cyrnus trimaculatus*.
- 4.3.5** The number of taxa recorded at the Gatwick Stream site increased from 9 in the summer sample to 15 in the autumn. Three damselfly larvae from the family Calopteryx, recorded in the autumn were absent in the summer sample. Four additional caddisfly (Family: Hydropsychidae) were also recorded in the autumn sample.
- 4.3.6** The increased number of taxa and the presence of the four additional caddisfly larvae contributed to an increase in the BMWP score from 27 in the summer to 50 in the autumn (Table 7). This represents a shift, from the poor to the moderate pollution category. The presence of high scoring taxa such as the Hydropsychidae in the autumn sample also resulted in an increase in ASPT from 3.86 to 5.0.
- 4.3.7** LIFE scores ranged from 7.67 to 7.11 indicating slow/sluggish conditions and a slight reduction from summer to autumn. This supports the PSI readings as they range from slightly sedimented to moderately sediment following the reduction in flow. The CCI ranges from low conservation value to moderate. This is in contrast to the BMWP and ASPT scores which increase in the autumn sample.

*Fish*

**Table 8: Gatwick Stream fish species data collected through summer and autumn electrofishing surveys.**

Summer					
Overall					
Species	Latin name	Abundance	Mean Size (mm)	Min size (mm)	Max size (mm)
Dace	<i>Leuciscus leuciscus</i>	67	146.7164179	80	190
Chub	<i>Squalius cephalus</i>	73	271.4383562	100	470
Barbel	<i>Barbus barbus</i>	11	290.4545455	190	480
Roach	<i>Rutilus rutilus</i>	33	139.1515152	70	235
Perch	<i>Perca fluviatilis</i>	4	241.25	130	330
Gudgeon	<i>Gobio gobio</i>	32	115.3125	70	160
Autumn					
Overall					
Species	Latin name	Abundance	Mean Size (mm)	Min size (mm)	Max size (mm)
Dace	<i>Leuciscus leuciscus</i>	53	142.3584906	80	180
Chub	<i>Squalius cephalus</i>	134	200.9253731	100	450
Barbel	<i>Barbus barbus</i>	5	246	200	270
Roach	<i>Rutilus rutilus</i>	9	160	120	200
Perch	<i>Perca fluviatilis</i>	8	164.375	100	260
Gudgeon	<i>Gobio gobio</i>	28	116.7857143	100	150
Pike	<i>Esox lucius</i>	3	483.3333333	300	650
Bullhead	<i>Cottus gobio</i>	4	100	100	100

- 4.3.8** The results from the summer and autumn fish surveys (Table 8) cannot be directly compared, due to the disparities in the survey methodology. Only one run was completed on the Gatwick Stream in summer compared with two in autumn. The total number of fish captured during three runs (244) in autumn was similar to that recorded in a single run in the summer (220). This suggests that there were lower numbers of fish present in the watercourse in autumn, although this cannot be concluded definitively without further sampling.
- 4.3.9** The fish community is typical of a slow flowing freshwater system with the presence of roach (*Rutilus rutilus*) and pike (*Esox Lucius*). Additionally, species such as dace (*Leuciscus leuciscus*) and chub (*Squalius cephalus*) favour the shaded habitat provided by bankside scrub and trees, and marginal macrophytes.
- 4.3.10** One adult Signal crayfish (*Pacifastacus leniusculus*) was caught in the Gatwick Stream during the autumn fish survey. Signal crayfish is an invasive non-native species (INNS) which competes with the native, white-clawed crayfish (*Austropotamobius pallipes*) for food and shelter and carries a crayfish plague (*Aphanomyces astaci*) which causes mortality in white-clawed crayfish. They also burrow into riverbank faces causing instability and increasing the rate in which siltation occurs. The presence of multiple holes in banks within the survey site of the Gatwick Stream indicate that this is a problem which could result in undermining and collapse if not managed.



- 4.3.11 The size of the fish has remained fairly constant when compared to the 2020 data, but the chub has increased in the summer months from a mean size of 194.5mm in summer 2020 to 271.4mm in summer 2022. There has been a reduction in numbers of fish such as the gudgeon for which 57 individuals were recorded in summer and 36 in autumn 2020; reducing to 32 in summer and 28 in autumn in 2022. Similar reductions were noted in the numbers of chub,
- 4.3.12 The species community is fairly similar between survey seasons with the addition of pike and bullhead (*Cottus gobio*) in the autumn sample. Bullhead are on Annex II of the EU Habitats Directive and IUCN red list. The pike could have been introduced to the area as the stream is stocked by the EA.
- 4.3.13 There are some differences in the species composition of the Gatwick Stream in 2022 when compared to the survey conducted in 2020. The main differences are the absence of bream *Abramis brama* and stone loach *Barbatula barbatula* and the presence of the barbel *Barbus barbus* in 2022

#### 4.4 River Mole

##### Water Quality

Table 9: Water quality at River Mole

Season	Temperature (°C)	DO (%)	DO (mg/L)	Conductivity (µS/cm)	pH	ORP	Turbidity (NTU)
Summer	18.3	14	1.31	554	7.2	11.51	1.43
Autumn	11.2	56.4	6.18	314.6	7.32	207	40.76

- 4.4.1 Water temperatures in the River Mole during the summer visit were slightly lower than the Gatwick Stream (18.3°C compared with 19.7°C), most likely due to the larger volumes of water in the channel and partial shading provided by the scrub and macrophytes (Table 9). Water temperature dropped by around 7 °C to 11.2°C. in the autumn
- 4.4.2 The DO increased notably between the summer and autumn survey from 14% in August to 56.4% in October. Such low DO in the summer is likely due to the high-water temperatures (in which oxygen is less soluble) and possibly the impacts of a Category 1 impact level pollution incident involving organic chemicals (surfactants and detergents) several months before the survey.
- 4.4.3 The River Mole also saw a significant decrease in conductivity by 239.4 µS/cm, which is the opposite of the increase observed in the 2020 data. Alongside this, there was a dramatic increase in turbidity from summer to autumn from 1.43NTU to 40.76NTU, the largest increase of any sites. The ORP also revealed a large increase from 11.51 to 207.

- 4.4.4** The study stretch on the River Mole lies within open floodplain grassland with no shading from trees but excessive shading from tall reeds around the edges and especially in narrow parts of the stretch. This means that water temperatures, and therefore dissolved oxygen, fluctuate considerably. Bacterial activity associated with organic pollution also depletes dissolved oxygen levels, and therefore macroinvertebrate taxa which occur in organically polluted conditions are tolerant of low dissolved oxygen conditions. Both factors are likely to be influencing the macroinvertebrate community at the River Mole site. However, such low DO may also be due to a Category 1 impact level pollution incident involving organic chemicals (surfactants and detergents) in the months before the survey.
- 4.4.5** Extensive stands of macrophytes covered approximately 90% of the channel surface, including submerged and emergent species such as Arrow heads (*Sagittaria sagittifolia*), Common Duckweed (*Lemna minor*). Although this channel vegetation will have contributed dissolved oxygen to the water during the summer through photosynthesis, their decay in autumn will contribute to organic pollution in reducing dissolved oxygen. Significant increases in conductivity such as those seen on the River Mole from spring to autumn are likely attributed to the decay of macrophytes and the release of ions such as phosphorous.

**Macroinvertebrates**

**Table 10: Macroinvertebrate biotic indices at River Mole**

Biotic Index	Summer	Autumn
	02/08/2022	11/10/2022
BMWP (TL1)	55	52
NTAXA (TL1)	15	13
ASPT (TL1)	3.67	4.00
LIFE (TL5)	5.80	5.77
PSI (TL5)	2.56	0.00
EPSI (TL5)	0.00	0.00
CCI (TL5)	4.71	5.75
WHPT (TL2)	57.1	51.0
NTAXA (TL2)	17	15
ASPT (TL2)	3.36	3.40

- 4.4.6** The River Mole had the greatest mean number of species (22.5 across the summer and autumn season) of the three sites (i.e. River Mole, Gatwick Stream and Pond F). Taxa varied between seasons, for example bivalve molluscs were found in summer, some of which in large numbers such as the *Sphaerium corneum* and none in autumn, and autumn had large numbers of 108 of water flea *Cladocera*, a species that was not present in the summer sample. This was the same for the 2020 data set in the Gatwick Stream and River Mole. It also experienced a slight increase in the ASPT from 3.67 to 4.00.
- 4.4.7** The BMWP remained at ‘poor’ with a slight decline from summer to autumn (Table 10), this is likely influenced by the large presence of species low scoring such as the water flea *Cladocera* in autumn and the reduction of high scoring taxa such as the mayfly nymph *Cloeon dipterum*.

This was coupled with the decline of the PSI from 2.56 to 0 showing the site to be heavily sedimented. This was supported by the low flow revealed by the low score and decline in LIFE score from 5.80 to 5.77. The River Mole, like the other 2 sites, demonstrated an increase in the CCI from low conservation value to moderate conservation value, this is likely due to the increase of water beetle *Coleoptera* species and the decline of low scoring taxa such as the water louse *Asellus aquaticus*.

- 4.4.8 The change in PSI scores between summer and autumn is likely to be primarily a result of low flow conditions and extensive macrophyte coverage (e.g. bulrush (*Typha* spp.)), detritus and, in stream netting trapping debris. The PSI readings correlate with the LIFE readings, with slower velocities we see increased sediment tolerant species such as worms and Black fly larvae (*Simuliidae*).
- 4.4.9 The biotic indices are fairly similar when compared to the 2020 data, with slight reduction in the BMWP in autumn from 49 in 2020 to 40 in 2022. The PSI has also worsened but still remains in the same sedimented category. The ASPT and the CCI has actually improved in 2022 when compared to the 2020 data, this could be due to the presence of taxa such as the mayfly *Ephemeroptera* in large numbers over both seasons, which is a high BMWP scoring taxa of 10.

### Fish

**Table 11: River Mole fish species data collected during the autumn electrofishing survey.**

Autumn					
Overall					
Species	Latin name	Abundance	Mean Size (mm)	Min size (mm)	Max size (mm)
Dace	<i>Leuciscus leuciscus</i>	18	127.1428571	100	160
Chub	<i>Squalius cephalus</i>	12	247.5	120	440
Roach	<i>Rutilus rutilus</i>	35	139.3103448	100	210
Gudgeon	<i>Gobio gobio</i>	2	<100	<100	<100
Pike	<i>Esox lucius</i>	3	243.3333333	150	410
Minnow	<i>Phoxinus phoxinus</i>	1	<100	<100	<100

- 4.4.10 Fish surveys on the River Mole were only undertaken during autumn on the River Mole due to a major pollution event (defined by the EA as a Category 1 pollution event) that occurred on the watercourse during the summer of 2022. As for the Gatwick Stream, the fish community is typical of a slow flowing freshwater stream due to the presence of pike and roach (Table 11). The River Mole is also rich in macrophyte communities so provides a suitable environment for chub and dace which favour shaded habitat.
- 4.4.11 The range of sizes indicates a diverse age range for species such as the chub and pike, with smaller classes for the dace, roach and gudgeon, especially when compared to the Gatwick Stream.
- 4.4.12 Although fish abundance at the River Mole site is lower than that of the Gatwick Stream, it is similar to the autumn results of the 2020 survey in which five species were recorded (chub, roach, tench, pike and perch) and the total abundance was 28 individuals. The results from the

autumn 2022 visit suggest that there has been some recovery from the pollution event which occurred prior to the summer visit. Tench (*Tinca tinca*) was absent from the River Mole sample site during autumn 2022 despite having been recorded in both spring and autumn surveys in 2020.

**4.4.13** There are some species composition changes when compared to the 2020 data. Most notably the absence of perch, bream and rudd. However, there was a significant reduction or total loss in the autumn 2020 of these species so they likely struggled to repopulate. The River Mole provides optimal conditions for species such as the bream as there is a stable food source of worms and other macroinvertebrates and they are tolerant to low oxygen levels and sedimentation. The rudd and perch however are less tolerant to poor water quality so is likely to be a contributing factor to the diminishing population.

#### 4.5 Pond F

##### Water Quality

Table 12: Water quality at Pond F

Season	Temperature (°C)	DO (%)	DO (mg/L)	Conductivity (µS/cm)	pH	ORP	Turbidity (NTU)
Summer							
Autumn	14.1	84	8.6	324	7.64	156.7	20

**4.5.1** Pond F underwent a large variation in temperature between summer and autumn of 8.1°C, experiencing the highest temperature of the 4 sites in summer at 22.2 °C (Table 12). The pond is an artificial feature with no significant inflow or outfall, and therefore mixing is limited. The prolonged period of no rainfall combined with exceptionally high temperatures during in summer 2022 will have reduced water levels from evaporation and resulted in heating of the remaining water. The warming effect would have been exacerbated by the lack of shading.

**4.5.2** Despite high water temperatures DO were relatively high during the summer visit (77.2%) and increased to 84% in the autumn. That may have been due to oxygenation of the water column by green algae (although chlorophyll levels were not measured) Additionally, turbidity increased from 12.1NTU to 20NTU which may reflect die off of algae in the autumn. There was a drop in conductivity from 520 µS/cm to 324 µS/cm. Similarly to the River Mole, the ORP levels at Pond F saw a significant increase from -30.1 to 146.7, an decrease would be expected due to die back of macrophytes and algae.

*Macroinvertebrates*

**Table 13: Macroinvertebrate biotic indices at Pond F**

Biotic Index	Summer	Autumn
	02/08/2022	11/10/2022
BMWP (TL1)	55	40
NTAXA (TL1)	13	11
ASPT (TL1)	4.23	3.64
LIFE (TL5)	6.14	5.50
PSI (TL5)	0.00	0.00
EPSI (TL5)	0.00	17.67
CCI (TL5)	8.33	11.25
WHPT (TL2)	42.4	47.2
NTAXA (TL2)	13	13
ASPT (TL2)	3.26	3.63

- 4.5.3** The main composition changes between the two seasons taxa are the presence of water flea (*Cladocera*), 2 species of water beetle (*Halipus ruficollis*), biting midges (*Ceratopogonidae*), meniscus midges (*Dixella* sp.), smooth rams horn snail (*Gyraulus laevis*) and worms (*Oligochaeta*). Whereas the summer had different taxa that was not present in autumn such as non-biting midges *Chironomini Gen sp.*, mayfly *Baetidae*, Ramshorn snails *Planorbidae*, leech species *Glossiphonia complanate*, common blue dragonfly *Enallagma cyathigerum*, and ruddy darter dragonfly *Sympetrum sanguineum*.
- 4.5.4** Macroinvertebrate abundance was significantly greater in the autumn (690 individuals compared with 242 in the summer) although diversity remained relatively consistent between the seasons (16 taxa in summer compared with 17 in autumn). The increase in abundance in the autumn is largely due to the presence of 470 common water fleas (*Cladocera*).
- 4.5.5** There was a significant decline in BMWP score between the summer and autumn sample from 55 to 40 taking it from moderate to poor (Table 13). Alongside this there was a reduction in the LIFE score which was also the greatest reduction (from 6.14 to 5.50) amongst the 3 sites, remaining in sluggish conditions. However, LIFE score is not considered to be important in the context of a standing water body as it is based on the sensitivity of macroinvertebrate taxa to low flow conditions. The PSI didn't change and remained at the poorest level of 0 indicating sedimented conditions. The CCI however, improved from summer to autumn increasing from the second highest score of 8.33, indicating moderate conservation value to the greatest score of 11.25, indicating fairly high conservation value.
- 4.5.6** Reduction of the BMWP of Pond F from moderate (55) to poor (40) can be as a result of the loss of high scoring taxa from summer to autumn such as the Ruddy Darter larvae *Libellulidae*, the loss of which is likely due to the 0 PSI score, revealing sluggish conditions, further supported by the LIFE score. Additionally, as a result in the decline of water quality indicated by the ASPT from 4.23 to 3.64, showing levels from poor to very poor.

**4.5.7** Despite the BMWP reducing, there was an increase in the CCI between summer and autumn, this could be due to the establishment of a sediment tolerant community and the nationally scarce, smooth ramshorn snail *Gyraulus laevis*. Additionally, the presence of *Sigara limitata*, a species of water boatman, contributed to a slightly higher score of 11.25 at Pond F in Autumn from the previous moderate score.

**4.6 River Mole confluence**

*Water Quality*

**Table 14: Water quality at the River Mole confluence**

Season	Temperature (°C)	DO (%)	DO (mg/L)	Conductivity (µS/cm)	pH	ORP	Turbidity (NTU)
Summer	19.9	77.3	7.02	884	7.38	84.8	6.82

**4.6.1** The water temperatures at the River Mole confluence were high which is reflective of the high air temperatures experienced in the summer (Table 14). The conductivity and pH is typical of freshwater conditions. The turbidity is fairly low at 6.82 which is expected as the two sites (Gatwick Stream and the River Mole) that feed into this site, also had low turbidity readings for the summer.

*Macroinvertebrates*

**Table 15: Macroinvertebrate biotic indices at the River Mole confluence**

Biotic Index	Summer
	<i>02/08/2022</i>
BMWP (TL1)	52
NTAXA (TL1)	12
ASPT (TL1)	4.33
LIFE (TL5)	7.25
PSI (TL5)	35.29
EPSI (TL5)	47.51
CCI (TL5)	8.57
WHPT (TL2)	51.0
NTAXA (TL2)	12
ASPT (TL2)	4.25

**4.6.2** There was a total of 16 different taxa found in the summer sample at the River Mole confluence with the dominant species being freshwater shrimp *Gammaridae*, worms *Oligochaeta* and non-biting midge larvae *Chironomini*.

**4.6.3** The BMWP of 52 suggests a moderately impacted waterbody, this is likely influenced by the large numbers of mid scoring freshwater shrimp (Table 15).

- 4.6.4** A PSI of 35.29 indicates sedimented conditions, with the presence of tolerant taxa such as worms and freshwater shrimp underlying the score. This compliments the sluggish flow conditions reflected by the LIFE score of 7.25.

---

## 5. Discussion

- 5.1.1** High water temperatures were recorded at all three sites in summer, and notably on the Gatwick Stream where a temperature of 19.7°C was recorded. According to the Met Office, the UK recorded the warmest summer in 2022 and the driest year since 1976. Summer water temperatures in the Gatwick Stream were three degrees higher than for the same period in 2020. Although the sampling sites were in different locations both sites were shaded by bank side trees which suggests that the differences were due to a combination of low flow conditions and exceptionally high ambient temperatures. High dissolved oxygen concentrations during both summer and autumn are unexpected given that oxygen is less soluble in warm water. This may have been due to hydromorphological features in the vicinity of the site, such as a riffle, which creates turbulence and introduces oxygen.
- 5.1.2** Although high dissolved oxygen may suggest good water quality, based on both the 2020 and 2022 data the Gatwick Stream supported a macroinvertebrate assemblage indicative of moderate to poor water quality. Slightly higher scores were obtained from the upstream sampling site in 2020, suggesting that there may be a pollution source such as a storm overflow between the two sampling sites. The sampling location for the 2022 survey was located in Riverside Park approximately 650m downstream of the 2020 upstream site and 450m upstream from the confluence with the River Mole. There was an increase in diversity and abundance of macroinvertebrate taxa between the summer and the autumn sample at this site, with a consequential increase in biotic scores. This was considered to be due to an autumn re-charge of the watercourse resulting in higher flow velocities and dissolved oxygen concentrations. Average Score Per Taxon (ASPT) scores from the August 2022 sample were lower (3.86) than either the up or downstream site in July 2020 (4.53 and 3.91 respectively). This may be due to discharges affecting water quality throughout the reach, combined with the effects of low flows and higher water temperatures in 2022. An ASPT of 5 was recorded for the October 2022 sample compared with 4.13 and 2.88 in September 2020, possibly indicating that a recovery in macroinvertebrate diversity increases through the autumn period.
- 5.1.3** The record of bullhead from the autumn sampling round on the Gatwick Stream is notable since the species is listed on both Annex II of the EU Habitats Directive and the IUCN Red list. Although Annex II species are qualifying features for the designation of sites as Special Area of Conservation (SAC) the presence of bullhead, a species that is relatively widespread in rivers in England, does not confer special status on the watercourse. However, measures to enhance the channel for fish as part of the development masterplan would be desirable.
- 5.1.4** The reductions in abundance of chub and gudgeon could have been due to the introduction of predators such as the pike, increasing the competition for prey. Additionally, the presence of a European Mink (*Mustela lutreola*) was noted whilst conducting the survey, this could have contributed to the reduction of the larger species as they are having to compete for food.
- 5.1.5** Burrowing by Signal Crayfish may be contributing to sedimentation of the channel, with bank stabilisation presenting a potential increased flood risk. However, fewer individuals were recorded during 2022 compared with 2020 which suggests that the population may be stable.



---

Additionally, there were 6 records in autumn and 8 recordings in summer of the invasive New Zealand mud snail (*Potamopyrgus antipodarum*), a decline from the 12 and 40 recorded in 2020.

- 5.1.6** In both 2020 and 2022 there was a reduction in PSI scores on the River Mole between summer and autumn, likely reflecting low flow conditions leading to sediment accretion. Autumn die back of the dense in-stream macrophyte beds on the River Mole combined with in-stream netting trapping debris and silt. Nevertheless, the Community Conservation Index for the River Mole indicated moderate conditions, possibly due to the presence of water beetle species (Coleoptera).
- 5.1.7** An environmental pollution incident categorised by the Environment Agency as 'Category 1', major incident, occurred on the River Mole during summer 2022 (ID number 106775). The impact level only affected the water through the presence of surfactants and detergents. Detergents and surfactants affect aquatic biota by causing the breakdown of mucus membranes that coats fish, making them more susceptible to parasites and bacteria. They also reduce the surface tension of the water, making it easier for aquatic biota to absorb pesticides and other pollutants in the water. This is particularly important as this stretch of the Mole failed the chemical status in the most recent assessment (2019) by the EA. Concentrations at high volumes (15ppm) can cause fish mortality and lower concentrations (5ppm) can cause fish eggs to die, there is also potential for chronic and sublethal effects to occur when left untreated. Finally, the phosphates within the detergents can contribute to oxygen depletion through algal blooms in the waterways which has already been exacerbated by the high temperatures that we have recorded.
- 5.1.8** No Tench (*Tinca tinca*) were recorded in the River Mole, a species that was present during the 2020 survey. This is unusual due to their diverse diets and hardy nature allowing them to be more resilient to aspects such as low DO concentrations, high turbidity and pollution events. Tench are hardy species that can tolerate high levels of pollution and sedimentation, so it is surprising that no Tench were found in the 2022 surveys, especially given that the dense macrophyte stands and variable depths offer ideal breeding habitat. However, the small sizes of Tench in the 2020 surveys could indicate that none were of breeding size and as the numbers were so low, no fry were produced. Predation by fish such as pike may also have impacted the population.

---

## 6. Conclusions and recommendations

### 6.1 Conclusions

- 6.1.1 The macroinvertebrate assemblages recorded on the Gatwick Stream and River Mole during summer and autumn 2022 indicate moderate to poor water quality and are therefore consistent with WFD classification of the waterbodies by the Environment Agency.
- 6.1.2 Exceptionally hot weather and drought conditions during summer 2022 combined with a pollution incident in the months prior to the summer survey prevented sampling for fish on the River Mole. There appeared to be no acute or chronic impacts from the pollution event on the macroinvertebrate community. An increase in biotic scores between summer and autumn 2022 is likely to be as a result of a recovery in flow conditions during autumn.
- 6.1.3 The record of bullhead from the Gatwick Stream is notable and measures to safeguard the population will need to be incorporated into the mitigation strategy for the Gatwick airport expansion development scheme.
- 6.1.4 Siltation was identified as a problem on the River Mole based on LIFE and PSI scores from data collected in both 2020 and 2022. This will need to be considered in the detailed design of the new river channel. The new river diversion should be designed to maximise flow velocity.

### 6.2 Recommendations

- 6.2.1 The design of the outfall structure from the proposed new water treatment plant discharging into the Gatwick Stream should take account of the presence of bullhead. This should include minimising the loss of bankside and channel habitat. The outfall should also be designed to minimise scour from flows discharged through the outfall.
- 6.2.2 The new river diversion should be designed to increase habitat heterogeneity through variable flow patterns and the creation of new channel features such as pools, side bars and points bars. Any planting should avoid bulrush which tends to dominate the vegetation community and choke the channel, particularly under slow flowing and eutrophic conditions.
- 6.2.3 There should be a plan for the management of the river diversion and associated floodplain of the River Mole. Management of bulrush stands would need to be informed by an understanding of the distribution of water voles on the site. To minimise impacts on water voles management of the bulrush would need to be implemented on a careful rotation ensuring that vegetation was always retained on one side of the channel.
- 6.2.4 There should also be a plan for the management of invasive species on both the River Mole and Gatwick Stream. Management of the Signal crayfish population on the Gatwick Stream will be essential since burrowing by the species is currently contributing to the sedimentation of the Gatwick Stream and has the potential to cause bank instability and increase flood risk.

---

## 7. References

- 7.1.1** BS EN 27828:1994, ISO 7828-1985 Water quality. Methods for biological testing. Methods of biological sampling: guidance on hand-net sampling of aquatic benthic macro-invertebrates;
- 7.1.2** British Standards (2003) BS EN 14011:2003 Water Quality: Sampling of Fish with Electricity.
- 7.1.3** Chadd, R. Extence, C. (2004) The conservation of freshwater macroinvertebrate populations: A community based classification scheme. Aquatic Conservation Marine and Freshwater Ecosystems. Vol 14. Pp. 597-624.
- 7.1.4** Clarke, R. (2011) Hydroseral habitat requirement of the endangered Shining Ramshorn Snail *Segmentina nitida*. Aquatic Conserv: Mar. Freshw. Ecosyst. 14: 597-624
- 7.1.5** Department for Environment Food and Rural Affairs; undated. [England | Catchment Data Explorer](#).
- 7.1.6** Environment Agency and Pond Action (2002) A guide to monitoring the ecological quality of ponds and canals using PSYM. PSYM Manual.
- 7.1.7** Extence, C.A, Balbi, D.M. Chadd, R.P. (1999) River Flow Indexing Using British Macroinvertebrates: A Framework For Setting Hydroecological Objectives. Regulated Rivers: Research and Management 15. pp543-574
- 7.1.8** Extence, C.A. Chadd, R.P. England, J. Naura, M. Pickwell, A.G.G. (2017) Application of the Proportion of Sediment-Sensitive Invertebrates (psi) Biomonitoring Index. River Research and Applications. Vol 33. Issue 10. December 2017. Pp. 1596 - 1605.
- 7.1.9** Paisley, M.F. Trigg, D.J. Walley, W.J. (2013) Revision of the Biological Monitoring Working Party (BMWP) score system: Derivation of present-only and abundance-related scores from field data. <https://onlinelibrary.wiley.com/doi/abs/10.1002/rra.2686>
- 7.1.10** Suffolk Biological Information Service (2003) Shining Ram's Horn Snail (*Segmentina nitida*) Biodiversity Action Plan. <https://www.suffolkbis.org.uk/sites/default/files/biodiversity/priorityspecieshabitats/actionplans/shiningramshornsnail.pdf>
- 7.1.11** Water Framework Directive - United Kingdom Advisory Group. UKTAG River Assessment Method. Benthic Invertebrate Fauna. Invertebrates (General Degradation): Whalley, Hawkes, Paisley & Trigg (WHPT) metric in River Invertebrate Classification Tool (RICT). July 2014. ISBN: 978-1-906934-49-1.
- 7.1.12** UK TAG (2021) UKTAG River Assessment Method Benthic Invertebrate Fauna: Invertebrates (General Degradation): Walley, Hawkes, Paisley & Trigg (WHPT) metric in River Invertebrate Classification Tool (RICT). WFD UK Technical Advisory Group.

## Appendix 1 - Taxa lists

Gatwick Stream			Summer	Autumn
Order	Family	Taxa Name	Abundance	Abundance
AMPHIPODA	CRANGONYCTIDAE	Crangonyx pseudogracilis/floridanus agg.		15
AMPHIPODA	GAMMARIDAE	Gammarus pulex/fossarum agg.	5	
AMPHIPODA	GAMMARIDAE	Gammarus pulex	4	2
BIVALVIA	SPHAERIIDAE	Euglesa sp.		
CLADOCERA	N/A	Cladocera		
COLEOPTERA	DYTISCIDAE	Hygrotus inaequalis		
COLEOPTERA	DYTISCIDAE	Agabus/Ilybius sp.		
COLEOPTERA	HALIPLIDAE	Halipus ruficollis		
COLEOPTERA	HALIPLIDAE	Halipus ruficollis group		
COLEOPTERA	HALIPLIDAE	Halipus sp.		
CYCLOPOIDA	N/A	Cyclopoida		
DIPTERA	CERATOPOGONIDAE	Ceratopogonidae		
DIPTERA	CHIRONOMIDAE	Chironomini Gen. sp.	6	
DIPTERA	CHIRONOMIDAE	Chironomidae	20	54
DIPTERA	CHIRONOMIDAE	Tanypodinae		
DIPTERA	CHIRONOMIDAE	Tanytarsini		1
DIPTERA	CULICIDAE	Anopheles sp.		
DIPTERA	DIXIDAE	Dixella sp.		
DIPTERA	SIMULIIDAE	Simulium sp.		11
EPHEMEROPTERA	BAETIDAE	Cloeon dipterum		
GASTROPODA	PHYSIDAE	Physella acuta/heterostropha		
GASTROPODA	PLANORBIDAE	Gyraulus laevis		
GASTROPODA	TATEIDAE	Potamopyrgus antipodarum	8	6
HEMIPTERA	CORIXIDAE	Callicorixa praeusta		
HEMIPTERA	CORIXIDAE	Corixidae		
HEMIPTERA	CORIXIDAE	Sigara sp.		
HEMIPTERA	NOTONECTIDAE	Notonecta glauca		
HIRUDINEA	ERPOBDELLIDAE	Erpobdella testacea		
HIRUDINEA	GLOSSIPHONIIDAE	Helobdella stagnalis		
HIRUDINEA	GLOSSIPHONIIDAE	Theromyzon tessulatum		
ISOPODA	ASELLIDAE	Asellus aquaticus	1	2
MEGALOPTERA	SIALIDAE	Sialis lutaria		
N/A	N/A	Ostracoda		
ODONATA	CALOPTERYGIDAE	Calopteryx sp.		2
ODONATA	CALOPTERYGIDAE	Calopteryx splendens		2
ODONATA	CALOPTERYGIDAE	Calopteryx virgo		1
OLIGOCHAETA	N/A	Oligochaeta	76	96
TRICHOPTERA	GLOSSOSSOMATIDAE	Agapetus fuscipes	1	
TRICHOPTERA	HYDROPSYCHIDAE	Hydropsyche angustipennis	31	10
TRICHOPTERA	HYDROPSYCHIDAE	Hydropsyche pellucidula		9
TRICHOPTERA	LEPTOCERIDAE	Mystacides sp.		1
TRICHOPTERA	POLYCENTROPODIDAE	Cyrnus trimaculatus		1
TRICLADIDA	DENDROCOELIDAE	Dendrocoelum lacteum		
TRICLADIDA	DUGESIIDAE	Schmidtea lugubris/polychroa		
TRICLADIDA	DUGESIIDAE	Girardia tigrina		
TRICLADIDA	PLANARIIDAE	Polycelis felina		

River Mole			Summer	Autumn
Order	Family	Taxa Name	Abundance	Abundance
AMPHIPODA	CRANGONYCTIDAE	Crangonyx pseudogracilis/floridanus agg.	5	22
AMPHIPODA	GAMMARIDAE	Gammarus pulex	1	
BIVALVIA	SPHAERIIDAE	Sphaerium corneum	25	
BIVALVIA	SPHAERIIDAE	Euglesa subtruncata	4	
BIVALVIA	SPHAERIIDAE	Euglesa sp.	3	
CLADOCERA	N/A	Cladocera		108
COLEOPTERA	DYTISCIDAE	Hygrotus inaequalis	5	1
COLEOPTERA	DYTISCIDAE	Agabus/Ilybius sp.		3
COLEOPTERA	HALIPLIDAE	Haliphus ruficollis	1	2
COLEOPTERA	HALIPLIDAE	Haliphus ruficollis group		20
COLEOPTERA	HALIPLIDAE	Haliphus sp.	2	6
CYCLOPOIDA	N/A	Cyclopoida		1
DIPTERA	CERATOPOGONIDAE	Ceratopogonidae		
DIPTERA	CHIRONOMIDAE	Chironomidae	3	12
DIPTERA	CHIRONOMIDAE	Tanypodinae	1	
DIPTERA	CHIRONOMIDAE	Tanytarsini		
DIPTERA	CULICIDAE	Anopheles sp.		1
DIPTERA	DIXIDAE	Dixella sp.		
DIPTERA	SIMULIIDAE	Simulium sp.		
EPHEMEROPTERA	BAETIDAE	Cloeon dipterum	35	27
GASTROPODA	PHYSIDAE	Physella acuta/heterostropha		
GASTROPODA	PLANORBIDAE	Gyraulus laevis		
GASTROPODA	PLANORBIDAE	Hippeutis complanatus	1	
GASTROPODA	TATEIDAE	Potamopyrgus antipodarum		
GASTROPODA	VALVATIDAE	Valvata piscinalis	8	
HEMIPTERA	CORIXIDAE	Callicorixa praeusta		1
HEMIPTERA	CORIXIDAE	Corixidae	1	
HEMIPTERA	CORIXIDAE	Sigara dorsalis	1	
HEMIPTERA	CORIXIDAE	Sigara sp.	6	1
HEMIPTERA	NOTONECTIDAE	Notonecta glauca		6
HIRUDINEA	ERPOBDELLIDAE	Erpobdella testacea		5
HIRUDINEA	GLOSSIPHONIIDAE	Helobdella stagnalis	3	2
HIRUDINEA	GLOSSIPHONIIDAE	Theromyzon tessulatum		1
ISOPODA	ASELLIDAE	Asellus aquaticus	196	59
MEGALOPTERA	SIALIDAE	Sialis lutaria	3	
N/A	N/A	Ostracoda		
ODONATA	CALOPTYRIDAE	Calopteryx sp.		
ODONATA	CALOPTYRIDAE	Calopteryx splendens		
ODONATA	CALOPTYRIDAE	Calopteryx virgo		
ODONATA	COENAGRIONIDAE	Enallagma cyathigerum		
ODONATA	LIBELLULIDAE	Sympetrum sanguineum		
OLIGOCHAETA	N/A	Oligochaeta	35	24
TRICHOPTERA	HYDROPSYCHIDAE	Hydropsyche angustipennis		
TRICHOPTERA	HYDROPSYCHIDAE	Hydropsyche pellucidula		
TRICHOPTERA	LEPTOCERIDAE	Mystacides sp.		
TRICHOPTERA	POLYCENTROPODIDAE	Cynus trimaculatus		
TRICLADIDA	DENDROCOELIDAE	Dendrocoelum lacteum		1
TRICLADIDA	DUGESIIDAE	Schmidtea lugubris/polychroa		4
TRICLADIDA	DUGESIIDAE	Girardia tigrina	2	
TRICLADIDA	PLANARIIDAE	Polycelis felina	5	29
TROMBIDIFORMES	N/A	Hydracarina	2	

Pond F			Summer 2022	Autumn 2022
Order	Family	Taxa Name	Abundance	Abundance
AMPHIPODA	CRANGONYCTIDAE	Crangonyx pseudogracilis/floridanus agg.	3	48
AMPHIPODA	GAMMARIDAE	Gammarus pulex		
BIVALVIA	SPHAERIIDAE	Euglesa sp.	2	17
CLADOCERA	N/A	Cladocera		470
COLEOPTERA	DYTISCIDAE	Hygrotus inaequalis		
COLEOPTERA	DYTISCIDAE	Agabus/llybius sp.		
COLEOPTERA	HALIPLIDAE	Halipus ruficollis		1
COLEOPTERA	HALIPLIDAE	Halipus ruficollis group		
COLEOPTERA	HALIPLIDAE	Halipus sp.		6
CYCLOPOIDA	N/A	Cyclopoida		
DIPTERA	CERATOPOGONIDAE	Ceratopogonidae		30
DIPTERA	CHIRONOMIDAE	Chironomini Gen. sp.	65	
DIPTERA	CHIRONOMIDAE	Chironomidae	115	24
DIPTERA	CHIRONOMIDAE	Tanypodinae	13	2
DIPTERA	CHIRONOMIDAE	Tanytarsini		
DIPTERA	CULICIDAE	Anopheles sp.		
DIPTERA	DIXIDAE	Dixella sp.		1
DIPTERA	SIMULIIDAE	Simulium sp.		
EPHEMEROPTERA	BAETIDAE	Cloeon dipterum	1	
GASTROPODA	PLANORBIDAE	Planorbidae	1	
GASTROPODA	PHYSIDAE	Physella acuta/heterostropha	2	21
GASTROPODA	PLANORBIDAE	Gyraulus laevis		16
GASTROPODA	TATEIDAE	Potamopyrgus antipodarum		
HEMIPTERA	CORIXIDAE	Callicorixa praeusta		
HEMIPTERA	CORIXIDAE	Corixidae	1	1
HEMIPTERA	CORIXIDAE	Sigara sp.		
HEMIPTERA	NOTONECTIDAE	Notonecta glauca		
HIRUDINEA	ERPOBDELLIDAE	Erpobdella testacea		
HIRUDINEA	GLOSSIPHONIIDAE	Glossiphonia complanata	1	
HIRUDINEA	GLOSSIPHONIIDAE	Helobdella stagnalis		
HIRUDINEA	GLOSSIPHONIIDAE	Theromyzon tessulatum		
ISOPODA	ASELLIDAE	Asellus aquaticus	1	3
MEGALOPTERA	SIALIDAE	Sialis lutaria	3	1
N/A	N/A	Ostracoda	29	8
ODONATA	CALOPTERYGIDAE	Calopteryx sp.		
ODONATA	CALOPTERYGIDAE	Calopteryx splendens		
ODONATA	COENAGRIONIDAE	Enallagma cyathigerum	2	
ODONATA	LIBELLULIDAE	Sympetrum sanguineum	2	
ODONATA	CALOPTERYGIDAE	Calopteryx virgo		
OLIGOCHAETA	N/A	Oligochaeta		13
TRICHOPTERA	HYDROPSYCHIDAE	Hydropsyche angustipennis		
TRICHOPTERA	HYDROPSYCHIDAE	Hydropsyche pellucidula		
TRICHOPTERA	LEPTOCERIDAE	Mystacides sp.		
TRICHOPTERA	POLYCENTROPODIDAE	Cyrnus trimaculatus		
TRICLADIDA	DENDROCOELIDAE	Dendrocoelum lacteum		
TRICLADIDA	DUGESIIDAE	Schmidtea lugubris/polychroa		
TRICLADIDA	DUGESIIDAE	Girardia tigrina	1	28
TRICLADIDA	PLANARIIDAE	Polycelis felina		



**Gatwick Shining Ramshorn  
Snail (*Segmentina nitida*)**

**Aquatic mollusc survey 2022**

For

**RPS Ltd**

Project No.: RPS001-022

August 2022



**London & South East**

Compass House  
Surrey Research Park  
Guildford  
GU2 7AG, UK  
t: +44 (0)1483 466 000

**North East**

Brooklands Court Business Centre  
Tunstall Road  
Leeds  
LS11 5HL, UK  
t: +44 (0)113 518 2121

**North West**

St James Tower  
7 Charlotte Street  
Manchester  
M1 4DZ  
t: +44 (0)113 518 2121

**Wales & South West**

Sophia House  
28 Cathedral Road  
Cardiff  
CF11 9LJ UK  
t: +44 (0)292 066 0180

**Midlands**

605 Holly Court, Holly Farm Business Park  
Honiley, Kenilworth  
Warwickshire  
CV8 1NP  
t: +44 (0) 121 827 3180

**Enquiries**

e: [enquiries@thomsonec.com](mailto:enquiries@thomsonec.com)

w: [www.thomsonec.com](http://www.thomsonec.com)



Project Number	Report No.
RPS001-022-001	002

Revision No.	Date of Issue	Author	Reviewer	Approver
001	31/01/2023	Toby Abrehart	Alexander Charlesworth	Tessa Harding
002	Date			
003	Date			

**Disclaimer:**

Copyright Thomson Environmental Consultants Limited. All rights reserved.

No part of this report may be copied or reproduced by any means without prior written permission from Thomson Environmental Consultants Limited. If you have received this report in error, please destroy all copies in your possession or control and notify Thomson Environmental Consultants Limited.

This report has been prepared for the exclusive use of the commissioning party and unless otherwise agreed in writing by Thomson Environmental Consultants Limited, no other party may use, make use of or rely on the contents of the report. No liability is accepted by Thomson Environmental Consultants Limited for any use of this report, other than for the purposes for which it was originally prepared and provided.

Opinions and information provided in the report are on the basis of Thomson Environmental Consultants Limited using due skill, care and diligence in the preparation of the same and no explicit warranty is provided as to their accuracy. It should be noted and it is expressly stated that no independent verification of any of the documents or information supplied to Thomson Environmental Consultants Limited has been made.

## Contents

1. Summary and Recommendations .....	5
1.1 Summary .....	5
1.2 Recommendations.....	5
2. Introduction .....	6
2.1 Development Background .....	6
2.2 Ecology Background.....	6
2.3 The Brief and Objectives .....	6
3. Methodology .....	7
3.1 Sample point locations .....	7
3.2 Sampling method.....	8
3.3 Biosecurity .....	8
3.4 Limitations .....	8
4. Results.....	9
5. References .....	10
Appendix 1 - Sample data .....	11
Table 3-1: Sample site locations .....	8

Figure 1 Site locations, survey area and species locations

## 1. Summary and Recommendations

### 1.1 Summary

- 1.1.1** Thompson Environmental Consultants in association with Abrehart Ecology Ltd was commissioned by RPS Ltd to undertake an aquatic mollusc survey of River Mole, running west of the main Gatwick Airport site.
- 1.1.2** The study aimed to update the records and status of the possible presence of Shining Ramshorn Snail (*Segmentina nitida*) along the stretch of the River Mole which lies within the development area for the proposed expansion at Gatwick airport. Additionally, the survey was used to evaluate the conservation value and the overall condition of the site. Samples were taken in July 2022.
- 1.1.3** The site comprised a single water course, the River Mole, running approximately north south, with a meandering channel that heads east. The river had been cleared in the past year with the material placed on the banks.
- 1.1.4** The channel was steep sided with abundant emergent and floating vegetation; all habitats were in early succession.
- 1.1.5** Ten samples were collected from both sides of the channel. Fourteen species of aquatic mollusc were found and identified. No species of conservation significance were found during the survey, all were common species of water courses.
- 1.1.6** No Shining Ramshorn Snails were found in any of the samples and the habitat appeared to currently be unsuitable to support it. Only the similar looking Flat Ram's-horn snail (*Hippeutis complanata*) was recorded.
- 1.1.7** Three invasive species were found across the survey areas: the common freshwater shrimp *Crangonyx pseudogracilis*, the nationally common mollusc *Potamopyrgus antipodarum*, and the invasive plant Himalayan Balsam, which was abundant along the river margins.

### 1.2 Recommendations

- 1.2.1** The results of the survey provide evidence that the Shining Ram's Horn snail is absent from this reach of the River Mole. No further surveys are required and the species does not need to be considered in the Environmental Impact Assessment for the proposed Gatwick expansion scheme.

## 2. Introduction

### 2.1 Development Background

2.1.1 The River Mole will be directly affected by proposals to expand operations at Gatwick airport. The project proposes alterations to the existing northern runway, and development of a range of infrastructure and facilities to increase passenger numbers and aircraft movements. Specifically in relation to the River Mole the development will include creation of additional flood storage to the west of Gatwick Airport within the flood plain of the River Mole. An existing, approximately 300m long straightened section of the River Mole immediately north of the airport will be re-meandered to create a natural sinuous course.

### 2.2 Ecology Background

2.2.1 A single record record of Shining Ramshorn Snail (*Segmentina nitida*), dating back to 2013, was returned in desk study data received from Sussex Biodiversity Records Centre supporting macroinvertebrate and fish surveys undertaken by Thomson Environmental Consultants for the project in 2020. The Shining Ramshorn Snail is nationally scarce<sup>1</sup>, a UK Priority Species under the UK Post 2010 Biodiversity Framework, and listed on the Sussex Rare Species Inventory.

2.2.2 A recommendation was made from the 2020 study to undertake a survey to establish the presence or absence of Shining Ramshorn Snail. It was recommended that the survey should focus initially on the site of the 2013 record, and if found to be present, extended to incorporate the whole study section. It was also recommended that surveys should be scoped and undertaken by a specialist mollusc ecologist.

### 2.3 The Brief and Objectives

2.3.1 RPS commissioned Thomson Environmental Consultants on 16/06/2022 to undertake Shining Ram's Horn Snail surveys on the River Mole in the vicinity of the 2013 record. The survey was undertaken by specialist mollusc consultants, Abrehart Ecology Ltd. The brief was to:

- Determine the presence/absence of Shining Ramshorn Snail in the vicinity of the 2013 record, and if found to be present extend the survey to include the whole length of the section that will be affected by the scheme.
- Provide a report on the survey giving the methods and results of the survey, with recommendations, including, if necessary, measures to mitigate for the species.

2.3.2 This report presents the findings of the survey.

---

<sup>1</sup> Occurring in 16-100 hectads in Great Britain. Excludes rare species qualifying under the main IUCN criteria. This category replaces Notable, Notable A and Notable B.

### 3. Methodology

#### 3.1 Sample point locations

**3.1.1** Sampling points were chosen once on site to allow the most diverse habitats to be sampled, rather than be restricted to previous sample locations. Data and sample collection were undertaken by one experienced on-site mollusc surveyor (Toby Abrehart FLS MCIEEM). All the sampling was undertaken during July 2022.

**3.1.2** The study area encompassed the banks, aquatic margins, and localised areas of floating and submerged vegetation of the River Mole (Figure 3-1). Ten sampling locations were selected and one sample taken from each (Table 3-1). The sample sites were identified by Toby Abrehart as suitable to survey and verified in the field.



**Figure 3-1: Map showing sample point locations**

**Table 3-1: Sample site locations**

Sample site	National Grid Reference location
1	TQ2562340908
2	TQ2561940901
3	TQ2563540903
4	TQ2563340900
5	TQ2560640857
6	TQ2556240794
7	TQ2553540733
8	TQ2551440697
9	TQ2555440620
10	TQ2553740560

### 3.2 Sampling method

**3.2.1** Samples were collected using thirty-second sweeps with a net with 0.5mm mesh. Sweeps were repeated three times in different sections of the waterbody profile, that is, floating vegetation (where present), the benthic layer, and the submerged edge of the nearside bank. Once collected species were identified in the field, with a focus on finding Shining Ramshorn Snail. A selection of the samples was placed into a 1-litre bucket and preserved in 99% Iso-propyl alcohol for confirmation and quality control within the Abrehart Ecology lab.

**3.2.2** For identification, all molluscs were separated from the retained sediment, detritus, and vegetation under x10 head mounted magnifiers and, in the laboratory, using a stereo binocular microscope. Where possible, all specimens were identified to species level.

### 3.3 Biosecurity

**3.3.1** All equipment used was subject to the check, clean, and dry methods prior to use on site. However, should surveying include the movement between systems then prior to entering a new waterbody, the net, and trays from one site would be washed in a solution of Virkon and left to dry. A clean and dry set was then used in the new waterbody. This prevented species or pathogens being transmitted from one area to another. On return to the laboratory the nets were washed again in Virkon solution and left to dry for at least one day before being taken into the field.

### 3.4 Limitations

**3.4.1** The survey was carried out in optimal conditions.

**3.4.2** Some of the marginal habitats were very dry at the time of the survey. It was apparent that the river had been cleared in the past year.

## 4. Results

- 4.1.1** The sampling sites were associated with the emergent, submerged, and floating aquatic vegetation within the river channel. The river ran through the valley bottom marshes; running to the east and west of these marshes in the north of the site, before joining halfway through the marshes to proceed south through the centre of the last marsh with a more meandering course.
- 4.1.2** The emergent vegetation within the channel was dominated by Arrowhead (*Sagittaria sagittifolia*), Flowering Rush (*Butomus umbellatus*), Branched Bur-reed (*Sparganium erectum*), Reed Canary Grass (*Phalaris arundinacea*), Common Water Plantain (*Alisma plantago-aquatica*), Hemlock Water Dropwort (*Oenanthe crocata*) and Purple Loosestrife (*Lythrum salicaria*). Floating aquatic vegetation was dominated by Yellow Water Lily (*Nuphar lutea*), Common Duckweed (*Lemna minor*). Submerged macrophytes included Canadian pond-weed (*Elodea canadensis*). The water was deep and clear with little to no successional habitat. The river had been recently cleared with the debris scattered over the floodplain. All successional habitat was surveyed if present; however, this was only found in Samples 1-4 in the north of the survey area.
- 4.1.3** No Shining Ram's Horn Snail were found in any of the samples, though thirteen species of mollusc and bivalve were identified in the samples (full results in Appendix A). The mollusc Flat Ram's-horn snail (*Hippeutis complanate*) was found in four samples. This species is superficially similar to *Segmentina nitida* and could have been mistaken for it in the past.
- 4.1.4** No other species of interest were found. The habitat appeared unsuitable to support Shining Ram's Horn Snail but was suitable for the more catholic Flat Ram's-horn snail, which was present in low to moderate numbers (as seen in Appendix A).
- 4.1.5** It appears possible that the single specimen found during the Water Quality Survey in 2013 may have been mis-identified or the habitat is no longer suitable to support it.



## 5. References

- 5.1.1 Kerans, B.L. et al. 2005. Potamopyrgus antipodarum: distribution, density, and effects on native macroinvertebrate assemblages in the Greater Yellowstone Ecosystem. *Journal of the North American Benthological Society* 24(1): 123-
- 5.1.2 Kerney, M., 1999. Atlas of the Land and Freshwater Molluscs of Britain and Ireland. The Conchological Society of Great Britain and Ireland. Harley Books, Great Horkesly.
- 5.1.3 Killeen, I., Aldridge, D. & Oliver, G., 2004. Freshwater Bivalves of Britain and Ireland. FSC Publications, Telford.
- 5.1.4 Reynoldson, T.B. & Young, J.O., 2000. A key to the triclads of Britain and Ireland with notes on their ecology. *Freshwater Biological Association Scientific Publication* 58. Freshwater Biological Association, Ambleside.

## Appendix 1 - Sample data

Species	1	2	3	4	5	6	7	8	9	10
<i>Acroloxus lacustris</i>	2	1	1		3					
<i>Ambullaceana balthica</i>	8	1	1	1	2			1		2
<i>Bathymphalis contortus</i>		1				1				
<i>Gyraulus albus</i>	12	3	4				1			
<i>Hipppeutis complanatus</i>	12	6	3		1					
<i>Pisidium henslowianum</i>	16	2	1							
<i>Planorbarius corneus</i>	2	1		13	19	12	9	6	9	11
<i>Planorbis planorbis</i>	4									
<i>Potamopyrgus antipodarum</i>			2							
<i>Segmentina nitida</i>	0	0	0	0	0	0	0	0	0	0
<i>Sphaerium corneum</i>	31	51	18	21	47	42	26	14	12	19
<i>Succinia putris</i>			1	3			1			1
<i>Valvata cristata</i>							2		1	1
<i>Valvata piscinalis</i>	3	16	12	21	38	8	12	4	2	5