

Gatwick Airport Northern Runway Project

Environmental Statement Appendix 9.6.3: Bat Trapping and Radio Tracking Surveys – Part 1

Book 5

VERSION: 1.0

DATE: JULY 2023

Application Document Ref: 5.3

PINS Reference Number: TR020005



Table of Contents		Table 4.2.4: Importance of foraging areas outside Proje	ct Area
1 Introduction	1	12	
2 Methodology	2	Table 7.1.1: Glossary of Terms	13
3 Results	5		
4 Discussion	10		
5 Conclusions	13		
6 References	13		
7 Glossary	13		
Annex			
Annex 1 Trapping data			
Annex 2 Home Range Analysis			
Annex 3 Figures			
Annex 4 2019 Survey Report			
Annex 5 Bat Collision Risk			
Tables			
Table 2.2.1: Trapping Locations	3		
Table 3.1.1: Trapping results by species	5		
Table 3.1.2: Trapping results by location	5		
Table 3.2.1: Radio-tagged bats	5		
Table 3.1.1: Roost Location, Count and Characterisation	8		
Table 4.2.1: Importance of roosts within Project Area	11		
Table 4.2.2: Importance of roosts outside Project Area	11		
Table 4.2.3: Importance of foraging areas within Project Are	ea 12		



1 Introduction

1.1 Overview

- 1.1.1 This document forms Appendix 9.6.3 of the Environmental Statement (ES) prepared on behalf of Gatwick Airport Limited (GAL) for the proposal to make best use of Gatwick Airport's existing runways and infrastructure (referred to within this report as 'the Project').
- 1.1.2 This report provides details of bat surveys undertaken on land within and around Gatwick Airport, Horley, West Sussex to inform the design of the Project, as described in **ES Chapter 5: Project Description** (Doc Ref. 5.1).
- 1.1.3 RPS contracted The Ecology Consultancy/Temple to provide specialist bat consultancy services, which included Advanced Bat Survey Techniques for the Project, including trapping and subsequent radio-tracking of target bat species.
- 1.1.4 Surveys undertaken in 2019 identified individuals from a population of Bechstein's bats that utilise the Project Area. Data for these surveys are presented in Annex 4. Due to restrictions to access within the wider landscape surrounding the Project Area during the surveys undertaken in 2019, it was considered likely that the evaluation of the importance of the site for bats was constrained by the lack of knowledge of the status of the Bechstein's bat population in the landscape surrounding the Project Area. Therefore, it was recommended that a landscape level population assessment of Bechstein's bats was undertaken in and adjacent to the Project Area to inform any mitigation proposals. The results of this assessment are presented in this report.
- 1.1.5 In addition, work was undertaken to examine the existing risk of bat collision within the airport in order to inform the requirements for any mitigation/avoidance measures from the Project. Due to the COVID-19 pandemic and associated subsequent reductions in air travel, work to parameterise the model was only partially completed. However, data and modelling to inform this work are presented in Annex 5.

1.2 Relevant legislation and planning policy

1.2.1 The following key pieces of nature conservation legislation are relevant to this assessment:

- The Conservation of Habitats and Species Regulations 2017 (as amended);
- The Wildlife and Countryside Act 1981 (as amended); and
- Natural Environment and Rural Communities Act 2006.
- 1.2.2 All native UK bat species are protected under Schedule 5 of the Wildlife and Countryside Act 1981 (as amended) and the Conservation of Habitats and Species Regulations 2017 (as amended). Under this combined legislation it is an offence to:
 - deliberately capture, injure or kill a bat;
 - intentionally or recklessly disturb a bat whilst occupying a place of shelter or protection;
 - possess or advertise/sell/exchange a bat (dead or alive) or any part of a bat; and
 - deliberately disturb a bat species.
- 1.2.3 Disturbance of the species includes in particular any disturbance which is likely to impair their ability to survive, to breed or reproduce, or to rear or nurture their young, or in the case of animals of a hibernating or migratory species, to hibernate or migrate. Under the legislation it is therefore an offence to;
 - damage or destroy a bat breeding site or resting place of a bat; and
 - intentionally or recklessly obstruct access to any structure or place used by a bat for shelter or protection.
- 1.2.4 In addition, certain rarer species of bat including Bechstein's bat *Myotis bechsteinii* and barbastelle *Barbastella barbastellus* are listed on Annex II of the EC Habitats Directive (Council Directive 92/43/EEC).
 - In accordance with Section 41 of the Natural Environment and Rural Communities Act (2006), several bat species are also identified on the England Biodiversity List as Species of Principal Importance. The presence of bats represents a material consideration in the planning process.

Biological records

1.2.5

1.3

1.3.1

1.3.2

- Biological records were obtained from the Gatwick Biodiversity Action Plan Five Year Review 2012-2017 (Gatwick Airport, 2018b).
 - A total of 12 species of bat have been recorded within the Project Area including Bechstein's bats, Brandt's bat *Myotis brandtii*, brown long-eared bat *Plecotus auritus*, common pipistrelle *Pipistrellus*, Daubenton's bat *Myotis daubentonii*, Leisler's bat

Nyctalus leislerii, Nathusius' pipistrelle Pipistrellus nathusii, Natterer's bat Myotis nattereri, noctule Nyctalus noctula, serotine Eptesicus serotinus, soprano pipistrelle Pipistrellus pygmaeus and whiskered bat Myotis mystacinus (Gatwick Airport, 2018b).

- 1.3.3 During a five-year monitoring programme of bat boxes on site undertaken by Surrey Bat Group (2012-2017) the species recorded occupying boxes included Bechstein's bat, Natterer's bat, soprano pipistrelle and brown long-eared bat (Gatwick Airport, 2018b).
- 1.3.4 A whiskered bat maternity roost was recorded at Charlwood Park Farmhouse in 2016 and 2017. This building was also previously occupied by common and soprano pipistrelles (Gatwick Airport, 2018b).
- 1.3.5 A previous trapping and radio-tracking project undertaken in 2014 identified Bechstein's bat roosts in several dead trees in the northern part of Brockley Wood including ash *Fraxinus excelsior*, alder *Alnus glutinosa* and pedunculate oak *Quercus robur* (Gatwick Airport, 2018b).

Previous surveys undertaken in the Project Area

1.3.6

1.3.7

1.3.8

- RPS were commissioned by Gatwick Airport to undertake advanced bat survey techniques in the form of bat trapping and a follow up radio-tracking survey in 2019, to inform the development of potential masterplan scenarios (Annex 4). Surveys were limited to land owned by GAL and within the Project Area and therefore not undertaken in the wider landscape. Trapping, and subsequent radio-tracking of target species from the woodland assemblage of bats (including Bechstein's bat, Brandt's bat, brown long-eared bat, Daubenton's bat, Natterer's bat and whiskered bat), was undertaken across 22 locations within the Project Area over three survey periods between May and September 2019 which were aligned with key stages of the annual bat life cycle.
- A total of 154 bats of nine species were captured over nine nights between 28th May 2019 and 4th September 2019. The species captured included Bechstein's bat, Brandt's bat, brown longeared bat, common pipistrelle, Daubenton's bat, Natterer's bat, noctule, soprano pipistrelle, and whiskered bat.
- Breeding females of seven species were trapped during the survey. These species comprised Brandt's bat, brown long-eared bat, common pipistrelle, Daubenton's bat, Natterer's bat, soprano pipistrelle, and whiskered bat. No female breeding Bechstein's bats were captured within the Project Area, but the presence of



juvenile males and females indicated there was likely to be a colony of breeding females in the wider landscape that was functionally connected to the Project Area.

Radio-tags were fitted to 20 bats of six species of woodland bat including Rechstein's bat Brandt's hat brown long-eared bat

- 1.3.9 Radio-tags were fitted to 20 bats of six species of woodland bat including Bechstein's bat, Brandt's bat, brown long-eared bat, Daubenton's bat, Natterer's bat and whiskered bat. The radio-tracking survey identified roosting locations for all species. As a result, 19 roosting locations were subsequently identified; nine of which were located within the Project Area.
- 1.3.10 Core foraging areas for Bechstein's bats were identified within and adjacent to the Project Area including the Aviation Museum, Charlwood Place Farm, woodland strip to the west of Brockley Wood, River Mole, woodland to the east of Shangri-La and south of Brook Farm, woodland strip to the south-west of the Project Area north of Charlwood Road, Riverside Park, Upper Pickett's Wood, and woodland to the north of sewage works.
- 1.3.11 Bechstein's bats were also recorded foraging in woodland to the east of Bonnett's Lane and Hyder's Farm, River Mole to the south of Charlwood Road, land to the east of Charlwood, Horleyland Wood, Lower Pickett's Wood, woodland strip to the south of Povey Cross Road and River Mole, Man's Brook, Great Burlands woodland, Prestwood Copse, and Gatwick Airport runway.

1.4 Requirement for surveys

- 1.4.1 Bat survey work is required within and adjacent to the Project Area to help inform any future changes to the airport. Surveys undertaken in the area include automated static bat detector surveys and walked activity surveys. However, advanced bat surveys were required because the data on woodland species including Bechstein's bats, which are a cryptic species, cannot be reliably obtained using standard survey techniques (such as activity surveys and/or automated surveys) alone due to overlapping call parameters within the genus *Myotis*. The advanced bat surveys detailed in this report were undertaken under Natural England Project Licence (Dr Stephanie Murphy: 2020-47826-SCI-SCI) and involved trapping, attaching radiotransmitter tags and ringing target species.
- 1.4.2 Surveys were required to develop a more comprehensive understanding of how Bechstein's bats are using the Project Area and surrounding landscape, and how they may be affected by the Project.
- 1.4.3 The surveys were required to provide information on roost locations, status of identified roosts (based on the number of bats

in roost), flightlines and foraging areas for target species (Annex II Bechstein's and barbastelle bats).

1.5 Project purpose

- 1.5.1 The purpose of the surveys was to gather information on the bat assemblage in the area within and adjacent to the Project Area.
- 1.5.2 The surveys encompassed a range of techniques including:
 - trapping bats using harp traps with the assistance of an acoustic lure within land permitted accessible by Gatwick Airport Limited (GAL), Woodland Trust and private landowners, and radio-tagging target species including barbastelle and Bechstein's bat;
 - non-invasive DNA analysis on small Myotis bats (Alcathoe/ /Brandt's bat/whiskered bat) in order to differentiate and confirm the presence of these species in and adjacent to the Project Area;
 - radio-tracking of barbastelle and Bechstein's bats to determine foraging areas, commuting routes and roost locations and counts; and
 - emergence surveys and roost counts undertaken on roosts of all radio-tagged bats (access permitting).
- 1.5.3 It was proposed to radio-tag and radio-track up to a maximum of 30 barbastelle and Bechstein's bats over the survey period.
- 1.5.4 This comprised adult females (both parous and non-parous), males and juveniles (providing they are of an appropriate weight and in good health condition).
- 1.5.5 In order to obtain a robust assessment of the bats' movements and roosts during pre-maternity, post-maternity and autumnal dispersal seasons and across the Project Area it was considered necessary to radio-tag the numbers of bats prescribed above.

2 Methodology

2.1.1 The methods generally followed the standard best practice (Collins, 2016 and Natural England, 2013). Any specific deviations due to objectives of the surveys are described where necessary.

2.2 Trapping surveys

2.2.1 Trapping surveys were undertaken during three key periods which corresponded with key stages of the annual life cycle of a

bat and included surveys between 13th 16th July 2020 (post-maternity), 7th and 10th September 2020 (autumnal dispersal) and 3rd and 6th May 2021 (maternity).

- 2.2.2 Trapping focused more on parts of the Project Area and adjacent wider landscape that may be of particular importance to bats, such as locations of known roosts and areas of high suitability foraging/commuting habitat. Trapping locations are shown in Figure 3.2.1 and Table 2.2.1.
- 2.2.3 Trapping surveys were carried out using a combination of triple and double bank harp traps including woodland habitat, hedgerows and watercourses, which are likely to be habitat features used by commuting and / or foraging bats.
- 2.2.4 At each of the locations, one harp trap (Austbat) was set up.
 Each trap was fitted with an acoustic lure (Model: Sussex
 Austbat) to increase the likelihood of catching bats present within the vicinity of the traps.
- 2.2.5 Trapping commenced at sunset and continued for a maximum of six to eight hours per night dependant on the conditions, weather, capture success and general bat activity. If the weather became unsuitable during a survey (ie temperatures fell below 8°C and/or heavy rain and/or strong wind), the survey was terminated to avoid captured bats becoming torpid. Where unseasonably cold or poor conditions completely curtailed a survey, the survey was rescheduled where possible.

2.2.6

When bats were captured in a harp trap they were removed by a suitably experienced and qualified person (under the direction of the licence holder or accredited agent) and transferred to a clean calico cloth bag. Biometric data was be recorded for each bat including weight, sex, breeding status and forearm measurements. Droppings from small Myotis species were collected and transferred to a sterile tube for possible DNA analysis. Bats not selected for radio-tracking were released at the site of capture immediately after biometric data was recorded.



Table 2.2.1: Trapping Locations

Trapping location ¹	Easting	Northing	Description
1	522591	140332	Glover's Wood south – Beside large oak. Hazel understory
2	522442	140472	Glover's Wood south – Beside large oak on narrow footpath
3	522522	140542	Glover's Wood south – Beside large oak, path overlooking small valley
4	522815	140731	Glover's Wood north – By pond
5	523077	141047	Glover's Wood north – Ancient deciduous woodland
6	523034	140711	Glover's Wood north – Ancient deciduous woodland
7	523544	142326	Edolph's Copse – Under large oak at pond edge
8	523461	142376	Edolph's Copse – Beside footpath next to large oak
9	523579	142520	Edolph's Copse – Woodland interior next to med/large oak
10	523713	142145	Edolph's Copse – Close to open meadow area
11	523754	142294	Edolph's Copse – Woodland interior next to med oak
12	525744	140813	Brockley Wood - Next to med/large oak
13	525776	140837	Brockley Wood - Next to med/large oak
14	525809	140892	Brockley Wood - Next to med/large oak
15	525457	140631	Strip west of Brockley Wood
16	525492	140681	Strip west of Brockley Wood
17	525600	139752	Strip of woodland to the south west of the Project Area, north of Charlwood Road – west

Trapping location ¹	Easting	Northing	Description
18	525862	139810	Strip of woodland to the south west of the Project Area, north of Charlwood Road – central
19	525928	139818	Strip of woodland to the south west of the Project Area, north of Charlwood Road – east
20	521870	138624	Orltons Copse - Gulley with some standing water
21	521898	138831	Orltons Copse - Woodland edge beside large oak
22	522184	138752	Orltons Copse - Beside med/large oak woodland interior
23	522673	138816	Scragg Copse
24	522610	139058	Scragg Copse
25	522716	138986	Scragg Copse
26	522669	138383	Mount Wood - Ancient deciduous woodland; located in uncluttered western section
27	522788	138370	Mount Wood - Ancient deciduous woodland; 6m mist net located in shrub-layer gap at boundary
28	522899	138298	Mount Wood - Ancient deciduous woodland; triple bank located in more cluttered southern section near mature oaks

2.3 Radio-tracking survey

2.3.1 Bechstein's bats were selected for radio-tagging on the basis of their species and apparent health and body condition. No underweight bats were selected for radio-tagging. The weight of the radio-tag was always less than 5% of the bat's weight. Female bats, and in particular reproductive females (avoiding

heavily pregnant bats), were radio-tagged in preference to male bats to enable the location of breeding colonies.

Lotek radio-tags were attached to the bats using Skin-Bond® (Pfizer Inc) to the area between the shoulder blades from which fur had been clipped.

The bats that were fitted with radio-tags were released on the night of capture at the location of capture and held only for the minimum time necessary to obtain the required data and to be fitted with rings and radio-tags as necessary.

The roost locations of radio-tagged bats were located by teams of a minimum of two surveyors using radio-tracking equipment. The radio-tracking equipment comprised of a Biotrack Sika radio-receiver unit coupled with a Biotrack Yagi radio-antenna which allowed the surveyors to tune into the frequency of the radio-tag on each radio-tagged bat. By moving through the landscape on foot or by vehicle, surveyors were able to locate radio-tagged bat roost locations by following the direction and strength of the radio signal emitted by the radio-tag.

The movements of active bats at dusk, night and at dawn, were ascertained by multiple teams of surveyors using the radiotracking equipment described above. Each team was positioned in separate locations within the landscape with their exact location recorded (eastings, northings recorded with GPS unit). Each team was in constant communication with each other via handheld two-way radios or mobile telephones. When bats commenced emergence from their roost location and / or their radio-tag was picked up whilst in flight, the surveyor teams took close approach estimates of location (if the bat was in the immediate vicinity). If bats were located some distance between the surveyors and / or surveyors' access was not permitted to the area in which the bat was situated simultaneous compass bearings on the direction of the strongest radio signal were taken. Through triangulating simultaneous bearings (using LOAS software; version 2.12, Ecological Software Solutions), surveyor teams were able to record 'fixes' (the point at which two or more separate team's bearings cross) of the bats' locations within the landscape, (ie locations at which simultaneous bearings crossed).

¹ Trapping locations were surveyed during each session with the following exclusions; Orltons Copse was not surveyed in September 2020 or May 2021, Scragg Copse was not surveyed in July 2020, Mount Wood was not surveyed in July 2020 or September 2020. Trapping locations

within woodlands varied slightly between sessions due to the condition of the woodland and suitability for trapping. More details are provided in Section 3.5.

Radio-tracking data can include outliers, especially with fast

moving animals such as bats.



2.3.6	The positions of the radio-tagged bats were received at intervals after leaving the roost and were used to identify flightlines, which		simply under-recorded. Bats are highly mobile animals and can move roost sites both within and between years.		Radio-tracking survey (including roost count and emergence surveys)
	are important connectivity features between the roosting locations and foraging areas and between separate foraging areas.	2.5.2	Data from bat surveys should be considered to be valid for a period of 18 months, unless there are any meaningful changes to	2.5.8	The positions of the radio-tagged bats are received at intervals after leaving the roost and used to identify flightlines between the
2.3.7	The cumulative home range size was plotted against the number of successive locations for all bats radio-tracked in order to		the buildings or other habitats within the site.		roost and foraging areas. Therefore, flightlines are partially estimated based on best available data. This is considered as an
	determine the point at which the foraging area reached an asymptote, indicating that sufficient data has been gathered from		Trapping surveys		acceptable limitation as assumptions on direction and use of
	each bat. Estimates of range size for this analysis is based on 100% minimum convex polygons, 95% kernels and 50% kernels,	2.5.3	Where identification to species level was not possible in the field, particularly for cryptic species such as the small <i>Myotis</i> genus,		landscape features can be made based on known bat behavioural ecology.
	obtained using BIOTAS software (version 2.0 Alpha, Ecological Software Solutions).		droppings of trapped bats were collected and sent for DNA analysis. As bats were captured for the minimum amount of time necessary, it was not always possible to collect a sample from	2.5.9	It was not GAL policy to obtain ad-hoc access for roost counts outside the Project Area or in private properties. Therefore, roost
2.3.8	Each bat fitted with a radio-tag was followed for a minimum of		each bat had they not provided a dropping in the calico cloth bag after one hour. Due to animal welfare priorities, bat droppings	0.7.40	counts were not possible in these areas.
	three nights and a maximum of seven nights, depending on the results obtained from the estimates of home range analysis (ie whether sufficient data had been collected). Bats were radio-		collected for DNA analysis was therefore not possible with all captured target species.	2.5.10	All radio-tracking was carried out from land where access was permitted, public roads and public rights of way. Restricted access to land affected the accuracy of the bearings taken during
	tracked concurrently.	2.5.4	The species of three whiskered/Brandt's/Alcathoe bats were not		radio-tracking. If a radio-tagged bat was foraging in an area of land where the strength of the signal was weak (as a
2.4	Roost count surveys		confirmed in July 2020 and one in May 2021 due to lack of droppings.		consequence of topography) and/or had disappeared, then it was not possible to ascertain the precise location of the bat. These
2.4.1	Roosts were identified by locating the radio-tagged bats during the day (as detailed above) and subsequent counts of the	2.5.5	Precise trapping locations varied slightly over the trapping survey		constraints were overcome using the following methods:
	number of bats emerging was carried out, where access was permitted by GAL.		period. The locations varied slightly as the condition of the woodland differed between seasons due to vegetation growth. This was not considered to be a significant limitation to the		 if the bat was likely to be roosting in land where access was prohibited, multiple bearings were taken from a variety of
2.4.2	The surveys were carried out in appropriate weather conditions following standard guidelines (Collins, 2016) and when bats are		findings of the study as the woodland was thoroughly surveyed during each session.		different locations to get a better estimate of likely roosting locations; and
	likely to be active. The dusk surveys commenced approximately	2.5.6	Orltons Copse was not surveyed in September 2020 or May		 where radio-tracking data on bats was deficient, subsequent radio-tracking night(s) concentrated effort on those bats for
	15 minutes before sunset and continued for a minimum of at least 90 minutes; the optimum time for bats to emerge from a roost.	2.0.0	2021. This was due to access restrictions in September 2020. The decision was made to not survey this area in May 2021 due		which data was limited.
2.4.3	An infra-red camera (such as the Canon XA-20) equipped with		to higher quality habitats for target species (Bechstein's bats and	2.5.11	Fast moving traffic on roads within and adjacent to the Project Area meant that surveyors could not survey from the most
	infra-red lamps (such as the IR Lab Outdoor IR Illuminator LIR-IC88) was used to aid accurate counts of bats emerging from the		barbastelles) being available. Scragg Copse was not surveyed in July 2020 or May 2021 due to access restrictions. Mount Wood		appropriate areas due to health and safety concerns.
	roost. Full spectrum bat detectors, such as Elekon Batlogger M,		was not surveyed in July 2020 or September 2020 due to access restrictions. This was not considered to be a significant limitation	2.5.12	Surrey Bat Group were also involved in data collection as
	were used to detect bat echolocation calls of any emerging bats and identify species where possible. Both video and ultrasonic		to the findings of the study as the wider area around the Project		information was shared on radio-transmitter frequencies.
	audio recordings were subsequently analysed to ensure accurate species identification and roost count. Flightlines from roosts		Area was covered thoroughly.	2.5.13	Due to the limited access to areas outside the Project Area, radio-tracking data was difficult to obtain. As mentioned above,
	were determined by roost count survey teams taking bearings on emerging bats.	2.5.7	Bechstein's bats radio-tagged in May 2021 were inactive for part of the radio-tracking study period due to sub-optimal		multiple bearings were taken from a variety of different locations when bats were recorded in inaccessible land. However, when
2.5	Data validity and limitations		temperatures after they were radio-tagged. The Surrey bat group continued to monitor these roosts after the radio-transmitters		bats moved between locations, surveyors moved to locate the bat but due to the access restrictions, the signal was often lost by the
2.5.1	It is important to note that even where data are held, a lack of		were no longer active and once the weather conditions improved and provided the data of emergence counts to the Project.		time they had reached a suitable surveying location.

records for a defined geographical area does not necessarily

mean that there is a lack of ecological interest; the area may be



- 2.5.15 Radio-tracking data on Bats 3J, 4J and 1S was obtained for two nights. Surveyors searched the Project Area for these bats over at least three nights, but the bats were not found, indicating that they were likely foraging outside the Project Area or the tag had failed.
- 2.5.16 Bat 5S was not found during the radio-tracking period. It is considered likely that the radio-tag failed on this bat or the bat was outside the Project Area. Bats are highly mobile species and consequently any bats trapped and tagged within the survey area may not always be identified roosting or in flight in the survey area on subsequent survey days.

3 Results

3.1 Trapping surveys

3.1.1 A total of 98 bats of nine species were captured over nine trapping nights between 13th July 2020 and 7th May 2021 in 28 different locations. Figure 2.2.1 shows the trapping locations, detailed trapping data is shown in Annex 1 and trapping results by species are summarised in Table 3.1.1 below.

Table 3.1.1: Trapping results by species

Species	Number of bats
Barbastelle	1
Bechstein's bat	16
Daubenton's bat	1
Whiskered bat	10
Whiskered bat/Brandt's bat ²	4
Natterer's bat	14
Noctule	1
Brown long-eared bat	45
Common pipistrelle	4

Species	Number of bats
Soprano pipistrelle	2

3.1.2 The average number of bats caught across each trapping area is shown in Table 3.1.2 below.

Table 3.1.2: Trapping results by location

	Number of	Number of	Number of bats			
Location	bats	nights	per night			
Glover's Wood	32	4	8			
Edolph's Copse	29	4	7.25			
Brockley Wood	15	3	5			
Strip west of Brockley Wood	5	3	1.67			
Strip of woodland to the south west of the Project Area, north of Charlwood	0	3	0			
Orltons Copse	10	1	10			
Scraggs Copse	5	1	5			
Mount Wood	2	1	2			

DNA analysis

3.1.3 Droppings were obtained from three of the trapped small *Myotis* bats, which were all sent for DNA analysis. Results were pending when this report was produced and will be provided in the ES.

3.2 Radio-tracking

3.2.1 Fourteen of the sixteen trapped Bechstein's bats were selected for radio-tracking. The sex, breeding status and bat identification numbers are shown in Table 3.2.1 below.

Table 3.2.1: Radio-tagged bats

Bat identification number ³	Trapping location	Species	Sex	Breeding status
1J	1	Bechstein's bat	F	Lactating
2J	6	Bechstein's bat	F	Lactating
3J	7	Bechstein's bat	F	Lactating/post- lactating
4J	11	Bechstein's bat	F	Post-lactating
5J	10	Bechstein's bat	F	Post-lactating
6J	13	Bechstein's bat	М	Adult
7J	12	Bechstein's bat	М	Adult
8J	14	Bechstein's bat	М	Adult
1S	4	Bechstein's bat	М	N/A – Juvenile
2S	4	Bechstein's bat	F	N/A – Juvenile
3S	4	Barbastelle	М	N/A – Juvenile
4S	9	Bechstein's bat	F	N/A – Juvenile
58	14	Bechstein's bat	F	N/A – Juvenile
1M	6	Bechstein's bat	F	Adult – Non- parous
2M	4	Bechstein's bat	F	Adult - Parous

² DNA analysis not possible

³ The letter after the number indicates month of capture; J=July 2020, S=September 2020, M=May2021



Flightlines

3.2.2 No flightlines were recorded from bat roosts to foraging areas as the majority of bats were recorded close to their roosting locations.

Roosts, Home ranges and foraging areas

- 3.2.3 A total of 10 confirmed roosting locations were identified from nine radio-tagged bats. Additionally, nine estimated roosting locations were identified. Dusk emergence surveys were undertaken on seven of the confirmed roosts. The location of these roosts and counts of the roosts are shown in Table 3.1.1 and Figure 3.2.1. The emergence survey section below provides information on roost counts and characterisation.
- 3.2.4 The fixes obtained during the radio-tracking study bats were analysed with BIOTAS software to calculate the maximum home range (100% MCP), the peripheral foraging areas (95% KDE) and the core foraging areas (50% KDE) of each tracked Bechstein's bat. The calculated 100% MCP, 95% KDE and 50% KDE for each tracked bat are shown in Annex 2. Figures 3.2.2-3.2.5 display visual representations of the home ranges and foraging areas for the radio-tracked bats.

Bechstein's bat

- 3.2.5 Figure 3.2.2 shows the Bechstein's bats that were recorded within and adjacent to the Project Area in July 2020. This only included bats tagged in the post-maternity session (Bats 3J, 4J, 6J, 7J and 8J). Two of the bats were post-lactating females (Bat 3J and 4J) and the remainder were adult males.
- 3.2.6 Bat 1J, a lactating female Bechstein's bat, was caught in Glover's Wood and recorded roosting in an inaccessible roost within the same woodland. This bat was radio-tracked for four nights. It was recorded foraging within Glover's Wood only. It was not recorded within the Project Area.
- 3.2.7 Bat 2J, a lactating female Bechstein's bat, was caught in Glover's Wood and recorded roosting in an inaccessible roost on a treeline east of Ifield Road, Woodcote Estate. This bat was radio-tracked for two nights. It was recorded foraging in Glover's Wood. It was not recorded within the Project Area.
- 3.2.8 Bat 3J, a lactating female Bechstein's bat, was caught in Edolph's Copse and was recorded roosting in two separate inaccessible roosts between Glover's Wood and Edolph's Copse. The bat was radio-tracked for five nights. It was recorded foraging within Glover's Wood and Edolph's Copse, and the woodland blocks

between the two, as well as within the Project Area along the River Mole north of Brockley Wood.

3.2.9

3.2.13

- Bat 4J, a post-lactating female Bechstein's bat, was caught in Edolph's Copse. Three roosts were recorded for this bat; two in a tree line east of Edolph's Copse (one of which had 5J also roosting in it) and one in a tree line off Norwood hill Road/Rectory Lane/Stan Hill (peak count of six bats emerging from the roost). This bat was radio-tracked for two nights. It was recorded foraging within Edolph's Copse, the woodland blocks to the north east of Glover's Wood and the River Mole within the Project Area, northwest of Brockley Wood.
- 3.2.10 Bat 5J, a post-lactating female Bechstein's bat, was caught in Edolph's Copse. Two separate roosts were recorded for this bat; one in a tree line east of Edolph's Copse and one in a tree line off Norwood hill Road/Rectory Lane/Stan Hill (peak count of six bats emerging from the roost). Bat 4J was also recorded roosting in both of these trees. This bat was radio-tracked for three nights. It was recorded foraging within Edolph's Copse and the woodland blocks to the north east of Glover's Wood. It was not recorded within the Project Area.
- 3.2.11 Bat 6J, an adult male Bechstein's bat, was caught in Brockley Wood and was recorded roosting in an estimated location within Brockley Wood. This bat was radio-tracked for three nights. It was recorded foraging within the Project Area along Man's Brook to the northwest of Brockley Wood, as well as outside the Project area in Edolph's Copse and the woodland blocks and hedgerows between Edolph's Copse and the northwest of the Project Area.
- 3.2.12 Bat 7J, an adult male Bechstein's bat, was caught in Brockley Wood and was recorded roosting in an estimated location within Brockley Wood as well as a confirmed ash tree roost within Brockley Wood where it was the only bat recorded emerging from the tree during an emergence survey. This bat was radio-tracked for three nights. It was recorded foraging within the Project Area along Man's Brook, River Mole and woodland adjacent to the runway, as well as pockets of woodland between the Project Area and Glover's Wood to the west and north west.
 - Bat 8J, an adult male Bechstein's bat, was caught in Brockley Wood. Bat 8J was recorded roosting in an inaccessible roost within a woodland block to the north east of Glover's Wood. This bat was radio-tracked for three nights. It was recorded foraging within Charlwood, along Man's Brook to the east of Charlwood and within woodland blocks between Glover's Wood and Edolph's Copse.

- 3.2.14 Figure 3.2.3 details the Bechstein's bats that were recorded within and adjacent to the Project area in September 2020. Bat 1S, a juvenile male Bechstein's bat, was caught in Glover's Wood and was recorded roosting within an ash tree on the eastern edge of Glover's Wood. A peak count of eight bats were recorded roosting within the roost. This bat was radio-tracked for two nights and was recorded foraging in Glover's Wood only. It was not recorded within the Project Area.
- 3.2.15 Bat 2S, a juvenile female Bechstein's bat, was caught in Glover's Wood and was recorded roosting within southeast Glover's Wood. Two emergence surveys were undertaken on the tree which recorded counts of 19 and 20 bats. This bat was radiotracked for four nights and was recorded foraging in Glover's Wood, Edolph's Copse and nearby areas. It was not recorded within the Project Area.
- 3.2.16 Bat 4S, a juvenile male Bechstein's bat, was caught in Edolph's Copse and was recorded roosting in an inaccessible location, estimated to be off Partridge Lane, Cudworth. This bat was radiotracked for three nights and was recorded foraging in Glover's Wood, Edolph's Copse, around Beam Brook in Toogoolawah, Cudworth, and pockets of woodland between Edolph's Copse and Toogoolawah. It was not recorded within the Project Area.
- 3.2.17 Bat 5S, a juvenile female Bechstein's bat, was caught in Brockley Wood but was not located by radio-telemetry following capture.
- 3.2.18 Figure 3.2.4 shows Bechstein's bats that were recorded in the Project area in May 2021. Bat 1M, an adult non-breeding female Bechstein's bat, was caught in Glover's Wood. This bat, along with Bat 2M was recorded roosting in two roosts within Glover's Wood; one of which had a peak count of three bats (including both Bat 1M and 2M) and the other had a peak count of 15 bats emerging from the roost. This bat was radio-tracked for five nights. This bat was recorded foraging exclusively in Glover's Wood during surveys. It was not recorded within the Project Area.
 - Bat 2M, an adult breeding female Bechstein's bat, was caught in Glover's Wood. The details of the Bat 2M roosts are shown in the paragraph above. This bat was radio-tracked for five nights. This bat was recorded foraging exclusively in Glover's Wood during surveys. It was not recorded within the Project Area.

Barbastelle bat

3.2.19

3.2.20 Figure 3.2.5 Bat 3S, shows an adult male barbastelle, was caught in Glover's Wood. This bat was radio-tracked for five nights. Core foraging areas for the one barbastelle (Bat 3S) were identified to





the south east of Glover's Wood, around Russ Hill Farm and adjacent to Man's Brook to the west of Prestwood Copse. No core foraging areas were recorded within the Project Area for barbastelles.

3.2.21 The barbastelle bat was also recorded foraging within Faygate Forest, Prestwood Copse and adjacent watercourses, and a block of woodland and Bewbush Brook in Bewbush, west of Crawley. No peripheral foraging areas were recorded within the Project Area for this species.



Table 3.1.1: Roost Location, Count and Characterisation

Roost locations	Emergence surveys		Roost Characterisation					
Bat identification number	Species	Estimated/confirmed roost	Easting	Northing	Description	Emergence date	Roost	Likely roost type
1J	Bechstein's bat	Estimated	523084	140706	Glover's Wood			Maternity ⁴
2J	Bechstein's bat	Estimated	523853	140217	Tree line E of Ifield Road, Woodcote Estate			Maternity ⁵
3J	Bechstein's bat	Estimated	523344	141522	Woodland block north east of Glover's Wood			Maternity ⁶
3J	Bechstein's bat	Estimated	523376	141489	In woodland between Edolph's Copse and Glover's Wood			Maternity ⁷
4J	Bechstein's bat	Confirmed	524085	142066	On tree line east of Edolph's Copse	17/07/2020	0	Day
4J and 5J	Bechstein's bat	Estimated	524031	142105	Hedge/tree line east of Edolph's Copse			Maternity ⁸
4J and 5J	Bechstein's bat	Confirmed	523900	141446	Ash tree in tree line off Norwood Hill Road/Rectory Lane/Stan Hill. Two woodpecker holes 10m in height on southern elevation	15/07/2020	6	Maternity
6J	Bechstein's bat	Estimated	525830	140917	Brockley Wood - unknown location			Unknown
7J	Bechstein's bat	Estimated	525766	140712	Brockley Wood - unknown location			Unknown
7J	Bechstein's bat	Confirmed	525771	140717	Ash tree within Brockley Wood	17/07/2020	1	Day
8J	Bechstein's bat	Estimated	523470	141513	Woodland block north east of Glover's Wood			
1S	Bechstein's bat	Confirmed	523171	140839	Eastern edge of Glover's Wood in branch of Ash tree 4-5m on SE elevation	08/09/2020	8	Maternity
2S	Bechstein's bat	Confirmed	523197	140875	SE Glover's Wood within stem at base of second major fork	10/09/2020 11/09/2020	19 20	Maternity
3S	Barbastelle	Confirmed	522088	133294	Faygate Forest - downy birch, lifted bark			Unknown

⁴ Although a roost count was not undertaken, it is considered likely that this roost was a maternity roost due to the breeding status of the tagged bat within the roost ⁵ Although a roost count was not undertaken, it is considered likely that this roost was a maternity roost due to the breeding status of the tagged bat within the roost ⁶ Although a roost count was not undertaken, it is considered likely that this roost was a maternity roost due to the breeding status of the tagged bat within the roost ⁷ Although a roost count was not undertaken, it is considered likely that this roost was a maternity roost due to the breeding status of the tagged bat within the roost

Although a roost count was not undertaken, it is considered likely that this roost was a maternity roost due to the breeding status of the tagged bat within the roost 8 Although a roost count was not undertaken, it is considered likely that this roost was a maternity roost due to the breeding status of the tagged bat within the roost



Roost locations	Roost locations							
Bat identification number	Species	Estimated/confirmed roost	Easting	Northing	Description	Emergence date	Roost	Likely roost type
3S	Barbastelle	Confirmed	523190	139589	NE section of Prestwood Copse in dead oak under lifted bark 3-4m on eastern elevation			Unknown
48	Bechstein's bat	Estimated	521655	142192	Off Partridge Lane, Cudworth			Unknown
48	Bechstein's bat	Confirmed	521592	142301	Tree along Beam Brook E of Partridge Lane			Unknown
1M and 2M	Bechstein's bat	Confirmed	522441	140507	Glover's Wood - split feature in oak main stem at 10m height. Split has been enlarged by woodpecker. Northern aspect	07/05/2021	3	Satellite maternity
1M and 2M	Bechstein's bat	Confirmed	522768	149621	Glover's Wood - oak tree 4x multistem. Woodpecker hole at 12m south west aspect	09/05/2021 14/05/2021 ⁹	14 15	Maternity

⁹ Emergence survey undertaken by Martyn Cooke of Surrey Bat Group



	Emergence surveys		Due to the breeding status of the bats this roost was considered to be a satellite maternity roost for Bechstein's bats.		this species' population and range is unknown its habitat status is considered to be in decline (Matthews <i>et al.</i> , 2018).
3.2.22	A total of nine emergence surveys were undertaken on seven of the confirmed roosts for barbastelle and Bechstein's bats.	3.2.30	On 9 th May 2021 an emergence survey was undertaken on a separate roost for Bat 1M and 2M in Glover's Wood. A total of 14	4.1.5	No barbastelle bats were captured within the Project Area during surveys in 2019.
3.2.23	One emergence survey was undertaken within the Project Area for Bat 7J that was roosting within an ash tree in Brockley Wood. The tagged Bat 7J was the only bat to be recorded emerging from the tree. Due to the number of bats recorded emerging from this roost, this was considered to be a day roost for Bochstein's		bats were recorded emerging from the roost, including the two tagged bats. A repeat emergence survey of this tree was undertaken on 10 th May 2021 which recorded 15 bats emerging from the roost. Due to the number of bats recorded emerging	4.1.6	The sections below summarise the results obtained for the trapping and radio-tracking surveys undertaken in 2020 and 2021.
	this roost, this was considered to be a day roost for Bechstein's bats.		from the roost this was considered to be a maternity roost for Bechstein's bats.	4.2	Field survey
3.2.24	The other seven roosts were located outside the Project Area, within Glover's Wood, Faygate Forest, in a treeline off Norwood Hill Road/Rectory Lane/Stan Hill and in a treeline east of Edolph's	4	Discussion	4.2.1	Trapping A total of 98 bats from a minimum of nine species were captured over nine nights between 13 th July 2020 and 7 th May 2021.
	Copse.	4.1	Species		Species included barbastelle, Bechstein's bat, brown long-eared
3.2.25	On 15 th July 2020 an emergence survey was undertaken on the bat 4J and 5J roost in an ash tree in the tree line off Norwood Hill		Bechstein's bat		bat, common pipistrelle, Daubenton's bat, Natterer's bat, noctule, soprano pipistrelle, whiskered/Brandt's bat and whiskered bat.
	Road/Rectory Lane/Stan Hill. Six bats were recorded emerging from the roost including the two tagged bats. Due to the number of bats recorded emerging from this roost, and the breeding status of the tagged bats, this was considered to be a small maternity roost for Bechstein's bats.	4.1.1	This species is listed in Annex II of the EU Habitat Directive, categorised as Near Threatened on the IUCN Red List of Threatened Species (Hutson and Paunovic, 2016), is a Species of Principal Importance in England, and is considered to be very rare at a national (Bat Conservation Trust, 2010) and local level (Sussex Bat Group, 2019 and Surrey Bat Group, 2019). Although	4.2.2	The most commonly caught bat species was brown long-eared bat with a total of 45 bats captured, followed by Bechstein's bats (16 bats). The least commonly caught bats were barbastelle, noctule and Daubenton's bats with just one of each species caught during the survey period.
3.2.26	On 17 th July 2020 an emergence survey was undertaken on the Bat 4J roost in the tree line to the east of Edolph's Copse. No bats were recorded emerging from the roost. As no bats were recorded emerging from this roost, this tree was considered to be		this species' range is considered to be stable, the future prospects for this species' habitat are considered to be in decline (Matthews <i>et al.</i> , 2018).	4.2.3	Trapping locations were split into eight distinct areas across the Project Area; Glover's Wood, Edolph's Copse, Brockley Wood, strip west of Brockley Wood, strip of woodland to the south west
3.2.27	a day roost for Bechstein's bats. On 8 th September 2020 an emergence survey was undertaken on the Bat 1S roost located in an ash tree on the eastern edge of Glover's Wood. Eight bats were recorded emerging from the roost. Due to the number of bats recorded emerging from the	4.1.2	Trapping surveys in 2019 captured a total of seven Bechstein's bats within the Project Area within the woodland strip west of Brockley Wood/River Mole, Brockley Wood, the strip of woodland to the south west of the Project area north of Charlwood Road, Riverside Park and Horleyland Wood.		of the Project Area, Orltons Copse, Scraggs Copse and Mount Wood. The highest number of average bats caught per night was 10 and this was recorded within Orltons Copse, however none of the target species were captured at this location. The location with the second highest number of bats caught per night was Glover's Wood, with an average of eight bats per night.
	roost this was considered to be a maternity roost for Bechstein's bats.	4.1.3	No breeding individuals were recorded for this species during surveys within the Project Area in 2019; captured bats were all		Radio-tracking
3.2.28	On 10 th and 11 th September 2020 an emergence survey was		male or non-breeding females. However, juvenile male and		Roosts
0.2.20	undertaken on the Bat 2S roost in south east Glover's Wood. A total of 19 and 20 bats were recorded emerging respectively. Eight bats were recorded emerging from the roost. Due to the		females were recorded present in September indicating that there is likely to be a breeding colony close to the Project Area. Barbastelle bat	4.2.4	Seventeen Bechstein's bat roosts and two barbastelle roosts were identified during the radio-tracking surveys. Emergence surveys were undertaken on seven of these roosts. Of these
	number of bats recorded emerging from the roost this was considered to be a maternity roost for Bechstein's bats.		This species is listed in Annex II of the EU Habitat Directive, categorised as Near Threatened on the IUCN Red List of		roosts, three were located within the Project Area. An emergence survey was undertaken on one of these roosts which identified
3.2.29	On 7 th May 2021 an emergence survey was undertaken on the Bat 1M and 2M roost in Glover's Wood. A total of three bats were recorded emerging from the roost, including the two tagged bats.		Threatened Species (Hutson and Paunovic, 2016), is a Species of Principal Importance in England, and is considered to be very rare at a national (Bat Conservation Trust, 2010) and local level (Sussex Bat Group, 2019 and Surrey Bat Group, 2019). Although		one bat (Bat 7J) emerging from the roost. As the bats recorded roosting within the Project Area were all adult males it is considered unlikely that the roosts are roosts of high conservation significance for Bechstein's bats eg maternity roosts are likely to be present in the Project Area.



4.2.5 Of the remaining 14 Bechstein's bat roosts located outside the Project Area, four were confirmed as maternity roosts from roost counts, five were considered likely to be maternity roosts due to the breeding status of the females within the roost (lactating/post-lactating), one as a satellite maternity roost, one as a day roost, and three unknown roost types. Bechstein's bats are typically associated with fission fusion roost dynamics where social groups change over time, sometimes growing to breed or splitting up to forage, for example (Kerth et al, 2006, Kerth et al 2014) It is likely that these bats use a number of trees in close proximity to one another throughout the pre & post breeding period.

Foraging areas

- 4.2.6 Core foraging areas for radio-tracked Bechstein's bats within the Project Area were identified within the following areas:
 - Brockley Wood;
 - Man's Brook: and
 - River Mole.
- 4.2.7 Although two post-lactating female Bechstein's bats were recorded within the Project Area (Bats 3J and 4J), core foraging areas for these bats were not located within the Project Area. No core foraging areas were recorded within the Project Area for barbastelles.
- 4.2.8 Core foraging areas for radio-tracked Bechstein's bats outside the Project Area were identified within the following areas:
 - Glover's Wood;
 - Mountnoddy Wood;
 - Edolph's Copse;
 - woodlands around Toogoolawah and Beam Brook; and
 - woodlands and waterbodies adjacent to Biltam Farm Nurseries.
- 4.2.9 Core foraging areas for the barbastelle were identified in the following areas:
 - south east of Glover's Wood;
 - around Russ Hill Farm; and
 - adjacent to Man's Brook to the west of Prestwood Copse.
- 4.2.10 Bechstein's bats were also recorded foraging around the following areas (peripheral foraging areas):
 - Beggars Gill;
 - Pagewood;
 - Charlwood;

- woodland blocks and hedgerows north east of Charlwood; and
- Man's Brook and adjacent arable fields to the west of the Project Area.
- 4.2.11 Barbastelle bats were recorded foraging around the following areas (peripheral foraging areas):
 - within Faygate Forest;
 - Prestwood Copse and adjacent watercourses; and
 - block of woodland and Bewbush Brook in Bewbush, west of Crawley.
- 4.2.12 The bats recorded foraging within the Project Area included two post-lactating females and three adult male Bechstein's bats. No core foraging areas for breeding females were identified within the Project Area, with data for these two bats limited to a relatively low number of fixes along the River Mole and Man's Brook, with the majority of their foraging (core foraging areas) recorded outside the Project Area in Edolph's Copse and woodland block to the north east of Glover's Wood.

Assessment of importance

- 4.2.13 This section aims to assess the importance of the Bechstein's bat and barbastelle populations within and adjacent to the Project Area. The importance of the areas is assessed at a geographical level eq local, regional, national.
- 4.2.14 Paragraphs 4.1.1 and 4.1.4 above detail the protection afforded to Bechstein's and Barbastelle bats, rarity of the species within the UK, Sussex, and Surrey, and future prospects of the species.
- 4.2.15 The value of roosts and foraging areas for both Bechstein's bats and barbastelles have been calculated using Wray *et al.* (2010) for both within the Project Area and within the wider landscape, as shown in Table 4.2.1-Table 4.2.4 below.

Table 4.2.1: Importance of roosts within Project Area

Species	Rarity	Roost type present (highest conservation significance listed)	Geographic frame of reference
Barbastelle	Rarest	None	N/A

Species	Rarity	Roost type present (highest conservation significance listed)	Geographic frame of reference
Bechstein's bat	Rarest	Day roost	County

Table 4.2.2: Importance of roosts outside Project Area

Species	Rarity	Roost type present (highest conservation significance listed)	Geographic frame of reference
Barbastelle	Rarest	Likely day roost	County
Bechstein's bat	Rarest	Maternity	National/UK

- 4.2.16 As barbastelle have not been recorded within the Project Area, this area is not considered to be of importance to this species. However, the areas in the wider landscape are considered to be of County/Regional importance for this species.
- 4.2.17 The population of Bechstein's bats is considered to be of County/Regional importance within the Project Area. However, the importance of the areas in the wider landscape are considered to be of Regional/National Importance.



Table 4.2.3: Importance of foraging areas within Project Area

Species	Species rarity	Number of bats	Roosts/potential roosts nearby	Type/Complexity of linear features	Total Score
Barbastelle	Rarest (20)	No bats	None	N/A	0 – Not important
Bechstein's	Rarest (20)	Individual bats (5)	Small number of	Isolated woodland patches, less intensive	31 – Regional importance
bat	(20)	ilidividuai bats (5)	roosts (3)	arable and/or small towns and villages (3)	31 – Negional importance

Table 4.2.4: Importance of foraging areas outside Project Area

Species	Species rarity	Number of bats	Roosts/potential roosts nearby	Type/Complexity of linear features	Total Score
Barbastelle	Rarest (20)	Individual bats (5)	Small number of roosts (3)	Larger or connected woodland blocks, mixed agriculture, and small villages/hamlets (4)	32 – Regional importance
Bechstein's bat	Rarest (20)	Small number of bats (10)	Small number of roosts (3)	Larger or connected woodland blocks, mixed agriculture, and small villages/hamlets (4)	37 – Regional importance



5 Conclusions

- 5.1.1 The highest rates of bat captures were recorded outside the Project Area in Orltons Copse, Glover's Wood and Edolph's Copse.
- 5.1.2 Three roosts were identified within the Project Area, all for male Bechstein's bats. The roosts were considered to be day roosts, or of unknown status. Fourteen roosts for Bechstein's bats were recorded outside the Project Area. Four of these roosts were confirmed as maternity roosts from roost counts, five were considered likely to be maternity roosts due to the breeding status of the females within the roost (lactating/post-lactating), one as a satellite maternity roost, one as a day roost, and three unknown roost types. Two barbastelle roosts of unknown type were also recorded outside the Project Area.
- 5.1.3 Glovers Wood and Edolphs Copse were used as core foraging areas for multiple Bechstein's bats with the majority of activity recorded in these areas being from breeding female Bechstein's bats.
- 5.1.4 Core foraging areas identified for Bechstein's bats within the Project Area included Brockley Wood, the River Mole and Man's Brook. These core foraging areas were for male Bechstein's bats.
- 5.1.5 The Project Area is considered to not to be of importance for barbastelle bats due to the lack of records but is considered to be of County/Regional importance for Bechstein's bats.
- 5.1.6 The area outside the Project is considered to be of County/Regional importance for barbastelle bats and Regional/National importance for Bechstein's bats.

6 References

Bat Conservation Trust (2010) *Species Factsheet* https://www.bats.org.uk/about-bats/what-are-bats/uk-bats last accessed 07 November 2019.

British Standards Institution (2015) *BS8956 Surveying for Bats in Trees and Woodland*. BSI London.

CIEEM (2018) Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, Freshwater and Coastal, 2nd edition. Chartered Institute of Ecology and Environmental Management, Winchester. Collins (2016) Bat Surveys for Professional Ecologists Good Practice Guidelines 3rd Edition. Bat Conservation Trust, London.

Fleischmann D, and Kerth G (2014) Roosting behavior and group decision making in 2 syntopic bat species with fission–fusion societies. Behavioral Ecology, Volume 25, Issue 5, September-October 2014, Pages 1240–1247,

Gatwick Airport (2018a) Gatwick Airport Draft Master Plan 2018.

Gatwick Airport (2018b) *Gatwick Biodiversity Action Plan Five* Year Review 2012-2017. Gatwick Airport Unpublished Report.

Harris & Yalden (2008). *Mammals of the British Isles: Handbook,* 4th Edition. The Mammal Society, Southampton.

Hutson, A.M. and Paunovic, M. (2016) *Myotis alcathoe*. The IUCN Red List of Threatened Species 2016 https://www.iucnredlist.org/species/136680/518740 last accessed 07 November 2019

Institute of Lighting Professionals (2018) Guidance Note 08/18: Bats and artificial lighting in the UK.

https://www.theilp.org.uk/documents/guidance-note-8-bats-and-artificial-lighting/ last accessed 27 November 2019

Kerth G., Ebert C. and Schmidtke C (2006) *Group decision making in fission–fusion societies: evidence from two-field experiments in Bechstein's bats.* Proceedings of the Royal Society Volume 273 Issue 1602.

Matthews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C.A., McDonald, R.A., Shore, R.F. (2018) A Review of the Population and Conservation Status of British Mammals. A report by the Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage. Natural England, Peterborough. ISBN 978-1-78354-494-3.

Ministry of Housing, Communities and Local Government (2021) National Planning Policy Framework (NPPF). [Online] Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1005759/NPPF_July_2021.pdf

Mitchell-Jones, A.J. & McLeish, A.P. (2004) *The Bat Workers' Manual 3rd Edition*. Joint Nature Conservation Committee, Peterborough.

Natural England (2013) Guidance on the capture and marking of bats under the authority of a Natural England Licence (WML-G39(05/13)).

Surrey Bat Group (2019) Bats In Surrey. Unpublished.

Sussex Bat Group (2019) *Bats In Sussex*http://www.sussexbatgroup.org.uk/batsinsussex last accessed 07
November 2019

Wray, S., Wells, D., Long, E., Mitchell-Jones, T. (2010) Valuing Bats in Ecological Impact Assessment. In Practice, December 2010 pp 23-25.

7 Glossary

7.1 Glossary of Terms

Table 7.1.1: Glossary of Terms

Term	Description
EIA	Environmental Impact Assessment
ES	Environmental Statement
EU	European Union
GAL	Gatwick Airport Limited
IUCN	International Union for Conservation of Nature
PEIR	Preliminary Environmental Information Report



Annex 1

Trapping data

Bat No.	Date	Time (24hr)	Trap No.	Easting	Northing	Species	Extracted by	Measured by	Determined by	Sex - M/F	Age - Juv/ Imm/ Ad	Breeding status (male) - Testes size 0-2	Breeding status (female) - Par/ NP/ Preg/ Lact	Forearm (mm)	Total weight (g)	Bag weight(g)	Bat weight (g)	Tagged bat? Include bat number	Sample taken?	ID features, damage, parasites photos and samples collected
1	13/07/2020	22:16	1	522591	140332	M.bec	DH	DH	DH	F	Ad	N/A	Lact	39.7	25.9	17.1	8.7	1J	N	
2	13/07/2020	22:58	3	522522	140542	M.mys	DH	DH	DH	М	Ad	0	N/A	33.9	22.7	17.1	5.6	N/A	N	
3	13/07/2020	23:47	1	522591	140332	M.mys	DH	DH	DH	М	Ad	1	N/A	35.1	22.7	17.8	4.9	N/A	N	
4	13/07/2020	23:58	3	522522	140542	P.pip	DH	DH	DH	M	Ad	2	N/A	33	22.2	17.2	5	N/A	N	
5	13/07/2020	22:30	6	523034	140711	M.bec	KH	KH	RM	F	Ad	N/A	Lactating	39.3	44.5	35.5	9	2J	N	
6	13/07/2020	22:30	6	523034	140711	M.mys	KH	RM	RM	М	Juv	0	N/a	36	21.5	15.5	6	N/A	N	
7	13/07/2020	23:45	6	523034	140711	P.aur	KH	KH	KH	F	Ad	N/A	Lactating	35.8	46	38	8	N/A	N	
8	13/07/2020	00:00	5	523077	141047	M.mys/bra	KH	KH	KH	F	Ad	N/A	NP	31.4	24.5	19.5	5	N/A	Υ	Straight tragus
9	14/07/2020	22:12	9	523579	142520	M.nat	DH	DH	DH	М	Ad	0	N/A	38.5	26.1	19.9	6.2	N/A	N	
10	14/07/2020	22:22	7	523544	142326	P.aur	DH	DH	DH	М	Ad	1	N/A	39	24.3	16.8	7.5	N/A	N	
11	14/07/2020	22:22	7	523544	142326	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	36.9	22.8	14.8	8	N/A	N	
12	14/07/2020	22:22	7	523544	142326	M.bec	DH	DH	DH	F	Ad	N/A	Lac/post	42.5	25.7	16.8	8.9	3J	N	
13	14/07/2020	23:09	8	523461	142376	P.aur	DH	DH	DH	M	Ad	0	N/A	37.1	26.4	19	7.4	N/A	N	
14	14/07/2020	23:12	7	523544	142326	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	39	25.7	17.5	8.2	N/A	N	
15	14/07/2020	23:47	9	523579	142520	P.aur	DH	DH	DH	F	Ad	N/A	NP	39.6	26.5	17.5	9	N/A	N	
16	14/07/2020	00:30	7	523544	142326	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	38	28.2	18.9	9.3	N/A	N	
17	14/07/2020	01:27	9	523579	142520	P.aur	DH	DH	DH	F	Ad	N/A	Post lact	39.9	24	16.3	7.7	N/A	N	
18	14/07/2020	01:31	8	523461	142376	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	39.6	25.7	17.7	8	N/A	N	
19	14/07/2020	01:31	8	523461	142376	P.pip	DH	DH	DH	F	Ad	N/A	NP	31.5	19.8	14.8	5	N/A	N	
20	14/07/2020	02:15	8	523461	142376	M.mys	DH	DH	DH	M	Ad	0	N/A	34.2	22.8	17.8	5	N/A	N	
21	14/07/2020	21:45	11	523754	142294	P.aur	SR	SR	RM	M	Ad	1	N/A	39	27	19.5	8.5	N/A	N	
22	14/07/2020	22:30	10	523713	142145	P.aur	SR	SR	SR	F	Ad	N/A	Post lact	38	26.5	18	8.5	N/A	N	
23	14/07/2020	22:30	10	523713	142145	M.bec	SR	SR	SR	M	Juv	0	N/A	41	27	18.5	8.5	N/A	N	
24	14/07/2020	22:40	11	523754	142294	M.bec	SR	SR	SR	F	Ad	N/A	Post lact	42	25.5	16	9.5	4J	N	
25	14/07/2020	23:55	11	523754	142294	P.aur	RM	RM	RM	F	Ad	N/A	NP	39	24	16.5	7.5	N/A	N	
26	14/07/2020	23:55	11	523754	142294	P.aur	RM	RM	RM	F	Ad	N/A	Post lact	39	26	19	7	N/A	N	
27	14/07/2020	00:10	10	523713	142145	M.mys	RM	RM	RM	M	Ad	1	N/A	38	20.5	15.5	5	N/A	N	
28	14/07/2020	00:10	10	523713	142145	M.bec	RM	RM	RM	F	Ad	N/A	Post lact	41	44	34.5	9.5	5J	N	
29	14/07/2020	00:10	10	523713	142145	M.mys/bra	RM	RM	RM	F	Juv	N/A	NP	40	50	44	6	N/A	N	No droppings left



Bat No.	Date	Time (24hr)	Trap No.	Easting	Northing	Species	Extracted by	Measured by	Determined by	Sex - M/F	Age - Juv/ Imm/ Ad	Breeding status (male) - Testes size 0-2	Breeding status (female) - Par/ NP/ Preg/ Lact	Forearm (mm)	Total weight (g)	Bag weight(g)	Bat weight (g)	Tagged bat? Include bat number	Sample taken?	ID features, damage, parasites photos and samples collected
30	14/07/2020	01:10	10	523713	142145	M.bec	SR	RM	RM	M	Juv	0	N/A	41	46	38.5	7.5	N/A	N	
31	14/07/2020	01:10	10	523713	142145	P.aur	SR	RM	RM	М	Juv	0	N/A	37	43.5	36.5	8	N/A	N	
32	15/07/2020	22:24	13	525776	140837	M.bec	DH	DH	DH	М	Ad	0	N/A	40	25	17.6	7.4	6J	N	
33	15/07/2020	23:45	12	525744	140813	M.bec	DH	DH	DH	М	Ad	0	N/A	42.1	25.8	16.8	9	7J	N	
34	15/07/2020	23:54	13	525776	140837	M.nat	DH	DH	DH	F	Juv	N/A	NP	39.5	21.6	14.7	6.9	N/A	N	
35	15/07/2020	01:20	14	525809	140892	M.bec	DH	DH	DH	М	Ad	0	N/A	41.5	25.4	16.8	8.6	8J	N	
36	15/07/2020	01:54	13	525776	140837	M.nat	DH	DH	DH	F	Juv	N/A	NP	N/A	N/A	N/A	N/A	N/A	N	Released at trap
37	15/07/2020	02:15	13	525776	140837	P.aur	DH	DH	DH	М	Ad	0	N/A	34.9	24.9	17.8	7.1	N/A	N	
38	15/07/2020	22:45	16	525492	140681	M.mys	TE	TE	TE	F	Ad	N/A	NP	34.6			5.48	N/A	Y	Likely young of last year. 2x ticks on face. Pale skin
39	15/07/2020	01:35	15	525457	140631	M.mys	TE	TE	TE	М	Juv	0	N/A	34.6			5.31	N/A	Υ	Wing mite E = black
40	15/07/2020	01:40	16	525492	140681	M.mys/bra	TE	TE	TE	F	Ad	N/A	Lact	35.4			5.9	N/A	N	Likely whiskered
41	16/07/2020	23:34	20	521870	138624	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	40.7	25.4	16.6	8.8	N/A	N	
42	16/07/2020	23:43	21	521898	138831	M.nat	DH	DH	DH	F	Ad	N/A	NP	37.6	26.6	18.9	7.7	N/A	N	
43	16/07/2020	00:24	20	521870	138624	N.noc	DH	DH	DH	М	Ad	2	N/A	55.6	49.5	17.1	32.4	N/A	N	
44	16/07/2020	00:36	22	522184	138752	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	39.5	29.1	20	9.1	N/A	N	
45	16/07/2020	00:36	22	522184	138752	P.aur	DH	DH	DH	М	Ad	0	N/A	39.2	26.6	18.9	7.7	N/A	N	
46	16/07/2020	00:36	22	522184	138752	P.aur	DH	DH	DH	F	Ad	N/A	Lac/post	38.7	24.2	16.3	7.9	N/A	N	
47	16/07/2020	00:36	22	522184	138752	P.aur	DH	DH	DH	F	Ad	N/A	NP	40.3	23.3	14.9	8.4	N/A	N	
48	16/07/2020	02:10	21	521898	138831	P.pip	DH	DH	DH	F	Ad	N/A	Lac/post	32.4	23.2	17.8	5.4	N/A	N	
49	16/07/2020	02:10	21	521898	138831	P.pyg	DH	DH	DH	F	Ad	N/A	Lac/post	31.2	22.1	16.8	5.3	N/A	N	
50	16/07/2020	02:25	22	522184	138752	P.aur	DH	DH	DH	М	Ad	1	N/A	36.6	24.3	16.9	7.4	N/A	N	
51	07/09/2020	20:51	2	522442	140472	P.aur	DH	DH	DH	F	Ad	N/A	Post lact	39.5	24.1	16.4	7.7	N/A	N	
52	07/09/2020	22:21	2	522442	140472	P.aur	DH	DH	DH	F	Juv	N/A	NP	38	25.7	17.7	8	N/A	N	
53	07/09/2020	23:01	2	522442	140472	M.nat	DH	DH	DH	F	Ad	N/A	NP	40.9	24.3	17.3	7	N/A	N	
54	07/09/2020	00:14	2	522442	140472	P.aur	DH	DH	DH	F	Ad	N/A	NP	39.7	24.7	16.4	8.3	N/A	N	
55	07/09/2020	00:15	2	522442	140472	P.aur	DH	DH	DH	F	Juv	N/A	NP	40.2	26.2	16.8	9.4	N/A	N	
56	07/09/2020	00:51	2	522442	140472	P.aur	DH	DH	DH	F	Juv	N/A	NP	39.3	24.6	16.4	8.2	N/A	N	
57	07/09/2020	21:00	4	523053	140706	P.aur	KH	KH	KH	F	Ad	N/A	NP	37	28	19.5	8.5	N/A	N	
58	07/09/2020	21:00	4	523053	140706	M.bec	KH	KH	KH	М	Juv	0	N/A	37.1	24	15.5	9.5	1S	N	
59	07/09/2020	21:00	4	523053	140706	M.bec	KH	KH	KH	F	Juv	N/A	NP	38.2	25	19	6	2S	N	
60	07/09/2020	22:30	4	523053	140706	M.nat	KH	KH	KH	М	Juv	1	N/A	35.8	24	16.5	7.5	N/A	N	
61	07/09/2020	22:30	4	523053	140706	P.aur	RM	RM	RM	М	Juv	1	N/A	34.7	22.5	15.5	7	N/A	N	
62	07/09/2020	22:30	4	523053	140706	B.bar	KH	KH	RM	М	Juv	1	N/A	35.8	25.5	18.5	7	3S	N	Parasites
63	07/09/2020	22:45	6	523081	141049	P.aur	RM	RM	RM	М	Juv	1	N/A	35	42.5	36	6.5	N/A	N	
64	07/09/2020	22:45	6	523081	141049	P.aur	RM	RM	RM	F	Juv	N/A	NP	33.5	25.5	18.5	7	N/A	N	Parasites



Bat No.	Date	Time (24hr)	Trap No.	Easting	Northing	Species	Extracted by	Measured by	Determined by	Sex - M/F	Age - Juv/ Imm/ Ad	Breeding status (male) - Testes size 0-2	Breeding status (female) - Par/ NP/ Preg/ Lact	Forearm (mm)	Total weight (g)	Bag weight(g)	Bat weight (g)	Tagged bat? Include bat number	Sample taken?	ID features, damage, parasites photos and samples collected
65	07/09/2020	22:45	6	523081	141049	P.aur	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	Ecaped
66	07/09/2020	01:15	4	523053	140706	P.aur	KH	KH	KH	М	Juv	1	N/A	35	38	30.5	7.5	N/A	N	
67	07/09/2020	01:15	4	523053	140706	M.nat	KH	KH	KH	F	Ad	N/A	NP	36.5	26.5	18.5	8	N/A	N	Parasites
68	08/09/2020	21:32	7	523544	142326	P.aur	DH	DH	DH	M	Ad	0	N/A	37.4	23.1	14.7	8.4	N/A	N	
69	08/09/2020	21:32	7	523544	142326	P.aur	DH	DH	DH	F	Ad	N/A	NP	38.1	22.1	14.8	7.3	N/A	N	
70	08/09/2020	22:18	7	523544	142326	P.aur	DH	DH	DH	F	Juv	N/A	NP	36.3	22.1	14.6	7.5	N/A	N	
71	08/09/2020	00:00	7	523544	142326	P.aur	DH	DH	DH	М	Juv	N/A	N/A	36.1	21.7	14.8	6.9	N/A	N	
72	08/09/2020	00:37	9	523579	142520	M.bec	DH	DH	DH	F	Juv	N/A	NP	41.7	25.7	17.2	8.5	4S	N	
73	08/09/2020	22:55	10	523544	142326	P.pyg	KH	KH	KH	F	Juv	N/A	NP	28.5	24	19.5	4.5	N/A	N	
74	09/09/2020	20:57	13	525776	140837	M.nat	DH	DH	DH	F	Juv	N/A	NP	40.1	24	17.2	6.8	N/A	N	
75	09/09/2020	21:41	13	525776	140837	P.pip	DH	DH	DH	М	Juv	0	N/A	30.2	23.7	19.8	3.9	N/A	N	
76	09/09/2020	22:15	12	525744	140813	M.nat	DH	DH	DH	F	Ad	N/A	Post lact	37.6	23.7	16.9	6.8	N/A	N	
77	09/09/2020	22:15	12	525744	140813	M.mys	DH	DH	DH	M	Ad	1	N/A	34.5	25	19.8	5.2	N/A	N	
78	09/09/2020	22:52	12	525744	140813	M.nat	DH	DH	DH	F	Juv	N/A	NP	39.9	24.7	17.5	7.2	N/A	N	
79	09/09/2020	23:37	14	525809	140892	M.bec	DH	DH	DH	F	Juv	N/A	NP	40.3	25.7	17.6	8.1	5S	N	
80	09/09/2020	00:19	14	525809	140892	M.mys	DH	DH	DH	М	Juv	0	N/A	31.4	20.6	16.3	4.3	N/A	N	
81	09/09/2020	00:49	14	525809	140892	M.nat	DH	DH	DH	M	Ad	0	N/A	37.2	21.7	14.7	7	N/A	N	
82	09/09/2020	01:20	14	525744	140813	P.aur	DH	DH	DH	F	Ad	N/A	NP	40	25.3	17.5	7.8	N/A	N	
83	09/09/2020	23:20	16	525492	140681	M.mys	TE	TE	TE	F	Ad	N/A	Par	34.1	N/A	N/A	5.6	N/A	N	Small hole in right wing. No cusp on P4
84	09/09/2020	23:50	15	525457	140631	P.aur	TE	TE	TE	F	Ad	N/A	Par	40.3	N/A	N/A	9	N/A	N	No droppings
85	10/09/2020	22:05	24	522610	139058	P.aur	DH	DH	DH	М	Juv	N/A	N/A	36.1	27.6	19.8	7.8	N/A	N	
86	10/09/2020	22:05	24	522610	139058	P.aur	DH	DH	DH	F	Ad	N/A	Post lact	39.8	22.4	17.6	4.8	N/A	N	
87	10/09/2020	22:10	25	522716	138986	P.aur	DH	DH	DH	М	Ad	1	N/A	37.6	27.8	19.4	8.4	N/A	N	
88	10/09/2020	22:59	23	522673	138816	M.nat	DH	DH	DH	F	Ad	N/A	NP	38.2	26.4	19.4	7	N/A	N	
89	10/09/2020	01:28	25	522716	138986	P.aur	DH	DH	DH	F	Ad	N/A	NP	38.2	25.7	17.7	8	N/A	N	
90	04/05/2021	21:23	6	522543	140426	M.bec	DH	DH	DH	F	Ad	N/A	NP	42.9	25.1	17.2	7.9	1M	N	
91	04/05/2021	21:27	4	522458	140462	P.aur	DH	DH	DH	F	Ad	N/A	NP	39.5	27.8	21.1	6.7	N/A	N	
92	04/05/2021	22:17	6	522543	140426	M.mys/bra	DH	DH	DH	F	Ad	N/A	Par	33.5	20.8	17.1	3.7	N/A	N	
93	04/05/2021	22:20	4	522458	140462	M.bec	DH	DH	DH	F	Ad	N/A	Par	42.5	25.4	17.2	3.2	2M	N	
94	04/05/2021	22:20	4	522458	140462	M.nat	DH	DH	DH	F	Ad	N/A	Par	40.5	23.6	16.4	7.2	N/A	N	
95	06/05/2021	22:15	26	522669	138383	M. nat	TE	TE	TE	М	Imm	0	N/A	38.45	N/A	N/A	7.33	N/A	N	
96	06/05/2021	23:50	27	522788	138370	M. dau	TE	JL	TE	М	Α	0	N/A	37.9	N/A	N/A	8.08	N/A	N	
97	07/05/2021	22:00	1	523034	140733	P. aur	TE	JL	TE	F	Ad	N/A	Par	39.2	N/A	N/A	6.57	N/A	N	Minor wing damage (hole with diameter <2mm), generally calm





Bat No.	Date	Time (24hr)	Trap No.	Easting	Northing	Species	Extracted by	Measured by	Determined by	Sex - M/F	Age - Juv/ Imm/ Ad	status (male) - Testes	Breeding status (female) - Par/ NP/ Preg/ Lact	Forearm (mm)	Total weight (g)	Bag weight(g)	Bat weight (g)	Tagged bat? Include bat number	Sample taken?	ID features, damage, parasites photos and samples collected
																				Bald patch on abdomen
98	07/05/2021	23:00	1	523034	140733	P. aur	TE	JL	TE	F	Ad	N/A	Par	40	N/A	N/A	7.59	N/A	N	indicative of mating, very
																				active



Annex 2

Home Range Analysis

Table 8.1.2 Home Range Analysis

Bat number	50% KDE (ha)	95% KDE (ha)	100% MCP (ha)	
1J	0.09	2.21	28.37	
2J	4.47	36.35	32.71	
3J	0.48	15.61	183.49	
4J	0.22	17.12	507.07	
5J	4.03	40.87	58.77	
6J	0.82	37.74	289.73	
7J	5.51	64.76	388.46	
8J	2.40	51.93	262.23	
1S	N/A	N/A	5.04	
2S	2.64	36.74	286.85	
3S	6.01	107.70	1470.98	
4S	1.67	23.40	290.99	
5S	N/A	N/A	N/A	
1M	1.29	11.31	10.39	
2M	0.87	9.44	11.09	





Annex 3

Figures











