

# IMMINGHAM EASTERN RO-RO TERMINAL



Environmental Statement: Volume 3  
Appendix 9.1: Benthic Ecology Survey  
Document Reference: 8.4.9 (a)

APFP Regulations 2009 – Regulation 5(2)(a) and 5(2)(e)  
PINS Reference – TR030007




December 2022

# Associated British Ports

## Immingham Eastern RoRo Terminal

### Environmental Statement: Appendix 9.1: Benthic Ecology Survey

December 2022



Innovative Thinking - Sustainable Solutions

Page intentionally left blank

# Immingham Eastern RoRo Terminal

## Environmental Statement:

### Appendix 9.1: Benthic Ecology Survey

December 2022



## Document Information

Document History and Authorisation		
<b>Title</b>	Immingham Eastern RoRo Terminal	
	Environmental Statement Appendix 9.1: Benthic Ecology Survey	
<b>Commissioned by</b>	Associated British Ports	
<b>Issue date</b>	December 2022	
<b>Document ref</b>	R.3742	
<b>Project no</b>	R/5035/4	
Date	Version	Revision Details
13/12/2022	1	Issued for client use

Prepared (PM)	Approved (QM)	Authorised (PD)
A J Pearson	E San Martin	N J Frost

### Suggested Citation

ABPmer, (2022). Immingham Eastern RoRo Terminal, Environmental Statement Appendix 9.1: Benthic Ecology Survey ABPmer Report No. R.3742. A report produced by ABPmer for Associated British Ports, December 2022.

### Authors

ABPmer

# Contents

1	Introduction.....	1
2	Methodology.....	1
	2.1 Intertidal benthic sampling.....	1
	2.2 Subtidal benthic sampling.....	1
	2.3 Laboratory analysis.....	2
3	Results.....	4
	3.1 Introduction.....	4
	3.2 Intertidal ecology survey results.....	4
	3.3 Subtidal benthic survey results.....	9
4	Summary.....	17
5	References.....	17
6	Abbreviations/Acronyms.....	18

## Annexes

A	Laboratory Macrofauna Results
B	Laboratory Sediment Results
	B.1 Particle Size Analysis
	B.2 Total Organic Carbon

## Images

Image 1.	Station IMM 04 (sandy mud with bird footprints).....	8
Image 2.	Pacific oysters <i>Crassostrea gigas</i> and barnacles colonising piles.....	8
Image 3.	Station IMM 14 sample (sandy mud).....	10
Image 4.	Station HU060 06 (sand).....	11
Image 5.	HU056 05 (Gravelly sand).....	12

## Figures

Figure 1.	The location of intertidal core and subtidal grab sample stations.....	3
-----------	--	---

## Tables

Table 1.	Intertidal benthic survey results.....	5
Table 2.	Size classes of key bird prey species.....	9
Table 3.	Subtidal benthic survey results.....	13

# 1 Introduction

- 1.1.1 This report has been prepared by ABPmer for Associated British Ports (ABP) and provides a summary of the results of intertidal and subtidal benthic sampling undertaken in September 2021 as part of the Immingham Eastern RoRo Terminal Project (also referred to as the proposed development).
- 1.1.2 The survey methodologies are detailed in Section 2 and the results of the intertidal and subtidal surveys are presented in Sections 3. Overall summary conclusions are then provided in Section 4.

## 2 Methodology

### 2.1 Intertidal benthic sampling

- 2.1.1 The intertidal sampling survey was undertaken on the 7 September 2021 using a dedicated two-person hovercraft to access the shoreline safely.
- 2.1.2 Ten stations were successfully sampled (Figure 1). At each of these stations, a sample was collected using a 0.01 m<sup>2</sup> hand-held corer (to a depth of approximately 15 cm) and analysed for macrofaunal analysis (faunal composition, abundance and biomass). An additional core sample was also be collected at each station for Particle Size Analysis (PSA) and Total Organic Carbon (TOC).
- 2.1.3 Field notes were also made about the nature of the habitats at each of the sampling points. Information recorded included details on the sediment type, evidence of bird feeding (e.g. footprints), the specific characteristics of the habitat at the precise point where the samples were retrieved and general characteristics of the wider habitat. A photographic record of the sediment type and the broader habitat appearance was also taken.
- 2.1.4 All infaunal samples were immediately delivered to the laboratory once the survey was complete, where samples were sieved (using a 0.5 mm sieve) and fixed ahead of analysis.
- 2.1.5 Anecdotal ornithology observations were also recorded during the survey.

### 2.2 Subtidal benthic sampling

- 2.2.1 The subtidal grab sampling survey was undertaken on the 10-11 September 2021. The sampling methods followed the established and recognised procedures outlined in the Recommended Operational Guidelines (ROG) for Grab Sampling and Sorting and Treatment of Samples (Guerra and Freitas, 2013) and the Marine Monitoring Handbook, Procedural Guideline No 3-9 (Thomas, 2000).

- 2.2.2 The survey was undertaken to characterise the benthic fauna within the following areas:
- Immingham Eastern RoRo Terminal: Ten stations were sampled with the location of these stations shown in Figure 1.
  - Disposal sites (HU060/HU056): In total, twelve stations were sampled (four within each of the proposed disposal sites and two nearby to each of the disposal sites) as shown in Figure 1.
- 2.2.3 At each station, a benthic sample was collected using a 0.1 m<sup>2</sup> Day Grab for macrofauna analysis (faunal composition, abundance and biomass). An additional sample was also taken at each station for determination of PSA and TOC.
- 2.2.4 At each site, up to three attempts were made to retrieve a suitable sample (i.e. a grab containing sufficient volume of sediment for analysis). The sediment depths within the grab which were used for sample acceptance were a minimum of 7 cm for muddy or soft sediments and 5 cm for hard packed or coarse sediments. Anything less than these values was only retained if no other viable sample was collected.
- 2.2.5 Each grab sample was photographed upon successful retrieval and transferred into a labelled plastic bucket. All infaunal samples were immediately delivered to the laboratory once the survey was complete, where samples were sieved (using a 0.5 mm sieve) and fixed ahead of analysis.
- 2.2.6 Anecdotal ornithology and marine mammal observations were also recorded during the survey.

## 2.3 Laboratory analysis

- 2.3.1 The benthic macrofaunal analysis was undertaken by Hull Marine Laboratory. The laboratory is Marine Biological Analytical Quality Control Scheme (NMBAQC) accredited. The PSA and TOC elements of the benthic ecology analysis were conducted by the ABPmer in-house NMBAQC accredited laboratory.
- 2.3.2 Faunal samples were sorted from the sieve residue using low power binocular microscopes. All of the macroinfaunal specimens were identified to species level (where practicable) and enumerated. This work was undertaken in adherence with ISO 16665 standards and the NMBAQC Scheme Guidelines.
- 2.3.3 The PSA sample analysis was undertaken by ABPmer using the NMBAQC standardised methodology. The analysis was carried out using a Mastersizer laser diffractor which produces detailed sedimentary profiles for fine sediments (clay, sand and silts). The TOC analysis was carried out using an elemental analyser.






Figure 1. The location of intertidal core and subtidal grab sample stations

## 3 Results

### 3.1 Introduction

- 3.1.1 The results of the intertidal and subtidal sampling are described in the following sections, including key characterising species and any species of particular note (such as protected species or non-native species)<sup>1</sup>.
- 3.1.2 The laboratory results are presented in Annex 1 (faunal analysis) and Annex 2 (PSA and TOC). Summary information on the sedimentary and ecological conditions at the sites based on these results is presented in Table 1.

### 3.2 Intertidal ecology survey results

- 3.2.1 The sediment in samples collected in this area consisted predominantly of sandy mud (Table 1 and Image 1). The TOC in the samples ranged between approximately 1 % and 3 % (Table 1). Overall, the number of taxa found in the samples was variable and ranged from four (Station IMM 1 and IMM 3) to 15 (Station IMM 7). The number of individuals was also highly variable and ranged from 1,100 organisms per m<sup>2</sup> (Station IMM 1) to 40,600 organisms per m<sup>2</sup> (Station IMM 7). The range in total species biomass in the samples was between 1 gram per m<sup>2</sup> at Station IMM 3 and 190 grams per m<sup>2</sup> at Station IMM 7 (which was primarily attributed to the ragworm *Hediste diversicolor* and the peppery furrow shell *Scrobicularia plana*).
- 3.2.2 The infaunal samples were predominantly characterised by nematodes, the oligochaetes *Tubificoides benedii* and *Enchytraeidae* spp., the mud shrimp *Corophium volutator*, the mudsnail *Peringia ulvae*, Baltic tellin *Limecola balthica* as well as the polychaetes *Hediste diversicolor* and *Pygospio elegans*. These characterising species dominated the assemblage and contributed almost entirely to the total abundances of organisms recorded at most of the sites. All the species recorded from the samples in this area were considered commonly occurring in the region and not protected.
- 3.2.3 During the surveys, the non-native Pacific oyster *Crassostrea gigas* and barnacles were recorded attached to piles on jetties in the area (Image 2).
- 3.2.4 The assemblage recorded is considered typical of the community recorded on mudflats in the nearby area (ABPmer, 2009; IECS, 2010 Able UK Limited, 2021). For example, intertidal surveys at North Killingholme (located approximately 3 km from the proposed development) in 2015 and 2016 also recorded a benthic assemblage characterised by species such as *Corophium volutator*, *Tubificoides benedii*, *Pygospio elegans*, *Hediste diversicolor*, *Limicola balthica* and nematodes with a broadly similar total number of individuals in the samples (up to around 50,000 organisms per m<sup>2</sup>) (Able UK Limited, 2021).

---

<sup>1</sup> As is standard practice for benthic analysis, the data in the report is presented as m<sup>2</sup> rather than per 0.1m<sup>2</sup> grab.

**Table 1. Intertidal benthic survey results**


Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
IMM 1	Mud	3.65	4	1,100	6.29	<i>Nematoda</i> (400) <i>Limecola balthica</i> (300) <i>Tubificoides benedii</i> (300) <i>Nephtys</i> (100)
IMM 2	Sandy Mud	3.32	14	15,400	105.76	<i>Peringia ulvae</i> (4,600) <i>Nematoda</i> (2,400) <i>Enchytraeidae</i> (2,100) <i>Hediste diversicolor</i> (1,500) <i>Tubificoides benedii</i> (1,400) <i>Pygospio elegans</i> (1,100) <i>Abra tenuis</i> (500)
IMM 3	Sandy Mud	2.99	4	1,300	1.13	<i>Nematoda</i> (500) <i>Limecola balthica</i> (500) <i>Tubificoides benedii</i> (200) <i>Tharyx</i> (100)
IMM 4	Sandy Mud	2.92	9	20,700	31.14	<i>Tubificoides benedii</i> (14,400) <i>Corophium volutator</i> (3,600) <i>Nematoda</i> (800) <i>Limecola balthica</i> (700) <i>Tellinoidea</i> (600) <i>Pygospio elegans</i> (300)
IMM 5	Sandy Mud	3.05	6	1,600	6.16	<i>Tubificoides benedii</i> (900) <i>Limecola balthica</i> (300) <i>Nematoda</i> (100) <i>Enchytraeidae</i> (100) <i>Corophium volutator</i> (100) <i>Tellinoidea</i> (100)

Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
IMM 6	Sandy Mud	2.90	11	30,300	58.07	<i>Enchytraeidae</i> (5,400) <i>Peringia ulvae</i> (5,400) <i>Tubificoides benedii</i> (5,000) <i>Nematoda</i> (4,900) <i>Hediste diversicolor</i> (2,700) <i>Limecola balthica</i> (2,500) <i>Abra tenuis</i> (2,000)
IMM 7	Sandy Mud	3.36	15	40,600	189.77	<i>Tubificoides benedii</i> (13,800) <i>Enchytraeidae</i> (5,700) <i>Nematoda</i> (5,100) <i>Limecola balthica</i> (3,500) <i>Pygospio elegans</i> (3,400) <i>Hediste diversicolor</i> (3,300) <i>Peringia ulvae</i> (1,800)
IMM 8	Sandy Mud	3.05	14	4,100	15.87	<i>Nematoda</i> (800) <i>Limecola balthica</i> (700) <i>Tubificoides benedii</i> (600) <i>Peringia ulvae</i> (400) <i>Hediste diversicolor</i> (300)
IMM 9	Sandy Mud	3.73	14	21,600	47.98	<i>Hediste diversicolor</i> (6,800) <i>Nematoda</i> (3,200) <i>Abra tenuis</i> (2,000) <i>Enchytraeidae</i> (1,600) <i>Peringia ulvae</i> (1,500) <i>Tubificoides benedii</i> (1,400) <i>Limecola balthica</i> (1,200)

Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
IMM 10	Sandy Mud	2.71	8	26,800	57.37	<i>Corophium volutator</i> (16,400) <i>Tubificoides benedii</i> (4,800) <i>Nematoda</i> (2,100) <i>Limecola balthica</i> (1,800) <i>Tellinoidea</i> (1,100) <i>Eteone longa</i> (400)



**Image 1. Station IMM 04 (sandy mud with bird footprints)**



**Image 2. Pacific oysters *Crassostrea gigas* and barnacles colonising piles**

3.2.5 Many of the species recorded in the samples are considered prey species for coastal waterbirds such as polychaetes, Baltic tellin *Limecola balthica*, mudsnail *Peringia* spp. and mudshrimp *Corophium* spp. (Stillman *et al.*, 2005; Woodward *et al.*, 2014). The species and size of the prey taken varies between different coastal waterbirds. Larger waders are typically capable of consuming larger invertebrate prey items than smaller species. In order to better understand prey size in the samples collected, prey species were assigned to different size classes based on a size class classification supplied by the laboratory which has been used by Natural England and the Environment Agency in previous studies. The results are summarised in Table 2. The benthic prey recorded in the surveys were typically small size classes that are consumed by both smaller and larger wading bird species.

**Table 2. Size classes of key bird prey species**

Species Group	Species	Size Class	Abundance (per sample)	% (per sample)
Polychaetes	<i>Eteone longa</i>	<25mm	14	100
		>25mm	0	0
	<i>Hediste diversicolor</i>	<25mm	113	77
		25-50mm	34	23
	<i>Nephtys spp</i>	<25mm	3	100
		>25mm	0	0
	<i>Pygospio elegans</i>	<25mm	68	100
		>25mm	0	0
	<i>Streblospio shrubsolii</i>	<25mm	12	100
		>25mm	0	0
	<i>Tharyx</i>	<25mm	3	100
		>25mm	0	0
<i>Manayunkia aestuarina</i>	<25mm	22	100	
	>25mm	0	0	
Crustacean	<i>Corophium volutator</i>	<3mm	142	65
		>3mm	75	35
Gastropod	<i>Peringia ulvae</i>	<3mm	136	99
		3-5mm	1	1
Bivalves	<i>Limecola balthica</i>	<9mm	117	98
		9-15mm	2	2
	<i>Abra tenuis</i>	<5mm	51	100
		>5mm	0	0
	<i>Scrobicularia plana</i>	20-25mm	2	100
<b>Size classes used:</b>				
<i>Hediste diversicolor</i> + other polychaetes:		<25 mm, 25-50 mm, 50-75 mm, 75-100 mm, >100 mm		
<i>Corophium volutator</i> + other corophiid species:		<3 mm, >3 mm		
<i>Peringia ulvae</i> :		<3 mm, 3-5 mm, >5 mm		
<i>Macoma balthica</i> :		<9 mm, 9-15 mm, 15-20 mm, >20 mm		
Other bivalve species:		< 5 mm, 5-10 mm, 10-15 mm, 15-20 mm		

## Ornithology observations

3.2.6 Species such as Curlew, Black-tailed Godwit, Redshank and Dunlin were recorded foraging within the Proposed Development area approximately 150 waders using this area in total. Approximately 30 Turnstone and 50 Herring Gull were present roosting on the outfall and the concrete structures.

## 3.3 Subtidal benthic survey results


3.3.1 To summarise the key findings from the subtidal sampling, the survey area has been divided into three broad areas with the results for each area summarised.

3.3.2 These areas are as follows:

- Immingham Eastern RoRo Terminal (Stations IMM 11 to IMM 20);
- Disposal site HU060 (Stations HU060 1 to HU060 6); and
- Disposal site HU056 (Stations HU056 1 to HU056 6).

### Immingham Eastern RoRo Terminal

3.3.3 The sediment from samples collected from the area of the Immingham Eastern RoRo Terminal consisted of mud and sandy mud (Image 3). The TOC in the samples ranged between approximately 3 % and 13 % (Table 3). Overall, the number of taxa found in the samples ranged from two (Station IMM 15) to 17 (Station IMM 14), and the number of individuals from 20 organisms per m<sup>2</sup> (Station IMM 15) to 37,540 organisms per m<sup>2</sup> (Station IMM 13). However, most stations were relatively impoverished (<10 taxa and <10,000 organisms per m<sup>2</sup>). The range in total species biomass in the samples was between >1 and 14 grams per m<sup>2</sup>.



**Image 3. Station IMM 14 sample (sandy mud)**

3.3.4 The faunal samples were predominantly characterised by nematodes, the mudsnail *Corophium volutator*, polychaetes (such as *Streblospio shrubsolii*, *Polydora cornuta*, *Tharyx* spp. and *Nephtys* spp.), oligochaetes *Tubificoides* spp. and barnacle *Amphibalanus improvises*. All the species recorded from the samples in this area were considered commonly occurring in the region and not protected.

3.3.5 The faunal assemblage recorded is considered characteristic of subtidal habitats in this section of the Humber Estuary. For example, subtidal benthic surveys undertaken in the Immingham area in 2009, 2010 and 2016 predominantly recorded mud or muddy sand habitat which was generally impoverished (with a low number of taxa occurring at the majority of sites). The most commonly recorded infaunal species (generally recorded in low



abundances) were the polychaetes *Capitella capitata*, *Streblospio shrubsolii*, *Pygospio elegans*, *Polydora cornuta*, oligochaetes *Tubificoides* spp., mud shrimp *Corophium volutator*, and nematodes (ABPmer, 2009; IECS, 2010; Able UK Limited, 2021).

### Disposal site HU060

- 3.3.6 The sediment in samples collected in this area consisted predominately of sand (Image 4) with TOC between approximately <1 and 3 % at all stations (Table 3).
- 3.3.7 Most stations were considered impoverished (<7 taxa and <121 organisms per m<sup>2</sup>). However, 16 taxa were recorded at both Station HU060 4 and HU060 6 with 1,880 and 4,030 organisms per m<sup>2</sup> respectively at each of these stations. Biomass ranged from 0 to 3.37 grams per m<sup>2</sup>.
- 3.3.8 The samples were characterised by a wide range of species but typically in low abundances including nematodes, barnacle *Amphibalanus improvises*, polychaetes (such as *Pygospio elegans* and *Arenicola* spp.) and the amphipod *Corophium volutator*. All the species recorded from the samples in this area were considered commonly occurring in the region and not protected.
- 3.3.9 Benthic surveys undertaken in 2008 within and near to Clay Huts disposal sites also recorded a community characterised by the polychaetes *Arenicola marina* and *Pygospio elegans* as well as nematodes and amphipods (ABPmer, 2009).



Image 4. Station HU060 06 (sand)

### Disposal site HU056

- 3.3.10 The sediment in samples collected in this area consisted of sand, gravelly sand and sandy gravel (Image 5) with TOC ranging from between

approximately 1 % and 3 % (Table 3). The stations were considered highly impoverished (with 0 to 2 taxa and 0 to 30 organisms per m<sup>2</sup> recorded). The samples were characterised by low abundances of a few species (the amphipod *Corophium volutator*, mysid shrimp *Gastrosaccus spinifer*, bryozoan *Electra monostachys* and springtails *Collembola* spp.).

3.3.11 Benthic monitoring in 2017 at disposal site HU056 recorded commonly occurring estuarine species generally in low abundances such as the polychaetes *Polydora cornuta*, *Pygospio elegans* *Arenicola marina* and *Capitella* spp., bivalve *Limecola balthica*, mysid shrimps and amphipods (ABPmer, 2017).

3.3.12 The impoverished assemblage recorded is considered typical of scoured subtidal habitats in the Humber Estuary (which are subject to very strong tidal currents). No protected species were recorded.



**Image 5. HU056 05 (Gravelly sand)**

### **Marine mammal and ornithology observations**

3.3.13 Approximately 10 to 15 grey seals were recorded hauling out on mudflat at Sunk Island (on the north bank of the Humber Estuary), approximately 4 km from the proposed development. In addition, a harbour porpoise was recorded foraging in the channel (approximately 2 km from the proposed development).

3.3.14 A juvenile auk species (a Razorbill or Guillemot) was recorded foraging mid channel between Grimsby and Immingham.

**Table 3. Subtidal benthic survey results**

Area	Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
Immingham Eastern RoRo Terminal	IMM 11	Mud	3.83	12	11,740	8.32	<i>Corophium volutator</i> (8,910) <i>Tubificoides benedii</i> (1,570) <i>Streblospio shrubsolii</i> (420) <i>Nematoda</i> (250) <i>Tharyx</i> (240) <i>Limecola balthica</i> (130) <i>Tubificoides swirencoides</i> (100)
	IMM 12	Sandy Mud	4.63	16	12,270	1.44	<i>Nematoda</i> (9,830) <i>Streblospio shrubsolii</i> (1,210) <i>Amphibalanus improvises</i> (450) <i>Polydora cornuta</i> (440) <i>Corophium volutator</i> (110) <i>Mytilus edulis</i> (90) <i>Tharyx</i> (60)
	IMM 13	Sandy Mud	13.01	4	37,540	14.13	<i>Corophium volutator</i> (33,130) <i>Polydora cornuta</i> (4,170) <i>Nematoda</i> (230) <i>Tubificoides benedii</i> (10)
	IMM 14	Sandy Mud	4.03	17	22,480	3.34	<i>Streblospio shrubsolii</i> (13,790) <i>Nematoda</i> (7,150) <i>Amphibalanus improvisus</i> (520) <i>Polydora cornuta</i> (340) <i>Tharyx</i> (210) <i>Tubificoides benedii</i> (210) <i>Corophium volutator</i> (70)
	IMM 15	Sandy Mud	13.01	2	20	0.10	<i>Nephtys hombergii</i> (10) <i>Amphibalanus improvisus</i> (10)

Area	Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
	IMM 16	Sandy Mud	4.03	5	250	1.19	<i>Tubificoides benedii</i> (120) <i>Nephtys</i> (50) <i>Nematoda</i> (40) <i>Limecola balthica</i> (40)
	IMM 17	Sandy Mud	3.98	4	80	0.09	<i>Nephtys</i> (30) <i>Nematoda</i> (20) <i>Diastylis rathkei</i> (20) <i>Corophium volutator</i> (10)
	IMM 18	Sandy Mud	3.69	5	9,580	6.30	<i>Corophium volutator</i> (9,550) <i>Tubificoides benedii</i> (10) <i>Enchytraeidae</i> (10) <i>Limecola balthica</i> (10)
	IMM 19	Mud	4.23	8	300	0.57	<i>Streblospio shrebsolii</i> (110) <i>Nematoda</i> (50) <i>Nephtys hombergii</i> (50) <i>Tubificoides benedii</i> (30) <i>Tharyx</i> (20) <i>Limecola balthica</i> (20) <i>Diastylis rathkei</i> (10)
	IMM 20	Sand	4.22	9	5,130	4.91	<i>Corophium volutator</i> (4,950) <i>Streblospio shrebsolii</i> (70) <i>Nematoda</i> (30) <i>Nephtys</i> (30) <i>Limecola balthica</i> (20) <i>Diastylis rathkei</i> (10) <i>Austrominius modestus</i> (10) <i>Tubificoides benedii</i> (10)

Area	Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
Disposal site HU060	HU060 1	Sand	4.04	6	40	0.004	<i>Nematoda</i> (10) <i>Pygospio elegans</i> (10) <i>Arenicola</i> (10) <i>Bathyporeia elegans</i> (10)
	HU060 2	Sand	0.38	0	0	0.00	
	HU060 3	Slightly Gravelly Muddy Sand	0.92	6	60	0.01	<i>Scoloplos armiger</i> (20) <i>Eteone longa</i> (10) <i>Tharyx</i> (10) <i>Corophium volutator</i> (10) <i>Tellinoidea</i> (10)
	HU060 4	Sand	1.69	16	1,880	3.37	<i>Amphibalanus improvisus</i> (1,800) <i>Nototropis guttatus</i> (20) <i>Jaera</i> ( <i>Jaera</i> ) <i>albifrons</i> (20) <i>Scoloplos armiger</i> (10) <i>Tubificoides benedii</i> (10) <i>Corophium volutator</i> (10) <i>Limecola balthica</i> (10)
	HU060 5	Sand	2.51	3	120	0.01	<i>Protodriloides chaetifer</i> (90) <i>Mytilus edulis</i> (20) <i>Tubificoides benedii</i> (10)
	HU060 6	Sand	3.04	16	4,030	0.56	<i>Nematoda</i> (2,170) <i>Pygospio elegans</i> (900) <i>Arenicola</i> (590) <i>Polydora cornuta</i> (80) <i>Ampharete cf. acutifrons</i> (80) <i>Austrominius modestus</i> (50) <i>Corophium volutator</i> (50)
	HU056 1	Sand	2.01	1	30	0.001	<i>Corophium volutator</i> (30)

Area	Station	Sediment Type	TOC (%)	No. of Taxa (per m <sup>2</sup> )	No. of Individuals (per m <sup>2</sup> )	Total Biomass (g per m <sup>2</sup> )	Key Characterising Species (Number per m <sup>2</sup> Shown in Brackets)
Disposal site HU056	HU056 2	Slightly Gravelly Muddy Sand	2.84	2	0	0.001	<i>Corophium volutator</i> (P) <i>Electra monostachys</i> (P)
	HU056 3	Muddy Gravel	1.05	1	10	0.002	<i>Corophium volutator</i> (10)
	HU056 4	Gravelly Mud	1.01	0	0	0.00	
	HU056 5	Gravelly Sand	1.40	0	0	0.00	
	HU056 6	Muddy Gravel	1.03	2	20	0.12	<i>Gastrosaccus spinifer</i> (10) <i>Collembola</i> (10)

## 4 Summary

- 4.1.1 The intertidal benthic samples consisted predominantly of sandy mud and were characterised by a range of commonly occurring species of oligochaetes, polychaetes, amphipods, molluscs and nematodes which are considered characteristic of intertidal mudflat in this area of the Humber Estuary. Many of the dominant species recorded are also considered important prey items for overwintering birds. Intertidal mudflat is a qualifying interest features of the Humber Estuary Special Area of Conservation (SAC) and Ramsar site. This feature is also supporting habitat of the Humber Estuary Special Protection Area (SPA). In addition, intertidal mudflat is a Habitat of Principal Importance in England under the Natural Environment and Rural Communities (NERC) Act 2006 Section 41. No intertidal species which are considered nationally rare or protected were recorded in the surveys.
- 4.1.2 The subtidal benthic samples consisted predominantly of mud or sandy mud, with gravelly sediment and sand mainly recorded at the disposal sites. Samples were typically impoverished and characterised by commonly occurring polychaetes, oligochaetes, nematodes, crustaceans and bivalves. No subtidal species considered nationally rare or protected were recorded, with the assemblages observed considered characteristic of estuarine communities found more widely in the Humber Estuary.

## 5 References

- Able UK Limited. (2021). Able Marine Energy Park (Material Change 2 – Tr030006). Updated Environmental Statement: Chapter 10: Aquatic Ecology.
- ABPmer. (2009). Humber Estuary: Environmental Management and Monitoring Plan: Data 2009. R. 1587.
- ABPmer (2017). Disposal Site HU056 Benthic Survey Data (Unpublished Data Collected for ABP).
- Institute of Estuarine and Coastal Studies (IECS). South Humber Channel Marine Studies: Intertidal and Subtidal Benthic & Fish Surveys 2010: Report to Yorkshire Forward.
- Guerra, M.T and Freitas, R. (2013). Recommended Operational Guidelines (ROG) for grab sampling and sorting and treatment of samples. Mesh Atlantic.
- ISO 16665: (2014) Water quality -- Guidelines for quantitative sampling and sample processing of marine soft-bottom macrofaunal. International Organization for Standardization.

JNCC (2015). The Marine Habitat Classification for Britain and Ireland Version 15.03 [Online]. [Date accessed]. Available from: [jncc.defra.gov.uk/MarineHabitatClassification](http://jncc.defra.gov.uk/MarineHabitatClassification)

Stillman, R.A., West, A.D., Goss-Custard, J.D., McGrorty, S., Frost, N.J., Morrissey, D.J., Kenny, A.J. and Drewitt, A.L. (2005). Predicting site quality for shorebird communities: a case study on the Humber estuary, UK. *Marine Ecological Progress Series*, 305, pp.203–217.

Thomas, N.S. (2000). Procedural Guideline No. 3-9 Quantitative sampling of sublittoral sediment biotopes and species using remote-operated grabs. In Davies J., Baxter J., Bradley M., Connor D., Khan J., Murray E., Sanderson W., Turnbull C. & Vincent M. 2001. *Marine Monitoring Handbook*, 405 pp.

Woodward, I.D., Calbrade, N.A and Holt., C.A. (2014). Humber Estuary Bird Decline Investigation 2014. BTO Research Report No. 668. Report of work carried out by The British Trust for Ornithology under contract to Natural England.

Wyn, G. and Brazier, P. (2001). Procedural Guideline No. 3-1 - In situ intertidal biotope recording. In Davies J., Baxter J., Bradley M., Connor D., Khan J., Murray E., Sanderson W., Turnbull C. & Vincent M. 2001. *Marine Monitoring Handbook*, 405 pp.

Wyn, G., Brazier, D. P. and McMath A. J., (2000). CCW handbook for marine intertidal Phase 1 survey and mapping. CCW Marine Sciences Report: 00/06/0.

## 6 Abbreviations/Acronyms




ABP	Associated British Ports
BAP	Biodiversity Action Plan
CCW	Countryside Council for Wales
GIS	Geographical Information System
GPS	Global Positioning System
ISO	International Organization for Standardization
JNCC	Joint Nature Conservation Committee
MHCBI	Marine Habitat Classification for Britain & Ireland
NERC	Natural Environment and Rural Communities
NMBAQC	National Marine Biological Analytical Quality Control Scheme
PSA	Particle Size Analysis
ROG	Recommended Operational Guidelines
TOC	Total Organic Carbon
UK	United Kingdom

Cardinal points/directions are used unless otherwise stated.

SI units are used unless otherwise stated.



# Annexes



Innovative Thinking - Sustainable Solutions

# A Laboratory Macrofauna Results





# **B Laboratory Sediment Results**


## **B.1 Particle Size Analysis**

# Particle Size Analysis Report

Sample Name: Imm\_01 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 21.08 %  
 Sample Collected:      Weighted Residual: 0.728 %


d(0.1): 3.212 μm      d(0.5): 12.745 μm      d(0.9): 60.020 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	24 %	36 %	27 %	7 %	3 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_02 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 13.23 %  
 Sample Collected:      Weighted Residual: 0.685 %


d(0.1): 3.383 μm      d(0.5): 18.132 μm      d(0.9): 68.935 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	20 %	30 %	34 %	13 %	0 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_03 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 11.64 %  
 Sample Collected:      Weighted Residual: 0.825 %


d(0.1): 4.335 μm      d(0.5): 31.757 μm      d(0.9): 93.194 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	13 %	21 %	40 %	22 %	2 %	0 %

## Frequency Curve






# Particle Size Analysis Report

Sample Name: Imm\_04 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 13.26 %  
 Sample Collected:      Weighted Residual: 0.883 %


d(0.1): 3.886 μm      d(0.5): 25.980 μm      d(0.9): 96.161 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	17 %	25 %	32 %	23 %	1 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_05 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 12.03 %  
 Sample Collected:      Weighted Residual: 0.914 %


d(0.1): 4.304 μm      d(0.5): 32.134 μm      d(0.9): 99.160 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	14 %	23 %	34 %	27 %	0 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_06 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 13.25 %  
 Sample Collected:      Weighted Residual: 0.861 %


d(0.1): 3.809 μm      d(0.5): 26.688 μm      d(0.9): 86.764 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	16 %	24 %	38 %	19 %	2 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_07 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 14.72 %  
 Sample Collected:      Weighted Residual: 0.883 %


d(0.1): 3.772 μm      d(0.5): 24.883 μm      d(0.9): 92.111 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	17 %	25 %	35 %	18 %	2 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_08 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 12.57 %  
 Sample Collected:      Weighted Residual: 1.038 %


d(0.1): 4.074 μm      d(0.5): 36.169 μm      d(0.9): 112.040 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	15 %	20 %	31 %	30 %	2 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_09 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 10.54 %  
 Sample Collected:      Weighted Residual: 0.860 %


d(0.1): 3.272 μm      d(0.5): 21.181 μm      d(0.9): 95.817 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	20 %	26 %	31 %	18 %	2 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_10 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 12.45 %  
 Sample Collected:      Weighted Residual: 0.775 %


d(0.1): 3.205 μm      d(0.5): 18.700 μm      d(0.9): 90.354 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	22 %	26 %	28 %	21 %	0 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_11 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 13.90 %  
 Sample Collected:      Weighted Residual: 0.599 %


d(0.1): 2.728 μm      d(0.5): 9.389 μm      d(0.9): 57.901 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
4 %	30 %	37 %	19 %	8 %	1 %	0 %

## Frequency Curve






# Particle Size Analysis Report

Sample Name: Imm\_12 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 13.83 %  
 Sample Collected:      Weighted Residual: 0.796 %


d(0.1): 3.076 μm      d(0.5): 12.737 μm      d(0.9): 98.640 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	25 %	33 %	21 %	15 %	3 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_13 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 15.93 %  
 Sample Collected: Rhizomes >1mm present in sample      Weighted Residual: 0.926 %


d(0.1): 5.339 μm      d(0.5): 27.691 μm      d(0.9): 171.668 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
1 %	11 %	29 %	28 %	22 %	8 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_14 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 14.61 %  
 Sample Collected:      Weighted Residual: 0.842 %


d(0.1): 3.359 μm      d(0.5): 16.579 μm      d(0.9): 96.798 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	21 %	30 %	25 %	18 %	3 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_15 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 15.70 %  
 Sample Collected:      Weighted Residual: 0.721 %


d(0.1): 3.278 μm      d(0.5): 15.313 μm      d(0.9): 82.670 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	22 %	31 %	27 %	16 %	1 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_16 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 14.33 %  
 Sample Collected:      Weighted Residual: 0.738 %


d(0.1): 3.391 μm      d(0.5): 15.508 μm      d(0.9): 79.285 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
2 %	22 %	33 %	28 %	15 %	1 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_17 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 14.16 %  
 Sample Collected:      Weighted Residual: 0.664 %


d(0.1): 2.795 μm      d(0.5): 10.761 μm      d(0.9): 75.386 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
4 %	28 %	32 %	22 %	13 %	1 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_18 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 21.83 %  
 Sample Collected:      Weighted Residual: 0.700 %


d(0.1): 2.884 μm      d(0.5): 12.797 μm      d(0.9): 89.358 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
4 %	25 %	31 %	23 %	14 %	3 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: Imm\_19 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 13.92 %  
 Sample Collected:      Weighted Residual: 0.642 %


d(0.1): 2.984 μm      d(0.5): 10.503 μm      d(0.9): 57.480 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
3 %	28 %	38 %	22 %	9 %	1 %	0 %

## Frequency Curve






# Particle Size Analysis Report

Sample Name: Imm\_20 - Average      Measured by: IDavidson      on 24 September 2021  
 Sample Source: Sugar      Obscuration: 12.27 %  
 Sample Collected:      Weighted Residual: 0.435 %


d(0.1): 81.836 μm      d(0.5): 155.361 μm      d(0.9): 257.761 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	2 %	3 %	1 %	66 %	27 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU56\_01 - Average      Measured by: IDavidson      on 30 September 2021  
 Sample Source: Sugar      Obscuration: 12.86 %  
 Sample Collected:      Weighted Residual: 0.773 %


d(0.1): 108.455 μm      d(0.5): 158.002 μm      d(0.9): 229.834 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	0 %	0 %	0 %	79 %	21 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU56\_02 Measured by: IDavidson  
 Sample Source: Sugar  
 Sample Collected: 11/09/2021


d(0.1): 22.3  $\mu\text{m}$  d(0.5): 188.7  $\mu\text{m}$  d(0.9): 461.9  $\mu\text{m}$

## Cumulative Frequency Plot



Clay (%)	Silt (%)			Sand (%)			Gravel (%)			Cobble (%)
	Fine	Medium	Coarse	Fine	Medium	Coarse	Fine	Medium	Coarse	
0.3	3.0	6.2	4.7	43.3	36.5	4.3	0.4	1.2	0.0	0.0

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU56\_03 Measured by: IDavidson  
 Sample Source: Sugar  
 Sample Collected: 11/09/2021


d(0.1): 7.7  $\mu\text{m}$  d(0.5): 86.6  $\mu\text{m}$  d(0.9): 10531.9  $\mu\text{m}$

## Cumulative Frequency Plot



Clay (%)	Silt (%)			Sand (%)			Gravel (%)			Cobble (%)
	Fine	Medium	Coarse	Fine	Medium	Coarse	Fine	Medium	Coarse	
0.9	6.8	17.0	21.4	10.2	3.1	3.5	17.0	20.2	0.0	0.0

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU56\_04 Measured by: IDavidson  
 Sample Source: Sugar  
 Sample Collected: 11/09/2021


d(0.1): 3.4 μm d(0.5): 18.0 μm d(0.9): 13348.1 μm

## Cumulative Frequency Plot



Clay (%)	Silt (%)			Sand (%)			Gravel (%)			Cobble (%)
	Fine	Medium	Coarse	Fine	Medium	Coarse	Fine	Medium	Coarse	
3.3	17.9	32.1	17.9	8.5	1.4	2.5	2.6	13.7	0.0	0.0

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU56\_05 Measured by: IDavidson  
 Sample Source: Sugar  
 Sample Collected: 11/09/2021


d(0.1): 158.8  $\mu\text{m}$  d(0.5): 729.4  $\mu\text{m}$  d(0.9): 4659.8  $\mu\text{m}$

## Cumulative Frequency Plot



Clay (%)	Silt (%)			Sand (%)			Gravel (%)			Cobble (%)
	Fine	Medium	Coarse	Fine	Medium	Coarse	Fine	Medium	Coarse	
0.0	0.2	0.6	0.4	15.3	29.1	35.8	15.3	3.4	0.0	0.0

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU56\_06 Measured by: IDavidson  
 Sample Source: Sugar  
 Sample Collected: 11/09/2021


d(0.1): 7.5 μm d(0.5): 75.9 μm d(0.9): 10610.9 μm

## Cumulative Frequency Plot



Clay (%)	Silt (%)			Sand (%)			Gravel (%)			Cobble (%)
	Fine	Medium	Coarse	Fine	Medium	Coarse	Fine	Medium	Coarse	
0.9	7.0	17.5	22.0	10.5	3.2	3.9	9.9	25.1	0.0	0.0

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU60\_01 - Average      Measured by: IDavidson      on 30 September 2021  
 Sample Source: Sugar      Obscuration: 10.66 %  
 Sample Collected:      Weighted Residual: 0.787 %


d(0.1): 158.985 μm      d(0.5): 229.115 μm      d(0.9): 329.772 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	0 %	0 %	0 %	32 %	68 %	0 %

## Frequency Curve






# Particle Size Analysis Report

Sample Name: HU60\_02 - Average      Measured by: IDavidson      on 30 September 2021  
 Sample Source: Sugar      Obscuration: 12.29 %  
 Sample Collected:      Weighted Residual: 0.635 %


d(0.1): 158.062 μm      d(0.5): 226.015 μm      d(0.9): 322.084 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	0 %	0 %	0 %	33 %	67 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU60\_03 Measured by: IDavidson  
 Sample Source: Sugar  
 Sample Collected: 11/09/2021


d(0.1): 7.8  $\mu\text{m}$  d(0.5): 148.4  $\mu\text{m}$  d(0.9): 395.9  $\mu\text{m}$

## Cumulative Frequency Plot



Clay (%)	Silt (%)			Sand (%)			Gravel (%)			Cobble (%)
	Fine	Medium	Coarse	Fine	Medium	Coarse	Fine	Medium	Coarse	
0.9	6.9	14.7	15.1	27.1	32.5	2.4	0.4	0.0	0.0	0.0

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU60\_04 - Average      Measured by: IDavidson      on 30 September 2021  
 Sample Source: Sugar      Obscuration: 11.45 %  
 Sample Collected:      Weighted Residual: 0.729 %


d(0.1): 165.268 μm      d(0.5): 230.698 μm      d(0.9): 319.948 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	0 %	0 %	0 %	29 %	71 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU60\_05 - Average      Measured by: IDavidson      on 30 September 2021  
 Sample Source: Sugar      Obscuration: 12.32 %  
 Sample Collected:      Weighted Residual: 0.592 %


d(0.1): 141.521 μm      d(0.5): 203.339 μm      d(0.9): 290.934 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	0 %	0 %	0 %	48 %	52 %	0 %

## Frequency Curve




# Particle Size Analysis Report

Sample Name: HU60\_06 - Average      Measured by: IDavidson      on 30 September 2021  
 Sample Source: Sugar      Obscuration: 11.71 %  
 Sample Collected:      Weighted Residual: 0.780 %


d(0.1): 149.550 μm      d(0.5): 222.455 μm      d(0.9): 328.328 μm

## Cumulative Frequency Plot



Clay	Silt			Sand		
	Fine	Medium	Coarse	Fine	Medium	Coarse
0 %	0 %	0 %	0 %	37 %	63 %	0 %

## Frequency Curve



## **B.2 Total Organic Carbon**

Crucible ID	Sample	Weight of crucible (g)	Weight of crucible + wet sample (g)	Weight of crucible + dry sample @105 (g)	Weight of crucible + dry sample @475 (g)	Weight of dry material @105 (g)	Weight of dry material @475 (g)	LOI475
1	Imm_01	36.18	101.73	76.19	74.73	40.01	38.55	3.65
2	Imm_02	36.61	90.18	71.59	70.43	34.98	33.82	3.32
3	Imm_03	37.08	83.12	63.46	62.67	26.38	25.59	2.99
4	Imm_04	36.40	91.10	60.74	60.03	24.34	23.63	2.92
5	Imm_05	36.60	81.62	63.51	62.69	26.91	26.09	3.05
6	Imm_06	34.93	102.73	78.79	77.52	43.86	42.59	2.90
7	Imm_07	36.62	88.33	69.70	68.59	33.08	31.97	3.36
8	Imm_08	36.14	69.83	52.53	52.03	16.39	15.89	3.05
9	Imm_09	35.23	76.73	70.04	68.74	34.81	33.51	3.73
10	Imm_10	36.12	91.78	68.59	67.71	32.47	31.59	2.71
11	Imm_11	35.74	71.89	54.03	53.33	18.29	17.59	3.83
12	Imm_12	35.07	87.43	63.15	61.85	28.08	26.78	4.63
13	Imm_13	36.21	67.98	48.66	47.04	12.45	10.83	13.01
14	Imm_14	36.64	78.16	57.24	56.41	20.60	19.77	4.03
15	Imm_15	36.27	75.53	56.11	55.32	19.84	19.05	3.98
16	Imm_16	36.29	83.26	62.02	61.07	25.73	24.78	3.69
17	Imm_17	36.78	84.01	62.79	61.69	26.01	24.91	4.23
18	Imm_18	36.54	91.83	67.32	66.02	30.78	29.48	4.22
19	Imm_19	35.60	82.17	60.37	59.37	24.77	23.77	4.04
20	Imm_20	36.30	94.61	81.04	80.87	44.74	44.57	0.38
01	HU56_01	36.62	65.71	59.41	59.20	22.79	22.58	0.92
05	HU56_02	36.21	66.62	59.89	59.49	23.68	23.28	1.69
04	HU56_03	36.16	68.58	61.30	60.67	25.14	24.51	2.51
06	HU56_04	36.62	71.16	64.21	63.37	27.59	26.75	3.04
02	HU56_05	36.59	70.99	64.02	63.47	27.43	26.88	2.01
10	HU56_06	36.13	67.43	59.01	58.36	22.88	22.23	2.84
12	HU60_01	37.08	67.98	61.80	61.54	24.72	24.46	1.05
08	HU60_02	36.40	71.09	63.29	63.02	26.89	26.62	1.00
03	HU60_03	35.23	65.65	58.42	58.06	23.19	22.83	1.55
11	HU60_04	36.53	61.16	55.42	55.23	18.89	18.70	1.01
07	HU60_05	34.93	67.02	60.57	60.21	25.64	25.28	1.40
09	HU60_06	36.28	67.01	60.51	60.26	24.23	23.98	1.03

## Contact Us

ABPmer

Quayside Suite,

Medina Chambers

Town Quay, Southampton

SO14 2AQ

T +44 (0) 23 8071 1840

F +44 (0) 23 8071 1841

E [enquiries@abpmer.co.uk](mailto:enquiries@abpmer.co.uk)

